

Uncertainty grows as irrigation modelling advances

R code

Arnald Puy

Contents

1	Preliminary functions	2
2	ISIMIP Data	3
2.1	Historical data	3
2.2	Predictions	6
2.3	ANOVA	9
3	Khan et al dataset	12
4	Bibliographical study	15
5	Session information	21

1 Preliminary functions

```
# PRELIMINARY FUNCTIONS #####

sensobol::load_packages(c("openxlsx", "data.table", "tidyverse", "cowplot",
                          "benchmarkme", "parallel", "wesanderson", "scales", "ncdf4",
                          "countrycode", "rworldmap", "sp", "doParallel", "here", "lme4",
                          "microbenchmark"))

# Create custom theme -----

theme_AP <- function() {
  theme_bw() +
    theme(panel.grid.major = element_blank(),
          panel.grid.minor = element_blank(),
          legend.background = element_rect(fill = "transparent",
                                            color = NA),
          legend.key = element_rect(fill = "transparent",
                                     color = NA),
          strip.background = element_rect(fill = "white"),
          legend.margin = margin(0.5, 0.1, 0.1, 0.1),
          legend.box.margin = margin(0.2, -4, -7, -7),
          plot.margin = margin(3, 4, 0, 4),
          legend.text = element_text(size = 7.3),
          axis.title = element_text(size = 10),
          legend.key.width = unit(0.4, "cm"),
          legend.key.height = unit(0.4, "cm"),
          legend.title = element_text(size = 7.8),
          axis.text.x = element_text(size = 7),
          axis.text.y = element_text(size = 7),
          axis.title.x = element_text(size = 7.3),
          axis.title.y = element_text(size = 7.3),
          strip.text.x = element_text(size = 7.4))
}

# Select color palette -----

selected.palette <- "Darjeeling1"

# SOURCE ALL R FUNCTIONS NEEDED FOR THE STUDY #####

# Source all .R files in the "functions" folder -----

r_functions <- list.files(path = here("functions"), pattern = "\\\\.R$", full.names = TRUE)
lapply(r_functions, source)
```

2 ISIMIP Data

2.1 Historical data

```
# RETRIEVE DATA FROM ISIMIP #####

# Create vector with list of files -----

list.of.files <- list.files("./files/isimip")
model.names <- sub("^(.*)_.*", "\\1", list.of.files)
climate.scenarios <- sapply(strsplit(list.of.files, "_"), function(x) x[2])
social.scenarios <- sapply(strsplit(list.of.files, "_"), function(x) x[which(x == "co2") - 1])
files.directory <- paste("./files/isimip", list.of.files, sep = "/")
start_year <- 1971

# Create parallel cluster -----

numCores <- detectCores() * 0.75
cl <- makeCluster(numCores)
registerDoParallel(cl)

# Run for loop -----

isimip.hist <- foreach(i = 1:length(files.directory),
                      .packages = c("data.table", "countrycode", "tidyverse",
                                    "sp", "rworldmap", "ncdf4")) %dopar% {

    get_isimip_fun(nc_file = files.directory[i],
                  variable = "airrww",
                  start_year = start_year)

}

# Stop the cluster after the computation -----

stopCluster(cl)

# ARRANGE DATA #####

# Number of files -----

list.of.files

## [1] "dbh_gswp3_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [2] "dbh_princeton_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [3] "dbh_watch_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2001.nc4"
## [4] "dbh_watch-wfdei_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [5] "dbh_wfdei_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc"
## [6] "h08_gswp3_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [7] "h08_gswp3_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
```

```
## [8] "h08_princeton_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2012.nc4"
## [9] "h08_princeton_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2012(1).nc4"
## [10] "h08_princeton_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2012.nc4"
## [11] "h08_watch_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2001.nc4"
## [12] "h08_watch_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2001.nc4"
## [13] "h08_watch-wfdei_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [14] "h08_watch-wfdei_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [15] "h08_wfdei_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc"
## [16] "lpjml_gswp3_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [17] "lpjml_gswp3_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [18] "lpjml_princeton_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2012.nc4"
## [19] "lpjml_princeton_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2012.nc4"
## [20] "lpjml_watch_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2001.nc4"
## [21] "lpjml_watch_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2001.nc4"
## [22] "lpjml_watch-wfdei_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [23] "lpjml_watch-wfdei_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [24] "lpjml_wfdei_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc"
## [25] "pcr-globwb_gswp3_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [26] "pcr-globwb_gswp3_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [27] "pcr-globwb_princeton_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2012.nc4"
## [28] "pcr-globwb_princeton_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2012.nc4"
## [29] "pcr-globwb_watch_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2001.nc4"
## [30] "pcr-globwb_watch_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2001.nc4"
## [31] "pcr-globwb_watch-wfdei_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [32] "pcr-globwb_watch-wfdei_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [33] "pcr-globwb_wfdei_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc"
## [34] "vic_gswp3_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [35] "vic_gswp3_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [36] "vic_princeton_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [37] "vic_princeton_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [38] "vic_watch_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2001.nc4"
## [39] "vic_watch_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2001.nc4"
## [40] "vic_watch-wfdei_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc4"
## [41] "vic_watch-wfdei_nobc_hist_varsoc_co2_airrww_global_monthly_1971_2010.nc4"
## [42] "vic_wfdei_nobc_hist_pressoc_co2_airrww_global_monthly_1971_2010.nc"
```

```
# Name the slots -----
```

```
names(isimip.hist) <- paste(model.names, climate.scenarios, social.scenarios, sep = "/")
```

```
# Clean and bind dataset -----
```

```
isimip.dt <- rbindlist(isimip.hist, idcol = "model") %>%
  na.omit() %>%
  .[, model := factor(model)] %>%
  .[, c("model", "climate", "social") := tstrsplit(model, "/")]

fwrite(isimip.dt, "isimip.dt.csv")
```


```
# Pressoc: constant human impacts in the form of dams and reservoirs
# varsoc: variable human impacts.
```

2.1.1 Plot data

```
# PLOT ISIMIP #####
```


```
# Continental level -----
```

```
isimip.dt[, sum(V1, na.rm = TRUE), .(Continent, model, year, climate, social)] %>%
  ggplot(., aes(year, V1, group = interaction(climate, model), color = model,
        linetype = climate)) +
  facet_wrap(social~Continent, scales = "free_y", ncol = 5) +
  geom_line() +
  scale_x_continuous(breaks = breaks_pretty(n = 3)) +
  labs(x = "Year", y = bquote("IWW (km"3 * ")")) +
  theme_AP() +
  guides(color = guide_legend(nrow = 2)) +
  guides(linetype = guide_legend(nrow = 2)) +
  theme(legend.position = "top")
```



```
# Global level -----
```

```
isimip.dt[, sum(V1, na.rm = TRUE), .(year, model, climate, social)] %>%
  ggplot(., aes(year, V1, group = interaction(climate, model), color = model)) +
  geom_line() +
  facet_wrap(~social) +
  labs(x = "Year", y = bquote("IWW (km"3 * ")")) +
  theme_AP() +
  theme(legend.position = "top")
```



2.2 Predictions

```
# RETRIEVE PROJECTIONS FROM ISIMIP #####

# Create vector with list of files -----

path.projections <- "./files/isimip_future"
list.of.files.projections <- list.files(path.projections)
files.directory.projections <- paste(path.projections, list.of.files.projections, sep = "/")
variable <- "airrww"
start_year <- 2006

# Create parallel cluster -----

numCores <- detectCores() * 0.75
cl <- makeCluster(numCores)
registerDoParallel(cl)

# Run for loop -----

isimip.future <- foreach(i = 1:length(files.directory.projections),
                        .packages = c("data.table", "countrycode", "tidyverse",
                                      "sp", "rworldmap", "ncdf4")) %dopar% {

    get_isimip_fun(nc_file = files.directory.projections[i],
                  variable = variable,
                  start_year = start_year)

}

# Stop the cluster after the computation -----

stopCluster(cl)
```

```

# ARRANGE DATA #####

# Number of files -----

list.of.files.projections

## [1] "h08_miroc5_ewembi_rcp26_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [2] "h08_miroc5_ewembi_rcp26_rcp26soc_co2_airrww_global_monthly_2006_2099.nc4"
## [3] "h08_miroc5_ewembi_rcp60_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [4] "h08_miroc5_ewembi_rcp60_rcp60soc_co2_airrww_global_monthly_2006_2099.nc4"
## [5] "h08_miroc5_ewembi_rcp85_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [6] "lpjml_miroc5_ewembi_rcp26_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [7] "lpjml_miroc5_ewembi_rcp26_rcp26soc_co2_airrww_global_monthly_2006_2099.nc4"
## [8] "lpjml_miroc5_ewembi_rcp60_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [9] "lpjml_miroc5_ewembi_rcp85_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [10] "mpi-hm_miroc5_ewembi_picontrol_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [11] "mpi-hm_miroc5_ewembi_rcp26_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [12] "mpi-hm_miroc5_ewembi_rcp60_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [13] "pcr-globwb_miroc5_ewembi_rcp26_2005soc_co2_airrww_global_monthly_2006_2099.nc4"
## [14] "pcr-globwb_miroc5_ewembi_rcp60_2005soc_co2_airrww_global_monthly_2006_2099.nc4"

# Arrange names -----

model.names <- sub("^(.*?)_.*", "\\1", list.of.files.projections)
pattern <- "ewembi_(.*?)soc"
climate <- sub(".*ewembi_(.*?)soc.*", "\\1", list.of.files.projections)
names(isimip.future) <- paste(model.names, climate, sep = "/")

# Clean and bind dataset -----

isimip.future.dt <- rbindlist(isimip.future, idcol = "model") %>%
  na.omit() %>%
  .[, model:= factor(model)] %>%
  .[, year:= as.numeric(year)]

isimip.future.dt[, c("model", "climate") := tstrsplit(model, "/")]

# Export -----

fwrite(isimip.future.dt, "isimip.future.dt.csv")

# PLOT ISIMIP #####

# Continental level -----


isimip.future.dt[, sum(V1, na.rm = TRUE), .(year, Continent, model, climate)] %>%
  .[, climate:= gsub("_", "\\_\\_", climate)] %>%
  ggplot(., aes(year, V1, group = climate, color = climate)) +

```

```

facet_wrap(model~Continent, scales = "free_y", ncol = 5) +
geom_line() +
labs(x = "Year", y = bquote("IWW (km"3 * ")")) +
theme_AP() +
scale_x_continuous(breaks = breaks_pretty(n = 3)) +
theme(legend.position = "top")

```



```

# PLOT ISIMIP MERGED #####

```

```

a <- isimip.future.dt[, sum(V1, na.rm = TRUE), .(year, Continent, model, climate)] %>%
ggplot(., aes(year, V1, group = interaction(climate, model), color = model)) +
facet_wrap(~Continent, scales = "free_y", ncol = 5) +
geom_line() +
scale_color_manual(name = "", values = wes_palette(name = selected.palette)) +
labs(x = "Year", y = bquote("IWW (km"3 * ")")) +
scale_x_continuous(breaks = breaks_pretty(n = 3)) +
theme_AP() +
theme(legend.position = "top")

b <- isimip.future.dt[, sum(V1, na.rm = TRUE), .(year, Continent, model, climate)] %>%
ggplot(., aes(year, V1, group = interaction(climate, model), color = climate)) +
facet_wrap(~Continent, scales = "free_y", ncol = 5) +
geom_line() +

```




```

labs(x = "Year", y = bquote("IWW (km3 * ")")) +
scale_x_continuous(breaks = breaks_pretty(n = 3)) +
theme_AP() +
theme(legend.position = "top")

plot_grid(a, b, ncol = 1, labels = "auto")

```



2.3 ANOVA

```

# ANOVA #####

# Arrange ISIMIP datasets -----

isimip.full <- isimip.dt[social == "varsoc"][, context:= "historic"] %>%
  rbind(., isimip.future.dt[, context:= "prediction"], fill = TRUE) %>%
  .[, social:= NULL]

isimip.anova <- isimip.full[, .(estimation = sum(V1)),
  .(Continent, climate, context, model, year)]

# ARRANGE DATA #####

columns_to_factor <- c("Continent", "climate", "model")
isimip.full[, (columns_to_factor):= lapply(.SD, as.factor), .SDcols = (columns_to_factor)]
isimip.anova[, (columns_to_factor):= lapply(.SD, as.factor), .SDcols = (columns_to_factor)]

```

```
# RUN MODEL AND ANALYSIS OF VARIANCE #####
```

```
results.dt <- isimip.anova[, analysis_variance_fun(.SD), .(Continent, context)]
```

```
## boundary (singular) fit: see help('isSingular')
```

```
## boundary (singular) fit: see help('isSingular')
```

```
results.dt
```

```
##      Continent      context climate_variance model_variance random_variance
##      <fctr>      <char>          <num>          <num>          <num>
##  1:      Asia    historic      0.0182441856      0.9815439      1.604121e-04
##  2:     Europe    historic      0.0265735831      0.9732386      7.928135e-05
##  3:      Africa    historic      0.0046293623      0.9952289      7.974990e-05
##  4:   Americas    historic      0.0015875370      0.9983346      4.897266e-05
##  5:    Oceania    historic      0.0003011393      0.9996366      2.836314e-05
##  6:      Asia prediction      0.0144443043      0.9855396      1.802974e-21
##  7:     Europe prediction      0.0188199322      0.9811568      9.455137e-07
##  8:      Africa prediction      0.0847272814      0.9151935      1.015636e-22
##  9:   Americas prediction      0.0070916322      0.9928739      2.351915e-06
## 10:    Oceania prediction      0.0099009112      0.9899272      2.436002e-05
##      residual_variance
##      <num>
##  1:      5.146166e-05
##  2:      1.085044e-04
##  3:      6.196443e-05
##  4:      2.885478e-05
##  5:      3.387542e-05
##  6:      1.606501e-05
##  7:      2.232385e-05
##  8:      7.922237e-05
##  9:      3.208885e-05
## 10:      1.475610e-04
```

```
# PLOT RESULTS #####
```

```
a <- isimip.full[, .(estimation = sum(V1)), .(model, Continent, climate, year, context)] %>%
  ggplot(., aes(year, estimation, color = model, group = interaction(climate, model))) +
  geom_line() +
  facet_wrap(context~Continent, scale = "free", ncol = 5) +
  scale_x_continuous(breaks = breaks_pretty(n = 3)) +
  theme_AP() +
  guides(colour = guide_legend(nrow = 2)) +
  labs(x = "Year", y = bquote("IWW (km"3 * ")")) +
  theme(legend.position = "top")
```


```
b <- melt(results.dt, measure.vars = paste(c("climate", "model", "random", "residual"),
      "variance", sep = "_")) %>%
```

```

[, variance:= tstrsplit(variable, "_", fixed = TRUE)[[1]]] %>%
ggplot(., aes(Continent, value, fill = variance)) +
geom_bar(stat = "identity") +
facet_wrap(~context, ncol = 1) +
labs(x = "", y = "Fraction of variance") +
scale_fill_manual(name = "", values= wes_palette(selected.palette, n = 4)) +
scale_x_discrete(guide = guide_axis(n.dodge = 2)) +
guides(fill = guide_legend(nrow = 2)) +
theme_AP() +
theme(legend.position = "top")

plot_grid(a, b, ncol = 2, labels = "auto", rel_widths = c(0.72, 0.28))

```



COUNT COMBINATIONS OF MODEL AND CLIMATE

```

unique(isimip.full[, .(model, climate, context)]) %>%
ggplot(., aes(x = model, y = climate, fill = context)) +
geom_tile(color = "white", size = 0.5) +
scale_fill_manual(values = c("historic" = "steelblue", "prediction" = "orange")) +
theme_minimal() +
theme(axis.text.x = element_text(angle = 45, hjust = 1)) +
labs(x = "Model", y = "Climate", fill = "Context") +
scale_x_discrete(guide = guide_axis(n.dodge = 2)) +
theme_AP() +
theme(legend.position = "top")


```

```

## Warning: Using `size` aesthetic for lines was deprecated in ggplot2 3.4.0.
## i Please use `linewidth` instead.
## This warning is displayed once every 8 hours.
## Call `lifecycle::last_lifecycle_warnings()` to see where this warning was

```

generated.



3 Khan et al dataset

```
# KHAN ET AL 2023 DATASET #####

path.projections <- "./files/khan_et_al_2023"
list.of.files <- list.files(path.projections, pattern = "\\*.csv$")
combinations <- lapply(list.of.files, function(x) strsplit(x, "_")[[1]][1:4]) %>%
  do.call(rbind, .) %>%
  data.frame()
colnames(combinations) <- c("SSP", "RCP", "Climate", "Use")

# READ FILES IN PARALLEL #####

# Create parallel cluste -----

numCores <- detectCores() * 0.75
cl <- makeCluster(numCores)
registerDoParallel(cl)

# Run for loop -----

result <- foreach(i = 1:length(list.of.files),
  .combine = "rbind",
  .packages = c("data.table", "countrycode",
    "sp", "rworldmap")) %dopar% {

  out <- fread(paste("./files/khan_et_al_2023/", list.of.files
    out[, `:=`(SSP = combinations[i, 1],
      RCP = combinations[i, 2],
      Climate = combinations[i, 3],
      Use = combinations[i, 4])]
```

```

Country <- coords2country(out[1:nrow(out), 2:3])

df <- cbind(Country, out)

df[, Continent := countrycode(Country, origin = "country.name")]

df[, Dataset := list.of.files[i]]

df
}

# Stop the cluster after the computation -----
stopCluster(cl)

# ARRANGE DATA #####

numeric_cols <- grep("^([0-9])+$", names(result), value = TRUE)
khan.dt <- melt(result, measure.vars = numeric_cols, variable.name = "Year") %>%
  .[, Year:= as.numeric(as.character(Year))] %>%
  .[, model:= "GCAM"] %>%
  na.omit()

# EXPORT DATA #####

khan.dt.continent <- khan.dt[, .(estimation = sum(value)),
  .(Year, Continent, Use, RCP, SSP, Climate, Dataset, model)] %>%
  .[, climate:= paste(Climate, RCP, SSP, sep = "_")]

fwrite(khan.dt.continent, "khan.dt.continent.csv")

# PLOT #####

# Continental -----

plot.khan.continental <- khan.dt.continent %>%
  ggplot(., aes(Year, estimation, color = Continent, group = interaction(Dataset, Continent)))
  geom_line(alpha = 0.3) +
  facet_wrap(~Use) +
  theme_AP() +
  theme(legend.position = "top") +
  labs(x = "", y = bquote("km"^3))

plot.khan.continental

# PLOT #####

# Global -----

```

```

plot.khan.global <- khan.dt[, sum(value), .(Year, Use, Dataset)] %>%
  ggplot(., aes(Year, V1, group = Dataset)) +
  geom_line(alpha = 0.3) +
  facet_wrap(~Use) +
  theme_AP() +
  theme(legend.position = "top") +
  labs(x = "Year", y = bquote("km"^3))

```

```
plot.khan.global
```

```
# MERGE KHAN ET AL DATASETS #####
```

```

plot_grid(plot.khan.continental, plot.khan.global, ncol = 1, labels = "auto",
  rel_heights = c(0.53, 0.47))

```

```
# PLOT SSPs VS RCPS #####
```

```

khan.dt[, sum(value), .(Year, Use, Dataset, RCP, SSP)] %>%
  ggplot(., aes(Year, V1, group = Dataset, color = Use)) +
  geom_line() +
  facet_grid(RCP~SSP) +
  theme_AP() +
  theme(legend.position = "top") +
  labs(x = "Year", y = bquote("km"^3))

```

```
# MERGE KHAN ET AL DATA WITH ISIMIP #####
```

```
# Arrange data -----
```

```
khan.dt.continent <- fread("khan.dt.continent.csv")
```

```

khan.dt2 <- khan.dt.continent[Use == "withdrawals", .(model, Continent, climate, Year, estimation)]
  setnames(., "Year", "year")

```

```
# Extract prediction data from ISIMIP -----
```

```

isimip.full2 <- isimip.full[context == "prediction" & year >= 2010,
  .(estimation = sum(V1)), .(model, Continent, climate, year, context)] %>%
  .[, context:= NULL]

```

```
# Merge and plot -----
```


```
merged.dt <- rbind(khan.dt2, isimip.full2)
```

```

ggplot(merged.dt, aes(year, estimation, group = interaction(climate, model), color = model)) +
  geom_line(alpha = 0.4) +
  facet_wrap(~Continent, scale = "free_y", ncol = 5) +
  theme_AP() +

```

```
scale_x_continuous(breaks = breaks_pretty(n = 3)) +
theme(legend.position = "top") +
labs(x = "Year", y = bquote("km"^3))
```



Calculate the min and max in 2030-2050 given uncertainty and the global level -----

```
merged.dt[year %in% c(2030, 2040, 2050),
           .(min = min(estimation), max = max(estimation)), .(Continent, year)] %>%
  .[, .(sum_min = sum(min), sum_max = sum(max)), year]
```

```
##      year  sum_min  sum_max
##      <num>    <num>    <num>
## 1:  2030  272.8320 2529.235
## 2:  2040  281.8063 2958.560
## 3:  2050  278.4169 3188.283
```

4 Bibliographical study

NAOMI DATASET

```
references.projected <- data.table(read.xlsx("references_projection.xlsx")) %>%
  .[, focus:= "projected"]
```

```
references.current <- data.table(read.xlsx("references_current.xlsx")) %>%
  .[, focus:= "current"]
```

```
references.full.dt <- rbind(references.projected, references.current) %>%
  .[, study:= paste(author, model, climate.scenario, sep = ".")]
```

CLEAN THE DATASET

```
colnames_vector <- c("title", "author", "region")
```

Remove leading and trailing spaces -----

```
references.full.dt[, (colnames_vector):= lapply(.SD, trimws), .SDcols = (colnames_vector)]
references.full.dt[, (colnames_vector):= lapply(.SD, str_squish), .SDcols = (colnames_vector)]
```

```

# Lowercaps -----
references.full.dt[, (colnames_vector):= lapply(.SD, tolower), .SDcols = (colnames_vector)]

# Remove multiple spaces -----
references.full.dt[, (colnames_vector):= lapply(.SD, function(x)
  gsub("\\s+", " ", x)), .SDcols = (colnames_vector)]

# Correct America -----
references.full.dt[, region:= ifelse(region == "america", "americas", region)]

# Extract the publication year -----
references.full.dt[, publication.date:= str_extract(author, "\\d{4}")] %>%
  .[, publication.date:= as.numeric(publication.date)]

# FEATURES OF THE DATASET #####

references.full.dt[, range.estimation:= ifelse(publication.date >= 1990 & publication.date < 20
  ifelse(publication.date >= 2000 & publication.date < 2010,
    "2010-2024"))]

# RANGE ESTIMATION #####

references.full.dt[, range.estimation.year:= ifelse(estimation.year >= 1990 & estimation.year < 2010,
  ifelse(estimation.year >= 2010 & publication.date < 2024,
    "2050-2100"))]

# FEATURES OF THE DATASET #####

# Name of different studies -----

sort(unique(references.full.dt$title))

## [1] "a global hydrological simulation to specify the sources of water used by humans"
## [2] "a global water scarcity assessment under shared socio-economic pathways - part 2: wat
## [3] "a pathway of global food supply adaptation in a world with increasingly constrained g
## [4] "agricultural green and blue water consumption and its influence on the global water s
## [5] "agriculture, bioenergy, and water implications of constrained cereal trade and climat
## [6] "an estimation of global virtual water flow and sources of water withdrawal for major c
## [7] "an integrated assessment of global and regional water demands for electricity generat
## [8] "an interpreted model for the assessment of global water resources - part 2: applicati
## [9] "appraisal and assessment of world water resources"
## [10] "aquastat: fao's global information system on water and agriculture"
## [11] "climate change impacts on irrigation water requirements: effects of mitigation, 1990-2

```


[12] "climate policy implications for agricultural water demand"
 ## [13] "cooling water sufficiency in a warming world: projection using an integrated assessment"
 ## [14] "future long-term changes in global water resources driven by socio-economic and climate change"
 ## [15] "global and regional evaluation of energy for water"
 ## [16] "global impacts of conversions from natural to agricultural ecosystems on water resources"
 ## [17] "global irrigation characteristics and effects simulated by fully coupled land surface models"
 ## [18] "global irrigation water demand: variability and uncertainties arising from agricultural practices"
 ## [19] "global modeling of irrigation water requirements"
 ## [20] "global monthly sectoral water use for 2010-2100 at 0.5° resolution across alternative scenarios"
 ## [21] "global water demand and supply projections"
 ## [22] "globwat - a global water balance model to assess water use in irrigated agriculture"
 ## [23] "groundwater use for irrigation- a global inventory"
 ## [24] "how can we cope with the water resources situation by the year 2050?"
 ## [25] "human appropriation of renewable fresh water"
 ## [26] "human-induced changes in the global water cycle"
 ## [27] "impact of climate change and variability on irrigation requirements: a global perspective"
 ## [28] "impact of climate forcing uncertainty and human water use on global and continental water resources"
 ## [29] "impact of water withdrawals from groundwater and surface water on continental water resources"
 ## [30] "implementation and evaluation of irrigation techniques in the community land model"
 ## [31] "incorporating anthropogenic water regulation modules into a land surface model"
 ## [32] "incorporation of groundwater pumping in a global land surface model with the representation of soil moisture"
 ## [33] "isimip database"
 ## [34] "long-term global water projections using six socioeconomic scenarios in an integrated assessment model"
 ## [35] "modelling global water stress of the recent past: on the relative importance of trends and variability"
 ## [36] "pcr-globwb 2: a 5 arcmin global hydrological and water resources model"
 ## [37] "physical impacts of climate change on water resources"
 ## [38] "planetary boundaries: guiding human development on a changing planet"
 ## [39] "present-day irrigation mitigates heat extremes"
 ## [40] "projected water consumption in future global agriculture: scenarios and related impacts"
 ## [41] "projection of future world water resources under sres scenarios: water withdrawal"
 ## [42] "recent global cropland water consumption constrained by observations"
 ## [43] "reconciling irrigated food production with environmental flows for sustainable development"
 ## [44] "reconstructing 20th century global hydrography: a contribution to the global terrestrial water cycle"
 ## [45] "reconstructing 20th century global hydrography: a contribution to the global terrestrial water cycle"
 ## [46] "sustainability of global water use: past reconstruction and future projections"
 ## [47] "the number of people exposed to water stress in relation to how much water is reserved for agriculture"
 ## [48] "the state of the world's land and water resources for food and agriculture"
 ## [49] "the world's water, 2000-2001: the biennial report on freshwater resources"
 ## [50] "united nations world water development report 2020: water and climate change"
 ## [51] "water and sustainability: a global outlook"
 ## [52] "water and sustainability. global pattern and long-range problems"
 ## [53] "water resources for sustainable development"
 ## [54] "water sector assumptions for the shared socioeconomic pathways in an integrated model"
 ## [55] "world agriculture towards 2030/2050"
 ## [56] "world agriculture towards 2030/2050: the 2012 revision"
 ## [57] "world agriculture towards 2030/2053"
 ## [58] "world agriculture towards 2030/2054"
 ## [59] "world agriculture towards 2030/2055"

```
## [60] "world water in 2025 - global modeling and scenario analysis for the world commission o
## [61] "world water resources and water use: modern assessment and outlook for future"
```

```
# Name of different studies per variable -----
```

```
references.full.dt[, unique(title), variable] %>%
  .[, .N, variable]
```

```
##      variable      N
##      <char> <int>
## 1:      iww      36
## 2:      tww      34
## 3:      iwc      14
## 4:      twc       7
## 5:      iwr       3
```


```
# Cumulative sum of published studies -----
```

```
dt <- references.full.dt[, .(title, publication.date)] %>%
  .[!duplicated(.)] %>%
  setorder(., publication.date) %>%
  .[, .N, publication.date] %>%
  .[, cumulative_sum := cumsum(N)]
```

```
ggplot(dt, aes(publication.date, cumulative_sum)) +
  geom_line() +
  geom_point(size = 0.7) +
  theme_AP() +
  labs(x = "Year", y = "N° studies")
```

```
## Warning: Removed 1 row containing missing values or values outside the scale range
## (`geom_line()`).
```

```
## Warning: Removed 1 row containing missing values or values outside the scale range
## (`geom_point()`).
```



```
# Only irrigation water withdrawal studies -----
```

```
references.full.dt[, .(title, publication.date, variable)] %>%
  .[variable == "iww"] %>%
```

```


[!duplicated(.)] %>%
setorder(., publication.date) %>%
[, .N, publication.date] %>%
[, cumulative_sum := cumsum(N)] %>%
ggplot(., aes(publication.date, cumulative_sum)) +
geom_line() +
geom_point(size = 0.7) +
theme_AP() +
labs(x = "Year", y = "N° studies")

```

```

## Warning: Removed 1 row containing missing values or values outside the scale range
## (`geom_line()`).
## Removed 1 row containing missing values or values outside the scale range
## (`geom_point()`).

```



```

# PLOT ALL ESTIMATIONS #####


```

```

def.alpha <- 0.3

references.full.dt[variable == "iww" & region == "global"] %>%
[, .(author, study, estimation.year, value)] %>%
na.omit() %>%
ggplot(., aes(estimation.year, value, color = author, group = study)) +
geom_point(alpha = def.alpha, size = 0.2) +
labs(x = "Year", y = bquote("Km"^3)) +
scale_color_discrete(name = "") +
geom_line(alpha = def.alpha) +
theme_AP()

```



5 Session information

```
# SESSION INFORMATION #####
```

```
sessionInfo()
```

```
## R version 4.3.3 (2024-02-29)
## Platform: aarch64-apple-darwin20 (64-bit)
## Running under: macOS Sonoma 14.2.1
##
## Matrix products: default
## BLAS: /Library/Frameworks/R.framework/Versions/4.3-arm64/Resources/lib/libRblas.0.dylib
## LAPACK: /Library/Frameworks/R.framework/Versions/4.3-arm64/Resources/lib/libRlapack.dylib;
##
## locale:
## [1] en_US.UTF-8/en_US.UTF-8/en_US.UTF-8/C/en_US.UTF-8/en_US.UTF-8
##
## time zone: Europe/London
## tzcode source: internal
##
## attached base packages:
## [1] parallel stats graphics grDevices utils datasets methods
## [8] base
##
## other attached packages:
## [1] microbenchmark_1.5.0 lme4_1.1-35.5 Matrix_1.6-5
## [4] here_1.0.1 doParallel_1.0.17 iterators_1.0.14
## [7] foreach_1.5.2 rworldmap_1.3-8 sp_2.1-4
## [10] countrycode_1.6.0 ncd4_1.23 scales_1.3.0
## [13] wesanderson_0.3.7 benchmarkme_1.0.8 cowplot_1.1.3
## [16] lubridate_1.9.3 forcats_1.0.0 stringr_1.5.1
## [19] dplyr_1.1.4 purrr_1.0.2 readr_2.1.5
## [22] tidyr_1.3.1 tibble_3.2.1 ggplot2_3.5.1
## [25] tidyverse_2.0.0 data.table_1.16.2 openxlsx_4.2.7.1
##
## loaded via a namespace (and not attached):
## [1] dotCall64_1.2 benchmarkmeData_1.0.4 gtable_0.3.6
## [4] spam_2.11-0 xfun_0.49 raster_3.6-30
## [7] lattice_0.22-6 tzdb_0.4.0 Rdpack_2.6.2
## [10] vctrs_0.6.5 tools_4.3.3 generics_0.1.3
## [13] fansi_1.0.6 pkgconfig_2.0.3 lifecycle_1.0.4
## [16] compiler_4.3.3 fields_16.3 munsell_0.5.1
## [19] terra_1.7-78 codetools_0.2-20 htmltools_0.5.8.1
## [22] maps_3.4.2.1 yaml_2.3.10 nloptr_2.1.1
## [25] pillar_1.9.0 MASS_7.3-60.0.1 boot_1.3-31
## [28] nlme_3.1-166 tidyselect_1.2.1 zip_2.3.1
## [31] digest_0.6.37 stringi_1.8.4 splines_4.3.3
## [34] sensobol_1.1.5 rprojroot_2.0.4 fastmap_1.2.0
```

```
## [37] grid_4.3.3           colorspace_2.1-1      cli_3.6.3
## [40] magrittr_2.0.3        utf8_1.2.4           withr_3.0.2
## [43] timechange_0.3.0      rmarkdown_2.29       httr_1.4.7
## [46] hms_1.1.3            evaluate_1.0.1       knitr_1.49
## [49] rbibutils_2.3         viridisLite_0.4.2    rlang_1.1.4
## [52] Rcpp_1.0.13-1         glue_1.8.0           minqa_1.2.8
## [55] rstudioapi_0.17.1     R6_2.5.1
```

```
## Return the machine CPU -----
```

```
cat("Machine:      "); print(get_cpu()$model_name)
```

```
## Machine:
```

```
## [1] "Apple M1 Max"
```

```
## Return number of true cores -----
```

```
cat("Num cores:    "); print(detectCores(logical = FALSE))
```

```
## Num cores:
```

```
## [1] 10
```

```
## Return number of threads -----
```

```
cat("Num threads: "); print(detectCores(logical = FALSE))
```

```
## Num threads:
```

```
## [1] 10
```