

Notes on Lattices

arnaucube

2026

Abstract

Notes taken while studying lattice-based cryptography.

Usually while reading books and papers I take handwritten notes in paper notebooks, this document contains some of them re-written to *LaTeX*.

Contents

1	Lattices	1
1.1	Hard problems	1
1.2	Dual lattices	2

1 Lattices

1.1 Hard problems

Definition 1.1 (SVP (Shortest Vector Problem)). find a shortest non-zero vector $x \in \mathcal{L}$, ie. that minimizes the Euclidean norm $\|x\|$, that is $\lambda_1(\mathcal{L}) = \|x\|$.

Definition 1.2 (CVP (Closest Vector Problem)). given a vector $y \in \mathbb{R}^n$ with $y \notin \mathcal{L}$, find a nonzero vector $x \in \mathcal{L}$ that is closest to y , ie. that minimizes the Euclidean norm $\|y - x\|$.

Definition 1.3 (NTRU key recovery problem). Given $h(x)$, find ternary polynomials $f, g \in \mathcal{T}$ satisfying

$$f(x) \cdot h(x) = g(x) \pmod{q} \in R_q$$

Where $R_q = \mathbb{Z}_q[X]/(X^n - 1)$.

Definition 1.4 (SIS (Short Integer Solution)). Given $q \in \mathbb{Z}$ and matrix $A \in \mathbb{Z}_q^{m \times n}$, find a non-zero short $z \in \mathbb{Z}^n$ s.th.

$$A \cdot z = 0 \pmod{q}$$

"short": $z \in \{B, B\}^n$ for some bound B , often $z \in \{-1, 0, 1\}^n$.

The set of all solutions to the SIS problem forms the (q-ary) lattice

$$\Lambda_q^\perp(A) = \{z \in \mathbb{Z}^n \mid Az = 0 \in \mathbb{Z}_q^m\} \supseteq \mathbb{Z}_q^n$$

where the SIS problem becomes the SVP for $\Lambda_q^\perp(A)$

- LLL algorithm is used to solve SVP and CVP.
- Both SVP and CVP become computationally difficult (NP hard) as the lattice dimension n grows.

Definition 1.5 (Ring SIS (Ring Short Integer Solution)). given a uniformly random matrix $A \in^R R_q^n$, with elements $a_i \in R_q$, find a non-zero integer vector $z \in R^n$ of norm $\|z\| \leq \beta$ such that

$$Az = \sum_i a_i \cdot z_i = 0 \in R_q$$

Definition 1.6 (MSIS (Module Short Integer Solution)). given a uniformly random matrix $A \in^R R_q^{m \times n}$, with columns $a_i \in R_q^m$, find a non-zero integer vector $z \in R^n$ of norm $\|z\| \leq \beta$ such that

$$Az = 0 \in R_q^m$$

$$\text{ie. } \sum_i a_{ij} \cdot z_j = 0 \in R_q, \quad \text{for } i = 1, \dots, m$$

Notice that Module-SIS is a Ring-SIS with $rank = 1$ ($m = 1$).

1.2 Dual lattices

From previous section, we have LWE-Lattice and SIS-Lattice:

Definition 1.7 (LWE Lattice). A LWE-Lattice $\mathcal{L}$ is defined

$$\begin{aligned} \mathcal{L} &= \{A^T s \mid s \in \mathbb{Z}_q^n\} + q\mathbb{Z}^m \\ &= \{A^T s + qz \mid s \in \mathbb{Z}_q^n, z \in \mathbb{Z}^m\} \end{aligned}$$

Definition 1.8 (SIS-Lattice). A SIS-Lattice $\mathcal{L}$ is defined

$$\mathcal{L}^\perp = \{z \in \mathbb{Z}^m \mid Az = 0 \in \mathbb{Z}_q^n\} \supseteq q\mathbb{Z}^m$$

Definition 1.9 (Dual of a lattice). For a lattice $\mathcal{L}$, its dual is

$$\mathcal{L}^* = \{y \in \mathbb{R}^m \mid \langle y, x \rangle \in \mathbb{Z} \forall x \in \mathcal{L}\}$$

Lemma 1.10.

$$\mathcal{L} = q\mathcal{L}^{\perp*} \text{ ie. } \frac{1}{q}\mathcal{L} = \mathcal{L}^{\perp*}$$

Proof. i. Take $y \in \frac{1}{q}\mathcal{L}$, then by 1.7, $\exists s \in \mathbb{Z}^n, z \in \mathbb{Z}^m$ such that

$$y = \frac{1}{q}(A^T s + qz) = \frac{1}{q}A^T s + z$$

Let $x \in \mathcal{L}^\perp$. By definition (1.8) $Ax = 0 \pmod{q}$, therefore $\exists k \in \mathbb{Z}^n$ such that $Ax = qk$.

Compute

$$\begin{aligned} \langle y, x \rangle &= \left\langle \frac{1}{q}A^T s + z, x \right\rangle \\ &= \frac{1}{q}\langle A^T s, x \rangle + \langle z, x \rangle \\ &= \frac{1}{q}\langle s, Ax \rangle + \langle z, x \rangle \\ &= \frac{1}{q}\langle s, qk \rangle + \langle z, x \rangle \\ &= \langle s, k \rangle + \langle z, x \rangle \end{aligned}$$

and $\langle s, k \rangle + \langle z, x \rangle \in \mathbb{Z}$, since each of $s, k, z, x \in \mathbb{Z}$ separately.

Thus $\langle y, x \rangle \in \mathbb{Z}$, hence by 1.9, $y \in \mathcal{L}^{\perp*}$.

We started from $y \in \frac{1}{q}\mathcal{L}$, so, now we have

$$y \in \mathcal{L}^{\perp*} \quad \forall y \in \frac{1}{q}\mathcal{L}$$

therefore

$$\mathcal{L}^{\perp*} \supseteq \frac{1}{q}\mathcal{L}$$

ii. Now we prove the reverse inclusion. Take $y \in \mathcal{L}^{\perp*}$.

Observe that $q\mathbb{Z}^m \subseteq \mathcal{L}^\perp$, because

$$\forall z \in \mathbb{Z}^m, A(qz) = qAz = q0 = 0 \pmod{q}$$

Since $y \in \mathcal{L}^{\perp*}$, we must have $\langle y, qz \rangle \in \mathbb{Z} \quad \forall z \in \mathbb{Z}^m$.

Let $e_i \in \mathbb{Z}^m$ be the i -th standard basis vector, ie. $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ (where one is at the i -th position).

We know that $\langle y, qz \rangle \in \mathbb{Z} \quad \forall z \in \mathbb{Z}^m$ (including $z = e_i$), thus, for $qe_i = (0, \dots, 0, q, 0, \dots, 0)$,

$$\langle y, qe_i \rangle = q\langle y, e_i \rangle = q \cdot y_i$$

since $\langle y, e_i \rangle = y_i$ (extracts the i -th position from y).

So, since $\langle y, qz \rangle \in \mathbb{Z} \quad \forall z \in \mathbb{Z}^m$, and $\langle y, qe_i \rangle = qy_i$, then $qy_i \in \mathbb{Z}$

This argument works $\forall i \in [m]$, that is, $qy_1, \dots, qy_m \in \mathbb{Z}$. Therefore

$$qy \in \mathbb{Z}^m \quad \forall y \in \mathcal{L}^{\perp*}$$

For $\bar{x} \in \ker(A) \subseteq \mathbb{Z}_q^m$, since $A\bar{x} = 0$, then $Ax = 0 \pmod{q}$, thus $x \in \mathcal{L}^{\perp}$ (by its definition 1.8).

Since $y \in \mathcal{L}^{\perp*}$, by the definition of $\mathcal{L}^{\perp*}$ (1.9), $\langle y, x \rangle \in \mathbb{Z}$. Multiply it by q ,

$$q\langle y, x \rangle = \langle qy, x \rangle \in q\mathbb{Z}$$

hence $\langle qy, x \rangle = 0 \pmod{q}$, thus $qy \pmod{q} \in \ker(A)^{\perp}$.

We'll now prove the following identity, valid for every integer $q \geq 2$, which we will use later:

$$(\ker A)^{\perp} = \text{im}(A^T)$$

where $A : \mathbb{Z}_q^m \rightarrow \mathbb{Z}_q^n$.

Let $y \in \text{im}(A^T)$. Then $\exists s \in \mathbb{Z}_q^n$ such that $y = A^T s$.

Then, $\forall x \in \ker(A)$, by definition of the kernel, $Ax = 0$. Want to prove that y is orthogonal to every $x \in \ker(A)$:

$$\langle y, x \rangle = \langle A^T s, x \rangle = \langle s, Ax \rangle = 0$$

thus, $\langle y, x \rangle = 0 \quad \forall x \in \ker(A)$, hence y is orthogonal to every x , that is, $y \in (\ker(A))^{\perp}$.

Therefore,

$$\text{im}(A^T) \subseteq (\ker(A))^{\perp}$$

For the reverse inclusion, it suffices to show that both sets have the same number of elements:

By the First Isomorphism Theorem,

$$|\text{im}(A)| = \frac{|\mathbb{Z}_q^m|}{|\ker(A)|} = \frac{q^m}{|\ker(A)|}$$

Notice that for every subgroup $K \subseteq \mathbb{Z}_q^m$, the dot product on $\mathbb{Z}_q^m$ is nondegenerate, therefore

$$|K^{\perp}| = \frac{q^m}{|K|}$$

so, applied to $\ker(A)$,

$$|\ker(A)^{\perp}| = \frac{q^m}{|\ker(A)|} = |\text{im}(A)|$$

which, by transposition properties, $|im(A)| = |im(A^T)|$, thus

$$|\ker(A)^\perp| = |im(A^T)|$$

hence,

$$(\ker(A)^\perp) = im(A^T)$$

Continuing from $qy \pmod q \in (\ker(A))^\perp$, we've just seen that $(\ker(A))^\perp = im(A^T)$, thus

$$\exists s \in \mathbb{Z}^n \text{ s.th. } qy = A^T s \pmod q$$

Therefore, for some

$$z \in \mathbb{Z}^m, \quad qy = A^T s + qz$$

which means that $qy \in \mathcal{L}$, by definition of $\mathcal{L}$ (1.7). Divide that by q , obtaining

$$y \in \frac{1}{q}\mathcal{L}$$

Thus, $\forall y \in \mathcal{L}^{\perp*}$, we have $y \in \frac{1}{q}\mathcal{L}$.

Hence $\mathcal{L}^{\perp*} \subseteq \frac{1}{q}\mathcal{L}$.

Putting together (i). $\mathcal{L}^{\perp*} \supseteq \frac{1}{q}\mathcal{L}$, and (ii). $\mathcal{L}^{\perp*} \subseteq \frac{1}{q}\mathcal{L}$, we have

$$\mathcal{L}^{\perp*} = \frac{1}{q}\mathcal{L}$$

□

References

- [1] Chris Peikert. A decade of lattice cryptography, 2016.
- [2] Matan Pasma. Lattices, an elementary and rigorous account, 2024.
- [3] Daniele Micciancio. Introduction to Lattices, 2024.
- [4] Daniele Micciancio and Oded Regev. Lattice-based Cryptography, 2008.
- [5] Alfred Menezes. A Gentle Introduction to Lattice-Based Cryptography, 2026.
- [6] Cynthia Dwork. Lattices and Their Application to Cryptography, 1998.