

Notes on Lattices

arnaucube

2026

Abstract

Notes taken while studying lattice-based cryptography.

Usually while reading books and papers I take handwritten notes in paper notebooks, this document contains some of them re-written to *LaTeX*.

Contents

1	Lattices	1
1.1	Hard problems	1
1.2	Dual lattices	2

1 Lattices

1.1 Hard problems

Definition 1.1 (SVP (Shortest Vector Problem)). find a shortest non-zero vector $x \in \mathcal{L}$, ie. that minimizes the Euclidean norm $\|x\|$, that is $\lambda_1(\mathcal{L}) = \|x\|$.

Definition 1.2 (CVP (Closest Vector Problem)). given a vector $y \in \mathbb{R}^n$ with $y \notin \mathcal{L}$, find a nonzero vector $x \in \mathcal{L}$ that is closest to y , ie. that minimizes the Euclidean norm $\|y - x\|$.

Definition 1.3 (NTRU key recovery problem). Given $h(x)$, find ternary polynomials $f, g \in \mathcal{T}$ satisfying

$$f(x) \cdot h(x) = g(x) \pmod{q} \in R_q$$

Where $R_q = \mathbb{Z}_q[X]/(X^n - 1)$.

Definition 1.4 (SIS (Short Integer Solution)). Given $q \in \mathbb{Z}$ and matrix $A \in \mathbb{Z}_q^{m \times n}$, find a non-zero short $z \in \mathbb{Z}^n$ s.th.

$$A \cdot z = 0 \pmod{q}$$

"short": $z \in \{B, B\}^n$ for some bound B , often $z \in \{-1, 0, 1\}^n$.

The set of all solutions to the SIS problem forms the (q-ary) lattice

$$\Lambda_q^\perp(A) = \{z \in \mathbb{Z}^n \mid A \cdot z = 0 \pmod{q}\}$$

where the SIS problem becomes the SVP for $\Lambda_q^\perp(A)$

- LLL algorithm is used to solve SVP and CVP.
- Both SVP and CVP become computationally difficult (NP hard*) as the lattice dimension n grows.

Definition 1.5 (Ring SIS (Ring Short Integer Solution)). given a uniformly random matrix $A \in^R R_q^n$, with elements $a_i \in R_q$, find a non-zero integer vector $z \in R^n$ of norm $\|z\| \leq \beta$ such that

$$Az = \sum_i a_i \cdot z_i = 0 \in R_q$$

Definition 1.6 (MSIS (Module Short Integer Solution)). given a uniformly random matrix $A \in^R R_q^{m \times n}$, with columns $a_i \in R_q^m$, find a non-zero integer vector $z \in R^n$ of norm $\|z\| \leq \beta$ such that

$$Az = 0 \in R_q^m$$

ie. $\sum_i a_{ij} \cdot z_j = 0 \in R_q, \text{ for } i = 1, \dots, m$

Notice that Module-SIS is a Ring-SIS with $rank = 1$ ($m = 1$).

1.2 Dual lattices

Definition 1.7 (Dual of a lattice). For a lattice $\mathcal{L}$, its dual is

$$\mathcal{L}^* = \{y \in \mathbb{R}^m \mid \langle y, x \rangle \in \mathbb{Z} \forall x \in \mathcal{L}\}$$

Lemma 1.8.

$$\mathcal{L} = q\mathcal{L}^{\perp*} \text{ ie. } \frac{1}{q}\mathcal{L} = \mathcal{L}^{\perp*}$$

Proof. i. Take $y \in \frac{1}{q}\mathcal{L}$, then $\exists s \in \mathbb{Z}^n, z \in \mathbb{Z}^m$ such that

$$y = \frac{1}{q}(A^T s + qz) = \frac{1}{q}A^T s + z$$

Let $x \in \mathcal{L}^\perp$. By definition $Ax = 0 \pmod{q}$, therefore $\exists k \in \mathbb{Z}^n$ such that $Ax = qk$.

Compute

$$\begin{aligned}
\langle y, x \rangle &= \left\langle \frac{1}{q} A^T s + z, x \right\rangle \\
&= \frac{1}{q} \langle A^T s, x \rangle + \langle z, x \rangle \\
&= \frac{1}{q} \langle s, Ax \rangle + \langle z, x \rangle \\
&= \frac{1}{q} \langle s, qk \rangle + \langle z, x \rangle \\
&= \langle s, k \rangle + \langle z, x \rangle
\end{aligned}$$

and $\langle s, k \rangle + \langle z, x \rangle \in \mathbb{Z}$, since each of $s, k, z, x \in \mathbb{Z}$.

Thus $\langle y, x \rangle \in \mathbb{Z}$, hence by 1.7, $y \in \mathcal{L}^{\perp*}$.

We started from $y \in \frac{1}{q}\mathcal{L}$, so, now we have

$$y \in \mathcal{L}^{\perp*} \quad \forall y \in \frac{1}{q}\mathcal{L}$$

therefore

$$\mathcal{L}^{\perp*} \supseteq \frac{1}{q}\mathcal{L}$$

ii. Now we prove the reverse inclusion. Take $y \in \mathcal{L}^{\perp*}$.

Observe that $q\mathbb{Z} \subseteq \mathcal{L}^{\perp}$, because

$$\forall z \in \mathbb{Z}^m, \quad A(qz) = qAz = q0 = 0 \pmod{q}$$

Since $y \in \mathcal{L}^{\perp*}$, we must have $\langle y, qz \rangle \in \mathbb{Z} \quad \forall z \in \mathbb{Z}^m$.

Let $e_i \in \mathbb{Z}^m$ be the i -th standard basis vector, ie. $e_i = (0, \dots, 0, 1, 0, \dots, 0)$ (where one is at the i -th position).

We know that $\langle y, qz \rangle \in \mathbb{Z} \quad \forall z \in \mathbb{Z}^m$ (including $z = e_i$), thus $qe_i = (0, \dots, 0, q, 0, \dots, 0)$. Therefore,

$$\langle y, qe_i \rangle = q\langle y, e_i \rangle = q \cdot y_i$$

since $\langle y, e_i \rangle = y_i$ (extracts the i -th position from y).

So, since $\langle y, qz \rangle \in \mathbb{Z} \quad \forall z \in \mathbb{Z}^m$, and $\langle y, qe_i \rangle = qy_i$, then $qy_i \in \mathbb{Z}$

This argument works $\forall i \in [m]$, that is, $qy_1, \dots, qy_m \in \mathbb{Z}$. Therefore

$$qy \in \mathbb{Z}^m \quad \forall y \in \mathcal{L}^{\perp*}$$

$qy \pmod{q}$ is orthogonal to $\ker(A)$.

Let $\hat{x} \in \ker(A) \subseteq \mathbb{Z}_q^m$. Since $A\hat{x} = 0$, then $Ax = 0 \pmod{q}$, thus $x \in \mathcal{L}^{\perp}$.

Since $y \in \mathcal{L}^{\perp*}$, $\langle y, x \rangle \in \mathbb{Z}$. Multiply it by q ,

$$\langle qy, x \rangle = q\langle y, x \rangle \in q\mathbb{Z}$$

hence $\langle qy, x \rangle = 0 \pmod{q}$, thus $qy \pmod{q} \in \ker(A)^{\perp}$.

Observe: $\ker(A)^\perp = \text{im}(A^T)$, proof: let $A : \mathbb{Z}_q^m \rightarrow \mathbb{Z}_q^n$, define

$$\ker(A) = \{x \in \mathbb{Z}_q^m \mid Ax = 0\} = \mathcal{L}^\perp(A)$$

Define the orthogonal complement

$$(\ker(A))^\perp = \{y \in \mathbb{Z}_q^n \mid \langle y, x \rangle = 0 \pmod{q} \ \forall x \in \ker(A)\}$$

$$\text{im}(A^T) = \{A^T s \mid s \in \mathbb{Z}_q^n\}$$

Then, $(\ker(A))^\perp = \text{im}(A^T)$.

Continuing from $qy \pmod{q} \in (\ker(A))^\perp$, we've just seen that $(\ker(A))^\perp = \text{im}(A^T)$, thus

$$\exists s \in \mathbb{Z}^n \text{ s.t. } qy = A^T s \pmod{q}$$

Therefore, for some

$$z \in \mathbb{Z}^m, \ qy = A^T s + qz$$

which means that $qy \in \mathcal{L}$ (by definition of $\mathcal{L}$). Divide that by q , obtaining

$$y \in \frac{1}{q}\mathcal{L}$$

Thus, $\forall y \in \mathcal{L}^{\perp*}$, we have $y \in \frac{1}{q}\mathcal{L}$.

Hence $\mathcal{L}^{\perp*} \subseteq \frac{1}{q}\mathcal{L}$.

Putting together (i). $\mathcal{L}^{\perp*} \supseteq \frac{1}{q}\mathcal{L}$, and (ii). $\mathcal{L}^{\perp*} \subseteq \frac{1}{q}\mathcal{L}$, we have

$$\mathcal{L}^{\perp*} = \frac{1}{q}\mathcal{L}$$

□

References

- [1] Chris Peikert. A decade of lattice cryptography, 2016.
- [2] Daniele Micciancio. Introduction to Lattices, 2024.
- [3] Daniele Micciancio and Oded Regev. Lattice-based Cryptography, 2008.
- [4] Matan Prasma. Lattices, an elementary and rigorous account, 2024.
- [5] Alfred Menezes. A Gentle Introduction to Lattice-Based Cryptography, 2026.
- [6] Cynthia Dwork. Lattices and Their Application to Cryptography, 1998.