

A variational derivation of the field equations of an action-dependent Einstein-Hilbert Lagrangian

Arnau Mas

Supervised by
Dr Jordi Gaset

Universitat Autònoma de Barcelona
June 27th 2021



- ① Introduction
- ② The Herglotz variational problem
- ③ Action-dependent Einstein gravity
- ④ Significance of the equations
- ⑤ Conclusions

- ➊ Introduction
- ➋ The Herglotz variational problem
- ➌ Action-dependent Einstein gravity
- ➍ Significance of the equations
- ➎ Conclusions

- Modified theories of gravity: $f(R)$, Brans-Dicke, etc. to answer questions from cosmology

- Modified theories of gravity: $f(R)$, Brans-Dicke, etc. to answer questions from cosmology
- Action-dependent Lagrangian mechanics: variational formulation of dissipative systems, applied in many other fields (thermodynamics, control theory).

- Modified theories of gravity: $f(R)$, Brans-Dicke, etc. to answer questions from cosmology
- Action-dependent Lagrangian mechanics: variational formulation of dissipative systems, applied in many other fields (thermodynamics, control theory).

Idea: what does an action-dependent version of Einstein gravity look like?

- Writing down the equations of action-dependent GR has two main challenges: it is a field theory and it is a *second order* field theory.

- Writing down the equations of action-dependent GR has two main challenges: it is a field theory and it is a *second order* field theory.
- An attempt was made in [Laz+18], but we claim the equations found there are not the correct ones. We will use an alternative formulation of the action-dependent theory which allows us to treat second order theories.

- ① Introduction
- ② The Herglotz variational problem
- ③ Action-dependent Einstein gravity
- ④ Significance of the equations
- ⑤ Conclusions

If we allow the Lagrangian to depend on the action, then we have

$$S[q^\mu](t) = \int_a^t L(q^\mu(\tau), \dot{q}^\mu(\tau), S[q^\mu](\tau)) \, d\tau$$

If we allow the Lagrangian to depend on the action, then we have

$$S[q^\mu](t) = \int_a^t L(q^\mu(\tau), \dot{q}^\mu(\tau), S[q^\mu](\tau)) \, d\tau$$

and differentiating

$$\dot{S}[q^\mu](t) = L(q^\mu(t), \dot{q}^\mu(t), S[q^\mu](t)).$$

If we allow the Lagrangian to depend on the action, then we have

$$S[q^\mu](t) = \int_a^t L(q^\mu(\tau), \dot{q}^\mu(\tau), S[q^\mu](\tau)) \, d\tau$$

and differentiating

$$\dot{S}[q^\mu](t) = L(q^\mu(t), \dot{q}^\mu(t), S[q^\mu](t)).$$

So, solve this ODE for any given trajectory and then find which paths are extrema of $S[q^\mu](a) - S[q^\mu](b)$.

If we allow the Lagrangian to depend on the action, then we have

$$S[q^\mu](t) = \int_a^t L(q^\mu(\tau), \dot{q}^\mu(\tau), S[q^\mu](\tau)) \, d\tau$$

and differentiating

$$\dot{S}[q^\mu](t) = L(q^\mu(t), \dot{q}^\mu(t), S[q^\mu](t)).$$

So, solve this ODE for any given trajectory and then find which paths are extrema of $S[q^\mu](a) - S[q^\mu](b)$. Not very elegant!

Consider expanded trajectories q^μ, z , so z tracks *some quantity* along the path.

Consider expanded trajectories q^μ, z , so z tracks *some quantity* along the path. If we want z to be the change in action then we require

$$\dot{z}(t) = L(q^\mu(t), \dot{q}^\mu(t), z(t)).$$

Consider expanded trajectories q^μ, z , so z tracks *some quantity* along the path. If we want z to be the change in action then we require

$$\dot{z}(t) = L(q^\mu(t), \dot{q}^\mu(t), z(t)).$$

So the action functional becomes

$$S[q^\mu, z] = z(b) - z(a).$$

Consider expanded trajectories q^μ, z , so z tracks *some quantity* along the path. If we want z to be the change in action then we require

$$\dot{z}(t) = L(q^\mu(t), \dot{q}^\mu(t), z(t)).$$

So the action functional becomes

$$S[q^\mu, z] = z(b) - z(a).$$

We want to find the extrema of this functional

Consider expanded trajectories q^μ, z , so z tracks *some quantity* along the path. If we want z to be the change in action then we require

$$\dot{z}(t) = L(q^\mu(t), \dot{q}^\mu(t), z(t)).$$

So the action functional becomes

$$S[q^\mu, z] = z(b) - z(a).$$

We want to find the extrema of this functional *subject to the constraint* that z actually be the action.

Turns out, finding the extrema subject to the constraint is equivalent to finding the extrema of

$$\tilde{S}[q^\mu, z, \lambda] = S[q^\mu, z] - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt$$

Turns out, finding the extrema subject to the constraint is equivalent to finding the extrema of

$$\begin{aligned}\tilde{S}[q^\mu, z, \lambda] &= S[q^\mu, z] - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt \\ &= \int_a^b \dot{z}(t) dt - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt\end{aligned}$$

Turns out, finding the extrema subject to the constraint is equivalent to finding the extrema of

$$\begin{aligned}\tilde{S}[q^\mu, z, \lambda] &= S[q^\mu, z] - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt \\ &= \int_a^b \dot{z}(t) dt - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt \\ &= \int_a^b (1 - \lambda(t)) \dot{z}(t) + \lambda(t) L(q^\mu(t), \dot{q}^\mu(t), z(t)) dt.\end{aligned}$$

Turns out, finding the extrema subject to the constraint is equivalent to finding the extrema of

$$\begin{aligned}\tilde{S}[q^\mu, z, \lambda] &= S[q^\mu, z] - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt \\ &= \int_a^b \dot{z}(t) dt - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt \\ &= \int_a^b (1 - \lambda(t)) \dot{z}(t) + \lambda(t) L(q^\mu(t), \dot{q}^\mu(t), z(t)) dt.\end{aligned}$$

Turns out, finding the extrema subject to the constraint is equivalent to finding the extrema of

$$\begin{aligned}\tilde{S}[q^\mu, z, \lambda] &= S[q^\mu, z] - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt \\ &= \int_a^b \dot{z}(t) dt - \int_a^b \lambda(t) [\dot{z}(t) - L(q^\mu(t), \dot{q}^\mu(t), z(t))] dt \\ &= \int_a^b (1 - \lambda(t)) \dot{z}(t) + \lambda(t) L(q^\mu(t), \dot{q}^\mu(t), z(t)) dt.\end{aligned}$$

The Euler-Lagrange equations of this functional are

$$\frac{\partial L}{\partial q^\mu} - \frac{d}{dt} \frac{\partial L}{\partial \dot{q}^\mu} + \frac{\partial L}{\partial \dot{q}^\mu} \frac{\partial L}{\partial z} = 0.$$

These are the Herglotz equations.

The harmonic oscillator with linear action coupling is

The harmonic oscillator with linear action coupling is

$$L(q, \dot{q}, z) = \frac{1}{2}m\dot{q}^2 - \frac{1}{2}m\omega^2 q^2 - \gamma z.$$

The harmonic oscillator with linear action coupling is

$$L(q, \dot{q}, z) = \frac{1}{2}m\dot{q}^2 - \frac{1}{2}m\omega^2 q^2 - \gamma z.$$

The equation of motion is

The harmonic oscillator with linear action coupling is

$$L(q, \dot{q}, z) = \frac{1}{2}m\dot{q}^2 - \frac{1}{2}m\omega^2 q^2 - \gamma z.$$

The equation of motion is

$$\ddot{q} + \gamma\dot{q} + \omega^2 q = 0,$$

which is the equation of a damped harmonic oscillator.

There is a subtlety when thinking of action-dependent Lagrangians for field theory. We must replace z by a 3-form $z = z^\mu dx_\mu$ which we call the action flux.

There is a subtlety when thinking of action-dependent Lagrangians for field theory. We must replace z by a 3-form $z = z^\mu dx_\mu$ which we call the action flux. The action flux does *not* transform like a vector.

There is a subtlety when thinking of action-dependent Lagrangians for field theory. We must replace z by a 3-form $z = z^\mu dx_\mu$ which we call the action flux. The action flux does *not* transform like a vector.

The Herglotz equations for field theory are

$$\frac{\partial L}{\partial \phi^a} - \partial_\mu \frac{\partial L}{\partial (\partial_\mu \phi^a)} + \frac{\partial L}{\partial z^\mu} \frac{\partial L}{\partial (\partial_\mu \phi^a)} = 0.$$

Mechanics

Field theory

	Mechanics	Field theory
Action flux	$z(t)$	$z^\mu(x^\nu) dx_\mu$

	Mechanics	Field theory
Action flux	$z(t)$	$z^\mu(x^\nu) dx_\mu$
Action functional	$S[q^\mu, z] = z(b) - z(a)$	$S[\phi^a, z^\mu] = \int_{\partial D} z$

	Mechanics	Field theory
Action flux	$z(t)$	$z^\mu(x^\nu) dx_\mu$
Action functional	$S[q^\mu, z] = z(b) - z(a)$	$S[\phi^a, z^\mu] = \int_{\partial D} z$
Constraint	$\dot{z} = L(q^\mu, \dot{q}^\mu, z)$	$dz = \mathcal{L}(\phi^a, \partial_\mu \phi^a, z^\mu)$

- ① Introduction
- ② The Herglotz variational problem
- ③ Action-dependent Einstein gravity
- ④ Significance of the equations
- ⑤ Conclusions

Recall the standard Einstein-Hilbert Lagrangian

$$\mathcal{L}_{\text{E-H}} = R\omega_g = R\sqrt{g} \, \mathrm{d}^4x$$

where

$$R = g^{ab} R_{ab} = g^{ab} (\partial_m \Gamma^m_{ab} - \partial_a \Gamma^m_{mb} + \Gamma^m_{mn} \Gamma^n_{ab} - \Gamma^m_{an} \Gamma^n_{mb}).$$

Recall the standard Einstein-Hilbert Lagrangian

$$\mathcal{L}_{\text{E-H}} = R\omega_g = R\sqrt{g} d^4x$$

where

$$R = g^{ab}R_{ab} = g^{ab}(\partial_m\Gamma^m_{ab} - \partial_a\Gamma^m_{mb} + \Gamma^m_{mn}\Gamma^n_{ab} - \Gamma^m_{an}\Gamma^n_{mb}).$$

It's second order in g !

Recall the standard Einstein-Hilbert Lagrangian

$$\mathcal{L}_{\text{E-H}} = R\omega_g = R\sqrt{g} d^4x$$

where

$$R = g^{ab}R_{ab} = g^{ab}(\partial_m\Gamma^m_{ab} - \partial_a\Gamma^m_{mb} + \Gamma^m_{mn}\Gamma^n_{ab} - \Gamma^m_{an}\Gamma^n_{mb}).$$

It's second order in g !

The Euler-Lagrange equations of this Lagrangian are the Einstein field equations:

$$R_{ab} - \frac{1}{2}g_{ab}R = 0$$

10 second order PDEs.

How do we couple the action-flux to the Lagrangian?

How do we couple the action-flux to the Lagrangian? Simplest is linear coupling,

$$\mathcal{L}_{\text{E-H}}(g_{ab}, g_{ab,\mu}, g_{ab,\mu\nu}, z^\mu) = R\omega_g - \theta \wedge z.$$

How do we couple the action-flux to the Lagrangian? Simplest is linear coupling,

$$\mathcal{L}_{\text{E-H}}(g_{ab}, g_{ab,\mu}, g_{ab,\mu\nu}, z^\mu) = R\omega_g - \theta \wedge z.$$

θ must be a covector, $\theta = \theta_\mu dx^\mu$, the *dissipation form*.

How do we couple the action-flux to the Lagrangian? Simplest is linear coupling,

$$\mathcal{L}_{\text{E-H}}(g_{ab}, g_{ab,\mu}, g_{ab,\mu\nu}, z^\mu) = R\omega_g - \theta \wedge z.$$

θ must be a covector, $\theta = \theta_\mu dx^\mu$, the *dissipation form*.

In coordinates

$$\theta \wedge z = \theta_\mu z^\mu d^4x.$$

How do we couple the action-flux to the Lagrangian? Simplest is linear coupling,

$$\mathcal{L}_{\text{E-H}}(g_{ab}, g_{ab,\mu}, g_{ab,\mu\nu}, z^\mu) = R\omega_g - \theta \wedge z.$$

θ must be a covector, $\theta = \theta_\mu dx^\mu$, the *dissipation form*.

In coordinates

$$\theta \wedge z = \theta_\mu z^\mu d^4x.$$

This means the constraint is

$$\partial_\mu z^\mu = R\sqrt{g} - \theta_\mu z^\mu.$$

Let's sketch the derivation of the equations.

Let's sketch the derivation of the equations. Write out the expanded action

$$\tilde{S}[g_{ab}, z^\nu] = \int_D [(1 - \lambda)\partial_\mu z^\mu + \lambda(R\sqrt{g} - \theta_\mu z^\mu)] d^4x.$$

Let's sketch the derivation of the equations. Write out the expanded action

$$\tilde{S}[g_{ab}, z^\nu] = \int_D [(1 - \lambda)\partial_\mu z^\mu + \lambda(R\sqrt{g} - \theta_\mu z^\mu)] d^4x.$$

Let's compute the variation of this:

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\nu] &= \int_D [(1 - \lambda)\delta\partial_\mu z^\mu + \lambda(\delta(R\sqrt{g}) - \theta_\mu\delta z^\mu)] d^4x \\ &= \int_D (1 - \lambda)\partial_\mu\delta z^\mu - \lambda\theta_\mu\delta z^\mu d^4x + \int_D \lambda\delta(R\sqrt{g}) d^4x.\end{aligned}$$

Let's sketch the derivation of the equations. Write out the expanded action

$$\tilde{S}[g_{ab}, z^\nu] = \int_D [(1 - \lambda)\partial_\mu z^\mu + \lambda(R\sqrt{g} - \theta_\mu z^\mu)] d^4x.$$

Let's compute the variation of this:

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\nu] &= \int_D [(1 - \lambda)\delta\partial_\mu z^\mu + \lambda(\delta(R\sqrt{g}) - \theta_\mu\delta z^\mu)] d^4x \\ &= \int_D (1 - \lambda)\partial_\mu\delta z^\mu - \lambda\theta_\mu\delta z^\mu d^4x + \int_D \lambda\delta(R\sqrt{g}) d^4x.\end{aligned}$$

Let's sketch the derivation of the equations. Write out the expanded action

$$\tilde{S}[g_{ab}, z^\nu] = \int_D [(1 - \lambda)\partial_\mu z^\mu + \lambda(R\sqrt{g} - \theta_\mu z^\mu)] d^4x.$$

Let's compute the variation of this:

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\nu] &= \int_D [(1 - \lambda)\delta\partial_\mu z^\mu + \lambda(\delta(R\sqrt{g}) - \theta_\mu\delta z^\mu)] d^4x \\ &= \int_D (1 - \lambda)\partial_\mu\delta z^\mu - \lambda\theta_\mu\delta z^\mu d^4x + \int_D \lambda\delta(R\sqrt{g}) d^4x.\end{aligned}$$

Let's sketch the derivation of the equations. Write out the expanded action

$$\tilde{S}[g_{ab}, z^\nu] = \int_D [(1 - \lambda)\partial_\mu z^\mu + \lambda(R\sqrt{g} - \theta_\mu z^\mu)] d^4x.$$

Let's compute the variation of this:

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\nu] &= \int_D [(1 - \lambda)\delta\partial_\mu z^\mu + \lambda(\delta(R\sqrt{g}) - \theta_\mu\delta z^\mu)] d^4x \\ &= \int_D (1 - \lambda)\partial_\mu\delta z^\mu - \lambda\theta_\mu\delta z^\mu d^4x + \int_D \lambda\delta(R\sqrt{g}) d^4x.\end{aligned}$$

Let's sketch the derivation of the equations. Write out the expanded action

$$\tilde{S}[g_{ab}, z^\nu] = \int_D [(1 - \lambda)\partial_\mu z^\mu + \lambda(R\sqrt{g} - \theta_\mu z^\mu)] d^4x.$$

Let's compute the variation of this:

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\nu] &= \int_D [(1 - \lambda)\delta\partial_\mu z^\mu + \lambda(\delta(R\sqrt{g}) - \theta_\mu\delta z^\mu)] d^4x \\ &= \int_D (1 - \lambda)\partial_\mu\delta z^\mu - \lambda\theta_\mu\delta z^\mu d^4x + \int_D \lambda\delta(R\sqrt{g}) d^4x.\end{aligned}$$

The variations of the action flux and the metric decouple.

Integrating by parts in the variation of the action flux,

Integrating by parts in the variation of the action flux,

$$\int_D (1 - \lambda) \partial_\mu \delta z^\mu - \lambda \theta_\mu \delta z^\mu d^4x$$

Integrating by parts in the variation of the action flux,

$$\begin{aligned} & \int_D (1 - \lambda) \partial_\mu \delta z^\mu - \lambda \theta_\mu \delta z^\mu \, d^4x \\ &= \int_D \partial_\mu ((1 - \lambda) \delta z^\mu) \, d^4x + \int_D (\partial_\mu \lambda - \lambda \theta_\mu) \delta z^\mu \, d^4x \end{aligned}$$

Integrating by parts in the variation of the action flux,

$$\begin{aligned} & \int_D (1 - \lambda) \partial_\mu \delta z^\mu - \lambda \theta_\mu \delta z^\mu \, d^4x \\ &= \int_D \cancel{\partial_\mu ((1 - \lambda) \delta z^\mu)} \, d^4x + \int_D (\partial_\mu \lambda - \lambda \theta_\mu) \delta z^\mu \, d^4x \end{aligned}$$

Integrating by parts in the variation of the action flux,

$$\begin{aligned} & \int_D (1 - \lambda) \partial_\mu \delta z^\mu - \lambda \theta_\mu \delta z^\mu \, d^4x \\ &= \int_D \cancel{\partial_\mu ((1 - \lambda) \delta z^\mu)} \, d^4x + \int_D (\partial_\mu \lambda - \lambda \theta_\mu) \delta z^\mu \, d^4x \end{aligned}$$

This must vanish for any variation of z^μ , so it must be

$$\partial_\mu \lambda = \lambda \theta_\mu.$$

The variation of the second integral term gives three terms

The variation of the second integral term gives three terms

$$\delta(R\sqrt{g}) = \delta(g^{ab}R_{ab}\sqrt{g}) = \delta g^{ab}R_{ab}\sqrt{g} + g_{ab}\delta R^{ab}\sqrt{g} + g^{ab}R_{ab}\delta\sqrt{g}$$

The variation of the second integral term gives three terms

$$\delta(R\sqrt{g}) = \delta(g^{ab}R_{ab}\sqrt{g}) = \delta g^{ab}R_{ab}\sqrt{g} + g_{ab}\delta R^{ab}\sqrt{g} + g^{ab}R_{ab}\delta\sqrt{g}$$

So

$$\int_D \lambda \delta(R\sqrt{g}) d^4x =$$

The variation of the second integral term gives three terms

$$\delta(R\sqrt{g}) = \delta(g^{ab}R_{ab}\sqrt{g}) = \delta g^{ab}R_{ab}\sqrt{g} + g_{ab}\delta R^{ab}\sqrt{g} + g^{ab}R_{ab}\delta\sqrt{g}$$

So

$$\begin{aligned} \int_D \lambda \delta(R\sqrt{g}) d^4x &= \\ &= \int_D \lambda \delta g^{ab} R_{ab} \sqrt{g} d^4x + \int_D \lambda g^{ab} \delta R_{ab} \sqrt{g} d^4x + \int_D \lambda R \delta \sqrt{g} d^4x \end{aligned}$$

The variation of the second integral term gives three terms

$$\delta(R\sqrt{g}) = \delta(g^{ab}R_{ab}\sqrt{g}) = \delta g^{ab}R_{ab}\sqrt{g} + g_{ab}\delta R^{ab}\sqrt{g} + g^{ab}R_{ab}\delta\sqrt{g}$$

So

$$\begin{aligned} \int_D \lambda \delta(R\sqrt{g}) d^4x &= \\ &= \int_D \lambda \delta g^{ab} R_{ab} \sqrt{g} d^4x + \int_D \lambda g^{ab} \delta R_{ab} \sqrt{g} d^4x + \int_D \lambda R \delta \sqrt{g} d^4x \\ &= \int_D \lambda R_{ab} \delta g^{ab} \sqrt{g} d^4x + \int_D \lambda g^{ab} \delta R_{ab} \sqrt{g} d^4x - \int_D \frac{1}{2} \lambda g_{ab} R \delta g^{ab} \sqrt{g} d^4x \end{aligned}$$

$$\begin{aligned} \int_D \lambda \delta(R\sqrt{g}) d^4x &= \\ &= \int_D \lambda R_{ab} \delta g^{ab} \sqrt{g} d^4x + \int_D \lambda g^{ab} \delta R_{ab} \sqrt{g} d^4x - \int_D \frac{1}{2} \lambda g_{ab} R \delta g^{ab} \sqrt{g} d^4x \end{aligned}$$

$$\begin{aligned} \int_D \lambda \delta(R\sqrt{g}) d^4x &= \\ &= \int_D \lambda R_{ab} \delta g^{ab} \sqrt{g} d^4x + \int_D \lambda g^{ab} \delta R_{ab} \sqrt{g} d^4x - \int_D \frac{1}{2} \lambda g_{ab} R \delta g^{ab} \sqrt{g} d^4x \end{aligned}$$

If λ weren't there, the second term would vanish and we would get the Einstein field equations. To deal with λ we have to integrate by parts twice.

We use

$$g^{ab}\delta R_{ab} = \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})$$

We use

$$g^{ab}\delta R_{ab} = \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})$$

so integrating by parts

$$\int_D \lambda g^{ab}\delta R_{ab}\sqrt{g}\,d^4x =$$

We use

$$g^{ab}\delta R_{ab} = \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})$$

so integrating by parts

$$\begin{aligned}\int_D \lambda g^{ab}\delta R_{ab}\sqrt{g}\,d^4x &= \\ &= \int_D \lambda \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})\sqrt{g}\,d^4x\end{aligned}$$

We use

$$g^{ab}\delta R_{ab} = \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})$$

so integrating by parts

$$\begin{aligned}\int_D \lambda g^{ab}\delta R_{ab}\sqrt{g}\,d^4x &= \\ &= \int_D \lambda \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})\sqrt{g}\,d^4x \\ &= \int_D \nabla_n \left(\lambda(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \right) \sqrt{g}\,d^4x\end{aligned}$$

We use

$$g^{ab}\delta R_{ab} = \nabla_n (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})$$

so integrating by parts

$$\begin{aligned} \int_D \lambda g^{ab} \delta R_{ab} \sqrt{g} \, d^4x &= \\ &= \int_D \lambda \nabla_n (g^{ab} \delta\Gamma^n_{ab} - g^{nb} \delta\Gamma^m_{mb}) \sqrt{g} \, d^4x \\ &= \int_D \nabla_n \left(\lambda (g^{ab} \delta\Gamma^n_{ab} - g^{nb} \delta\Gamma^m_{mb}) \right) \sqrt{g} \, d^4x \\ &\quad - \int_D (\nabla_n \lambda) (g^{ab} \delta\Gamma^n_{ab} - g^{nb} \delta\Gamma^m_{mb}) \sqrt{g} \, d^4x \end{aligned}$$

We use

$$g^{ab}\delta R_{ab} = \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})$$

so integrating by parts

$$\begin{aligned} \int_D \lambda g^{ab}\delta R_{ab} \sqrt{g} d^4x &= \\ &= \int_D \lambda \nabla_n (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \sqrt{g} d^4x \\ &= \int_D \nabla_n \left(\lambda (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \right) \sqrt{g} d^4x \\ &\quad - \int_D (\nabla_n \lambda) (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \sqrt{g} d^4x \\ &= - \int_D \lambda \theta_n (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \sqrt{g} d^4x \end{aligned}$$

We use

$$g^{ab}\delta R_{ab} = \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})$$

so integrating by parts

$$\begin{aligned} \int_D \lambda g^{ab}\delta R_{ab}\sqrt{g}\,d^4x &= \\ &= \int_D \lambda \nabla_n(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})\sqrt{g}\,d^4x \\ &= \int_D \nabla_n \left(\lambda (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \right) \sqrt{g}\,d^4x \\ &\quad - \int_D (\nabla_n \lambda)(g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})\sqrt{g}\,d^4x \\ &= - \int_D \lambda \theta_n (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})\sqrt{g}\,d^4x \end{aligned}$$

We use

$$g^{ab}\delta R_{ab} = \nabla_n (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb})$$

so integrating by parts

$$\begin{aligned} \int_D \lambda g^{ab}\delta R_{ab} \sqrt{g} d^4x &= \\ &= \int_D \lambda \nabla_n (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \sqrt{g} d^4x \\ &= \int_D \nabla_n \left(\lambda (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \right) \sqrt{g} d^4x \\ &\quad - \int_D (\nabla_n \lambda) (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \sqrt{g} d^4x \\ &= - \int_D \lambda \theta_n (g^{ab}\delta\Gamma^n_{ab} - g^{nb}\delta\Gamma^m_{mb}) \sqrt{g} d^4x \end{aligned}$$

We use

$$\delta\Gamma^a_{bc} = \frac{1}{2}g^{am}(\nabla_c\delta g_{bm} + \nabla_b\delta g_{mc} - \nabla_m\delta g_{bc}).$$

We use

$$\delta\Gamma^a_{bc} = \frac{1}{2}g^{am}(\nabla_c\delta g_{bm} + \nabla_b\delta g_{mc} - \nabla_m\delta g_{bc}).$$

so $g^{ab}\delta R_{ab}$ splits into 6 terms.

We use

$$\delta\Gamma^a_{bc} = \frac{1}{2}g^{am}(\nabla_c\delta g_{bm} + \nabla_b\delta g_{mc} - \nabla_m\delta g_{bc}).$$

so $g^{ab}\delta R_{ab}$ splits into 6 terms. We will go through one in detail

$$- \frac{1}{2} \int_D \lambda \theta_n g^{ab} g^{nk} \nabla_b \delta g_{ak} \sqrt{g} d^4x$$

We use

$$\delta\Gamma^a_{bc} = \frac{1}{2}g^{am}(\nabla_c\delta g_{bm} + \nabla_b\delta g_{mc} - \nabla_m\delta g_{bc}).$$

so $g^{ab}\delta R_{ab}$ splits into 6 terms. We will go through one in detail

$$\begin{aligned} & -\frac{1}{2}\int_D \lambda\theta_n g^{ab}g^{nk}\nabla_b\delta g_{ak}\sqrt{g}\,d^4x \\ & = -\frac{1}{2}\int_D \lambda\theta_n\nabla_b(g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x \end{aligned}$$

We use

$$\delta\Gamma^a_{bc} = \frac{1}{2}g^{am}(\nabla_c\delta g_{bm} + \nabla_b\delta g_{mc} - \nabla_m\delta g_{bc}).$$

so $g^{ab}\delta R_{ab}$ splits into 6 terms. We will go through one in detail

$$\begin{aligned} & -\frac{1}{2}\int_D \lambda\theta_n g^{ab}g^{nk}\nabla_b\delta g_{ak}\sqrt{g}\,d^4x \\ &= -\frac{1}{2}\int_D \lambda\theta_n\nabla_b(g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x \\ &= -\frac{1}{2}\int_D \nabla_b(\lambda\theta_n g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x + \frac{1}{2}\int_D \nabla_b(\lambda\theta_n)(g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x \end{aligned}$$

We use

$$\delta\Gamma^a_{bc} = \frac{1}{2}g^{am}(\nabla_c\delta g_{bm} + \nabla_b\delta g_{mc} - \nabla_m\delta g_{bc}).$$

so $g^{ab}\delta R_{ab}$ splits into 6 terms. We will go through one in detail

$$\begin{aligned} & -\frac{1}{2}\int_D \lambda\theta_n g^{ab}g^{nk}\nabla_b\delta g_{ak}\sqrt{g}\,d^4x \\ &= -\frac{1}{2}\int_D \lambda\theta_n\nabla_b(g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x \\ &= -\frac{1}{2}\int_D \nabla_b(\lambda\theta_n g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x + \frac{1}{2}\int_D \nabla_b(\lambda\theta_n)(g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x \\ &= -\int_D \lambda(\theta_b\theta_n + \nabla_b\theta_n)\delta g^{bn}\sqrt{g}\,d^4x. \end{aligned}$$

We use

$$\delta\Gamma^a_{bc} = \frac{1}{2}g^{am}(\nabla_c\delta g_{bm} + \nabla_b\delta g_{mc} - \nabla_m\delta g_{bc}).$$

so $g^{ab}\delta R_{ab}$ splits into 6 terms. We will go through one in detail

$$\begin{aligned} & -\frac{1}{2}\int_D \lambda\theta_n g^{ab}g^{nk}\nabla_b\delta g_{ak}\sqrt{g}\,d^4x \\ &= -\frac{1}{2}\int_D \lambda\theta_n\nabla_b(g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x \\ &= -\frac{1}{2}\int_D \nabla_b(\lambda\theta_n g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x + \frac{1}{2}\int_D \nabla_b(\lambda\theta_n)(g^{ab}g^{nk}\delta g_{ak})\sqrt{g}\,d^4x \\ &= -\int_D \lambda(\theta_b\theta_n + \nabla_b\theta_n)\delta g^{bn}\sqrt{g}\,d^4x. \end{aligned}$$

After some calculation, one finds

After some calculation, one finds

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\mu] = & \int_D (\partial_\mu \lambda - \lambda \theta_\mu) \delta z^\mu d^4x \\ & + \int_D \lambda (R_{ab} - \tfrac{1}{2} R g_{ab} - K_{ab} + K g_{ab}) \delta g^{ab} \sqrt{g} d^4x\end{aligned}$$

with $K_{ab} = \theta_a \theta_b + \nabla_{(a} \theta_{b)}$.

After some calculation, one finds

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\mu] = & \int_D (\partial_\mu \lambda - \lambda \theta_\mu) \delta z^\mu d^4x \\ & + \int_D \lambda (R_{ab} - \tfrac{1}{2} R g_{ab} - K_{ab} + K g_{ab}) \delta g^{ab} \sqrt{g} d^4x\end{aligned}$$

with $K_{ab} = \theta_a \theta_b + \nabla_{(a} \theta_{b)}$.

The stationary configurations will be those for which this vanishes for any variation, so we arrive at

After some calculation, one finds

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\mu] = & \int_D (\partial_\mu \lambda - \lambda \theta_\mu) \delta z^\mu d^4x \\ & + \int_D \lambda (R_{ab} - \tfrac{1}{2} R g_{ab} - K_{ab} + K g_{ab}) \delta g^{ab} \sqrt{g} d^4x\end{aligned}$$

with $K_{ab} = \theta_a \theta_b + \nabla_{(a} \theta_{b)}$.

The stationary configurations will be those for which this vanishes for any variation, so we arrive at

$$R_{ab} - \tfrac{1}{2} R g_{ab} - K_{ab} + K g_{ab} = 0.$$

After some calculation, one finds

$$\begin{aligned}\delta\tilde{S}[g_{ab}, z^\mu] = & \int_D (\partial_\mu \lambda - \lambda \theta_\mu) \delta z^\mu d^4x \\ & + \int_D \lambda (R_{ab} - \tfrac{1}{2} R g_{ab} - K_{ab} + K g_{ab}) \delta g^{ab} \sqrt{g} d^4x\end{aligned}$$

with $K_{ab} = \theta_a \theta_b + \nabla_{(a} \theta_{b)}$.

The stationary configurations will be those for which this vanishes for any variation, so we arrive at

$$R_{ab} - \tfrac{1}{2} R g_{ab} - K_{ab} + K g_{ab} = 0.$$

- ① Introduction
- ② The Herglotz variational problem
- ③ Action-dependent Einstein gravity
- ④ Significance of the equations
- ⑤ Conclusions

We call the object K_{ab} the *dissipation tensor*.

We call the object K_{ab} the *dissipation tensor*. Coordinate free we can write

$$K(X, Y) = (\theta \otimes \theta)(X, Y) + \frac{1}{2} [(\nabla_X \theta)Y + (\nabla_Y \theta)X]$$

which is a $(0, 2)$ tensor.

We call the object K_{ab} the *dissipation tensor*. Coordinate free we can write

$$K(X, Y) = (\theta \otimes \theta)(X, Y) + \frac{1}{2} [(\nabla_X \theta)Y + (\nabla_Y \theta)X]$$

which is a $(0, 2)$ tensor. This means the equations we have derived are covariant.

In [Laz+17], the derived equations are

$$R_{ab} + \tilde{K}_{ab} - \frac{1}{2}g_{ab}(R + \tilde{K}) = 0$$

where

$$\tilde{K}_{ab} = \theta_m \Gamma^m_{ab} - \frac{1}{2} (\theta_a \Gamma^m_{mb} + \theta_b \Gamma^m_{am}).$$

In [Laz+17], the derived equations are

$$R_{ab} + \tilde{K}_{ab} - \frac{1}{2}g_{ab}(R + \tilde{K}) = 0$$

where

$$\tilde{K}_{ab} = \theta_m \Gamma^m_{ab} - \frac{1}{2} (\theta_a \Gamma^m_{mb} + \theta_b \Gamma^m_{am}).$$

\tilde{K}_{ab} is not a tensor so they are *not* covariant!

In [Laz+17], the derived equations are

$$R_{ab} + \tilde{K}_{ab} - \frac{1}{2}g_{ab}(R + \tilde{K}) = 0$$

where

$$\tilde{K}_{ab} = \theta_m \Gamma^m_{ab} - \frac{1}{2} (\theta_a \Gamma^m_{mb} + \theta_b \Gamma^m_{am}).$$

\tilde{K}_{ab} is not a tensor so they are *not* covariant!

From mechanics, linear action couplings in second order theories lead to terms *quadratic* in the dissipation. K_{ab} is quadratic in θ_μ , but not \tilde{K}_{ab} .

- ① Introduction
- ② The Herglotz variational problem
- ③ Action-dependent Einstein gravity
- ④ Significance of the equations
- ⑤ Conclusions**

As a recap

As a recap

- The Herglotz problem can be turned into an unconstrained optimisation problem.

As a recap

- The Herglotz problem can be turned into an unconstrained optimisation problem.
- Because we can write down an action, we can take its variation and deal with second order theories.

As a recap

- The Herglotz problem can be turned into an unconstrained optimisation problem.
- Because we can write down an action, we can take its variation and deal with second order theories.
- We have derived the field equations for the Einstein-Hilbert Lagrangian with a linear action flux coupling.

As a recap

- The Herglotz problem can be turned into an unconstrained optimisation problem.
- Because we can write down an action, we can take its variation and deal with second order theories.
- We have derived the field equations for the Einstein-Hilbert Lagrangian with a linear action flux coupling.
- The result are a set of equations which are *covariant* and with the expected structure.

As a recap

- The Herglotz problem can be turned into an unconstrained optimisation problem.
- Because we can write down an action, we can take its variation and deal with second order theories.
- We have derived the field equations for the Einstein-Hilbert Lagrangian with a linear action flux coupling.
- The result are a set of equations which are *covariant* and with the expected structure.

- What do the solutions look like?

- What do the solutions look like?
- What happens when we add matter terms?

- What do the solutions look like?
- What happens when we add matter terms?
- What kind of cosmology do we get?

- What do the solutions look like?
- What happens when we add matter terms?
- What kind of cosmology do we get?
- What happens with different action flux couplings?

- What do the solutions look like?
- What happens when we add matter terms?
- What kind of cosmology do we get?
- What happens with different action flux couplings?

A variational derivation of the field equations of an action-dependent Einstein-Hilbert Lagrangian

Arnau Mas

Supervised by
Dr Jordi Gaset

Universitat Autònoma de Barcelona
June 27th 2021

