

$$r^3 + 3r^2 + 2r = 0$$

$$r(r^2 + 3r + 2) = 0$$

$$r = 0, -1, -2$$

$$y(t) = A + Be^{-t} + Ce^{-2t}$$

$$y'(t) = -Be^{-t} - 2Ce^{-2t}$$

$$y''(t) = Be^{-t} + 4Ce^{-2t}$$

$$y(0) = A + B + C = 2$$

$$y'(0) = -B - 2C = -1$$

$$y''(0) = B + 4C = 1$$

$$1 - 2C + 4C = 1 + 2C = 1$$

$$2C = 0$$

$$C = 0$$

$$-B = -1 \rightarrow B = 1$$

$$A = 1$$

$$y(t) = 1 + e^{-t}$$

$$y(1) = 1 + e^{-1}$$

Let $y(t)$ denote the unique solution to the initial value problem

$$y''' + 3y'' + 2y' = 0, \quad y(0) = 2, \\ y'(0) = -1, \quad y''(0) = 1.$$

What is the value of $y(1)$?

☐ (A) $1 + 2e^{-1} + e^{-2}$

☐ (B) $1 - e^{-2}$

☒ (C) $1 + e^{-1}$

☐ (D) $e + e^2$

☐ (E) $2 + e^{-1} + 2e^{-2}$

$$F(t) = m x'' + c x' + k x$$

$$0 = 4x'' + 4x' + 17x$$

$$4r^2 + 4r + 17 = 0$$

$$r^2 + r + \frac{17}{4} = 0$$

$$r^2 + r + \frac{1}{4} + 4 = 0$$

$$\left(r + \frac{1}{2}\right)^2 + 4 = 0$$

$$\left(r + \frac{1}{2}\right)^2 = -4$$

$$r + \frac{1}{2} = \pm 2j$$

$$r = \pm 2j - \frac{1}{2}$$

$$x(t) = e^{-\frac{1}{2}t} (A \cos 2t + B \sin 2t)$$

$$x'(t) = -\frac{1}{2}e^{-\frac{1}{2}t} (A \cos 2t + B \sin 2t) + e^{-\frac{1}{2}t} (-2A \sin 2t + 2B \cos 2t)$$

$$x(0) = A \cos 0 + B \sin 0 = A = 0$$

$$x'(0) = -\frac{1}{2}(0) + 2B = 2 \rightarrow B = 1$$

$$x_p(2) = \frac{\sin 4}{e}$$

A mass-spring-dashpot system with mass $m=4$, damping constant $c=4$, and spring constant $k=17$ is set in free motion with initial conditions $x(0)=0$ and $x'(0)=2$, where $x(t)$ is the displacement from the equilibrium position at time t . Find $x(2)$.

☒ (A) $e^{-1} \sin(4)$

☐ (B) $e^{-1} \cos(4)$

☐ (C) $e^{-1} \cos(4) + e^{-1} \sin(4)$

☐ (D) $e^{-2} \cos(1) + e^{-2} \sin(1)$

☐ (E) $e^{-2} \sin(1)$

$$y(t) = Ae^{4t} + Bte^{4t} + C$$

$$r = 4 \text{ m2, } 0$$

$$r(r-4)^2$$

$$= r(r^2 - 8r + 16)$$

$$= r^3 - 8r^2 + 16r$$

$$\hookrightarrow y''' - 8y'' + 16y' + C = 0$$

Which of the following differential equations has

$$y(t) = C_1 e^{4t} + C_2 t e^{4t} + C_3$$

as a general solution?

(A) $y'' - 8y + 16y = 0$

(B) $y^{(3)} - 8y'' + 16y' + y = 0$

(C) $y^{(3)} - 8y'' + 16y' = 0$

(D) $y^{(3)} - 8y'' + 24y' - 16y = 0$

(E) $y^{(3)} - 16y' = 0$

$$r^2 - 10r + 26 = 0$$

$$r^2 - 10r + 25 + 1 = 0$$

$$(r-5)^2 = -1$$

$$r-5 = \pm j$$

$$r = 5 \pm j$$

$$y(t) = e^{5t}(A \cos t + B \sin t)$$

$$y'(t) = 5e^{5t}(A \cos t + B \sin t) + e^{5t}(-A \sin t + B \cos t)$$

$$y(0) = A = 1$$

$$y'(0) = 5 + B = 4 \rightarrow B = -1$$

$$y(t) = e^{5t}(\cos t - \sin t)$$

Solve the initial value problem

$$y'' - 10y' + 26y = 0, y(0) = 1, \\ y'(0) = 4$$

(A) $y(t) = e^t(\cos 5t - \frac{2}{5} \sin 5t)$

(B) $y(t) = e^{5t}(\cos t + 4 \sin t)$

(C) $y(t) = e^t(\cos 5t + \frac{3}{5} \sin 5t)$

(D) $y(t) = e^t(\cos t + 2 \sin t)$

(E) $y(t) = e^{5t}(\cos t - \sin t)$

$$r^3 + 6r^2 + 9r = 0$$

$$r(r+3)^2 = 0$$

$$r = -3 \text{ m2, } 0$$

$$y(t) = A + Be^{-3t} + Cte^{-3t}$$

Find the general solution of $y''' + 6y'' + 9y' = 0$

☒ A $y(t) = c_1 + c_2e^{-3t} + c_3te^{-3t}$

☐ B $y(t) + c_1t + c_2e^{3t} + c_3te^{3t}$

☐ C $y(t) = c_1 + c_2 \cos(-3t) + c_3 \sin(-3t)$

☐ D $y(t) = c_1t + c_2e^{-3t} + c_3e^{3t}$

☐ E $y(t) = c_1t + c_2 \cos(3t) + c_3 \sin(3t)$

$$r^2 + r - 12 = 0$$

$$r = 3, -4$$

$$y(t) = Ae^{3t} + Be^{-4t}$$

$$y'(t) = 3Ae^{3t} - 4Be^{-4t}$$

$$y(0) = 3 = A + B$$

$$y'(0) = -4 = 3A - 4B$$

$$3(3 - B) - 4B = -4$$

$$13 = 7B$$

$$B = 13/7$$

$$A = 3 - \frac{B}{7} = \frac{21}{7} - \frac{13}{7} = \frac{8}{7}$$

$$y(t) = \frac{8}{7}e^{3t} + \frac{13}{7}e^{-4t}$$

Solve the initial value problem

$$y'' + y' - 12y = 0$$

$$y(0) = 3, y'(0) = -4$$

(A) $y(t) = 3e^{-4t} - 4e^{3t}$

(B) $y(t) = \frac{16}{7}e^{-3t} + \frac{5}{7}e^{4t}$

(C) $y(t) = \frac{13}{7}e^{-4t} + \frac{8}{7}e^{3t}$

(D) $y(t) = -\frac{5}{7}e^{-4t} - \frac{16}{7}e^{3t}$

(E) $y(t) = \frac{11}{7}e^{-3t} + \frac{10}{7}e^{4t}$

$$\frac{dy}{dx} = \frac{y}{x} + e^{y/x}$$

$$v + x \frac{dv}{dx} = v + e^v$$

$$x \frac{dv}{dx} = e^v$$

$$\int e^{-v} dv = \int x^{-1} dx$$

$$-e^{-v} = \ln x + C$$

$$-e^{-y/x} = \ln x + C$$

$$e^{-y/x} = -C - \ln x = C - \ln x$$

$$-\frac{y}{x} = \ln(C - \ln x)$$

$$y = -x \ln(C - \ln x)$$

$$v = y/x$$

$$y = xv$$

$$dy/dx = v + x dv/dx$$

Find the general solution to the homogenous differential equation using the substitution $v = y/x$. Assume $x > 0$.

$$x^2 y' = xy + x^2 e^{y/x}$$

☒ A $y = -x \ln(C - \ln x)$

☐ B $y = x(\ln x + C)$

☐ C $y = -\ln(C - \ln x)$

☐ D $y = x \ln\left(\frac{x^2}{2} + C\right)$

☐ E $y = -x \ln(C - x)$

let $C_1, C_2, C_3 = A, B, C$

$$y(x) = \underline{A}e^{2x} + \underline{e}^{0x} \left[B \cos \underline{2}x + C \sin \underline{2}x \right]$$

$0 \pm j2$

$$r = 2, \pm j2$$

$$(r-2)(r^2+4)=0$$

$$r^3 + 4r - 2r^2 - 8 = 0$$

$$y''' - 2y'' + 4y' - 8y = 0$$

Find a linear homogenous constant-coefficient differential equation with the general solution $y(x) = C_1 e^{2x} + C_2 \cos(2x) + C_3 \sin(2x)$.