

Complex Analysis and Differential Equations II

Course Code: MA 203
Programme: B.Tech.



Department of Mathematics
Indian Institute of Technology Indore

Contents

- 1 Syllabus
- 2 Evaluation
- 3 Complex number: Origin
- 4 Real vs complex
- 5 Applications of complex analysis
- 6 Complex number: Review
- 7 Points and sets
- 8 Complex functions
- 9 Analytic functions
- 10 Harmonic functions
- 11 Elementary functions
- 12 Complex Integration
- 13 Cauchy Integral Theorem
- 14 Cauchy-Goursat's theorem for multiply connected domain

Syllabus-Tentative Distribution of contents

Syllabus-Tentative Distribution of contents

Before mid-semester

Syllabus-Tentative Distribution of contents

Before mid-semester

1 Dr. Bapan Ghosh:

Syllabus-Tentative Distribution of contents

Before mid-semester

- 1 Dr. Bapan Ghosh: First part-complex Analysis

Syllabus-Tentative Distribution of contents

Before mid-semester

1 Dr. Bapan Ghosh: First part-complex Analysis

Definitions and properties of analytic functions. Cauchy -Riemann equations, harmonic functions. Power series and their properties, Elementary functions, Cauchy's theorem and its applications.

Syllabus-Tentative Distribution of contents

Before mid-semester

1 Dr. Bapan Ghosh: First part-complex Analysis

Definitions and properties of analytic functions. Cauchy -Riemann equations, harmonic functions. Power series and their properties, Elementary functions, Cauchy's theorem and its applications.

Syllabus-Tentative Distribution of contents

Before mid-semester

1 Dr. Bapan Ghosh: First part-complex Analysis

Definitions and properties of analytic functions. Cauchy-Riemann equations, harmonic functions. Power series and their properties, Elementary functions, Cauchy's theorem and its applications.

2 Dr. Santanu Manna:

Syllabus-Tentative Distribution of contents

Before mid-semester

1 Dr. Bapan Ghosh: First part-complex Analysis

Definitions and properties of analytic functions. Cauchy -Riemann equations, harmonic functions. Power series and their properties, Elementary functions, Cauchy's theorem and its applications.

2 Dr. Santanu Manna: Ordinary Differential Equations part

Syllabus-Tentative Distribution of contents

Before mid-semester

1 Dr. Bapan Ghosh: First part-complex Analysis

Definitions and properties of analytic functions. Cauchy-Riemann equations, harmonic functions. Power series and their properties, Elementary functions, Cauchy's theorem and its applications.

2 Dr. Santanu Manna: Ordinary Differential Equations part

Review of power series and series solutions of ODE's. Legendre equation and Legendre Polynomials. Regular and singular points, method of Frobenius. Bessel's equation and Bessel's functions. Strum-Liouville problems.

Syllabus-Tentative Distribution of contents

Syllabus-Tentative Distribution of contents

After mid-semester

Syllabus-Tentative Distribution of contents

After mid-semester

- 1 Dr. Debopriya Mukherjee:

Syllabus-Tentative Distribution of contents

After mid-semester

- 1 Dr. Debopriya Mukherjee: Partial Differential Equations part

Syllabus-Tentative Distribution of contents

After mid-semester

- 1 Dr. Debopriya Mukherjee: **Partial Differential Equations part**

Fourier series. D'Alembert solution to the wave equations. Classification of linear second order PDE's in two variables. Laplace, wave, and Heat equations using separation of variables. Vibration of a circular membrane. Heat equation in the half space.

Syllabus-Tentative Distribution of contents

After mid-semester

- 1 Dr. Debopriya Mukherjee: **Partial Differential Equations part**

Fourier series. D'Alembert solution to the wave equations. Classification of linear second order PDE's in two variables. Laplace, wave, and Heat equations using separation of variables. Vibration of a circular membrane. Heat equation in the half space.

- 2 Dr. Vinay Kumar:

Syllabus-Tentative Distribution of contents

After mid-semester

- 1 Dr. Debopriya Mukherjee: Partial Differential Equations part

Fourier series. D'Alembert solution to the wave equations. Classification of linear second order PDE's in two variables. Laplace, wave, and Heat equations using separation of variables. Vibration of a circular membrane. Heat equation in the half space.

- 2 Dr. Vinay Kumar: Second part-complex Analysis

Syllabus-Tentative Distribution of contents

After mid-semester

- 1 Dr. Debopriya Mukherjee: **Partial Differential Equations part**

Fourier series. D'Alembert solution to the wave equations. Classification of linear second order PDE's in two variables. Laplace, wave, and Heat equations using separation of variables. Vibration of a circular membrane. Heat equation in the half space.

- 2 Dr. Vinay Kumar: **Second part-complex Analysis**

Taylor series and Laurent expansion, Residues and Cauchy's residue formula, Evaluation of improper integrals, Conformal mappings, inversion of Laplace transformations

Tentative Evaluation Plan (**out of 100 Marks**)

- 1 Internal evaluation-Before mid-semester: 15 Marks

Tentative Evaluation Plan (**out of 100 Marks**)

1 Internal evaluation-Before mid-semester: 15 Marks

It can be any combination of Assignment, Class test, Viva, etc. based on the declaration by the instructors.

Tentative Evaluation Plan (**out of 100 Marks**)

1 Internal evaluation-Before mid-semester: 15 Marks

It can be any combination of Assignment, Class test, Viva, etc. based on the declaration by the instructors.

2 Mid-semester: 30 Marks

Tentative Evaluation Plan (out of 100 Marks)

1 Internal evaluation-Before mid-semester: 15 Marks

It can be any combination of Assignment, Class test, Viva, etc. based on the declaration by the instructors.

2 Mid-semester: 30 Marks

3 End-semester: 55 Marks

Complex Analysis

LECTURE 1

July 26, 2023

BOOKS

- 1 E. Kreyszig, *Advanced Engineering Mathematics*, John Wiley & Sons.

BOOKS

- 1 E. Kreyszig, **Advanced Engineering Mathematics**, John Wiley & Sons.
- 2 R.V. Churchill and J.W. Brown, **Complex Variables and Applications**, McGraw-Hill Inc.

BOOKS

- 1 E. Kreyszig, **Advanced Engineering Mathematics**, John Wiley & Sons.
- 2 R.V. Churchill and J.W. Brown, **Complex Variables and Applications**, McGraw-Hill Inc.
- 3 M.J. Ablowitz and A.S. Fokas, **Complex Variables: Introduction and Applications**, Cambridge University Press.

BOOKS

- 1 E. Kreyszig, **Advanced Engineering Mathematics**, John Wiley & Sons.
- 2 R.V. Churchill and J.W. Brown, **Complex Variables and Applications**, McGraw-Hill Inc.
- 3 M.J. Ablowitz and A.S. Fokas, **Complex Variables: Introduction and Applications**, Cambridge University Press.
- 4 J.M. Howie, **Complex Analysis**, Springer-Verlag.

BOOKS

- 1 E. Kreyszig, **Advanced Engineering Mathematics**, John Wiley & Sons.
- 2 R.V. Churchill and J.W. Brown, **Complex Variables and Applications**, McGraw-Hill Inc.
- 3 M.J. Ablowitz and A.S. Fokas, **Complex Variables: Introduction and Applications**, Cambridge University Press.
- 4 J.M. Howie, **Complex Analysis**, Springer-Verlag.
- 5 S. Ponnusamy, H. Silverman **Complex Variable with Applications**, Birkhäuser.

Complex number: Origin

- 1 Assume that we know the definition of integer number.

Complex number: Origin

- 1 Assume that we know the definition of integer number.
Does $5x = 3$ have an integer solution?

Complex number: Origin

- 1 Assume that we know the definition of integer number.
Does $5x = 3$ have an integer solution? NO

Complex number: Origin

- 1 Assume that we know the definition of integer number.
Does $5x = 3$ have an integer solution? NO
- 2 To obtain a solution of the above we need an extension of integer numbers.

Complex number: Origin

- 1 Assume that we know the definition of integer number.
Does $5x = 3$ have an integer solution? NO
- 2 To obtain a solution of the above we need an extension of integer numbers.
- 3 Consider the set \mathbb{R} of real number. Then the above equation has a solution.

Complex number: Origin

- 1 Assume that we know the definition of integer number.
Does $5x = 3$ have an integer solution? NO
- 2 To obtain a solution of the above we need an extension of integer numbers.
- 3 Consider the set \mathbb{R} of real number. Then the above equation has a solution.
- 4 Note that $\mathbb{R} = \{x : x^2 \geq 0\}$.

Complex number: Origin

- 1 Assume that we know the definition of integer number.
Does $5x = 3$ have an integer solution? NO
- 2 To obtain a solution of the above we need an extension of integer numbers.
- 3 Consider the set \mathbb{R} of real number. Then the above equation has a solution.
- 4 Note that $\mathbb{R} = \{x : x^2 \geq 0\}$.
- 5 Does $x^2 = -1$ have a solution on \mathbb{R} ?

Complex number: Origin

- 1 Assume that we know the definition of integer number.
Does $5x = 3$ have an integer solution? NO
- 2 To obtain a solution of the above we need an extension of integer numbers.
- 3 Consider the set \mathbb{R} of real number. Then the above equation has a solution.
- 4 Note that $\mathbb{R} = \{x : x^2 \geq 0\}$.
- 5 Does $x^2 = -1$ have a solution on \mathbb{R} ? NO

Complex number: Origin

- 1 Assume that we know the definition of integer number.
Does $5x = 3$ have an integer solution? NO
- 2 To obtain a solution of the above we need an extension of integer numbers.
- 3 Consider the set \mathbb{R} of real number. Then the above equation has a solution.
- 4 Note that $\mathbb{R} = \{x : x^2 \geq 0\}$.
- 5 Does $x^2 = -1$ have a solution on \mathbb{R} ? NO
- 6 Like before, we need to extend our number system to obtain a solution.

Complex number

- 1 Euler: was the first to use the symbol i for $\sqrt{-1}$.

Complex number

- 1 **Euler:** was the **first** to use the symbol i for $\sqrt{-1}$.
- 2 **Gauss (1831):** He provided a geometric representation of complex number $z = x + iy$ with the coordinate point (x, y) in the plane. This is what we use today.

Complex number

- 1 **Euler**: was the **first** to use the symbol i for $\sqrt{-1}$.
- 2 **Gauss (1831)**: He provided a geometric representation of complex number $z = x + iy$ with the coordinate point (x, y) in the plane. This is what we use today.
- 3 Gauss was in favor to call such number as "**complex number**".

Motivations: Real number vs complex number

Real number vs complex number

- 1 Note \mathbb{R} is the set of all **real numbers**.

Real number vs complex number

- 1 Note \mathbb{R} is the set of all **real numbers**.
- 2 \mathbb{C} is the set of all **complex number->**

Real number vs complex number

- 1 Note \mathbb{R} is the set of all **real numbers**.
- 2 \mathbb{C} is the set of all **complex number**-> xy-plane.

Real number vs complex number

- 1 Note \mathbb{R} is the set of all **real numbers**.
- 2 \mathbb{C} is the set of all **complex numbers** \rightarrow xy-plane.
- 3 **Result:** \mathbb{C} cannot be **totally ordered** in consistence with the usual order on \mathbb{R} .

Explanation

Real number vs complex number

Real number vs complex number

1 Note

$$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$$

is called the **extended real number system**.

Real number vs complex number

1 Note

$$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$$

is called the **extended real number system**.

2 $\hat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$ is called the **extended complex number system**.

Real number vs complex number

1 Note

$$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$$

is called the **extended real number system**.

- 2 $\hat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$ is called the **extended complex number system**.
- 3 There is **only one infinity** in the extended complex number system.

Real number vs complex number

1 Note

$$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$$

is called the **extended real number system**.

- 2 $\hat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$ is called the **extended complex number system**.
- 3 There is **only one infinity** in the extended complex number system.
- 4 We define it as $|z| \rightarrow \infty$ or $z \rightarrow \infty$.

Real number vs complex number

1 Note

$$\mathbb{R}^* := \mathbb{R} \cup \{-\infty, \infty\}$$

is called the **extended real number system**.

- 2 $\hat{\mathbb{C}} := \mathbb{C} \cup \{\infty\}$ is called the **extended complex number system**.
- 3 There is **only one infinity** in the extended complex number system.
- 4 We define it as $|z| \rightarrow \infty$ or $z \rightarrow \infty$.
- 5 Infinite in complex plane means that we are travelling **far away** from the origin **in any direction**.

Real function vs complex function

Real function vs complex function

- 1 We know real functions.

Real function vs complex function

- 1 We know real functions.
- 2 There are two terminologies for complex function:

Real function vs complex function

- 1 We know real functions.
- 2 There are two terminologies for complex function:
 - 1 Complex function with real variable

Real function vs complex function

- 1 We know real functions.
- 2 There are two terminologies for complex function:
 - 1 Complex function with real variable
 - 2 Example:

Real function vs complex function

- 1 We know real functions.
- 2 There are two terminologies for complex function:
 - 1 Complex function with real variable
 - 2 Example:
 - 3 Complex function with complex variable.

Real function vs complex function

- 1 We know real functions.
- 2 There are two terminologies for complex function:
 - 1 Complex function with real variable
 - 2 Example:
 - 3 Complex function with complex variable.
 - 4 Example:

Real function vs complex function

- 1 We know real functions.
- 2 There are two terminologies for complex function:
 - 1 Complex function with real variable
 - 2 Example:
 - 3 Complex function with complex variable.
 - 4 Example:
- 3 In this particular discussion we mean **complex function** by **Complex function with complex variable**.

Real function vs complex function

Real function vs complex function

- 1 Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then it could be bounded or unbounded.

Real function vs complex function

- 1 Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then it could be bounded or unbounded.
- (i) Bounded $f(x) = \sin x$, (ii) unbounded $f(x) = \exp(x)$.

Real function vs complex function

- 1 Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then it could be bounded or unbounded.
 - (i) Bounded $f(x) = \sin x$,
 - (ii) unbounded $f(x) = \exp(x)$.
- 2 Any $f : \mathbb{C} \rightarrow \mathbb{C}$ **non-constant complex** function and its derivatives of all order exist on \mathbb{C} , then it is **ALWAYS** unbounded.

Real function vs complex function

- 1 Let $f : \mathbb{R} \rightarrow \mathbb{R}$ be a **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then it could be bounded or unbounded.
 - (i) Bounded $f(x) = \sin x$,
 - (ii) unbounded $f(x) = \exp(x)$.
- 2 Any $f : \mathbb{C} \rightarrow \mathbb{C}$ **non-constant complex** function and its derivatives of all order exist on \mathbb{C} , then it is **ALWAYS** unbounded.
- 3 Here, boundedness of a complex function $f(z)$ means $|f(z)|$ is unbounded.

Real function vs complex function

- 1 If the **first derivative** of a **real** function exists on \mathbb{R} , we **CANNOT** claim that all higher order derivative exist on \mathbb{R} .

Real function vs complex function

- 1 If the **first derivative** of a **real** function exists on \mathbb{R} , we **CANNOT** claim that all higher order derivative exist on \mathbb{R} .

Example $f(x) = x^{3/2}$,

Real function vs complex function

- 1 If the **first derivative** of a **real** function exists on \mathbb{R} , we **CANNOT** claim that all higher order derivative exist on \mathbb{R} .

Example $f(x) = x^{3/2}$, first derivative exists,

Real function vs complex function

- 1 If the **first derivative** of a **real** function exists on \mathbb{R} , we **CANNOT** claim that all higher order derivative exist on \mathbb{R} .

Example $f(x) = x^{3/2}$, first derivative exists, but higher derivatives do not exist at $x = 0$.

Real function vs complex function

- 1 If the **first derivative** of a **real** function exists on \mathbb{R} , we **CANNOT** claim that all higher order derivatives exist on \mathbb{R} .

Example $f(x) = x^{3/2}$, first derivative exists, but higher derivatives do not exist at $x = 0$.

- 2 If the first derivative of any **complex** function $f : \mathbb{C} \rightarrow \mathbb{C}$ exists, then all order derivatives must exist on \mathbb{C} .

Domain, co-domain and range

We recall the above ideas from real function to discuss the next...

Domain, co-domain and range

Domain, co-domain and range

Real function vs complex function

- 1 A **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then the **range** of f can leave infinitely many values from \mathbb{R} .

Real function vs complex function

- 1 A **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then the **range** of f can leave infinitely many values from \mathbb{R} .
- 2 Example: Range of $f(x) = \exp(x)$ does not contain the **negative real axis** including zero.

Real function vs complex function

- 1 A **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then the **range** of f can leave infinitely many values from \mathbb{R} .
- 2 Example: Range of $f(x) = \exp(x)$ does not contain the **negative real axis** including zero.
- 3 The range of any **non-constant complex** function (derivatives of all order exist) on \mathbb{C} , can leave **at most one value**.

Real function vs complex function

- 1 A **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then the **range** of f can leave infinitely many values from \mathbb{R} .
- 2 Example: Range of $f(x) = \exp(x)$ does not contain the **negative real axis** including zero.
- 3 The range of any **non-constant complex** function (derivatives of all order exist) on \mathbb{C} , can leave **at most one value**. This is called **Picard's Little Theorem** (Emile Picard, at the age of 22, 1856-1941).

Real function vs complex function

- 1 A **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then the **range** of f can leave infinitely many values from \mathbb{R} .
- 2 Example: Range of $f(x) = \exp(x)$ does not contain the **negative real axis** including zero.
- 3 The range of any **non-constant complex** function (derivatives of all order exist) on \mathbb{C} , can leave **at most one value**. This is called **Picard's Little Theorem** (Emile Picard, at the age of 22, 1856-1941).
- 4 Examples: Polynomial
 $P(\mathbb{C}) = \sin(\mathbb{C}) = \cos(\mathbb{C}) = \sinh(\mathbb{C}) = \cosh(\mathbb{C}) = \mathbb{C}$

Real function vs complex function

- 1 A **real, non-constant, smooth** function (derivatives of all order exist) on \mathbb{R} , then the **range** of f can leave infinitely many values from \mathbb{R} .
- 2 Example: Range of $f(x) = \exp(x)$ does not contain the **negative real axis** including zero.
- 3 The range of any **non-constant complex** function (derivatives of all order exist) on \mathbb{C} , can leave **at most one value**. This is called **Picard's Little Theorem** (Emile Picard, at the age of 22, 1856-1941).
- 4 Examples: Polynomial
 $P(\mathbb{C}) = \sin(\mathbb{C}) = \cos(\mathbb{C}) = \sinh(\mathbb{C}) = \cosh(\mathbb{C}) = \mathbb{C}$
- 5 But $e^{(\mathbb{C})} = \mathbb{C} - \{0\}$.

Complex Analysis

LECTURE 2

July 27, 2023

Applications—Why do we study?

- 1 Of course, we need to utilize imaginary number to find zeros of polynomial.

Applications—Why do we study?

- 1 Of course, we need to **utilize imaginary number** to find zeros of polynomial.
- 2 We can easily determine many **complicated real integrations** using the concepts of complex integration techniques.

Applications—Why do we study?

- 1 Of course, we need to **utilize imaginary number** to find zeros of polynomial.
- 2 We can easily determine many **complicated real integrations** using the concepts of complex integration techniques. For instance,

$$\int_0^\infty \sin(t^2) dt = \frac{1}{2} \sqrt{\frac{\pi}{2}}.$$

Applications—Why do we study?

- 1 Of course, we need to **utilize imaginary number** to find zeros of polynomial.
- 2 We can easily determine many **complicated real integrations** using the concepts of complex integration techniques. For instance,

$$\int_0^\infty \sin(t^2) dt = \frac{1}{2} \sqrt{\frac{\pi}{2}}.$$

Applications—Why do we study?

- 1 We can determine the inverse Laplace transform using **complex inversion formula**.

Applications—Why do we study?

- 1 We can determine the inverse Laplace transform using **complex inversion formula**.
- 2 Generally, we memorize the Laplace transform $F(s)$ of a given function $f(t)$.

Applications—Why do we study?

- 1 We can determine the inverse Laplace transform using **complex inversion formula**.
- 2 Generally, we memorize the Laplace transform $F(s)$ of a given function $f(t)$. Then then we recall the inverse Laplace transform $f(t)$ for a given and known $F(s)$.

Applications—Why do we study?

- 1 We can determine the inverse Laplace transform using **complex inversion formula**.
- 2 Generally, we memorize the Laplace transform $F(s)$ of a given function $f(t)$. Then then we recall the inverse Laplace transform $f(t)$ for a given and known $F(s)$. But what would be the inverse Laplace transform of any arbitrary expression $F(s)$?

Applications—Why do we study?

- 1 We can determine the inverse Laplace transform using **complex inversion formula**.
- 2 Generally, we memorize the Laplace transform $F(s)$ of a given function $f(t)$. Then then we recall the inverse Laplace transform $f(t)$ for a given and known $F(s)$. But what would be the inverse Laplace transform of any arbitrary expression $F(s)$? We need to learn the above theorem.

Applications—Why do we study?

- 1 We can determine the inverse Laplace transform using **complex inversion formula**.
- 2 Generally, we memorize the Laplace transform $F(s)$ of a given function $f(t)$. Then then we recall the inverse Laplace transform $f(t)$ for a given and known $F(s)$. But what would be the inverse Laplace transform of any arbitrary expression $F(s)$? We need to learn the above theorem.
- 3 Many theorems (including **prime number theorem**) in number theory can be proved using complex analysis.

Complex number-Review

- 1 A *complex number* denoted by z is an **ordered pair** (x, y) with $x \in \mathbb{R}, y \in \mathbb{R}$.

Complex number-Review

- 1 A *complex number* denoted by z is an **ordered pair** (x, y) with $x \in \mathbb{R}, y \in \mathbb{R}$.
- 2 Also we write $z = x + iy$.

Complex number-Review

- 1 A *complex number* denoted by z is an **ordered pair** (x, y) with $x \in \mathbb{R}, y \in \mathbb{R}$.
- 2 Also we write $z = x + iy$.
- 3 For $z = (x, y)$, x and y are respectively called the **real** and **imaginary** part of z .

Complex number-Review

- 1 A *complex number* denoted by z is an **ordered pair** (x, y) with $x \in \mathbb{R}, y \in \mathbb{R}$.
- 2 Also we write $z = x + iy$.
- 3 For $z = (x, y)$, x and y are respectively called the **real** and **imaginary** part of z .
- 4 In symbolically, $\text{Re}(z) = x$ and $\text{Im}(z) = y$.

Complex number-Review

- 1 A *complex number* denoted by z is an **ordered pair** (x, y) with $x \in \mathbb{R}, y \in \mathbb{R}$.
- 2 Also we write $z = x + iy$.
- 3 For $z = (x, y)$, x and y are respectively called the **real** and **imaginary** part of z .
- 4 In symbolically, $\operatorname{Re}(z) = x$ and $\operatorname{Im}(z) = y$.
- 5 The set of all complex numbers is denoted by \mathbb{C} .

Operations on complex number

- 1 **Addition** of two complex numbers is defined as follows:

$$(a, b) + (c, d) := (a + c, b + d).$$

Operations on complex number

- 1 **Addition** of two complex numbers is defined as follows:

$$(a, b) + (c, d) := (a + c, b + d).$$

- 2 **Multiplication** of complex numbers is defined as

$$(a, b)(c, d) := (ac - bd, bc + ad).$$

Operations on complex number

1 **Addition** of two complex numbers is defined as follows:

$$(a, b) + (c, d) := (a + c, b + d).$$

2 **Multiplication** of complex numbers is defined as

$$(a, b)(c, d) := (ac - bd, bc + ad).$$

3 One can easily verify

- 1 $z_1 + z_2 = z_2 + z_1$.
- 2 $z_1 z_2 = z_2 z_1$.
- 3 $z_1(z_2 + z_3) = z_1 z_2 + z_1 z_3$.

Operations on complex number

- 1 **Inverse** of a **non-zero** complex number is defined as:

Operations on complex number

1 **Inverse** of a **non-zero** complex number is defined as:

$$\frac{1}{z} = \frac{1}{x+iy} = \frac{x-iy}{x^2+y^2}.$$

Operations on complex number

- 1 **Inverse** of a **non-zero** complex number is defined as:

$$\frac{1}{z} = \frac{1}{x+iy} = \frac{x-iy}{x^2+y^2}.$$

- 2 We can define **division** of complex numbers

Operations on complex number

1 **Inverse** of a **non-zero** complex number is defined as:

$$\frac{1}{z} = \frac{1}{x+iy} = \frac{x-iy}{x^2+y^2}.$$

2 We can define **division** of complex numbers

$$\frac{x_1 + iy_1}{x_2 + iy_2} = (x_1 + iy_1) \left(\frac{x_2 - iy_2}{x_2^2 + y_2^2} \right) = \frac{(x_1x_2 + y_1y_2) + i(x_2y_1 - x_1y_2)}{x_2^2 + y_2^2}.$$

Operations on complex number

- 1 **Equal complex numbers:** Two complex numbers (x_1, y_1) and (x_2, y_2) are said to be *equal* if both their real parts and imaginary parts are equal.

Operations on complex number

- 1 **Equal complex numbers:** Two complex numbers (x_1, y_1) and (x_2, y_2) are said to be *equal* if both their real parts and imaginary parts are equal.
- 2 Mathematically,

$$(x_1, y_1) = (x_2, y_2) \text{ if and only if } x_1 = x_2 \text{ and } y_1 = y_2.$$

Complex plane

- 1 For any complex number $z = x + iy$ there is a **unique** ordered pair (x, y) in the xy -plane.

Complex plane

- 1 For any complex number $z = x + iy$ there is a **unique** ordered pair (x, y) in the xy -plane.
- 2 Thus, there is a **one-to-one correspondence** between the complex number $z = x + iy$ and the point (x, y) .
- 3 We refer to xy -plane as the ***complex plane*** or the z -plane.

Complex plane

- 1 For any complex number $z = x + iy$ there is a **unique** ordered pair (x, y) in the **xy-plane**.
- 2 Thus, there is a **one-to-one correspondence** between the complex number $z = x + iy$ and the point (x, y) .
- 3 We refer to **xy-plane** as the **complex plane** or the **z -plane**.
- 4 x -axis is called the **real axis**.

Complex plane

- 1 For any complex number $z = x + iy$ there is a **unique** ordered pair (x, y) in the **xy-plane**.
- 2 Thus, there is a **one-to-one correspondence** between the complex number $z = x + iy$ and the point (x, y) .
- 3 We refer to **xy-plane as the *complex plane*** or the **z -plane**.
- 4 x -axis is called the **real axis**.
- 5 y -axis is called the **imaginary axis**.

Complex conjugate

- 1 The complex conjugate \bar{z} of a given complex number $z = x + iy$ is defined by

$$\bar{z} = x - iy.$$

Complex conjugate

- 1 The complex conjugate \bar{z} of a given complex number $z = x + iy$ is defined by

$$\bar{z} = x - iy.$$

- 2 \bar{z} is just the reflection of the point z wrt x-axis.
- 3 The following properties are easy to prove:
 - 1 $\operatorname{Re}(z) = \frac{1}{2}(z + \bar{z})$ and $\operatorname{Im}(z) = \frac{1}{2i}(z - \bar{z})$.
 - 2 $\bar{z_1 + z_2} = \bar{z_1} + \bar{z_2}$.
 - 3 $\bar{z_1 z_2} = \bar{z_1} \bar{z_2}$, and hence for a real number α , $\overline{\alpha z} = \alpha \bar{z}$.

Magnitude or modulus

- 1 We know $z = x + iy$ represents a point (x, y) in the plane.

Magnitude or modulus

- 1 We know $z = x + iy$ represents a point (x, y) in the plane.
- 2 Each point (x, y) determines a **two dimension vector** with $(0, 0)$ as the initial point and (x, y) as the terminal point.

Magnitude or modulus

- 1 We know $z = x + iy$ represents a point (x, y) in the plane.
- 2 Each point (x, y) determines a **two dimension vector** with $(0, 0)$ as the initial point and (x, y) as the terminal point.
- 3 Addition rule of vectors is also then valid for complex numbers.

Magnitude or modulus

- 1 We know $z = x + iy$ represents a point (x, y) in the plane.
- 2 Each point (x, y) determines a **two dimension vector** with $(0, 0)$ as the initial point and (x, y) as the terminal point.
- 3 Addition rule of vectors is also then valid for complex numbers.
- 4 Magnitude or modulus of z is defined as $|z| = \sqrt{x^2 + y^2}$

Magnitude or modulus

- 1 We know $z = x + iy$ represents a point (x, y) in the plane.
- 2 Each point (x, y) determines a **two dimension vector** with $(0, 0)$ as the initial point and (x, y) as the terminal point.
- 3 Addition rule of vectors is also then valid for complex numbers.
- 4 **Magnitude or modulus** of z is defined as $|z| = \sqrt{x^2 + y^2}$
- 5 We have **triangle inequalities**:

$$|z_1 + z_2| \leq |z_1| + |z_2|,$$

$$|z_1 - z_2| \geq ||z_1| - |z_2||.$$

Polar form of complex numbers

- 1 Ok, we know the magnitude of the vector z .

Polar form of complex numbers

- 1 Ok, we know the magnitude of the vector z .
- 2 Then what about its direction?

Polar form of complex numbers

- 1 Ok, we know the magnitude of the vector z .
- 2 Then what about its direction?
- 3 On the other hand, if we know the magnitude and direction, we'll be able to determine z uniquely.

Polar form of complex numbers

- 1 Ok, we know the magnitude of the vector z .
- 2 Then what about its direction?
- 3 On the other hand, if we know the magnitude and direction, we'll be able to determine z uniquely.
- 4 To find the direction we determine an angle corresponding to given z .

Polar form of complex numbers

- 1 Ok, we know the magnitude of the vector z .
- 2 Then what about its direction?
- 3 On the other hand, if we know the magnitude and direction, we'll be able to determine z uniquely.
- 4 To find the direction we determine an angle corresponding to given z .
- 5 To measure the angle we have to fix another vector.

Polar form of complex numbers

- 1 Ok, we know the magnitude of the vector z .
- 2 Then what about its direction?
- 3 On the other hand, if we know the magnitude and direction, we'll be able to determine z uniquely.
- 4 To find the direction we determine an angle corresponding to given z .
- 5 To measure the angle we have to fix another vector.
- 6 Conventionally, we fix positive x -axis as the initial vector.

Polar form of complex numbers

- 1 Let $z = x + iy$ be a **non-zero** complex number.

Polar form of complex numbers

- 1 Let $z = x + iy$ be a **non-zero** complex number.
- 2 Its **Cartesian** form is (x, y)

Polar form of complex numbers

- 1 Let $z = x + iy$ be a **non-zero** complex number.
- 2 Its **Cartesian** form is (x, y)
- 3 We know the **polar** representation of (x, y) as

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta$$

Polar form of complex numbers

- 1 Let $z = x + iy$ be a **non-zero** complex number.
- 2 Its **Cartesian** form is (x, y)
- 3 We know the **polar** representation of (x, y) as

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta$$

- 4 Our aim is to find r and θ for a give complex number.

Polar form of complex numbers

- 1 Let $z = x + iy$ be a **non-zero** complex number.
- 2 Its **Cartesian** form is (x, y)
- 3 We know the **polar** representation of (x, y) as

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta$$

- 4 Our aim is to find r and θ for a give complex number.
- 5 Easy to check that $r = \sqrt{x^2 + y^2}$ - a **unique value**

Polar form of complex numbers

- 1 Let $z = x + iy$ be a **non-zero** complex number.
- 2 Its **Cartesian** form is (x, y)
- 3 We know the **polar** representation of (x, y) as

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta$$

- 4 Our aim is to find r and θ for a give complex number.
- 5 Easy to check that $r = \sqrt{x^2 + y^2}$ - a **unique value**
- 6 Ok, let us encounter the **trouble** to determine θ .

Polar form of complex numbers

- 1 Let $z = x + iy$ be a **non-zero** complex number.
- 2 Its **Cartesian** form is (x, y)
- 3 We know the **polar** representation of (x, y) as

$$x = r \cos \theta \quad \text{and} \quad y = r \sin \theta$$

- 4 Our aim is to find r and θ for a give complex number.
- 5 Easy to check that $r = \sqrt{x^2 + y^2}$ - a **unique value**
- 6 Ok, let us encounter the **trouble** to determine θ .
- 7 Chose $z = 1 - i$.

Polar form of complex numbers

1 Here $r = \sqrt{2}$.

Polar form of complex numbers

- 1 Here $r = \sqrt{2}$.
- 2 We have to chose θ such that

Polar form of complex numbers

- 1 Here $r = \sqrt{2}$.
- 2 We have to chose θ such that

$$1 = \sqrt{2} \cos \theta \quad \text{and} \quad -1 = \sqrt{2} \sin \theta.$$

Polar form of complex numbers

- 1 Here $r = \sqrt{2}$.
- 2 We have to choose θ such that

$$1 = \sqrt{2} \cos \theta \quad \text{and} \quad -1 = \sqrt{2} \sin \theta.$$

- 3 What happens if we choose $\theta = \left(\frac{3\pi}{2} + \frac{\pi}{4}\right)$ and $-\frac{\pi}{4}$?

Polar form of complex numbers

- 1 Here $r = \sqrt{2}$.
- 2 We have to choose θ such that

$$1 = \sqrt{2} \cos \theta \quad \text{and} \quad -1 = \sqrt{2} \sin \theta.$$

- 3 What happens if we choose $\theta = \left(\frac{3\pi}{2} + \frac{\pi}{4}\right)$ and $-\frac{\pi}{4}$?
- 4 Both values work. **Very confusing** to fix θ uniquely.

Polar form of complex numbers




Figure: 3: Two angles.

Polar form of complex numbers

- 1 Which angle should we choose?.

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):
 - (i) $\text{Arg}(z)$

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):
 - (i) $\text{Arg}(z)$ —called **principal argument**

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):
 - (i) $\text{Arg}(z)$ —called **principal argument**— a unique value

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):
 - (i) $\text{Arg}(z)$ —called **principal argument**— a unique value
 - (ii) $\arg(z)$

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):
 - (i) $\text{Arg}(z)$ —called **principal argument**— a unique value
 - (ii) $\arg(z)$ —called **argument**

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):
 - (i) $\text{Arg}(z)$ —called **principal argument**— a unique value
 - (ii) $\arg(z)$ —called **argument**— **multi-valued function**

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):
 - (i) $\text{Arg}(z)$ —called **principal argument**— a unique value
 - (ii) $\arg(z)$ —called **argument**— **multi-valued function**
- 3 We define $\text{Arg}(z) = \theta \in (-\pi, \pi]$. This is a **unique measurement** for a given z

Polar form of complex numbers

- 1 Which angle should we choose?.
- 2 We now define two measure/quantities of angle (in **radian**):
 - (i) $\text{Arg}(z)$ —called **principal argument**— a unique value
 - (ii) $\arg(z)$ —called **argument**— **multi-valued function**
- 3 We define $\text{Arg}(z) = \theta \in (-\pi, \pi]$. This is a **unique measurement** for a given z
- 4 $\arg(z) = \theta + 2k\pi, k = 0, \pm 1, \pm 2, \dots$ (**Why $2k\pi$?**)

Problems

1 $\text{Arg}(-5i) = -\frac{\pi}{2}$

Problems

1 $\text{Arg}(-5i) = -\frac{\pi}{2}$

2 $\text{Arg}(-1 + i) = \frac{3\pi}{4}$

Problems

$$1 \quad \text{Arg}(-5i) = -\frac{\pi}{2}$$

$$2 \quad \text{Arg}(-1 + i) = \frac{3\pi}{4}$$

$$3 \quad \text{Arg}(-1 - i) = -\frac{3\pi}{4}$$

Problems

$$1 \quad \text{Arg}(-5i) = -\frac{\pi}{2}$$

$$2 \quad \text{Arg}(-1 + i) = \frac{3\pi}{4}$$

$$3 \quad \text{Arg}(-1 - i) = -\frac{3\pi}{4}$$

$$4 \quad \text{Arg}(-1 + \sqrt{3}i) = -\frac{2\pi}{3}$$

Problems

Let $z_k = r_k(\cos \theta_k + i \sin \theta_k)$, $k = 1, 2, \dots, n$. Show that

1 $z_1 z_2 = r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)]$.

Problems

Let $z_k = r_k(\cos \theta_k + i \sin \theta_k)$, $k = 1, 2, \dots, n$. Show that

1 $z_1 z_2 = r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)]$.

Very straight forward

Problems

Let $z_k = r_k(\cos \theta_k + i \sin \theta_k)$, $k = 1, 2, \dots, n$. Show that

1 $z_1 z_2 = r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)]$.

Very straight forward

2 $z_1 z_2 \cdots z_n = r_1 r_2 \cdots r_n [\cos(\theta_1 + \theta_2 + \cdots + \theta_n) + i \sin(\theta_1 + \theta_2 + \cdots + \theta_n)]$.

Problems

Let $z_k = r_k(\cos \theta_k + i \sin \theta_k)$, $k = 1, 2, \dots, n$. Show that

1 $z_1 z_2 = r_1 r_2 [\cos(\theta_1 + \theta_2) + i \sin(\theta_1 + \theta_2)]$.

Very straight forward

2 $z_1 z_2 \cdots z_n = r_1 r_2 \cdots r_n [\cos(\theta_1 + \theta_2 + \cdots + \theta_n) + i \sin(\theta_1 + \theta_2 + \cdots + \theta_n)]$.

Use method of induction

De Moiver's formula

1 De Moiver's formula:-1722—

De Moiver's formula

- 1 De Moiver's formula:-1722—If $z = r(\cos \theta + i \sin \theta)$, then any n (an integer),

$$z^n = [r(\cos \theta + i \sin \theta)]^n = r^n(\cos n\theta + i \sin n\theta).$$

De Moiver's formula

- 1 De Moiver's formula:-1722—If $z = r(\cos \theta + i \sin \theta)$, then any n (an integer),

$$z^n = [r(\cos \theta + i \sin \theta)]^n = r^n(\cos n\theta + i \sin n\theta).$$

De Moiver's formula

- 1 De Moiver's formula:-1722—if $z = r(\cos \theta + i \sin \theta)$, then any n (an integer),

$$z^n = [r(\cos \theta + i \sin \theta)]^n = r^n(\cos n\theta + i \sin n\theta).$$

- 2 Generally, De Moiver's formula is invalid for non integer n .
Give an example.

De Moiver's formula

- 1 De Moiver's formula:-1722—If $z = r(\cos \theta + i \sin \theta)$, then any n (an integer),

$$z^n = [r(\cos \theta + i \sin \theta)]^n = r^n(\cos n\theta + i \sin n\theta).$$

- 2 Generally, De Moiver's formula is invalid for non integer n .
Give an example.
- 3 Sometimes, De Moiver's formula is defined with $r = 1$.

Applications of De Moiver's theorem

- 1 Easily we can determine the **higher power** of a complex number.

Applications of De Moivre's theorem

- 1 Easily we can determine the **higher power** of a complex number.
- 2 Plays an important role in **finding explicit expressions for the n th roots of unity**.

Power of a complex number-Application of De Moiver's theorem




Figure: 4: Geometrical view of power of $2i$

Root finding

Problem: For a given complex number z_0 and $n \in \mathbb{N}$, determine **all values** of z satisfying $z^n = z_0$.

Check the following example and do by yourself:

Root finding

Determine all values of z such that $z^3 = -1$.

Root finding

Determine all values of z such that $z^3 = -1$.

1 $\operatorname{Arg}(-1) = \pi$.

Root finding

Determine all values of z such that $z^3 = -1$.

- 1 $\text{Arg}(-1) = \pi$.
- 2 Set $z = r(\cos \theta + i \sin \theta)$.

Root finding

Determine all values of z such that $z^3 = -1$.

- 1 $\text{Arg}(-1) = \pi$.
- 2 Set $z = r(\cos \theta + i \sin \theta)$.
- 3 $z^3 = r^3(\cos 3\theta + i \sin 3\theta)$
- 4 Thus, $z^3 = -1 = 1 \cdot \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$,
 $k = 0, \pm 1, \pm 2, \dots$

Root finding

Determine all values of z such that $z^3 = -1$.

- 1 $\text{Arg}(-1) = \pi$.
- 2 Set $z = r(\cos \theta + i \sin \theta)$.
- 3 $z^3 = r^3(\cos 3\theta + i \sin 3\theta)$
- 4 Thus, $z^3 = -1 = 1 \cdot \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$,
 $k = 0, \pm 1, \pm 2, \dots$
- 5 $r^3(\cos 3\theta + i \sin 3\theta) = \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$

Root finding

Determine all values of z such that $z^3 = -1$.

- 1 $\text{Arg}(-1) = \pi$.
- 2 Set $z = r(\cos \theta + i \sin \theta)$.
- 3 $z^3 = r^3(\cos 3\theta + i \sin 3\theta)$
- 4 Thus, $z^3 = -1 = 1 \cdot \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$,
 $k = 0, \pm 1, \pm 2, \dots$
- 5 $r^3(\cos 3\theta + i \sin 3\theta) = \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$

Root finding

Determine all values of z such that $z^3 = -1$.

- 1 $\text{Arg}(-1) = \pi$.
- 2 Set $z = r(\cos \theta + i \sin \theta)$.
- 3 $z^3 = r^3(\cos 3\theta + i \sin 3\theta)$
- 4 Thus, $z^3 = -1 = 1 \cdot \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$,
 $k = 0, \pm 1, \pm 2, \dots$
- 5 $r^3(\cos 3\theta + i \sin 3\theta) = \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$
- 6 $r = 1$ and $\theta = \frac{\pi + 2k\pi}{3}$, $k = 0, \pm 1, \pm 2, \dots$

Root finding

Determine all values of z such that $z^3 = -1$.

- 1 $\text{Arg}(-1) = \pi$.
- 2 Set $z = r(\cos \theta + i \sin \theta)$.
- 3 $z^3 = r^3(\cos 3\theta + i \sin 3\theta)$
- 4 Thus, $z^3 = -1 = 1 \cdot \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$,
 $k = 0, \pm 1, \pm 2, \dots$
- 5 $r^3(\cos 3\theta + i \sin 3\theta) = \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$
- 6 $r = 1$ and $\theta = \frac{\pi + 2k\pi}{3}$, $k = 0, \pm 1, \pm 2, \dots$
- 7 $z_k = \cos\left(\frac{\pi + 2k\pi}{3}\right) + i \sin\left(\frac{\pi + 2k\pi}{3}\right)$, $k = 0, 1, 2$.

Root finding

Determine all values of z such that $z^3 = -1$.

- 1 $\text{Arg}(-1) = \pi$.
- 2 Set $z = r(\cos \theta + i \sin \theta)$.
- 3 $z^3 = r^3(\cos 3\theta + i \sin 3\theta)$
- 4 Thus, $z^3 = -1 = 1 \cdot \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$,
 $k = 0, \pm 1, \pm 2, \dots$
- 5 $r^3(\cos 3\theta + i \sin 3\theta) = \cos(\pi + 2k\pi) + i \sin(\pi + 2k\pi)$
- 6 $r = 1$ and $\theta = \frac{\pi + 2k\pi}{3}$, $k = 0, \pm 1, \pm 2, \dots$
- 7 $z_k = \cos\left(\frac{\pi + 2k\pi}{3}\right) + i \sin\left(\frac{\pi + 2k\pi}{3}\right)$, $k = 0, 1, 2$.
- 8 We have only **three** distinct values of z .

Location of roots




Figure: 5: Geometrical view of the roots.

Finding nth roots of number z_0

- 1 Finding nth roots of number z_0 means we have to find the solution of $z^n = z_0$

Finding nth roots of number z_0

- 1 Finding nth roots of number z_0 means we have to find the solution of $z^n = z_0$
- 2 In notation, it is also written as $z = z_0^{1/n}$

Finding nth roots of number z_0

- 1 Finding nth roots of number z_0 means we have to find the solution of $z^n = z_0$
- 2 In notation, it is also written as $z = z_0^{1/n}$
- 3 If it is asked to find the values of $z_0^{1/n}$, then we basically find the nth roots of z_0

Finding nth roots of number z_0

- 1 Finding nth roots of number z_0 means we have to find the solution of $z^n = z_0$
- 2 In notation, it is also written as $z = z_0^{1/n}$
- 3 If it is asked to find the values of $z_0^{1/n}$, then we basically find the nth roots of z_0
- 4 I suggest to read (books) about the location of roots (geometrical view) on the complex plane.

Geometrical representation of sets or regions in \mathbb{C}

- 1 Sets are very important to **define domain, codomain, and range** of complex functions.

Geometrical representation of sets or regions in \mathbb{C}

- 1 Sets are very important to **define domain, codomain, and range** of complex functions.
- 2 Sets and points in sets also help to **define limit, continuity and differentiability**.

Geometrical representation of sets or regions in \mathbb{C}

- 1 Sets are very important to **define domain, codomain, and range** of complex functions.
- 2 Sets and points in sets also help to **define limit, continuity and differentiability**.
- 3 **Straight line:** $\operatorname{Re}z = \operatorname{Im}z$ represents the straight line $y = x$.

Geometrical representation of sets or regions in \mathbb{C}

- 1 Sets are very important to **define domain, codomain, and range** of complex functions.
- 2 Sets and points in sets also help to **define limit, continuity and differentiability**.
- 3 **Straight line:** $\operatorname{Re}z = \operatorname{Im}z$ represents the straight line $y = x$.
- 4 **General equation of straight line:** $\bar{a}z + a\bar{z} + b = 0$, where a is a **complex number** and b is a **real number**.

Geometrical representation of sets or regions in \mathbb{C}

- 1 Sets are very important to **define domain, codomain, and range** of complex functions.
- 2 Sets and points in sets also help to **define limit, continuity and differentiability**.
- 3 **Straight line:** $\operatorname{Re}z = \operatorname{Im}z$ represents the straight line $y = x$.
- 4 **General equation of straight line:** $\bar{a}z + a\bar{z} + b = 0$, where a is a **complex number** and b is a **real number**. **Establish it.**

Geometrical representation of sets or regions in \mathbb{C}


- 1 Sets are very important to **define domain, codomain, and range** of complex functions.
- 2 Sets and points in sets also help to **define limit, continuity and differentiability**.
- 3 **Straight line:** $\operatorname{Re}z = \operatorname{Im}z$ represents the straight line $y = x$.
- 4 **General equation of straight line:** $\bar{a}z + a\bar{z} + b = 0$, where a is a **complex number** and b is a **real number**. Establish it.
- 5 $|z - z_0| = 2 \implies \sqrt{(x - x_0)^2 + (y - y_0)^2} = 2$. The subset of points z consists of the points on the circle with center at $z_0 = (x_0, y_0)$ and radius $r = 2$.

Open disk

The collection of z satisfying $|z - z_0| < 2$ represents the subset consisting of the points (red points) **inside** the circle with center at $z_0 = (x_0, y_0)$ and radius $r = 2$.

Open disk

The collection of z satisfying $|z - z_0| < 2$ represents the subset consisting of the points (red points) **inside** the circle with center at $z_0 = (x_0, y_0)$ and radius $r = 2$.




Closed disk

The collection of z satisfying $|z - z_0| \leq 2$ represents the subset consisting of the points inside (red points) and on the circle (blue points) the circle with center at $z_0 = (x_0, y_0)$ and radius $r = 2$.

Closed disk

The collection of z satisfying $|z - z_0| \leq 2$ represents the subset consisting of the points inside (red points) and on the circle (blue points) the circle with center at $z_0 = (x_0, y_0)$ and radius $r = 2$.




Annulus

The collection of z satisfying $1 < |z - z_0| < 3$ represents an annulus.

Annulus

The collection of z satisfying $1 < |z - z_0| < 3$ represents an annulus.



Punctured disk

- 1 The collection of z satisfying $0 < |z - z_0| < 2$ represents an punctured disk

Punctured disk

- 1 The collection of z satisfying $0 < |z - z_0| < 2$ represents an punctured disk
- 2 The subset contains all points of $|z - z_0| < 2$ except z_0 .

Punctured disk

- 1 The collection of z satisfying $0 < |z - z_0| < 2$ represents an punctured disk
- 2 The subset contains all points of $|z - z_0| < 2$ except z_0 .
- 3 It has a hole at z_0 .

Complex Analysis

LECTURE 3

July 31, 2023

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define:

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set, interior points of a set,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set, interior points of a set, boundary points of a set,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set, interior points of a set, boundary points of a set, open set,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set, interior points of a set, boundary points of a set, open set, closed set,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set, interior points of a set, boundary points of a set, open set, closed set, bounded and unbounded set,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set, interior points of a set, boundary points of a set, open set, closed set, bounded and unbounded set, connected set,

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set, interior points of a set, boundary points of a set, open set, closed set, bounded and unbounded set, connected set, domain, and

Some topological definitions

- 1 Some of the definitions will be useful to categorize different sets.
- 2 We define: neighborhood of a point, deleted neighborhood of a point, limit points of a set, interior points of a set, boundary points of a set, open set, closed set, bounded and unbounded set, connected set, domain, and region.
- 3 We can apply these definitions to define limit, continuity and differentiability of functions.

Neighborhood of a point z_0

Neighborhood of a point z_0

Neighborhood of a point z_0

Neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $|z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.

Neighborhood of a point z_0

Neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $|z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $N(z_0; \epsilon)$.

Neighborhood of a point z_0

Neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $|z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $N(z_0; \epsilon)$.
- 3 $N(z_0; \epsilon) = \{z : |z - z_0| < \epsilon\}$.

Neighborhood of a point z_0

Neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $|z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $N(z_0; \epsilon)$.
- 3 $N(z_0; \epsilon) = \{z : |z - z_0| < \epsilon\}$.
- 4 $N(z_0; \epsilon)$ contains z_0 too.

Neighborhood of a point z_0

Neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $|z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $N(z_0; \epsilon)$.
- 3 $N(z_0; \epsilon) = \{z : |z - z_0| < \epsilon\}$.
- 4 $N(z_0; \epsilon)$ contains z_0 too.
- 5 Where do we apply?

Neighborhood of a point z_0

Neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $|z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $N(z_0; \epsilon)$.
- 3 $N(z_0; \epsilon) = \{z : |z - z_0| < \epsilon\}$.
- 4 $N(z_0; \epsilon)$ contains z_0 too.
- 5 Where do we apply? **Verifying continuity**.

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $0 < |z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $0 < |z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $\hat{N}(z_0; \epsilon)$.

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $0 < |z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $\hat{N}(z_0; \epsilon)$.
- 3 $\hat{N}(z_0; \epsilon) = \{z : 0 < |z - z_0| < \epsilon\}$.

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $0 < |z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $\hat{N}(z_0; \epsilon)$.
- 3 $\hat{N}(z_0; \epsilon) = \{z : 0 < |z - z_0| < \epsilon\}$.
- 4 $\hat{N}(z_0; \epsilon)$ does **NOT** contain z_0 .

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $0 < |z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $\hat{N}(z_0; \epsilon)$.
- 3 $\hat{N}(z_0; \epsilon) = \{z : 0 < |z - z_0| < \epsilon\}$.
- 4 $\hat{N}(z_0; \epsilon)$ does **NOT** contain z_0 .
- 5 $\hat{N}(z_0; \epsilon)$ is a **punctured disk**.

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $0 < |z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $\hat{N}(z_0; \epsilon)$.
- 3 $\hat{N}(z_0; \epsilon) = \{z : 0 < |z - z_0| < \epsilon\}$.
- 4 $\hat{N}(z_0; \epsilon)$ does **NOT** contain z_0 .
- 5 $\hat{N}(z_0; \epsilon)$ is a **punctured disk**.
- 6 Where do we apply?

Deleted neighborhood of a point z_0

Deleted neighborhood of a point z_0

- 1 Mathematically, neighborhood of a point z_0 is a **set of points z** such that $0 < |z - z_0| < \epsilon$, $z_0 \in \mathbb{C}$ and $\epsilon > 0$ is arbitrary small real.
- 2 We denote this set as $\hat{N}(z_0; \epsilon)$.
- 3 $\hat{N}(z_0; \epsilon) = \{z : 0 < |z - z_0| < \epsilon\}$.
- 4 $\hat{N}(z_0; \epsilon)$ does **NOT** contain z_0 .
- 5 $\hat{N}(z_0; \epsilon)$ is a **punctured disk**.
- 6 Where do we apply? **Verifying limit**.

Limit points of a set S

Limit points of a set S

Limit points of a set S

Limit point of a set S

Limit points of a set S

Limit point of a set S

- 1 Limit point is also called **accumulation point**.

Limit points of a set S

Limit point of a set S

- 1 Limit point is also called **accumulation point**.
- 2 Let S be a subset of \mathbb{C} .

Limit points of a set S

Limit point of a set S

- 1 Limit point is also called **accumulation point**.
- 2 Let S be a subset of \mathbb{C} .
- 3 Let $z_1 \in \mathbb{C}$.

Limit points of a set S


Limit point of a set S

- 1 Limit point is also called **accumulation point**.
- 2 Let S be a subset of \mathbb{C} .
- 3 Let $z_1 \in \mathbb{C}$.
- 4 z_1 is called a limit point of S if $\hat{N}(z_1; \epsilon)$ contains **a point** of S .

Limit points of a set S

Limit point of a set S

- 1 Limit point is also called **accumulation point**.
- 2 Let S be a subset of \mathbb{C} .
- 3 Let $z_1 \in \mathbb{C}$.
- 4 z_1 is called a limit point of S if $\hat{N}(z_1; \epsilon)$ contains **a point** of S .
- 5 Mathematically, $\hat{N}(z_1; \epsilon) \cap S \neq \emptyset$.

Limit points of a set S Figure: z_1, z_2 are limit points, but not z_3

Example: Limit points of a set S

- 1 Let $S = \{z : |z| < 1\}$.

Example: Limit points of a set S

- 1 Let $S = \{z : |z| < 1\}$.
- 2 All points of S are limit points of S .

Example: Limit points of a set S

- 1 Let $S = \{z : |z| < 1\}$.
- 2 All points of S are limit points of S .
- 3 In addition, all points on the circle $|z| = 1$ are limit points of S .

Example: Limit points of a set S

- 1 Let $S = \{z : |z| < 1\}$.
- 2 All points of S are limit points of S .
- 3 In addition, all points on the circle $|z| = 1$ are limit points of S .
- 4 Thus, limit points of S are $\{z : |z| \leq 1\}$.

Example: Limit points of a set S

- 1 Let $S = \{z : |z| < 1\}$.
- 2 All points of S are limit points of S .
- 3 In addition, all points on the circle $|z| = 1$ are limit points of S .
- 4 Thus, limit points of S are $\{z : |z| \leq 1\}$.
- 5 What are the limit points of the set $A = \{z : |z| \leq 1\}$?

Example: Limit points of a set S

- 1 Let $S = \{z : |z| < 1\}$.
- 2 All points of S are limit points of S .
- 3 In addition, all points on the circle $|z| = 1$ are limit points of S .
- 4 Thus, limit points of S are $\{z : |z| \leq 1\}$.
- 5 What are the limit points of the set $A = \{z : |z| \leq 1\}$?

Limit points of a set S

Problem: Let c_0 be a limit point of S . Then prove that every neighbourhood of c_0 contains infinitely many points of S .

Try yourself.

Ask me if you need help.

Interior points of a set

Interior points of a set

Interior points of a set

Interior points of a set

Interior points of a set

- 1 Let S be a subset of \mathbb{C} .

Interior points of a set

Interior points of a set

- 1 Let S be a subset of \mathbb{C} .
- 2 Let $z_1 \in S$.

Interior points of a set

Interior points of a set


- 1 Let S be a subset of \mathbb{C} .
- 2 Let $z_1 \in S$.
- 3 z_1 is called an interior point of S if it is **possible** to find a $N(z_1; \epsilon)$ such that $N(z_1; \epsilon) \subset S$.

Interior points of a set

Interior points of a set

- 1 Let S be a subset of \mathbb{C} .
- 2 Let $z_1 \in S$.
- 3 z_1 is called an interior point of S if it is **possible** to find a $N(z_1; \epsilon)$ such that $N(z_1; \epsilon) \subset S$.
- 4 In language, neighborhood of z_1 will be **completely inside** S .

Interior points of a set

Figure: z_1 is an interior point.

Scanned with CamScanner

Interior points of a set

Problem: Mathematically prove that all points of $S = \{z : |z - z_0| < 1\}$ are interior points.

Try yourself.

Boundary points of a set S

Boundary points of a set S

Boundary points of a set S

Boundary points of a set S

- 1 Let S be a subset of \mathbb{C} .

Boundary points of a set S

Boundary points of a set S

- 1 Let S be a subset of \mathbb{C} .
- 2 Let $z_1 \in \mathbb{C}$.

Boundary points of a set S

Boundary points of a set S

- 1 Let S be a subset of \mathbb{C} .
- 2 Let $z_1 \in \mathbb{C}$.
- 3 z_1 is called a boundary point of S if $N(z_1; \epsilon)$ contains at least one point of S and at least one point **not belonging** to S .

Example: Boundary points of a set S

- 1 Let $S = \{z : |z| < 1\}$ be a subset of \mathbb{C} .

Example: Boundary points of a set S

- 1 Let $S = \{z : |z| < 1\}$ be a subset of \mathbb{C} .
- 2 Then each point of $B = \{z : |z| = 1\}$ represents a boundary point.

Example: Boundary points of a set S

- 1 Let $S = \{z : |z| < 1\}$ be a subset of \mathbb{C} .
- 2 Then each point of $B = \{z : |z| = 1\}$ represents a boundary point.
- 3 **Boundary:** Collection of all boundary points is called the **boundary** of S .

Example: Boundary points of a set S

- 1 Let $S = \{z : |z| < 1\}$ be a subset of \mathbb{C} .
- 2 Then each point of $B = \{z : |z| = 1\}$ represents a boundary point.
- 3 **Boundary:** Collection of all boundary points is called the **boundary** of S .
- 4 Here B is the boundary of S

Open and closed sets

- 1 **Open set:** A set S be a subset of \mathbb{C} . S is said to be an open set if each point of S is an interior point of S .

Open and closed sets

- 1 **Open set:** A set S be a subset of \mathbb{C} . S is said to be an open set if each point of S is an interior point of S .
- 2 Example: Let $S = \{z : |z| < 1\}$

Open and closed sets

- 1 **Open set:** A set S be a subset of \mathbb{C} . S is said to be an open set if each point of S is an interior point of S .
- 2 Example: Let $S = \{z : |z| < 1\}$
- 3 **Closed set:** A set S be a subset of \mathbb{C} . S is said to be a closed set if it contains all of its boundary points.

Open and closed sets

- 1 **Open set:** A set S be a subset of \mathbb{C} . S is said to be an open set if each point of S is an interior point of S .
- 2 Example: Let $S = \{z : |z| < 1\}$
- 3 **Closed set:** A set S be a subset of \mathbb{C} . S is said to be a closed set if it contains all of its boundary points.
- 4 Example: Let $S = \{z : |z| \leq 1\}$.

Bounded and unbounded sets

- 1 **Bounded set:** A set S be a subset of \mathbb{C} is said to be bounded if $S \subseteq \{z : |z| < M\}$ for some real number M . Otherwise, it is called **unbounded**.

Bounded and unbounded sets

- 1 **Bounded set:** A set S be a subset of \mathbb{C} is said to be bounded if $S \subseteq \{z : |z| < M\}$ for some real number M . Otherwise, it is called **unbounded**.
- 2 $S = \{z : |z| < 1\}$ is **bounded set** with $M = 1$.

Bounded and unbounded sets

- 1 **Bounded set:** A set S be a subset of \mathbb{C} is said to be bounded if $S \subseteq \{z : |z| < M\}$ for some real number M . Otherwise, it is called **unbounded**.
- 2 $S = \{z : |z| < 1\}$ is **bounded set** with $M = 1$.
- 3 $S = \{z : |z| > 1\}$ is an **unbounded set**.

Bounded and unbounded sets


- 1 **Bounded set:** A set S be a subset of \mathbb{C} is said to be bounded if $S \subseteq \{z : |z| < M\}$ for some real number M . Otherwise, it is called **unbounded**.
- 2 $S = \{z : |z| < 1\}$ is **bounded set** with $M = 1$.
- 3 $S = \{z : |z| > 1\}$ is an **unbounded set**.
- 4 $S = \{z : y = x^2, x, y \in \mathbb{R}\}$ is an **unbounded set**.

Connected set

Connected set

A set S be a subset of \mathbb{C} is said to be connected if **any two points** in S can be joined by a polygonal line (pieces of lines joined end to end) or by any **continuous curve** which **lies entirely on S** .

Connected set

Figure: U, V are connected, but $U \cup V$ is not.

Examples

- 1 $S = \{z : |z| < 1\}$ is a connected set.

Examples

- 1 $S = \{z : |z| < 1\}$ is a connected set.
- 2 $S = \{z : |z| > 1\}$ is a connected set.

Examples

- 1 $S = \{z : |z| < 1\}$ is a connected set.
- 2 $S = \{z : |z| > 1\}$ is a connected set.
- 3 Neighborhood

Examples

- 1 $S = \{z : |z| < 1\}$ is a connected set.
- 2 $S = \{z : |z| > 1\}$ is a connected set.
- 3 Neighborhood and deleted neighborhood are connected sets.

Domain

An **open connected set** is called a domain.

Domain




Figure: S_1 is domain, but S_2, S_3 are not.

Examples

- 1 $S = \{z : |z| < 1\}$ is a domain.

Examples

- 1 $S = \{z : |z| < 1\}$ is a domain.
- 2 $S = \{z : |z| > 1\}$ is a domain.

Examples

- 1 $S = \{z : |z| < 1\}$ is a domain.
- 2 $S = \{z : |z| > 1\}$ is a domain.
- 3 Neighborhood

Examples

- 1 $S = \{z : |z| < 1\}$ is a domain.
- 2 $S = \{z : |z| > 1\}$ is a domain.
- 3 Neighborhood and deleted neighborhood are domain.

Region

A region is a **domain** together with all, some or none of its **boundary** points.

Examples

- 1 $S = \{z : |z| < 1\}$ is a region.

Examples

- 1 $S = \{z : |z| < 1\}$ is a region.
- 2 $S = \{z : |z| \leq 1\}$ is a region.

Examples

- 1 $S = \{z : |z| < 1\}$ is a region.
- 2 $S = \{z : |z| \leq 1\}$ is a region.
- 3 $S = \{z : |z| > 1\}$ is a region.

Examples

1 $S = \{z : |z| < 1\}$ is a region.

2 $S = \{z : |z| \leq 1\}$ is a region.

3 $S = \{z : |z| > 1\}$ is a region.

4 $S = \{z : |z| \geq 1\}$ is a region.

Examples

- 1 $S = \{z : |z| < 1\}$ is a region.
- 2 $S = \{z : |z| \leq 1\}$ is a region.
- 3 $S = \{z : |z| > 1\}$ is a region.
- 4 $S = \{z : |z| \geq 1\}$ is a region.
- 5 A **domain** is also a **region**, but the converse is not true.

Complex Analysis

LECTURE 4

August 3, 2023

Complex function

- 1 Let S be a subset of complex numbers in the complex plane.

Complex function

- 1 Let S be a subset of complex numbers in the complex plane.
- 2 For every point $z = x + iy \in S$, we specify the rule for assigning a corresponding complex number $w = u + iv$.
- 3 This defines a function of the complex variable z , and the function is denoted by

$$w = f(z).$$

Complex function

- 1 Geometrically, elements of S can be represented in the complex plane \mathbb{C} .

Complex function

- 1 Geometrically, elements of S can be represented in the complex plane \mathbb{C} .
- 2 The elements of $f(z)$ is presented in a different complex plane i.e., on the w -plane here.

Complex function

- 1 Geometrically, elements of S can be represented in the complex plane \mathbb{C} .
- 2 The elements of $f(z)$ is presented in a different complex plane i.e., on the w -plane here.
- 3 The set S is called the **domain of definition** of the function f or simply domain of f .

Complex function

- 1 Geometrically, elements of S can be represented in the complex plane \mathbb{C} .
- 2 The elements of $f(z)$ is presented in a different complex plane i.e., on the w -plane here.
- 3 The set S is called the **domain of definition** of the function f or simply domain of f .
- 4 The collection of all values of w is called the **range of f** .

Complex function

- 1 Geometrically, elements of S can be represented in the complex plane \mathbb{C} .
- 2 The elements of $f(z)$ is presented in a different complex plane i.e., on the w -plane here.
- 3 The set S is called the **domain of definition** of the function f or simply domain of f .
- 4 The collection of all values of w is called the **range of f** .
- 5 Mathematically, range denoted by $R(f) = \{f(z) : z \in S\}$.

Complex function

- 1 Geometrically, elements of S can be represented in the complex plane \mathbb{C} .
- 2 The elements of $f(z)$ is presented in a different complex plane i.e., on the w -plane here.
- 3 The set S is called the **domain of definition** of the function f or simply domain of f .
- 4 The collection of all values of w is called the **range of f** .
- 5 Mathematically, range denoted by $R(f) = \{f(z) : z \in S\}$.
- 6 **Domain of definition** and **domain (set)** are different terminology. "Domain of definition" could be a region as well.

Complex function

- 1 Geometrically, elements of S can be represented in the complex plane \mathbb{C} .
- 2 The elements of $f(z)$ is presented in a different complex plane i.e., on the w -plane here.
- 3 The set S is called the **domain of definition** of the function f or simply domain of f .
- 4 The collection of all values of w is called the **range of f** .
- 5 Mathematically, range denoted by $R(f) = \{f(z) : z \in S\}$.
- 6 **Domain of definition** and **domain (set)** are different terminology. "Domain of definition" could be a region as well.
- 7 Often we say "domain of a function". Note that "domain of a function" is not necessarily a "domain". Avoid confusion.

Complex function

- 1 A complex function can be thought of a pair of **real functions** in two-variable (**WHY?**).

Complex function

- 1 A complex function can be thought of a pair of **real functions** in two-variable (**WHY?**).
- 2 Thus, a complex function, **theoretically**, can be written as $w = f(z) = u + iv$, where

$$u = u(x, y), v = v(x, y)$$

Complex function

- 1 A complex function can be thought of a pair of **real functions** in two-variable (**WHY?**).
- 2 Thus, a complex function, **theoretically**, can be written as $w = f(z) = u + iv$, where

$$u = u(x, y), v = v(x, y)$$

- 3 One can also write it in polar form as:

$$u = u(r, \theta), v = v(r, \theta)$$

Complex function

- 1 For some complicated functions, it may be difficult to derive such u and v , but it is easy for most of the functions we are going to discuss.

Complex function

- 1 For some complicated functions, it may be difficult to derive such u and v , but it is easy for most of the functions we are going to discuss.
- 2 For instance, $f(z) = z^2 = (x + iy)^2 = x^2 - y^2 + 2ixy$.

Complex function

- 1 For some complicated functions, it may be difficult to derive such u and v , but it is easy for most of the functions we are going to discuss.
- 2 For instance, $f(z) = z^2 = (x + iy)^2 = x^2 - y^2 + 2ixy$.
- 3 Thus,

$$u(x, y) = x^2 - y^2, v(x, y) = 2ixy$$

Local analysis of functions

- 1 We now study some **local** properties of complex functions.

Local analysis of functions

- 1 We now study some **local** properties of complex functions.
- 2 Local properties **means**?

Local analysis of functions

- 1 We now study some **local** properties of complex functions.
- 2 Local properties **means**?
- 3 We determine the values of the function **at a point and nearby points only**, hence local.....

Local analysis of functions

- 1 We now study some **local** properties of complex functions.
- 2 Local properties **means**?
- 3 We determine the values of the function **at a point and nearby points only**, hence local.....
- 4 We investigate the following **at a point**:
 - 1 **Limit**

Local analysis of functions

- 1 We now study some **local** properties of complex functions.
- 2 Local properties **means**?
- 3 We determine the values of the function **at a point and nearby points only**, hence local.....
- 4 We investigate the following **at a point**:
 - 1 **Limit**
 - 2 **Continuity**

Local analysis of functions

- 1 We now study some **local** properties of complex functions.
- 2 Local properties **means**?
- 3 We determine the values of the function **at a point and nearby points only**, hence local.....
- 4 We investigate the following **at a point**:
 - 1 **Limit**
 - 2 **Continuity**
 - 3 **Differentiability**

Limit

Limit

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.

Limit

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.
- 2 Consider $D \subset \mathbb{C}$ is a **domain set** as of now, and let $z_0 \in \mathbb{C}$ be a **limit point** of D .

Limit

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.
- 2 Consider $D \subset \mathbb{C}$ is a **domain set** as of now, and let $z_0 \in \mathbb{C}$ be a **limit point** of D .
- 3 Define a function $f : D \rightarrow \mathbb{C}$.

Limit

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.
- 2 Consider $D \subset \mathbb{C}$ is a **domain set** as of now, and let $z_0 \in \mathbb{C}$ be a **limit point** of D .
- 3 Define a function $f : D \rightarrow \mathbb{C}$.
- 4 The function is said to have the limit $l \in \mathbb{C}$ as z approaches z_0 , if for every $\epsilon > 0$ (sufficiently small), there exists a $\delta > 0$ (depending on ϵ and z_0) such that

$$|f(z) - l| < \epsilon \text{ whenever } 0 < |z - z_0| < \delta \text{ with } z \in D.$$

Limit

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.
- 2 Consider $D \subset \mathbb{C}$ is a **domain set** as of now, and let $z_0 \in \mathbb{C}$ be a **limit point** of D .
- 3 Define a function $f : D \rightarrow \mathbb{C}$.
- 4 The function is said to have the limit $l \in \mathbb{C}$ as z approaches z_0 , if for every $\epsilon > 0$ (sufficiently small), there exists a $\delta > 0$ (depending on ϵ and z_0) such that

$$|f(z) - l| < \epsilon \text{ whenever } 0 < |z - z_0| < \delta \text{ with } z \in D.$$

- 5 NOTE: For finding limit, it is **NOT** required to define the function at z_0 .

Computing limit

Determine $\lim_{z \rightarrow 0} \frac{z}{\bar{z}}$, if it exists.

- 1 The limit l , if exists , must be unique. The value of l is independent of the direction along which $z \rightarrow z_0$.

Computing limit

Determine $\lim_{z \rightarrow 0} \frac{z}{\bar{z}}$, if it exists.

- 1 The limit l , if exists , must be unique. The value of l is independent of the direction along which $z \rightarrow z_0$.
- 2 If we move from the **positive of x-axis** towards the origin, we obtain $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x \rightarrow 0, y=0} \frac{x+iy}{x-iy} = 1$.

Computing limit

Determine $\lim_{z \rightarrow 0} \frac{z}{\bar{z}}$, if it exists.

- 1 The limit l , if exists, must be unique. The value of l is independent of the direction along which $z \rightarrow z_0$.
- 2 If we move from the **positive of x-axis** towards the origin, we obtain $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x \rightarrow 0, y=0} \frac{x+iy}{x-iy} = 1$.
- 3 If we move from the **positive of y-axis** towards the origin, we obtain $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x=0, y \rightarrow 0} \frac{x+iy}{x-iy} = -1$.

Computing limit

Determine $\lim_{z \rightarrow 0} \frac{z}{\bar{z}}$, if it exists.

- 1 The limit l , if exists, must be unique. The value of l is independent of the direction along which $z \rightarrow z_0$.
- 2 If we move from the **positive of x-axis** towards the origin, we obtain $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x \rightarrow 0, y=0} \frac{x+iy}{x-iy} = 1$.
- 3 If we move from the **positive of y-axis** towards the origin, we obtain $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x=0, y \rightarrow 0} \frac{x+iy}{x-iy} = -1$.
- 4 If we move **along $y = x$ line** towards the origin in the first quadrant, $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x \rightarrow 0} \frac{x+ix}{x-ix}$, which is **a complex number**.

Computing limit

Determine $\lim_{z \rightarrow 0} \frac{z}{\bar{z}}$, if it exists.

- 1 The limit l , if exists, must be unique. The value of l is independent of the direction along which $z \rightarrow z_0$.
- 2 If we move from the **positive of x-axis** towards the origin, we obtain $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x \rightarrow 0, y=0} \frac{x+iy}{x-iy} = 1$.
- 3 If we move from the **positive of y-axis** towards the origin, we obtain $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x=0, y \rightarrow 0} \frac{x+iy}{x-iy} = -1$.
- 4 If we move **along $y = x$ line** towards the origin in the first quadrant, $\lim_{z \rightarrow 0} \frac{z}{\bar{z}} = \lim_{x \rightarrow 0} \frac{x+ix}{x-ix}$, which is **a complex number**.
- 5 Thus, **limit does not exist**.

Computing limit

1 Here we were **lucky**. We found distinct values of f .

Computing limit

- 1 Here we were **lucky**. We found distinct values of f .
- 2 But what happens for other situations?

Computing limit

- 1 Here we were **lucky**. We found distinct values of f .
- 2 But what happens for other situations?
- 3 We have to use ϵ - δ definition.

Computing limit

- 1 Here we were **lucky**. We found distinct values of f .
- 2 But what happens for other situations?
- 3 We have to use ϵ - δ definition.
- 4 How do we apply the definition if **the existence of the limit is not ensured**?

Computing limit

- 1 Here we were **lucky**. We found distinct values of f .
- 2 But what happens for other situations?
- 3 We have to use ϵ - δ definition.
- 4 How do we apply the definition if **the existence of the limit is not ensured**? No answer.

Computing limit

- 1 Here we were **lucky**. We found distinct values of f .
- 2 But what happens for other situations?
- 3 We have to use ϵ - δ definition.
- 4 How do we apply the definition if **the existence of the limit is not ensured**? No answer.
- 5 Ok, compute a limit along a specific path. Then use the definition. If you are successful, then fine.

Use $\epsilon - \delta$ definition

Prove that $\lim_{z \rightarrow i} 2\left(\frac{z^2 + iz + 2}{z - i}\right) = 6i$.

Use $\epsilon - \delta$ definition

Show that if the function $f(z) = i\bar{z}/2$ in the open disk $|z| < 1$, then

$$\lim_{z \rightarrow 1} f(z) = i/2.$$

Note: We did not define the function for all z in a neighborhood of $z = 1$. Hence we should consider those z which belong to both the domain $|z| < 1$ and $\hat{N}(1, \delta)$.

Finding limit using the concept of multivariable calculus

- 1 Perhaps, you are **comfortable** to find limit of **real functions** in **two-variable**.

Finding limit using the concept of multivariable calculus

- 1 Perhaps, you are **comfortable** to find limit of **real functions** in **two-variable**.
- 2 Ok, good news. We provide such **a facility** to find limit of complex functions.

Finding limit using the concept of multivariable calculus

- 1 Perhaps, you are **comfortable** to find limit of **real functions** in **two-variable**.
- 2 Ok, good news. We provide such **a facility** to find limit of complex functions.
- 3 Follow the next theorem.

Theorem

Theorem

Theorem: Let $f(z) = u(x, y) + iv(x, y)$. Then limit of $f(z)$ exists at $z_0 = x_0 + iy_0$ if and only if limits of $u(x, y)$ and $v(x, y)$ exist at (x_0, y_0) .

Theorem

Theorem: Let $f(z) = u(x, y) + iv(x, y)$. Then limit of $f(z)$ exists at $z_0 = x_0 + iy_0$ if and only if limits of $u(x, y)$ and $v(x, y)$ exist at (x_0, y_0) .

Further,

$$\lim_{z \rightarrow z_0} f(z) = \lim_{(x,y) \rightarrow (x_0,y_0)} u(x, y) + i \lim_{(x,y) \rightarrow (x_0,y_0)} v(x, y)$$

Theorem

Theorem: Let $f(z) = u(x, y) + iv(x, y)$. Then limit of $f(z)$ exists at $z_0 = x_0 + iy_0$ if and only if limits of $u(x, y)$ and $v(x, y)$ exist at (x_0, y_0) .

Further,

$$\lim_{z \rightarrow z_0} f(z) = \lim_{(x,y) \rightarrow (x_0,y_0)} u(x, y) + i \lim_{(x,y) \rightarrow (x_0,y_0)} v(x, y)$$

It is now much easier to verify the existence of limit and computing it.

Proof

Proof

Proof

Continuity

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.

Continuity

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.
- 2 Unlike the case of limit, here, we assume $z_0 \in D \subset \mathbb{C}$.

Continuity

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.
- 2 Unlike the case of limit, here, we assume $z_0 \in D \subset \mathbb{C}$.
- 3 Define a functions $f : D \rightarrow \mathbb{C}$.

Continuity

- 1 Recall the usual **definition** you have studied for real function in single or two-variable.
- 2 Unlike the case of limit, here, we assume $z_0 \in D \subset \mathbb{C}$.
- 3 Define a functions $f : D \rightarrow \mathbb{C}$.
- 4 Concept of **deleted neighborhood will not work**. **WHY?**
- 5 The function is said to be continuous at z_0 if for every $\epsilon > 0$ (sufficiently small), there exists a $\delta > 0$ (depends on ϵ and the point z_0) such that

$$|f(z) - f(z_0)| < \epsilon \text{ whenever } |z - z_0| < \delta \text{ with } z \in D.$$

Remember

- 1 If the function is continuous at a point, limit of the function must exist at that point.

Remember

- 1 If the function is continuous at a point, limit of the function must exist at that point.
- 2 A complex function is said to be continuous in a **region** R if it is continuous at each point in R

Theorem

Theorem

Theorem: Let $f(z) = u(x, y) + iv(x, y)$. Then $f(z)$ is continuous at $z_0 = x_0 + iy_0$ if and only if $u(x, y)$ and $v(x, y)$ both are continuous at (x_0, y_0) .

Differentiability

- 1 Let $f : D \rightarrow \mathbb{C}$ be a function, and $z_0 \in D$ be a limit point of D .

Differentiability

- 1 Let $f : D \rightarrow \mathbb{C}$ be a function, and $z_0 \in D$ be a limit point of D .
- 2 Suppose that the limit

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0},$$

Differentiability

- 1 Let $f : D \rightarrow \mathbb{C}$ be a function, and $z_0 \in D$ be a limit point of D .
- 2 Suppose that the limit

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0},$$

- 3 or equivalently

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h},$$

exists. Note that h is a complex number.

Differentiability

- 1 Let $f : D \rightarrow \mathbb{C}$ be a function, and $z_0 \in D$ be a limit point of D .
- 2 Suppose that the limit

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0},$$

- 3 or equivalently

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h},$$

exists. Note that h is a complex number.

- 4 Then we say that f is **differentiable** at z_0 and the limit denoted by $f'(z_0)$ is called the **derivative** of f at z_0 .

Differentiability

Examine if the function $f(z) = |z|$ is differentiable at $z = 0$.

- 1 The following limit must exist for differentiability

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0},$$

Differentiability

Examine if the function $f(z) = |z|$ is differentiable at $z = 0$.

- 1 The following limit must exist for differentiability

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0},$$

- 2 Now

$$\lim_{x \rightarrow 0, y=0} \frac{|x| - 0}{x} = +1 (\text{when } x > 0) \text{ or } -1 (\text{when } x < 0)$$

Differentiability

Examine if the function $f(z) = |z|$ is differentiable at $z = 0$.

- 1 The following limit must exist for differentiability

$$\lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0},$$

- 2 Now

$$\lim_{x \rightarrow 0, y=0} \frac{|x| - 0}{x} = +1 (\text{when } x > 0) \text{ or } -1 (\text{when } x < 0)$$

- 3 The function is not differentiable at $z = 0$.

Differentiability

Determine the points at which the function $f(z) = |z|^2 = x^2 + y^2$ is differentiable.

- 1 The complex function is differentiable at $z = 0$.

Differentiability

Determine the points at which the function $f(z) = |z|^2 = x^2 + y^2$ is differentiable.

- 1 The complex function is differentiable at $z = 0$.
- 2 The complex function is NOT differentiable at any nonzero point (Verify).

Differentiability

Determine the points at which the function $f(z) = |z|^2 = x^2 + y^2$ is differentiable.

- 1 The complex function is differentiable at $z = 0$.
- 2 The complex function is NOT differentiable at any nonzero point (Verify).
- 3 OK, we can think the **above** function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ as a **real valued** function of two-variable.

Differentiability

Determine the points at which the function $f(z) = |z|^2 = x^2 + y^2$ is differentiable.

- 1 The complex function is differentiable at $z = 0$.
- 2 The complex function is NOT differentiable at any nonzero point (Verify).
- 3 OK, we can think the **above** function $f : \mathbb{R}^2 \rightarrow \mathbb{R}$ as a **real valued** function of two-variable.
- 4 Then the real valued function is differentiable (**Why?**) everywhere on \mathbb{R}^2 .

Complex Analysis

LECTURE 5

August 5, 2023

Condition for differentiability

Our questions:

Condition for differentiability

Our questions:

- 1 What are the **necessary conditions** for the differentiability of a complex function.

Condition for differentiability

Our questions:

- 1 What are the **necessary conditions** for the differentiability of a complex function.
- 2 If a complex function is differentiable, how to compute the derivative **without** using the limit definition?

Condition for differentiability

Our questions:

- 1 What are the **necessary conditions** for the differentiability of a complex function.
- 2 If a complex function is differentiable, how to compute the derivative **without** using the limit definition?
- 3 Is there any **sufficient condition** for the differentiability of a complex function?

Necessary condition for differentiability

Cauchy-Riemann equations (C-R equations):

Let $f(z) = u(x, y) + iv(x, y)$ be differentiable at $z_0 = x_0 + iy_0$.

Then the partial derivatives of u and v exist at the point (x_0, y_0) ,

Necessary condition for differentiability

Cauchy-Riemann equations (C-R equations):

Let $f(z) = u(x, y) + iv(x, y)$ be differentiable at $z_0 = x_0 + iy_0$.

Then the partial derivatives of u and v exist at the point (x_0, y_0) , and

$$u_x(x_0, y_0) = v_y(x_0, y_0) \quad \text{and} \quad v_x(x_0, y_0) = -u_y(x_0, y_0). \quad (1)$$

Necessary condition for differentiability

Cauchy-Riemann equations (C-R equations):

Let $f(z) = u(x, y) + iv(x, y)$ be differentiable at $z_0 = x_0 + iy_0$.

Then the partial derivatives of u and v exist at the point (x_0, y_0) , and

$$u_x(x_0, y_0) = v_y(x_0, y_0) \quad \text{and} \quad v_x(x_0, y_0) = -u_y(x_0, y_0). \quad (1)$$

Further,

$$f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0) = v_y(x_0, y_0) - iu_y(x_0, y_0). \quad (2)$$

C-R equations

C-R equations

1 The Equations

$$u_x(x_0, y_0) = v_y(x_0, y_0) \text{ and } v_x(x_0, y_0) = -u_y(x_0, y_0).$$

are called *Cauchy-Riemann equations* (C-R equations).

C-R equations

1 The Equations

$$u_x(x_0, y_0) = v_y(x_0, y_0) \text{ and } v_x(x_0, y_0) = -u_y(x_0, y_0).$$

are called *Cauchy-Riemann equations* (C-R equations).

2 The Equations in

$$f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0) = v_y(x_0, y_0) - iu_y(x_0, y_0).$$

represents the technique to find derivative **without** using the limit definition?

3 Note that the violating of the C-R equations implies the non-existence of derivative at the point.

Derive CR equations.

Derive CR equations.

Derive CR equations.

Applications of CR-Equations

Applications of CR-Equations

- 1 Apply the C-R equations to determine the points where $f(z) = \bar{z}$ are not differentiable.

Applications of CR-Equations

- 1 Apply the C-R equations to determine the points where $f(z) = \bar{z}$ are not differentiable.
- 2 Note that $u = x, v = -y$

Applications of CR-Equations

- 1 Apply the C-R equations to determine the points where $f(z) = \bar{z}$ are not differentiable.
- 2 Note that $u = x, v = -y$
- 3 Note that $u_x = 1, u_y = 0, v_x = 0, v_y = -1$

Applications of CR-Equations

- 1 Apply the C-R equations to determine the points where $f(z) = \bar{z}$ are not differentiable.
- 2 Note that $u = x, v = -y$
- 3 Note that $u_x = 1, u_y = 0, v_x = 0, v_y = -1$
- 4 C-R equations are not satisfied anywhere.

Applications of CR-Equations

- 1 Apply the C-R equations to determine the points where $f(z) = \bar{z}$ are not differentiable.
- 2 Note that $u = x, v = -y$
- 3 Note that $u_x = 1, u_y = 0, v_x = 0, v_y = -1$
- 4 C-R equations are not satisfied anywhere.
- 5 The function is not differentiable anywhere.

Applications of CR-Equations

Applications of CR-Equations

- 1 Apply the C-R equations to determine the points where $f(z) = |z|$ are not differentiable.

Applications of CR-Equations

- 1 Apply the C-R equations to **determine the points** where $f(z) = |z|$ are not differentiable.
- 2 We already know that the function is **not differentiable** at origin.

Applications of CR-Equations

- 1 Apply the C-R equations to **determine the points** where $f(z) = |z|$ are not differentiable.
- 2 We already know that the function is **not differentiable** at origin.
- 3 Note that $u = \sqrt{x^2 + y^2}$, $v = 0$

Applications of CR-Equations

- 1 Apply the C-R equations to **determine the points** where $f(z) = |z|$ are not differentiable.
- 2 We already know that the function is **not differentiable** at origin.
- 3 Note that $u = \sqrt{x^2 + y^2}$, $v = 0$
- 4 For nonzero point,
$$u_x = \frac{x}{\sqrt{x^2 + y^2}}, u_y = \frac{y}{\sqrt{x^2 + y^2}}, v_x = 0, v_y = 0.$$

Applications of CR-Equations

- 1 Apply the C-R equations to **determine the points** where $f(z) = |z|$ are not differentiable.
- 2 We already know that the function is **not differentiable** at origin.
- 3 Note that $u = \sqrt{x^2 + y^2}$, $v = 0$
- 4 For nonzero point,
$$u_x = \frac{x}{\sqrt{x^2 + y^2}}, u_y = \frac{y}{\sqrt{x^2 + y^2}}, v_x = 0, v_y = 0.$$
- 5 C-R equations are **not satisfied** for any nonzero z also.

Applications of CR-Equations

- 1 Apply the C-R equations to **determine the points** where $f(z) = |z|$ are not differentiable.
- 2 We already know that the function is **not differentiable** at origin.
- 3 Note that $u = \sqrt{x^2 + y^2}$, $v = 0$
- 4 For nonzero point,
$$u_x = \frac{x}{\sqrt{x^2 + y^2}}, u_y = \frac{y}{\sqrt{x^2 + y^2}}, v_x = 0, v_y = 0.$$
- 5 C-R equations are **not satisfied** for any nonzero z also.
- 6 The function is not differentiable anywhere.

Applications of CR-Equations

Applications of CR-Equations

- 1 Consider the functions $f(z) = |z|^2$.

Applications of CR-Equations

- 1 Consider the functions $f(z) = |z|^2$.
- 2 Note that $u = x^2 + y^2, v = 0$

Applications of CR-Equations

- 1 Consider the functions $f(z) = |z|^2$.
- 2 Note that $u = x^2 + y^2, v = 0$
- 3 Note that $u_x = 2x, u_y = 2y, v_x = 0, v_y = 0$.

Applications of CR-Equations

- 1 Consider the functions $f(z) = |z|^2$.
- 2 Note that $u = x^2 + y^2, v = 0$
- 3 Note that $u_x = 2x, u_y = 2y, v_x = 0, v_y = 0$.
- 4 Satisfy the C-R equations at origin only.

Applications of CR-Equations

- 1 Consider the functions $f(z) = |z|^2$.
- 2 Note that $u = x^2 + y^2, v = 0$
- 3 Note that $u_x = 2x, u_y = 2y, v_x = 0, v_y = 0$.
- 4 Satisfy the C-R equations at origin only.
- 5 In fact, $f'(0)$ exists (Verify).

Applications of CR-Equations

- 1 Consider the functions $f(z) = |z|^2$.
- 2 Note that $u = x^2 + y^2, v = 0$
- 3 Note that $u_x = 2x, u_y = 2y, v_x = 0, v_y = 0$.
- 4 Satisfy the C-R equations at **origin only**.
- 5 In fact, $f'(0)$ exists (Verify).
- 6 Now can we **claim** that the differentiability of the function is **due the satisfaction** of C-R equations?

CR-Equations are not sufficient for differentiability

CR-Equations are not sufficient for differentiability

- 1 Consider the function $f(z)$

$$f(z) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } z \neq 0 \\ 0, & \text{if } z = 0. \end{cases}$$

CR-Equations are not sufficient for differentiability

- 1 Consider the function $f(z)$

$$f(z) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } z \neq 0 \\ 0, & \text{if } z = 0. \end{cases}$$

- 2 Note that

$$u = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } (x, y) \neq 0 \\ 0, & \text{if } (x, y) = 0. \end{cases}$$

CR-Equations are not sufficient for differentiability

- 1 Consider the function $f(z)$

$$f(z) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } z \neq 0 \\ 0, & \text{if } z = 0. \end{cases}$$

- 2 Note that

$$u = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } (x, y) \neq 0 \\ 0, & \text{if } (x, y) = 0. \end{cases}$$

- 3 $v = 0$

CR-Equations are not sufficient for differentiability

- 1 Consider the function $f(z)$

$$f(z) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } z \neq 0 \\ 0, & \text{if } z = 0. \end{cases}$$

- 2 Note that

$$u = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } (x, y) \neq 0 \\ 0, & \text{if } (x, y) = 0. \end{cases}$$

- 3 $v = 0$

$$4 u_x(0, 0) = \lim_{h \rightarrow 0} \frac{u(h, 0) - u(0, 0)}{h} = \lim_{h \rightarrow 0} \frac{0 - 0}{h} = 0.$$

CR-Equations are not sufficient for differentiability

1 Similarly, $u_y = 0$

CR-Equations are not sufficient for differentiability

- 1 Similarly, $u_y = 0$
- 2 Thus, $u_x = u_y = v_x = 0, v_y = 0$.

CR-Equations are not sufficient for differentiability

- 1 Similarly, $u_y = 0$
- 2 Thus, $u_x = u_y = v_x = 0, v_y = 0$.
- 3 The C-R equations are satisfied at the origin.

CR-Equations are not sufficient for differentiability

- 1 Similarly, $u_y = 0$
- 2 Thus, $u_x = u_y = v_x = 0, v_y = 0$.
- 3 The C-R equations are satisfied at the origin.
- 4 Now we check differentiability of $f(z)$ at the origin.

CR-Equations are not sufficient for differentiability

- 1 Similarly, $u_y = 0$
- 2 Thus, $u_x = u_y = v_x = 0, v_y = 0$.
- 3 The C-R equations are satisfied at the origin.
- 4 Now we check differentiability of $f(z)$ at the origin.

5 In fact, $f'(0) = \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \frac{\frac{xy}{x^2 + y^2} - 0}{x + iy}$

CR-Equations are not sufficient for differentiability

- 1 Similarly, $u_y = 0$
- 2 Thus, $u_x = u_y = v_x = 0, v_y = 0$.
- 3 The C-R equations are satisfied at the origin.
- 4 Now we check differentiability of $f(z)$ at the origin.

$$5 \text{ In fact, } f'(0) = \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \frac{\frac{xy}{x^2 + y^2} - 0}{x + iy}$$

6 If $y = x$ and $x \rightarrow 0^+$, then $f'(0) \rightarrow \infty$

CR-Equations are not sufficient for differentiability

- 1 Similarly, $u_y = 0$
- 2 Thus, $u_x = u_y = v_x = 0, v_y = 0$.
- 3 The C-R equations are satisfied at the origin.
- 4 Now we check differentiability of $f(z)$ at the origin.

5 In fact, $f'(0) = \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \frac{\frac{xy}{x^2 + y^2} - 0}{x + iy}$

6 If $y = x$ and $x \rightarrow 0^+$, then $f'(0) \rightarrow \infty$

CR-Equations are not sufficient for differentiability

- 1 Similarly, $u_y = 0$
- 2 Thus, $u_x = u_y = v_x = 0, v_y = 0$.
- 3 The C-R equations are satisfied at the origin.
- 4 Now we check differentiability of $f(z)$ at the origin.

- 5 In fact, $f'(0) = \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \frac{\frac{xy}{x^2 + y^2}}{x + iy} - 0$
- 6 If $y = x$ and $x \rightarrow 0^+$, then $f'(0) \rightarrow \infty$ (complex infinity)
—derivative does not exist.

CR-Equations are not sufficient for differentiability

- 1 Similarly, $u_y = 0$
- 2 Thus, $u_x = u_y = v_x = 0, v_y = 0$.
- 3 The C-R equations are satisfied at the origin.
- 4 Now we check differentiability of $f(z)$ at the origin.

- 5 In fact, $f'(0) = \lim_{z \rightarrow 0} \frac{f(z) - f(0)}{z - 0} = \lim_{z \rightarrow 0} \frac{\frac{xy}{x^2 + y^2}}{x + iy} - 0$
- 6 If $y = x$ and $x \rightarrow 0^+$, then $f'(0) \rightarrow \infty$ (complex infinity)
—derivative does not exist.
- 7 Thus, we need alternative conditions for differentiability

Sufficient conditions for differentiability

Theorem:

Sufficient conditions for differentiability

Theorem: Suppose $f = u + iv$ is defined on some neighborhood

$N(z_0; \epsilon)$ of $z_0 = x_0 + iy_0$ such that u_x, u_y, v_x, v_y exist on $N(z_0; \epsilon)$ and are continuous at (x_0, y_0) . If u, v satisfies the C-R equations at (x_0, y_0) , then f' exists at z_0 and $f'(z_0) = u_x(x_0, y_0) + iv_x(x_0, y_0)$.

Application of Sufficient conditions

- 1 Consider $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$.

Application of Sufficient conditions

1 Consider $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$.

2

$$u_x(x, y) = -e^{-y} \sin x \quad u_y(x, y) = -e^{-y} \cos x$$

$$v_y(x, y) = -e^{-y} \sin x \quad v_x(x, y) = e^{-y} \cos x.$$

Application of Sufficient conditions

1 Consider $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$.

2

$$u_x(x, y) = -e^{-y} \sin x \quad u_y(x, y) = -e^{-y} \cos x$$

$$v_y(x, y) = -e^{-y} \sin x \quad v_x(x, y) = e^{-y} \cos x.$$

3 Therefore, u_x, u_y, v_x, v_y are all continuous on \mathbb{R}^2 , and u, v satisfies the C-R equations $u_x(x, y) = v_y(x, y)$,
 $u_y(x, y) = -v_x(x, y)$ at each point $(x, y) \in \mathbb{R}^2$.

Application of Sufficient conditions

- 1 Consider $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$.

2

$$\begin{aligned} u_x(x, y) &= -e^{-y} \sin x & u_y(x, y) &= -e^{-y} \cos x \\ v_y(x, y) &= -e^{-y} \sin x & v_x(x, y) &= e^{-y} \cos x. \end{aligned}$$

- 3 Therefore, u_x, u_y, v_x, v_y are all continuous on \mathbb{R}^2 , and u, v satisfies the C-R equations $u_x(x, y) = v_y(x, y)$, $u_y(x, y) = -v_x(x, y)$ at each point $(x, y) \in \mathbb{R}^2$.
- 4 The C-R equations **are satisfied** as well.

Application of Sufficient conditions

- 1 Consider $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$.

2

$$\begin{aligned} u_x(x, y) &= -e^{-y} \sin x & u_y(x, y) &= -e^{-y} \cos x \\ v_y(x, y) &= -e^{-y} \sin x & v_x(x, y) &= e^{-y} \cos x. \end{aligned}$$

- 3 Therefore, u_x, u_y, v_x, v_y are all continuous on \mathbb{R}^2 , and u, v satisfies the C-R equations $u_x(x, y) = v_y(x, y)$, $u_y(x, y) = -v_x(x, y)$ at each point $(x, y) \in \mathbb{R}^2$.
- 4 The C-R equations are satisfied as well.
- 5 Hence $f'(z) = u_x + iv_x = -e^{-y} \sin x + ie^{-y} \cos x$

Your comment

Your comment

- 1 Consider the function $f(z)$

$$f(z) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } z \neq 0 \\ 0, & \text{if } z = 0. \end{cases}$$

Your comment

- 1 Consider the function $f(z)$

$$f(z) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } z \neq 0 \\ 0, & \text{if } z = 0. \end{cases}$$

- 2 Check whether the sufficient conditions are satisfied at and in the neighborhood of origin.

Your comment

- 1 Consider the function $f(z)$

$$f(z) = \begin{cases} \frac{xy}{x^2 + y^2} & \text{if } z \neq 0 \\ 0, & \text{if } z = 0. \end{cases}$$

- 2 Check whether the sufficient conditions are satisfied **at and in the neighborhood** of origin.
- 3 What is your conclusion on the existence of $f'(z)$ at origin?

Complex Analysis

LECTURE 6

August 7, 2023

Your comment

Your comment

- 1 Ok, you may consider a relatively simple function $f(z) = |z|$

Your comment

- 1 Ok, you may consider a **relatively simple** function $f(z) = |z|$
- 2 Check whether the sufficient conditions are satisfied at and in the neighborhood of origin.

Your comment

- 1 Ok, you may consider a **relatively simple** function $f(z) = |z|$
- 2 Check whether the sufficient conditions are satisfied at and in the neighborhood of origin.
- 3 What is **your conclusion** on the existence of $f'(z)$ at the origin?

Our observations

Our observations

- 1 The functions $f(z) = |z|^2$ is differentiable **at the origin only**.
- 2 The function $f(z) = |z|^2$ is **NOT** differentiable in any **neighborhood** of origin.

Our observations

- 1 The functions $f(z) = |z|^2$ is differentiable at the origin only.
- 2 The function $f(z) = |z|^2$ is NOT differentiable in any neighborhood of origin.
- 3 The function $f(x + iy) = x^2 + iy^2$ is differentiable at all points on $y = x$ (HOW?).

Our observations

- 1 The functions $f(z) = |z|^2$ is differentiable **at the origin only**.
- 2 The function $f(z) = |z|^2$ is **NOT** differentiable in any **neighborhood** of origin.
- 3 The function $f(x + iy) = x^2 + iy^2$ is differentiable at all points on $y = x$ (**HOW?**).
- 4 But the above functions **too** is **NOT** differentiable in any **neighborhood** of any point on $y = x$.

Our observations

Our observations

- 1 OK, we **SEARCH** functions which are differentiable in some neighborhood of a point.

Our observations

- 1 OK, we **SEARCH** functions which are differentiable in some **neighborhood** of a point.
- 2 Consider the functions $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$ and $f(z) = z^2$ are differentiable at all points on \mathbb{C} .

Our observations

- 1 OK, we **SEARCH** functions which are differentiable in some **neighborhood** of a point.
- 2 Consider the functions $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$ and $f(z) = z^2$ are differentiable at all points on \mathbb{C} .
- 3 Therefore, the above functions **are** differentiable in any **neighborhood** of any point.

Our observations

- 1 OK, we **SEARCH** functions which are differentiable in some **neighborhood** of a point.
- 2 Consider the functions $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$ and $f(z) = z^2$ are differentiable at all points on \mathbb{C} .
- 3 Therefore, the above functions **are** differentiable in any **neighborhood** of any point.
- 4 Thus we obtain a spacial class of functions.

Our observations

- 1 OK, we **SEARCH** functions which are differentiable in some **neighborhood** of a point.
- 2 Consider the functions $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$ and $f(z) = z^2$ are differentiable at all points on \mathbb{C} .
- 3 Therefore, the above functions **are** differentiable in any **neighborhood** of any point.
- 4 Thus we obtain a spacial class of functions.
- 5 **Hence we present the concept** of analytic functions.

Analytic functions

Analytic functions

- 1 We first provide the definition of analytic functions **at a point**.

Analytic functions

- 1 We first provide the definition of analytic functions **at a point**.
- 2 Then we define analytic functions on a **set**.

Analytic functions

Analytic functions

- 1 **Analytic function at a point:** A function f is said to be analytic at a point z_0 if there exists a neighborhood $N(z_0, \epsilon)$ of z_0 , $\epsilon > 0$ (sufficiently small) such that f is differentiable at every point $z \in N(z_0, \epsilon)$.

Analytic functions

- 1 **Analytic function at a point:** A function f is said to be analytic at a point z_0 if there exists a neighborhood $N(z_0, \epsilon)$ of z_0 , $\epsilon > 0$ (sufficiently small) such that f is differentiable at every point $z \in N(z_0, \epsilon)$.
- 2 **Analytic function on a set:** A function f is said to be analytic (or **regular** or **holomorphic**) on a set D if it is differentiable at every point of **some open set containing D** .

Some interesting results

Some interesting results

- 1 If f is differentiable at all points of an **open set** D , then f is analytic on D .

Some interesting results

- 1 If f is differentiable at all points of an **open set** D , then f is analytic on D .
- 2 If f is **differentiable** at all points of a set D , then it **does not** mean that f will be analytic on D

Some interesting results

- 1 If f is differentiable at all points of an **open set** D , then f is analytic on D .
- 2 If f is **differentiable** at all points of a set D , then it **does not** mean that f will be analytic on D . Check the fact with $f(x + iy) = x^2 + iy^2$.

Some interesting results

- 1 If f is differentiable at all points of an **open set** D , then f is analytic on D .
- 2 If f is **differentiable** at all points of a set D , then it **does not** mean that f will be analytic on D . Check the fact with $f(x + iy) = x^2 + iy^2$.
- 3 If f is **analytic** at all points of a set D , then f is analytic on D .

Necessary and Sufficient Condition for Analyticity

Necessary and Sufficient Condition for Analyticity

Necessary and Sufficient Condition for Analyticity

A function $f = u + iv$ is analytic in a **domain** D if and only if u, v satisfies C-R equations in D , and u_x, u_y, v_x, v_y are continuous in D .

Problems

Problems

Apply the theorem to the function on \mathbb{C}

1 $f(z) = z$

Problems

Apply the theorem to the function on \mathbb{C}

1 $f(z) = z$

2 $f(x + iy) = x^2 + iy^2$

Problems

Apply the theorem to the function on \mathbb{C}

1 $f(z) = z$

2 $f(x + iy) = x^2 + iy^2$

- 1 Check each of the conditions for the above function.

Problems

Apply the theorem to the function on \mathbb{C}

1 $f(z) = z$

2 $f(x + iy) = x^2 + iy^2$

- 1 Check each of the conditions for the above function.
- 2 What is your conclusion about the analyticity of the function?

Entire functions

Entire functions

Entire function: A function analytic on the entire complex plane is called an entire function.

Entire functions

Entire function: A function analytic on the entire complex plane is called an entire function.

Example

Entire functions

Entire function: A function analytic on the entire complex plane is called an entire function.

Example

- 1 Any polynomial

Entire functions

Entire function: A function analytic on the entire complex plane is called an entire function.

Example

- 1 Any polynomial
- 2 $f(x + iy) = e^{-y} \cos x + ie^{-y} \sin x$

Problems

Problem: Suppose f is analytic in a domain D . If $f'(z) = 0$ for all $z \in D$, then f is constant on D .

Contd...

Problems

Problem: Suppose f is analytic in a domain D . If any of $\operatorname{Re} f$, $\operatorname{Im} f$ is constant in D , then f is constant in D .

Problems

Problem: Suppose f is analytic in a domain D such that $|f|$ is constant in D . Then show that f is a constant in D .

Problems

Problem: Let $f = u + iv$ be a non-constant function such that $\bar{f} = u - iv$ be analytic in a domain D . Show that f cannot be analytic in D .

Complex Analysis

LECTURE 7

August 9, 2023

Problems

Problem: Given an **analytic function**

$$w = f(z) = u(x, y) + iv(x, y), z = x + iy,$$

the equations $u(x, y) = \alpha$ and $v(x, y) = \beta$, α and β are constants, define two **families of curves** in the complex plane. Show that the two families are mutually orthogonal to each other.

The families of curves are called **level curves**.

Verify the orthogonality

Verify the orthogonality.

Draw the level curves too.

1 $f(z) = z$

Verify the orthogonality

Verify the orthogonality.

Draw the level curves too.

1 $f(z) = z$

2 $f(z) = z^2$

Verify the orthogonality

Verify the orthogonality.

Draw the level curves too.

1 $f(z) = z$

2 $f(z) = z^2$

3 $f(x + iy) = x^2 + iy^2$ (**Interesting problem**)

Harmonic functions

Harmonic functions

Definition:

A real-valued function $\phi(x, y)$ of two real variables x and y is said to be harmonic in a given **domain** D in the xy -plane if ϕ has **continuous partial derivatives up to the second order** in D

Harmonic functions

Definition:

A real-valued function $\phi(x, y)$ of two real variables x and y is said to be harmonic in a given **domain** D in the xy -plane if ϕ has **continuous partial derivatives up to the second order** in D **AND** satisfies the Laplace equation

$$\phi_{xx}(x, y) + \phi_{yy}(x, y) = 0, (x, y) \in D.$$

Example

Harmonic functions

Definition:

A real-valued function $\phi(x, y)$ of two real variables x and y is said to be harmonic in a given **domain** D in the xy -plane if ϕ has **continuous partial derivatives up to the second order** in D **AND** satisfies the Laplace equation

$$\phi_{xx}(x, y) + \phi_{yy}(x, y) = 0, (x, y) \in D.$$

Example

1 $\phi(x, y) = x$

Harmonic functions

Definition:

A real-valued function $\phi(x, y)$ of two real variables x and y is said to be harmonic in a given **domain** D in the xy -plane if ϕ has **continuous partial derivatives up to the second order** in D **AND** satisfies the Laplace equation

$$\phi_{xx}(x, y) + \phi_{yy}(x, y) = 0, (x, y) \in D.$$

Example

1 $\phi(x, y) = x$

2 $\psi(x, y) = y$

Harmonic functions

Definition:

A real-valued function $\phi(x, y)$ of two real variables x and y is said to be harmonic in a given **domain** D in the xy -plane if ϕ has **continuous partial derivatives up to the second order** in D **AND** satisfies the Laplace equation

$$\phi_{xx}(x, y) + \phi_{yy}(x, y) = 0, (x, y) \in D.$$

Example

- 1 $\phi(x, y) = x$
- 2 $\psi(x, y) = y$
- 3 $\phi(x, y) = x^2 - y^2$

Harmonic functions

Definition:

A real-valued function $\phi(x, y)$ of two real variables x and y is said to be harmonic in a given **domain** D in the xy -plane if ϕ has **continuous partial derivatives up to the second order** in D **AND** satisfies the Laplace equation

$$\phi_{xx}(x, y) + \phi_{yy}(x, y) = 0, (x, y) \in D.$$

Example

- 1 $\phi(x, y) = x$
- 2 $\psi(x, y) = y$
- 3 $\phi(x, y) = x^2 - y^2$
- 4 $\psi(x, y) = 2xy$

Conjugate Harmonic function

Conjugate Harmonic function

Definition:

If two harmonic functions $\phi(x, y), \psi(x, y)$ satisfy C-R equations, namely

$$\phi_x = \psi_y \text{ and } \phi_y = -\psi_x$$

in a domain D , then ψ is called conjugate harmonic function of ϕ .

Important note

Note that harmonic conjugacy is not a symmetric relation, that is, if ψ is conjugate harmonic function of ϕ , then it does not mean that ϕ will be conjugate harmonic function of ψ . This is due to the minus sign in the second Cauchy–Riemann relation.

Examples: Conjugate harmonic function

Examples: Conjugate harmonic function

Example

Examples: Conjugate harmonic function

Example

- 1 Set $\phi(x, y) = x, \psi(x, y) = y$

Examples: Conjugate harmonic function

Example

- 1 Set $\phi(x, y) = x, \psi(x, y) = y$
- 2 ψ is a conjugate Harmonic function of ϕ , but ϕ is NOT a conjugate Harmonic function of ψ .

Examples: Conjugate harmonic function

Example

- 1 Set $\phi(x, y) = x, \psi(x, y) = y$
- 2 ψ is a conjugate Harmonic function of ϕ , but ϕ is NOT a conjugate Harmonic function of ψ .
- 3 Again set $\phi(x, y) = x^2 - y^2, \psi(x, y) = 2xy$

Examples: Conjugate harmonic function

Example

- 1 Set $\phi(x, y) = x, \psi(x, y) = y$
- 2 ψ is a conjugate Harmonic function of ϕ , but ϕ is NOT a conjugate Harmonic function of ψ .
- 3 Again set $\phi(x, y) = x^2 - y^2, \psi(x, y) = 2xy$
- 4 ψ is a conjugate Harmonic function of ϕ , but ϕ is NOT a conjugate Harmonic function of ψ .

Examples: Conjugate harmonic function

Example

- 1 Set $\phi(x, y) = x, \psi(x, y) = y$
- 2 ψ is a conjugate Harmonic function of ϕ , but ϕ is NOT a conjugate Harmonic function of ψ .
- 3 Again set $\phi(x, y) = x^2 - y^2, \psi(x, y) = 2xy$
- 4 ψ is a conjugate Harmonic function of ϕ , but ϕ is NOT a conjugate Harmonic function of ψ .
- 5 Verify the statement by C-R equations.

Important observation

- 1 For $\phi(x, y) = x, \psi(x, y) = y,$
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z$ is **an analytic function.**

Important observation

- 1 For $\phi(x, y) = x, \psi(x, y) = y$,
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z$ is **an analytic function**.
- 2 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).
- 3 Again, for $\phi(x, y) = x^2 - y^2, \psi(x, y) = 2xy$,
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z^2$ is **an analytic function**.

Important observation

- 1 For $\phi(x, y) = x, \psi(x, y) = y$,
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z$ is **an analytic function**.
- 2 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).
- 3 Again, for $\phi(x, y) = x^2 - y^2, \psi(x, y) = 2xy$,
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z^2$ is **an analytic function**.
- 4 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).

Important observation

- 1 For $\phi(x, y) = x, \psi(x, y) = y$,
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z$ is **an analytic function**.
- 2 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).
- 3 Again, for $\phi(x, y) = x^2 - y^2, \psi(x, y) = 2xy$,
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z^2$ is **an analytic function**.
- 4 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).
- 5 Ohhhhh! something is happening.

Important observation

- 1 For $\phi(x, y) = x, \psi(x, y) = y,$
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z$ is **an analytic function**.
- 2 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).
- 3 Again, for $\phi(x, y) = x^2 - y^2, \psi(x, y) = 2xy,$
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z^2$ is **an analytic function**.
- 4 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).
- 5 **Ohhhhh!** something is happening.
- 6 Perhaps, special arrangement of $\phi(x, y)$ and $\psi(x, y)$ is
constructing an analytic function.

Important observation

- 1 For $\phi(x, y) = x, \psi(x, y) = y$,
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z$ is **an analytic function**.
- 2 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).
- 3 Again, for $\phi(x, y) = x^2 - y^2, \psi(x, y) = 2xy$,
 $f(x + iy) = \phi(x, y) + i\psi(x, y) = z^2$ is **an analytic function**.
- 4 But $f(x + iy) = \psi(x, y) + i\phi(x, y)$ is **NOT** (WHY??).
- 5 **Ohhhhh!** something is happening.
- 6 Perhaps, special arrangement of $\phi(x, y)$ and $\psi(x, y)$ is
constructing an analytic function.
- 7 Ok, go to the next theorem.

Next theorem

- 1 Next theorem give a relationship between analyticity and conjugacy.

Next theorem

- 1 Next theorem give a relationship between **analyticity** and **conjugacy**.
- 2 The theorem provides **conditions for analyticity** of a function.

Next theorem

- 1 Next theorem give a relationship between **analyticity** and **conjugacy**.
- 2 The theorem provides **conditions** for analyticity of a function.
- 3 Alternatively, it gives ideas to **construct** an analytic function from a given harmonic function.

Theorem

Theorem

A complex function $f(z) = u(x, y) + iv(x, y)$, $z = x + iy$, is analytic in a domain D if and only if v is a harmonic conjugate of u in D .

Theorem

Theorem

A complex function $f(z) = u(x, y) + iv(x, y)$, $z = x + iy$, is analytic in a domain D if and only if v is a harmonic conjugate of u in D .

Verification

- 1 Verify for $f(x + iy) = e^{-y}(\cos x + i \sin x)$.
- 2 Verify for $f(x + iy) = x^2 + iy^2$.

Uniqueness of harmonic conjugate functions

Result

Any two harmonic conjugates v, w of u in a domain D are **unique in the sense** that they differ by a constant, that is,

$$v(x, y) - w(x, y) = K$$

for all $(x, y) \in D$, where K is a **real** constant.

Uniqueness of harmonic conjugate functions

Result

Any two harmonic conjugates v, w of u in a domain D are **unique in the sense** that they differ by a constant, that is,

$$v(x, y) - w(x, y) = K$$

for all $(x, y) \in D$, where K is a **real** constant.

Proof:

Proof

Construction of harmonic conjugate function

Problem

Find a conjugate harmonic function of $u(x, y) = x^2 - y^2 - y$ in \mathbb{C} .

Construction of harmonic conjugate function

Problem

Find a conjugate harmonic function of $u(x, y) = x^2 - y^2 - y$ in \mathbb{C} .

Solution:

Construction of harmonic conjugate function

Construction of harmonic conjugate function

Answer: $v = 2xy + x + K, K \in \mathbb{R}$

Construction of harmonic conjugate function

Answer: $v = 2xy + x + K, K \in \mathbb{R}$

In addition, we can construct the analytic function

$$f(x + iy) = x^2 - y^2 - y + i(2xy + x + K) = z^2 + i(z + k)$$

Complex Analysis

LECTURE 8

August 10, 2023

Elementary functions

Elementary functions

Elementary functions

Elementary functions

1 Exponential function

Elementary functions

Elementary functions

- 1 Exponential function
- 2 Trigonometric functions

Elementary functions

Elementary functions

- 1 Exponential function
- 2 Trigonometric functions
- 3 Hyperbolic functions

Elementary functions

Elementary functions

- 1 Exponential function
- 2 Trigonometric functions
- 3 Hyperbolic functions
- 4 Logarithmic function

Complex exponential function

Complex exponential function

1 Complex exponential function e^z is defined by

$$e^z = e^x(\cos y + i \sin y).$$

Complex exponential function

- 1 Complex exponential function e^z is defined by

$$e^z = e^x(\cos y + i \sin y).$$

- 2 Euler's formula: When $x = 0$, we obtain

Complex exponential function

- 1 Complex exponential function e^z is defined by

$$e^z = e^x(\cos y + i \sin y).$$

- 2 Euler's formula: When $x = 0$, we obtain

$$e^{iy} = (\cos y + i \sin y).$$

Complex exponential function

- 1 Complex exponential function e^z is defined by

$$e^z = e^x(\cos y + i \sin y).$$

- 2 Euler's formula: When $x = 0$, we obtain

$$e^{iy} = (\cos y + i \sin y).$$

- 3 Euler's identity—"The most beautiful equation": When $y = \pi$ in Euler's formula, we obtain

Complex exponential function

- 1 Complex exponential function e^z is defined by

$$e^z = e^x(\cos y + i \sin y).$$

- 2 Euler's formula: When $x = 0$, we obtain

$$e^{iy} = (\cos y + i \sin y).$$

- 3 Euler's identity—"The most beautiful equation": When $y = \pi$ in Euler's formula, we obtain

$$e^{i\pi} + 1 = 0.$$

Complex exponential function

- 1 Complex exponential function e^z is defined by

$$e^z = e^x(\cos y + i \sin y).$$

- 2 Euler's formula: When $x = 0$, we obtain

$$e^{iy} = (\cos y + i \sin y).$$

- 3 Euler's identity—"The most beautiful equation": When $y = \pi$ in Euler's formula, we obtain

$$e^{i\pi} + 1 = 0.$$

- 4 What would be the form of De-Moivre's theorem in view of Euler's formula?

Properties exponential function

Properties exponential function

- 1 $|e^z| = e^x$, and $\arg e^z = y \pm 2n\pi$, ($n = 0, 1, 2, \dots$).

Properties exponential function

- 1 $|e^z| = e^x$, and $\arg e^z = y \pm 2n\pi$, ($n = 0, 1, 2, \dots$).
- 2 e^z is an entire function, and $\frac{d}{dz} e^z = e^z$.

Properties exponential function

- 1 $|e^z| = e^x$, and $\arg e^z = y \pm 2n\pi$, ($n = 0, 1, 2, \dots$).
- 2 e^z is an entire function, and $\frac{d}{dz} e^z = e^z$.
- 3 $e^{z_1+z_2} = e^{z_1} e^{z_2}$.

Properties exponential function

- 1 $|e^z| = e^x$, and $\arg e^z = y \pm 2n\pi$, ($n = 0, 1, 2, \dots$).
- 2 e^z is an entire function, and $\frac{d}{dz} e^z = e^z$.
- 3 $e^{z_1+z_2} = e^{z_1} e^{z_2}$.
- 4 $e^z \neq 0$ for all z .

Properties exponential function

- 1 $|e^z| = e^x$, and $\arg e^z = y \pm 2n\pi$, ($n = 0, 1, 2, \dots$).
- 2 e^z is an entire function, and $\frac{d}{dz} e^z = e^z$.
- 3 $e^{z_1+z_2} = e^{z_1}e^{z_2}$.
- 4 $e^z \neq 0$ for all z .
- 5 e^z is periodic with the **fundamental period** $2\pi i$ (Ohh! purely imaginary period),

$$\text{i.e., } e^{z+2k\pi i} = e^z, z \in \mathbb{C}, k \in \mathbb{Z}$$

Properties exponential function

- 1 $|e^z| = e^x$, and $\arg e^z = y \pm 2n\pi$, ($n = 0, 1, 2, \dots$).
- 2 e^z is an entire function, and $\frac{d}{dz} e^z = e^z$.
- 3 $e^{z_1+z_2} = e^{z_1}e^{z_2}$.
- 4 $e^z \neq 0$ for all z .
- 5 e^z is periodic with the **fundamental period** $2\pi i$ (Ohh! purely imaginary period),

$$\text{i.e., } e^{z+2k\pi i} = e^z, z \in \mathbb{C}, k \in \mathbb{Z}$$

Properties exponential function

- 1 $|e^z| = e^x$, and $\arg e^z = y \pm 2n\pi$, ($n = 0, 1, 2, \dots$).
- 2 e^z is an entire function, and $\frac{d}{dz} e^z = e^z$.
- 3 $e^{z_1+z_2} = e^{z_1}e^{z_2}$.
- 4 $e^z \neq 0$ for all z .
- 5 e^z is periodic with the **fundamental period** $2\pi i$ (Ohh! purely imaginary period),

$$\text{i.e., } e^{z+2k\pi i} = e^z, z \in \mathbb{C}, k \in \mathbb{Z}$$

- 6 e^z is NOT injective.

Properties exponential function

- 1 $|e^z| = e^x$, and $\arg e^z = y \pm 2n\pi$, ($n = 0, 1, 2, \dots$).
- 2 e^z is an entire function, and $\frac{d}{dz} e^z = e^z$.
- 3 $e^{z_1+z_2} = e^{z_1}e^{z_2}$.
- 4 $e^z \neq 0$ for all z .
- 5 e^z is periodic with the **fundamental period** $2\pi i$ (Oh! purely imaginary period),

$$\text{i.e., } e^{z+2k\pi i} = e^z, z \in \mathbb{C}, k \in \mathbb{Z}$$

- 6 e^z is NOT injective.
- 7 If $H = \{x + iy : -\pi < y \leq \pi\}$. Then e^z is bijective from H to $\mathbb{C} \setminus \{0\}$.

Trigonometric function

Trigonometric function

1 Euler formula yields

$$e^{iy} = \cos y + i \sin y \text{ and } e^{-iy} = \cos y - i \sin y.$$

Trigonometric function

1 Euler formula yields

$$e^{iy} = \cos y + i \sin y \text{ and } e^{-iy} = \cos y - i \sin y.$$

2 Solving the above equations, we obtain

$$\sin y = \frac{e^{iy} - e^{-iy}}{2i} \text{ and } \cos y = \frac{e^{iy} + e^{-iy}}{2}.$$

Trigonometric function

- 1 Euler formula yields

$$e^{iy} = \cos y + i \sin y \text{ and } e^{-iy} = \cos y - i \sin y.$$

- 2 Solving the above equations, we obtain

$$\sin y = \frac{e^{iy} - e^{-iy}}{2i} \text{ and } \cos y = \frac{e^{iy} + e^{-iy}}{2}.$$

- 3 Here sine and cosine functions are expressed in term of **complex exponential functions**.

Trigonometric function

Trigonometric function

- 1 Similarly, we define the **complex** sine and cosine functions in terms of the complex exponential functions e^{iz} and e^{-iz} below

Trigonometric function

- 1 Similarly, we define the **complex** sine and cosine functions in terms of the complex exponential functions e^{iz} and e^{-iz} below

2

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i} \text{ and } \cos z = \frac{e^{iz} + e^{-iz}}{2}.$$

Trigonometric function

- 1 Similarly, we define the **complex** sine and cosine functions in terms of the complex exponential functions e^{iz} and e^{-iz} below

2

$$\sin z = \frac{e^{iz} - e^{-iz}}{2i} \text{ and } \cos z = \frac{e^{iz} + e^{-iz}}{2}.$$

- 3 Now using sine and cosine we can define $\tan z$, $\sec z$, $\operatorname{cosec} z$ as in the real case.

Properties of sine and cosine functions

We have to prove all the properties:

Properties of sine and cosine functions

We have to prove all the properties:

- 1 $\sin z$ and $\cos z$ are entire functions.

Properties of sine and cosine functions

We have to prove all the properties:

- 1 $\sin z$ and $\cos z$ are entire functions.
- 2 $\frac{d}{dz} \sin z = \cos z$, and $\frac{d}{dz} \cos z = -\sin z$.

Properties of sine and cosine functions

We have to prove all the properties:

- 1 $\sin z$ and $\cos z$ are entire functions.
- 2 $\frac{d}{dz} \sin z = \cos z$, and $\frac{d}{dz} \cos z = -\sin z$.
- 3 Real and imaginary parts of $\sin z$ and $\cos z$ are:

$$\sin z = \sin x \cosh y + i \cos x \sinh y$$

$$\cos z = \cos x \cosh y - i \sin x \sinh y.$$

Properties of sine and cosine functions

We have to prove all the properties:

Properties of sine and cosine functions

We have to prove all the properties:

- 1 Modulii of $\sin z$ and $\cos z$ are:

$$\begin{aligned} |\sin z| &= \sqrt{\sin^2 x + \sinh^2 y} \\ |\cos z| &= \sqrt{\cos^2 x + \sinh^2 y}. \end{aligned}$$

Properties of sine and cosine functions

We have to prove all the properties:

- 1 Modulii of $\sin z$ and $\cos z$ are:

$$\begin{aligned} |\sin z| &= \sqrt{\sin^2 x + \sinh^2 y} \\ |\cos z| &= \sqrt{\cos^2 x + \sinh^2 y}. \end{aligned}$$

- 2 $\sin z$ and $\cos z$ are unbounded function in \mathbb{C} .

Properties of sine and cosine functions

We have to prove all the properties:

- 1 Modulii of $\sin z$ and $\cos z$ are:

$$\begin{aligned} |\sin z| &= \sqrt{\sin^2 x + \sinh^2 y} \\ |\cos z| &= \sqrt{\cos^2 x + \sinh^2 y}. \end{aligned}$$

- 2 $\sin z$ and $\cos z$ are unbounded function in \mathbb{C} .

What happens for real sine and cosine function in \mathbb{R} ?

Properties of sine and cosine functions

We have to prove all the properties:

- 1 Modulii of $\sin z$ and $\cos z$ are:

$$\begin{aligned} |\sin z| &= \sqrt{\sin^2 x + \sinh^2 y} \\ |\cos z| &= \sqrt{\cos^2 x + \sinh^2 y}. \end{aligned}$$

- 2 $\sin z$ and $\cos z$ are unbounded function in \mathbb{C} .

What happens for real sine and cosine function in \mathbb{R} ?

- 3 $\sin z = 0$ iff $z = k\pi$, and $\cos z = 0$ iff $z = k\pi + \frac{\pi}{2}$, where $k \in \mathbb{Z}$.

Trigonometric formula

We have to prove all the properties:

Trigonometric formula

We have to prove all the properties:

1 $\sin(-z) = -\sin z$ and $\cos(-z) = \cos z$.

Trigonometric formula

We have to prove all the properties:

- 1 $\sin(-z) = -\sin z$ and $\cos(-z) = \cos z$.
- 2 $\sin(z_1 + z_2) = \sin z_1 \cos z_2 + \cos z_1 \sin z_2$.

Trigonometric formula

We have to prove all the properties:

1 $\sin(-z) = -\sin z$ and $\cos(-z) = \cos z$.

2 $\sin(z_1 + z_2) = \sin z_1 \cos z_2 + \cos z_1 \sin z_2$.

3 $\sin 2z = 2 \sin z \cos z$.

Trigonometric formula

We have to prove all the properties:

1 $\sin(-z) = -\sin z$ and $\cos(-z) = \cos z$.

2 $\sin(z_1 + z_2) = \sin z_1 \cos z_2 + \cos z_1 \sin z_2$.

3 $\sin 2z = 2 \sin z \cos z$.

4 $\sin(z + \pi) = -\sin z$, $\sin(z + 2\pi) = \sin z$

Trigonometric formula

We have to prove all the properties:

1 $\sin(-z) = -\sin z$ and $\cos(-z) = \cos z$.

2 $\sin(z_1 + z_2) = \sin z_1 \cos z_2 + \cos z_1 \sin z_2$.

3 $\sin 2z = 2 \sin z \cos z$.

4 $\sin(z + \pi) = -\sin z$, $\sin(z + 2\pi) = \sin z$

5 $\cos(z_1 + z_2) = \cos z_1 \cos z_2 - \sin z_1 \sin z_2$.

Trigonometric formula

We have to prove all the properties:

1 $\sin(-z) = -\sin z$ and $\cos(-z) = \cos z$.

2 $\sin(z_1 + z_2) = \sin z_1 \cos z_2 + \cos z_1 \sin z_2$.

3 $\sin 2z = 2 \sin z \cos z$.

4 $\sin(z + \pi) = -\sin z$, $\sin(z + 2\pi) = \sin z$

5 $\cos(z_1 + z_2) = \cos z_1 \cos z_2 - \sin z_1 \sin z_2$.

6 $\cos 2z = \cos^2 z - \sin^2 z$.

Hyperbolic functions

Hyperbolic functions

Like real real hyperbolic sine and hyperbolic cosine, we define the **complex** hyperbolic sine and cosine functions as

Hyperbolic functions

Like real real hyperbolic sine and hyperbolic cosine, we define the **complex** hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \quad \text{and} \quad \cosh z = \frac{e^z + e^{-z}}{2}.$$

1 Properties

Hyperbolic functions

Like real real hyperbolic sine and hyperbolic cosine, we define the **complex** hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \text{ and } \cosh z = \frac{e^z + e^{-z}}{2}.$$

1 Properties

$$-i \sinh(iz) = \sin z \text{ and } \cosh(iz) = \cos(z)$$

$$-i \sin(iz) = \sinh z \text{ and } \cos(iz) = \cosh(z)$$

$$\sinh(-z) = -\sinh z \text{ and } \cosh(-z) = \cosh(z).$$

Hyperbolic functions

Like real real hyperbolic sine and hyperbolic cosine, we define the **complex** hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \text{ and } \cosh z = \frac{e^z + e^{-z}}{2}.$$

1 Properties

$$-i \sinh(iz) = \sin z \text{ and } \cosh(iz) = \cos(z)$$

$$\sinh(-z) = -\sinh z \text{ and } \cosh(-z) = \cosh(z).$$

Hyperbolic functions

Like real real hyperbolic sine and hyperbolic cosine, we define the **complex** hyperbolic sine and cosine functions as

$$\sinh z = \frac{e^z - e^{-z}}{2} \text{ and } \cosh z = \frac{e^z + e^{-z}}{2}.$$

1 Properties

$$-i \sinh(iz) = \sin z \text{ and } \cosh(iz) = \cos(z)$$

$$-i \sin(iz) = \sinh z \text{ and } \cos(iz) = \cosh(z)$$

$$\sinh(-z) = -\sinh z \text{ and } \cosh(-z) = \cosh(z).$$

Logarithmic functions

Logarithmic functions

Before studying the **complex logarithmic function** we define the following

- 1 $\ln |z|$ is called the **real** logarithm of $|z|$, $|z| \neq 0$.

Logarithmic functions

Before studying the **complex logarithmic function** we define the following

- 1 $\ln|z|$ is called the **real** logarithm of $|z|$, $|z| \neq 0$.
- 2 $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ —**punctured complex plane**.

Logarithmic functions

Before studying the **complex logarithmic function** we define the following

- 1 $\ln|z|$ is called the **real** logarithm of $|z|$, $|z| \neq 0$.
- 2 $\mathbb{C}^* = \mathbb{C} \setminus \{0\}$ —**punctured complex plane**.
- 3 $H = \{x + iy : -\pi < y \leq \pi\}$ - **a strip in the complex plane** (visualize it)

Logarithmic function

- 1 For $z \in \mathbb{C}^*$, we define

$$\log z = \ln |z| + i\arg(z).$$

Logarithmic function

- 1 For $z \in \mathbb{C}^*$, we define

$$\log z = \ln |z| + i\arg(z).$$

- 2 $\log z$ is called the **complex logarithm** of z .

Logarithmic function

- 1 For $z \in \mathbb{C}^*$, we define

$$\log z = \ln |z| + i\arg(z).$$

- 2 $\log z$ is called the **complex logarithm** of z .
- 3 $\log z$ is a **multi-valued function**. WHY?

Logarithmic function

- 1 For $z \in \mathbb{C}^*$, we define

$$\log z = \ln |z| + i\arg(z).$$

- 2 $\log z$ is called the **complex logarithm** of z .
- 3 $\log z$ is a **multi-valued function**. WHY?
- 4 By considering the **principle argument**, we define single-valued function as

$$\text{Log } z = \ln |z| + i\text{Arg}(z).$$

Logarithmic function

- 1 For $z \in \mathbb{C}^*$, we define

$$\log z = \ln |z| + i\arg(z).$$

- 2 $\log z$ is called the **complex logarithm** of z .
- 3 $\log z$ is a **multi-valued function**. WHY?
- 4 By considering the **principle argument**, we define single-valued function as

$$\text{Log } z = \ln |z| + i\text{Arg}(z).$$

- 5 Certainly, $\text{Log} : \mathbb{C}^* \rightarrow H$ is **well defined** i.e., **single-valued** function.

Example

1 When $z = 1 + i$,

$$\log(1+i) = \ln\sqrt{2} + i\arg(z) = \ln\sqrt{2} + i\left(\frac{\pi}{4} + 2k\pi\right), k \in \mathbb{Z}.$$

Example

1 When $z = 1 + i$,

$$\log(1+i) = \ln\sqrt{2} + i\arg(z) = \ln\sqrt{2} + i\left(\frac{\pi}{4} + 2k\pi\right), k \in \mathbb{Z}.$$

Also,

$$\text{Log}(1+i) = \ln\sqrt{2} + i\text{Arg}(z) = \ln\sqrt{2} + i\frac{\pi}{4}$$

Example

- 1 When $z = 1 + i$,

$$\log(1+i) = \ln\sqrt{2} + i\arg(z) = \ln\sqrt{2} + i\left(\frac{\pi}{4} + 2k\pi\right), k \in \mathbb{Z}.$$

Also,

$$\text{Log}(1+i) = \ln\sqrt{2} + i\text{Arg}(z) = \ln\sqrt{2} + i\frac{\pi}{4}$$

- 2 Complex logarithm for negative real exists. When $z = -1$,

$$\log(-1) = \ln 1 + i\arg(z) = i(\pi + 2k\pi), k \in \mathbb{Z}.$$

Example

- 1 When $z = 1 + i$,

$$\log(1+i) = \ln\sqrt{2} + i\arg(z) = \ln\sqrt{2} + i\left(\frac{\pi}{4} + 2k\pi\right), k \in \mathbb{Z}.$$

Also,

$$\text{Log}(1+i) = \ln\sqrt{2} + i\text{Arg}(z) = \ln\sqrt{2} + i\frac{\pi}{4}$$

- 2 Complex logarithm for negative real exists. When $z = -1$,

$$\log(-1) = \ln 1 + i\arg(z) = i(\pi + 2k\pi), k \in \mathbb{Z}.$$

Also,

$$\text{Log}(-1) = \ln 1 + i\text{Arg}(z) = i\pi$$

Complex Analysis

LECTURE 9

August 16, 2023

Properties

- 1 For $z \neq 0$, $e^{\operatorname{Log} z} = z$.

Properties

- 1 For $z \neq 0$, $e^{\operatorname{Log} z} = z$.
- 2 For $z \in H$, $\operatorname{Log} e^z = z$.

Properties

- 1 For $z \neq 0$, $e^{\operatorname{Log} z} = z$.
- 2 For $z \in H$, $\operatorname{Log} e^z = z$.
- 3 For $z \notin H$, $\operatorname{Log} e^z \neq z$.

Properties

- 1 For $z \neq 0$, $e^{\operatorname{Log} z} = z$.
- 2 For $z \in H$, $\operatorname{Log} e^z = z$.
- 3 For $z \notin H$, $\operatorname{Log} e^z \neq z$.
- 4 For real $x > 0$, $\operatorname{Log} x = \ln x$.

Properties

- 1 For $z \neq 0$, $e^{\operatorname{Log} z} = z$.
- 2 For $z \in H$, $\operatorname{Log} e^z = z$.
- 3 For $z \notin H$, $\operatorname{Log} e^z \neq z$.
- 4 For real $x > 0$, $\operatorname{Log} x = \ln x$.
- 5 $\operatorname{Log} z$ is not continuous on the negative real axis
 $\mathbb{R}^- = \{z = x + iy : x < 0, y = 0\}$.

Properties

- 1 Log z is analytic on the set $\mathbb{C}^* \setminus \mathbb{R}^-$.

Properties

1 Log z is analytic on the set $\mathbb{C}^* \setminus \mathbb{R}^-$.

2 $\frac{d}{dz} \text{Log } z = \frac{1}{z}$.

Properties

- 1 $\text{Log } z$ is analytic on the set $\mathbb{C}^* \setminus \mathbb{R}^-$.
- 2 $\frac{d}{dz} \text{Log } z = \frac{1}{z}$.
- 3 The identity $\text{Log}(z_1 z_2) = \text{Log } z_1 + \text{Log } z_2$ is true iff $\text{Arg } z_1 + \text{Arg } z_2 \in (-\pi, \pi]$.

Curves

Curves

1 Why do we need to study curves?

Curves

- 1 Why do we need to study curves?
- 2 A curve C in the complex plane \mathbb{C} is given by a function
(involving one parameter)

$$\gamma : [a, b] \rightarrow \mathbb{C}$$

Curves

- 1 Why do we need to study curves?
- 2 A curve C in the complex plane \mathbb{C} is given by a function
(involving one parameter)

$$\gamma : [a, b] \rightarrow \mathbb{C}$$

defined as

$$\gamma(t) = x(t) + iy(t),$$

with $x, y : [a, b] \rightarrow \mathbb{R}$ being continuous.

Examples

Examples

1 Circle:

Examples

1 Circle:

The function $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$ given by

$\gamma(t) = z_0 + r(\cos t + i \sin t)$, where z_0 is a fixed complex number, gives the circle with center at z_0 and radius r .

Examples

1 Circle:

The function $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$ given by

$\gamma(t) = z_0 + r(\cos t + i \sin t)$, where z_0 is a fixed complex number, gives the circle with center at z_0 and radius r .

2 Line segment:

Examples

1 Circle:

The function $\gamma : [0, 2\pi] \rightarrow \mathbb{C}$ given by

$\gamma(t) = z_0 + r(\cos t + i \sin t)$, where z_0 is a fixed complex number, gives the circle with center at z_0 and radius r .

2 Line segment:

The function $\gamma : [0, 1] \rightarrow \mathbb{C}$ given by

$$\gamma(t) = tz_1 + (1 - t)z_0$$

gives the line segment joining z_0 and z_1 .

Types of curves

1 Smooth curve

Types of curves

- 1 Smooth curve
- 2 Closed curve

Types of curves

1 Smooth curve

2 Closed curve

3 Contour

Types of curves

- 1 Smooth curve
- 2 Closed curve
- 3 Contour
- 4 Simple contour

Types of curves

- 1 Smooth curve
- 2 Closed curve
- 3 Contour
- 4 Simple contour
- 5 Simple closed contour

Types of curves

- 1 Smooth curve
- 2 Closed curve
- 3 Contour
- 4 Simple contour
- 5 Simple closed contour
- 6 Curve with reverse orientation

Smooth curve

- 1 A curve C defined for $t \in [a, b]$ is called a smooth curve if $\gamma'(t) = x'(t) + iy'(t)$ is continuous on $[a, b]$ and nonzero for all $t \in (a, b)$.

Smooth curve

- 1 A curve C defined for $t \in [a, b]$ is called a smooth curve if $\gamma'(t) = x'(t) + iy'(t)$ is continuous on $[a, b]$ and nonzero for all $t \in (a, b)$.
- 2 Geometrically this means that C has everywhere a continuously turning tangent.

Smooth curve

- 1 A curve C defined for $t \in [a, b]$ is called a smooth curve if $\gamma'(t) = x'(t) + iy'(t)$ is **continuous** on $[a, b]$ and **nonzero** for all $t \in (a, b)$.
- 2 Geometrically this means that C has everywhere a **continuously turning tangent**.
- 3 Alternatively, a smooth curve does not have any **sharp point or corner**.

Example

- 1 Circular arc and the circle are examples of smooth curves.

Example


- 1 Circular arc and the circle are examples of smooth curves.
- 2 $\gamma(t) = t^4 + it^5, t \in [-1, 1]$ is NOT a smooth curve.

Example

- 1 Circular arc and the circle are examples of smooth curves.
- 2 $\gamma(t) = t^4 + it^5, t \in [-1, 1]$ is NOT a smooth curve.
Explain mathematically??


Example

- 1 Circular arc and the circle are examples of smooth curves.
- 2 $\gamma(t) = t^4 + it^5, t \in [-1, 1]$ is NOT a smooth curve.
Explain mathematically??



Example

- 1 Circular arc and the circle are examples of smooth curves.
- 2 $\gamma(t) = t^4 + it^5, t \in [-1, 1]$ is NOT a smooth curve.
Explain mathematically??



Example

$\gamma(t) = 3 \cos^3 t + i3 \sin^3 t, t \in [0, 2\pi]$ is **NOT** a smooth curve.


Example

$\gamma(t) = 3 \cos^3 t + i3 \sin^3 t, t \in [0, 2\pi]$ is NOT a smooth curve.

Explain mathematically??

Example

$\gamma(t) = 3 \cos^3 t + i 3 \sin^3 t, t \in [0, 2\pi]$ is **NOT** a smooth curve.
Explain mathematically??



Closed curve

- 1 A curve C is called closed if its terminal point coincides with its initial point, that is $\gamma(a) = \gamma(b)$.

Closed curve

- 1 A curve C is called closed if its terminal point coincides with its initial point, that is $\gamma(a) = \gamma(b)$.
- 2 In this case, the curve may have property $\gamma(t_1) = \gamma(t_2)$ for some $t_1 \neq t_2, t_1, t_2 \in (a, b)$ too.

Contour

- 1 **Contour:** A contour is a curve that is obtained by joining finitely many **smooth curves** end to end.

Contour

- 1 **Contour:** A contour is a curve that is obtained by joining finitely many **smooth curves** end to end.
- 2 **Simple contour:** A contour is called simple if it does not cross itself (if initial point and the final point are same they are not considered as non simple).

Contour

- 1 **Contour:** A contour is a curve that is obtained by joining finitely many **smooth curves** end to end.
- 2 **Simple contour:** A contour is called simple if it does not cross itself (if initial point and the final point are same they are not considered as non simple).
- 3 **Examples:** A circle is simple, the curve of the shape U are examples of simple contour, but the curve of the shapes 8 and g are not simple

Simple closed contour

- 1 A contour C is called a simple closed contour if **ONLY** its terminal point coincides with its initial point, that is $\gamma(a) = \gamma(b)$.

Simple closed contour

- 1 A contour C is called a simple closed contour if **ONLY** its terminal point coincides with its initial point, that is $\gamma(a) = \gamma(b)$.
- 2 In this case, the contour **CANNOT** have property $\gamma(t_1) = \gamma(t_1)$ for some $t_1 \neq t_2, t_1, t_2 \in (a, b)$.

Simple closed contour

- 1 A contour C is called a simple closed contour if **ONLY** its terminal point coincides with its initial point, that is $\gamma(a) = \gamma(b)$.
- 2 In this case, the contour **CANNOT** have property $\gamma(t_1) = \gamma(t_1)$ for some $t_1 \neq t_2, t_1, t_2 \in (a, b)$.
- 3 **Examples:** A circle is simple.

Curve with Reverse Orientation

- 1 Let $C : \gamma : [a, b] \rightarrow \mathbb{C}$ be a curve. Then the curve with the reverse orientation, denoted as $-C : \gamma_- : [a, b] \rightarrow \mathbb{C}$ and is defined as

Curve with Reverse Orientation

- 1 Let $C : \gamma : [a, b] \rightarrow \mathbb{C}$ be a curve. Then the curve with the reverse orientation, denoted as $-C : \gamma_- : [a, b] \rightarrow \mathbb{C}$ and is defined as

$$\gamma_-(t) = \gamma(b + a - t).$$

Curve with Reverse Orientation

- 1 Let $C : \gamma : [a, b] \rightarrow \mathbb{C}$ be a curve. Then the curve with the reverse orientation, denoted as $-C : \gamma_- : [a, b] \rightarrow \mathbb{C}$ and is defined as

$$\gamma_-(t) = \gamma(b + a - t).$$

- 2 Thus for a contour C , the contour with the negative orientation $-C$ make sense.

Simply connected domain

- 1 A domain D is called simply connected if every simple closed contour within it encloses points of D only.

Simply connected domain

- 1 A domain D is called simply connected if every simple closed contour within it encloses points of D only.
- 2 A simply connected domain does not contain any **hole**.

Simply connected domain

- 1 A domain D is called simply connected if every simple closed contour within it encloses points of D only.
- 2 A simply connected domain does not contain any **hole**.
- 3 A domain D is called multiply connected if it is not simply connected.

Simply connected domain

- 1 A domain D is called simply connected if every simple closed contour within it encloses points of D only.
- 2 A simply connected domain does not contain any **hole**.
- 3 A domain D is called multiply connected if it is not simply connected.
- 4 A multiply connected domain contains at least **one** hole.

Simply connected domain

- 1 A domain D is called simply connected if every simple closed contour within it encloses points of D only.
- 2 A simply connected domain does not contain any **hole**.
- 3 A domain D is called multiply connected if it is not simply connected.
- 4 A multiply connected domain contains at least **one** hole.
- 5 A doubly connected domain contain **EXACTLY one** hole.

Simply connected domain

- 1 A domain D is called simply connected if every simple closed contour within it encloses points of D only.
- 2 A simply connected domain does not contain any **hole**.
- 3 A domain D is called multiply connected if it is not simply connected.
- 4 A multiply connected domain contains at least **one** hole.
- 5 A doubly connected domain contain **EXACTLY one** hole.
- 6 A triply connected domain contain **EXACTLY two** holes.

Integral of a complex valued function of a real variable

- 1 Understand the meaning of "**complex valued function of a real variable**".

Integral of a complex valued function of a real variable

- 1 Understand the meaning of "**complex valued function of a real variable**".
- 2 Let $f : [a, b] \rightarrow \mathbb{C}$ be a **piecewise** continuous function. Then $f(t) = u(t) + iv(t)$ where $u, v : [a, b] \rightarrow \mathbb{R}$. We then define

$$\int_a^b f(t) dt = \int_a^b u(t) dt + i \int_a^b v(t) dt.$$

Integral of a complex valued function of a real variable

- 1 Understand the meaning of "**complex valued function of a real variable**".
- 2 Let $f : [a, b] \rightarrow \mathbb{C}$ be a **piecewise** continuous function. Then $f(t) = u(t) + iv(t)$ where $u, v : [a, b] \rightarrow \mathbb{R}$. We then define

$$\int_a^b f(t) dt = \int_a^b u(t) dt + i \int_a^b v(t) dt.$$

- 3 The above expression is called integral of a complex valued function of a real variable.

Integral of a complex valued function of a real variable

- 1 Understand the meaning of "**complex valued function of a real variable**".
- 2 Let $f : [a, b] \rightarrow \mathbb{C}$ be a **piecewise** continuous function. Then $f(t) = u(t) + iv(t)$ where $u, v : [a, b] \rightarrow \mathbb{R}$. We then define

$$\int_a^b f(t) dt = \int_a^b u(t) dt + i \int_a^b v(t) dt.$$

- 3 The above expression is called integral of a complex valued function of a real variable.
- 4 Easy to calculate.

Important results

Result: Let $f : [a, b] \rightarrow \mathbb{C}$ be a piecewise continuous function.

Then

$$\left| \int_a^b f(t) dt \right| \leq \int_a^b |f(t)| dt.$$

Proof:

- 1 Let $\phi = \operatorname{Arg} \left(\int_a^b f(t) dt \right)$

Important results

Result: Let $f : [a, b] \rightarrow \mathbb{C}$ be a piecewise continuous function.
Then

$$\left| \int_a^b f(t) dt \right| \leq \int_a^b |f(t)| dt.$$

Proof:

1 Let $\phi = \operatorname{Arg} \left(\int_a^b f(t) dt \right)$

2

$$\begin{aligned} \left| \int_a^b f(t) dt \right| &= e^{-i\phi} \int_a^b f(t) dt \quad (\because |z| = e^{-i\operatorname{Arg} z} z) \\ &= \int_a^b e^{-i\phi} f(t) dt \end{aligned}$$

Proof

$$= \operatorname{Re} \int_a^b e^{-i\phi} f(t) dt \quad (\because \int_a^b e^{-i\phi} f(t) dt = \left| \int_a^b f(t) dt \right| \text{ is real})$$

$$= \int_a^b \operatorname{Re} (e^{-i\phi} f(t)) dt$$

[Note that this is Riemann integral of the real function $\operatorname{Re} (e^{-i\phi} f(t))$]

$$\leq \int_a^b |e^{-i\phi} f(t)| dt$$

[Using the property of Riemann integral with the fact that

$$\operatorname{Re} (e^{-i\phi} f(t)) \leq |e^{-i\phi} f(t)|$$

$$= \int_a^b |f(t)| dt.$$

Remark

ϕ is NOT defined if $\left| \int_a^b f(t) dt \right| = 0$. Then there is nothing to prove.

Important results

Result: For $\alpha \in \mathbb{R}$, show that

$$\int_a^b e^{i\alpha t} dt = \frac{e^{i\alpha b} - e^{i\alpha a}}{i\alpha}.$$

Complex Analysis

LECTURE 10

August 17, 2023

Important results

Result: Let $f : [a, b] \rightarrow \mathbb{C}$ be a **continuous** function. Then it is **NOT necessary** that there must exist $c \in (a, b)$ such that

$$\int_a^b f(t) dt = f(c)(b - a).$$

Important results

Result: Let $f : [a, b] \rightarrow \mathbb{C}$ be a **continuous** function. Then it is **NOT necessary** that there must exist $c \in (a, b)$ such that

$$\int_a^b f(t) dt = f(c)(b - a).$$

Example:

Important results

Result: Let $f : [a, b] \rightarrow \mathbb{C}$ be a **continuous** function. Then it is **NOT necessary** that there must exist $c \in (a, b)$ such that

$$\int_a^b f(t) dt = f(c)(b - a).$$

Example:

- 1 Set $f(t) = e^{it}$, $a = 0$, $b = 2\pi$.

Important results

Result: Let $f : [a, b] \rightarrow \mathbb{C}$ be a **continuous** function. Then it is **NOT necessary** that there must exist $c \in (a, b)$ such that

$$\int_a^b f(t) dt = f(c)(b - a).$$

Example:

1 Set $f(t) = e^{it}$, $a = 0$, $b = 2\pi$.

2 Then $\int_a^b f(t) dt = 0$

Important results

Result: Let $f : [a, b] \rightarrow \mathbb{C}$ be a **continuous** function. Then it is **NOT necessary** that there must exist $c \in (a, b)$ such that

$$\int_a^b f(t) dt = f(c)(b - a).$$

Example:

- 1 Set $f(t) = e^{it}$, $a = 0$, $b = 2\pi$.

- 2 Then $\int_a^b f(t) dt = 0$

- 3 But $f(t)$ is **zero nowhere**.

Integral of a complex valued function of complex variable

- 1 So far we have discussed about complex valued function of **a real variable**.

Integral of a complex valued function of complex variable

- 1 So far we have discussed about complex valued function of **a real variable**.
- 2 Understand the meaning of integration of complex valued function of **complex variable**.

Integral of a complex valued function of complex variable

- 1 So far we have discussed about complex valued function of **a real variable**.
- 2 Understand the meaning of integration of complex valued function of **complex variable**.
- 3 Now the domain of the integrand is in the complex plane.

Integral of a complex valued function of complex variable

- 1 So far we have discussed about complex valued function of **a real variable**.
- 2 Understand the meaning of integration of complex valued function of **complex variable**.
- 3 Now the domain of the integrand is in the complex plane.
- 4 The integration will be **taken over a contour or curve** in \mathbb{C} .

Integral of a complex valued function of complex variable

- 1 So far we have discussed about complex valued function of **a real variable**.
- 2 Understand the meaning of integration of complex valued function of **complex variable**.
- 3 Now the domain of the integrand is in the complex plane.
- 4 The integration will be **taken over a contour or curve** in \mathbb{C} .
- 5 Such an integration is called a **contour integration**.

Contour Integral

- 1 Let C be a contour contained in a domain D .

Contour Integral

- 1 Let C be a contour contained in a domain D .
- 2 Let $f : D \rightarrow \mathbb{C}$ be a piecewise continuous function defined on C .

Contour Integral

- 1 Let C be a contour contained in a domain D .
- 2 Let $f : D \rightarrow \mathbb{C}$ be a piecewise continuous function defined on C .
- 3 Then the contour integral of $f(z)$ along the contour C , denoted as $\int_C f(z) dz$.

Contour Integral

- 1 Let C be a contour contained in a domain D .
- 2 Let $f : D \rightarrow \mathbb{C}$ be a piecewise continuous function defined on C .
- 3 Then the contour integral of $f(z)$ along the contour C , denoted as $\int_C f(z) dz$.
- 4 We define the integration using Riemann sum.

Contour Integral-Riemann sum

- 1 Let C be a **contour** in \mathbb{C} with initial point A and terminal point B .

Contour Integral-Riemann sum

- 1 Let C be a **contour** in \mathbb{C} with initial point A and terminal point B .
- 2 We create a partition $P_n = \{a = z_0, z_1, z_2, \dots, z_{n-1}, z_n = B\}$ where all the points are **on the contour**.

Contour Integral-Riemann sum




Figure:

Contour Integral-Riemann sum

- 1 Let C be a **contour** in \mathbb{C} with initial point A and terminal point B .
- 2 We create a partition $P_n = \{a = z_0, z_1, \dots, z_{n-1}, z_n = B\}$ where all the points are **on the contour**.

Contour Integral-Riemann sum

- 1 Let C be a **contour** in \mathbb{C} with initial point A and terminal point B .
- 2 We create a partition $P_n = \{a = z_0, z_1, \dots, z_{n-1}, z_n = B\}$ where all the points are **on the contour**.
- 3 We define $\Delta z_k = z_{k+1} - z_k$

Contour Integral-Riemann sum

- 1 Let C be a **contour** in \mathbb{C} with initial point A and terminal point B .
- 2 We create a partition $P_n = \{a = z_0, z_1, \dots, z_{n-1}, z_n = B\}$ where all the points are **on the contour**.
- 3 We define $\Delta z_k = z_{k+1} - z_k$
- 4 We now select a point c_k on C lying between z_k and z_{k+1} .

Contour Integral-Riemann sum

- 1 Let C be a **contour** in \mathbb{C} with initial point A and terminal point B .
- 2 We create a partition $P_n = \{a = z_0, z_1, \dots, z_{n-1}, z_n = B\}$ where all the points are **on the contour**.
- 3 We define $\Delta z_k = z_{k+1} - z_k$
- 4 We now select a point c_k on C lying between z_k and z_{k+1} .
- 5 Then the Riemann sum for the partition P_n is

$$S(P_n) = \sum_{k=1}^n f(c_k) \Delta z_k$$

Contour Integral-Riemann sum

- 1 When every sequence $\{S(P_n)\}$ converges to a limit with maximum $|\Delta z_k| \rightarrow 0$, we say that the integration exists.

Contour Integral-Riemann sum

- 1 When every sequence $\{S(P_n)\}$ converges to a limit with maximum $|\Delta z_k| \rightarrow 0$, we say that the integration exists.
- 2 We now define

$$\int_C f(z) dz = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k) \Delta z_k$$

Example- Riemann sum

Use the Riemann sum to find an approximation $S(P_8)$ to the contour integral $\int_C e^z dz$ where C is the line joining the starting point $A = 0$ and terminal point $B = 2 + i\pi/4$.

Example- Riemann sum




Figure:

Example- Riemann sum

- 1 Develop $P_8 = \{z + k : z_k = \frac{k}{4} + i \frac{k\pi}{32}, k = 0, 1, 2, \dots, 8\}$.

Example- Riemann sum

- 1 Develop $P_8 = \{z + k : z_k = \frac{k}{4} + i\frac{k\pi}{32}, k = 0, 1, 2, \dots, 8\}$.
- 2 Then $\Delta z_k = z_{k+1} - z_k = \frac{1}{4} + i\frac{\pi}{32}, k = 1, 2, \dots, 8$

Example- Riemann sum

- 1 Develop $P_8 = \{z + k : z_k = \frac{k}{4} + i\frac{k\pi}{32}, k = 0, 1, 2, \dots, 8\}$.
- 2 Then $\Delta z_k = z_{k+1} - z_k = \frac{1}{4} + i\frac{\pi}{32}, k = 1, 2, \dots, 8$
- 3 We now select
 $c_k = \frac{z_{k+1} + z_k}{2} = \frac{2k - 1}{8} + i\frac{(2k - 1)\pi}{64}, k = 1, 2, \dots, 8$

Example- Riemann sum

1 Develop $P_8 = \{z + k : z_k = \frac{k}{4} + i \frac{k\pi}{32}, k = 0, 1, 2, \dots, 8\}$.

2 Then $\Delta z_k = z_{k+1} - z_k = \frac{1}{4} + i \frac{\pi}{32}, k = 1, 2, \dots, 8$

3 We now select

$$c_k = \frac{z_{k+1} + z_k}{2} = \frac{2k-1}{8} + i \frac{(2k-1)\pi}{64}, k = 1, 2, \dots, 8$$

4

$$\begin{aligned} S(P_8) &= \sum_{k=1}^8 \exp \left[\frac{2k-1}{8} + i \frac{(2k-1)\pi}{64} \right] \times \left(\frac{1}{4} + i \frac{\pi}{32} \right) \\ &\approx 4.23 + 5.20i \end{aligned}$$

Example- Riemann sum

- 1 Develop $P_8 = \{z + k : z_k = \frac{k}{4} + i \frac{k\pi}{32}, k = 0, 1, 2, \dots, 8\}$.
- 2 Then $\Delta z_k = z_{k+1} - z_k = \frac{1}{4} + i \frac{\pi}{32}, k = 1, 2, \dots, 8$
- 3 We now select
 $c_k = \frac{z_{k+1} + z_k}{2} = \frac{2k - 1}{8} + i \frac{(2k - 1)\pi}{64}, k = 1, 2, \dots, 8$
- 4

$$S(P_8) = \sum_{k=1}^8 \exp \left[\frac{2k - 1}{8} + i \frac{(2k - 1)\pi}{64} \right] \times \left(\frac{1}{4} + i \frac{\pi}{32} \right)$$

$$\approx 4.23 + 5.20i$$

- 5 Later we can calculate the analytical value

$$-1 + \frac{e^2}{\sqrt{2}} + i \frac{e^2}{\sqrt{2}} \approx 4.22485 + 4.22485i$$

Trouble with Riemann sum??

- 1 Ohhh, we cannot continue such **messy** calculations.

Trouble with Riemann sum??

- 1 Ohhh, we cannot continue such **messy** calculations.
- 2 In fact, we cannot bother **Riemann sum** as we **cannot visualize** it.

Trouble with Riemann sum??

- 1 Ohhh, we cannot continue such **messy** calculations.
- 2 In fact, we cannot bother **Riemann sum** as we **cannot visualize** it.
- 3 Need some **alternative approach**.

Trouble with Riemann sum??

- 1 Ohhh, we cannot continue such **messy** calculations.
- 2 In fact, we cannot bother **Riemann sum** as we **cannot visualize** it.
- 3 Need some **alternative approach**.
- 4 Ok, we can establish some **sophisticated** tools.....

Conversion to an integration of complex valued function with a real variable

- 1 Here we convert the complex integration of complex valued function with a **complex variable** into an integration of complex valued function with a **real variable**.

Conversion to an integration of complex valued function with a real variable

- 1 Here we convert the complex integration of complex valued function with a **complex variable** into an integration of complex valued function with a **real variable**.
- 2 Possibly, you would be happy now.

Conversion to an integration of complex valued function with a real variable

- 1 Here we convert the complex integration of complex valued function with a **complex variable** into an integration of complex valued function with a **real variable**.
- 2 Possibly, you would be happy now.
- 3 How to convert?

Procedure

- 1 Recall the parametric representation of the contour C

Procedure

- 1 Recall the parametric representation of the contour C
- 2 Mathematically, C is a one-parameter curve $\gamma(t)$ with

$$\gamma(t) = x(t) + iy(t), t \in [a, b].$$

Procedure

- 1 Recall the parametric representation of the contour C
- 2 Mathematically, C is a one-parameter curve $\gamma(t)$ with

$$\gamma(t) = x(t) + iy(t), t \in [a, b].$$

- 3 We can think $z(t) = x(t) + iy(t)$.

Procedure

- 1 Recall the parametric representation of the contour C
- 2 Mathematically, C is a one-parameter curve $\gamma(t)$ with

$$\gamma(t) = x(t) + iy(t), t \in [a, b].$$

- 3 We can think $z(t) = x(t) + iy(t)$.
- 4 Since C is a contour, $\gamma'(t) = z'(t) = x'(t) + iy'(t)$, $t \in [a, b]$ is (at least) continuous.

Procedure

- 1 Recall the parametric representation of the contour C
- 2 Mathematically, C is a one-parameter curve $\gamma(t)$ with

$$\gamma(t) = x(t) + iy(t), t \in [a, b].$$

- 3 We can think $z(t) = x(t) + iy(t)$.
- 4 Since C is a contour, $\gamma'(t) = z'(t) = x'(t) + iy'(t)$, $t \in [a, b]$ is (at least) continuous.
- 5 We can write

$$\int_C f(z) dz = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k) \Delta z_k$$

Procedure

- 1 Recall the parametric representation of the contour C
- 2 Mathematically, C is a one-parameter curve $\gamma(t)$ with

$$\gamma(t) = x(t) + iy(t), t \in [a, b].$$

- 3 We can think $z(t) = x(t) + iy(t)$.
- 4 Since C is a contour, $\gamma'(t) = z'(t) = x'(t) + iy'(t)$, $t \in [a, b]$ is (at least) continuous.
- 5 We can write

$$\int_C f(z) dz = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k) \Delta z_k = \lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k(t_k)) \frac{\Delta z_k}{\Delta t_k} \Delta t_k$$

Procedure

- 1 The **sum** can be written as

$$\int_C f(z) dz = \int_a^b f(\gamma(t)) \gamma'(t) dt$$

Procedure

- 1 The **sum** can be written as

$$\int_C f(z) dz = \int_a^b f(\gamma(t)) \gamma'(t) dt$$

- 2 We have two advantages

Procedure

- 1 The **sum** can be written as

$$\int_C f(z) dz = \int_a^b f(\gamma(t))\gamma'(t) dt$$

- 2 We have two advantages
- 3 Firstly, we can **visualize** the Riemann sum.

Procedure

- 1 The **sum** can be written as

$$\int_C f(z) dz = \int_a^b f(\gamma(t))\gamma'(t) dt$$

- 2 We have two advantages
- 3 Firstly, we can **visualize** the Riemann sum.
- 4 Secondly, $f(\gamma(t))\gamma'(t)$ is a complex valued function with **a real variable**. We know how to handle such integration.

Procedure

- 1 The **sum** can be written as

$$\int_C f(z) dz = \int_a^b f(\gamma(t))\gamma'(t) dt$$

- 2 We have two advantages
- 3 Firstly, we can **visualize** the Riemann sum.
- 4 Secondly, $f(\gamma(t))\gamma'(t)$ is a complex valued function with **a real variable**. We know how to handle such integration.
- 5 We solve many problems using the final expression.

Alternative representation of Riemann sum

- 1 Assume $f(z) = u + iv$ and $\Delta z = \Delta x + i\Delta y$.

Alternative representation of Riemann sum

- 1 Assume $f(z) = u + iv$ and $\Delta z = \Delta x + i\Delta y$.
- 2 In view of the above and after some egregious calculations, the Riemann sum

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k) \Delta z_k$$

can be written as

$$\int_C f(z) dz = \int_C (udx - vdy) + i \int_C (vdx + udy)$$

Alternative representation of Riemann sum

- 1 Assume $f(z) = u + iv$ and $\Delta z = \Delta x + i\Delta y$.
- 2 In view of the above and after some egregious calculations, the Riemann sum

$$\lim_{n \rightarrow \infty} \sum_{k=1}^n f(c_k) \Delta z_k$$

can be written as

$$\int_C f(z) dz = \int_C (udx - vdy) + i \int_C (vdx + udy)$$

- 3 Note the RHS expression could be remembered from the expansion of $\int_C (u + iv)(dx + idy)$

Importance of the last representation

- 1 The sum is represented in **Cartesian form**.
- 2 We have now **alternative form** to visualize the Riemann sum.

Importance of the last representation

- 1 The sum is represented in **Cartesian form**.
- 2 We have now **alternative form** to visualize the Riemann sum.
- 3 The last expression is nothing but **two ordinary line integrals**.
Easy to tackle as we know.

Importance of the last representation

- 1 The sum is represented in **Cartesian form**.
- 2 We have now **alternative form** to visualize the Riemann sun.
- 3 The last expression is nothing but **two ordinary line integrals**.
Easy to tackle as we know.
- 4 Note that, if we know the points $(x(t), y(t))$ on the contour,
it would yield

$$\int_C f(z) dz = \int_a^b \phi(t) dt + i \int_a^b \psi(t) dt$$

Importance of the last representation

- 1 The sum is represented in **Cartesian form**.
- 2 We have now **alternative form** to visualize the Riemann sum.
- 3 The last expression is nothing but **two ordinary line integrals**.
Easy to tackle as we know.
- 4 Note that, if we know the points $(x(t), y(t))$ on the contour,
it would yield

$$\int_C f(z) dz = \int_a^b \phi(t) dt + i \int_a^b \psi(t) dt$$

- 5 The last form is similar to the one obtained in the first procedure.

Importance of the last representation

- 1 The sum is represented in **Cartesian form**.
- 2 We have now **alternative form** to visualize the Riemann sun.
- 3 The last expression is nothing but **two ordinary line integrals**.
Easy to tackle as we know.
- 4 Note that, if we know the points $(x(t), y(t))$ on the contour,
it would yield

$$\int_C f(z) dz = \int_a^b \phi(t) dt + i \int_a^b \psi(t) dt$$

- 5 The last form is similar to the one obtained in the first procedure. What is the **great** in Cartesian ?

Importance of the last representation

- 1 The sum is represented in **Cartesian form**.
- 2 We have now **alternative form** to visualize the Riemann sun.
- 3 The last expression is nothing but **two ordinary line integrals**.
Easy to tackle as we know.
- 4 Note that, if we know the points $(x(t), y(t))$ on the contour, it would yield

$$\int_C f(z) dz = \int_a^b \phi(t) dt + i \int_a^b \psi(t) dt$$

- 5 The last form is similar to the one obtained in the first procedure. What is the **great** in Cartesian ?
- 6 Ok, wait a bit. We'll see the **beauty of the Cartesian form** soon.

Rules of contour integrals

Prove all:

$$1 \quad \int_C f(z) dz = - \int_{-C} f(z) dz.$$

Rules of contour integrals

Prove all:

1 $\int_C f(z) dz = - \int_{-C} f(z) dz.$

- 2 Let $C = C_1 + C_2 + \cdots + C_n$, where the terminal point of C_k coincides with the initial point of C_{k+1} , then

$$\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz + \cdots + \int_{C_n} f(z) dz.$$

Rules of contour integrals

Prove all:

1 $\int_C f(z) dz = - \int_{-C} f(z) dz.$

- 2 Let $C = C_1 + C_2 + \cdots + C_n$, where the terminal point of C_k coincides with the initial point of C_{k+1} , then

$$\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz + \cdots + \int_{C_n} f(z) dz.$$

- 3 $\int_C Kf(z) dz = K \int_C f(z) dz$, K is a complex constant.

Rules of contour integrals

Prove all:

1 $\int_C f(z) dz = - \int_{-C} f(z) dz.$

- 2 Let $C = C_1 + C_2 + \cdots + C_n$, where the terminal point of C_k coincides with the initial point of C_{k+1} , then

$$\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz + \cdots + \int_{C_n} f(z) dz.$$

- 3 $\int_C Kf(z) dz = K \int_C f(z) dz$, K is a complex constant.
4 $\int_C (f(z) + g(z)) dz = \int_C f(z) dz + \int_C g(z) dz.$

Example

Evaluate

$$I = \int_C z^2 dz,$$

where C is along x -axis from 0 to 1 and then along the line parallel to y -axis from 1 to $1 + 2i$.

Example-contd

- 1 Let C_1 be x -axis from 0 to 1 and C_2 be the line parallel to y -axis from 1 to $1 + 2i$. Then $C = C_1 + C_2$.

Example-contd

- 1 Let C_1 be x -axis from 0 to 1 and C_2 be the line parallel to y -axis from 1 to $1 + 2i$. Then $C = C_1 + C_2$.
- 2 Parametrizations of C_1 and C_2 are given by

$$C_1 : \quad \gamma_1(t) = t, \quad 0 \leq t \leq 1$$

$$\begin{aligned}C_2 : \quad \gamma_2(t) &= (1 - t) + t(1 + 2i) \\&= 1 + 2it, \quad 0 \leq t \leq 1\end{aligned}$$

Example-contd

- 1 Let C_1 be x -axis from 0 to 1 and C_2 be the line parallel to y -axis from 1 to $1 + 2i$. Then $C = C_1 + C_2$.
- 2 Parametrizations of C_1 and C_2 are given by

$$C_1 : \quad \gamma_1(t) = t, \quad 0 \leq t \leq 1$$

$$\begin{aligned}C_2 : \quad \gamma_2(t) &= (1 - t) + t(1 + 2i) \\&= 1 + 2it, \quad 0 \leq t \leq 1\end{aligned}$$

3

$$\int_{C_1} z^2 dz = \int_0^1 t^2 dt = \frac{1}{3}$$

Example-contd

- 1 Let C_1 be x -axis from 0 to 1 and C_2 be the line parallel to y -axis from 1 to $1 + 2i$. Then $C = C_1 + C_2$.
- 2 Parametrizations of C_1 and C_2 are given by

$$C_1 : \quad \gamma_1(t) = t, \quad 0 \leq t \leq 1$$

$$\begin{aligned}C_2 : \quad \gamma_2(t) &= (1 - t) + t(1 + 2i) \\&= 1 + 2it, \quad 0 \leq t \leq 1\end{aligned}$$

3

$$\int_{C_1} z^2 dz = \int_0^1 t^2 dt = \frac{1}{3}$$

4

$$\int_{C_2} z^2 dz = \int_0^1 (1 + 2it)^2 2i dt = -4 - \frac{2}{3}i.$$

Example-contd

$$\int_C z^2 dz = \int_{C_1} z^2 dz + \int_{C_2} z^2 dz = -\frac{11+2i}{3}.$$

Integration using Indefinite Integral

- 1 What do you mean by indefinite integral?

Integration using Indefinite Integral

- 1 What do you mean by indefinite integral?
- 2 Let $f(x)$ be a given function. If there exists anti-derivative $F(x)$ of $f(x)$, then $F(x)$ is called indefinite integral.

Integration using Indefinite Integral

- 1 What do you mean by indefinite integral?

- 2 Let $f(x)$ be a given function. If there exists anti-derivative $F(x)$ of $f(x)$, then $F(x)$ is called indefinite integral.

- 3 What is its role in integration?

Integration using Indefinite Integral

Theorem: Let f be a **continuous** function defined on a domain D and there exists a function F **defined on D** such that $F' = f$. Let $z_1, z_2 \in D$. Then for any contour C **lying** in D starting from z_1 , and ending at z_2 ,

Integration using Indefinite Integral

Theorem: Let f be a **continuous** function defined on a domain D and there exists a function F **defined on D** such that $F' = f$. Let $z_1, z_2 \in D$. Then for any contour C **lying** in D starting from z_1 , and ending at z_2 ,

$$\int_C f(z) dz = F(z_2) - F(z_1).$$

Integration using Indefinite Integral

Theorem: Let f be a **continuous** function defined on a domain D and there exists a function F **defined on D** such that $F' = f$. Let $z_1, z_2 \in D$. Then for any contour C **lying** in D starting from z_1 , and ending at z_2 ,

$$\int_C f(z) dz = F(z_2) - F(z_1).$$

Proof:

Integration using Indefinite Integral

Theorem: Let f be a **continuous** function defined on a domain D and there exists a function F **defined on D** such that $F' = f$. Let $z_1, z_2 \in D$. Then for any contour C **lying** in D starting from z_1 , and ending at z_2 ,

$$\int_C f(z) dz = F(z_2) - F(z_1).$$

Proof:

Integration using Indefinite Integral

Theorem: Let f be a **continuous** function defined on a domain D and there exists a function F **defined on D** such that $F' = f$. Let $z_1, z_2 \in D$. Then for any contour C **lying** in D starting from z_1 , and ending at z_2 ,

$$\int_C f(z) dz = F(z_2) - F(z_1).$$

Proof:

- 1 Suppose that that C is given by a map $\gamma : [a, b] \rightarrow \mathbb{C}$. Then

$$\frac{d}{dt} F(\gamma(t)) = F'(\gamma(t))\gamma'(t).$$

Proof-Contd.

Proof-Contd.

$$\begin{aligned}\int_C f(z) dz &= \int_a^b f(\gamma(t))\gamma'(t) dt \\&= \int_a^b F'(\gamma(t))\gamma'(t) dt \\&= \int_a^b \frac{d}{dt}F(\gamma(t)) dt \\&= F(\gamma(t))|_a^b = F(\gamma(b)) - F(\gamma(a)) \\&= F(z_2) - F(z_1).\end{aligned}$$

Proof-Contd.

$$\begin{aligned}\int_C f(z) dz &= \int_a^b f(\gamma(t))\gamma'(t) dt \\&= \int_a^b F'(\gamma(t))\gamma'(t) dt \\&= \int_a^b \frac{d}{dt}F(\gamma(t)) dt \\&= F(\gamma(t))|_a^b = F(\gamma(b)) - F(\gamma(a)) \\&= F(z_2) - F(z_1).\end{aligned}$$

- 1 Geometrically, it means that the integration does not depend on the path.

Proof-Contd.

$$\begin{aligned}\int_C f(z) dz &= \int_a^b f(\gamma(t))\gamma'(t) dt \\&= \int_a^b F'(\gamma(t))\gamma'(t) dt \\&= \int_a^b \frac{d}{dt}F(\gamma(t)) dt \\&= F(\gamma(t))|_a^b = F(\gamma(b)) - F(\gamma(a)) \\&= F(z_2) - F(z_1).\end{aligned}$$

- 1 Geometrically, it means that the integration does not depend on the path.
- 2 The integration depends on the end points only.

Application of the theorem

Example

Evaluate

$$I = \int_C z^2 dz,$$

where C is along x -axis from 0 to 1 and then along the line parallel to y -axis from 1 to $1 + 2i$.

Application of the theorem

Example

Evaluate

$$I = \int_C z^2 dz,$$

where C is along x -axis from 0 to 1 and then along the line parallel to y -axis from 1 to $1 + 2i$.

- 1 It is possible to construct a domain D such that contour C lies within the domain.

Application of the theorem

Example

Evaluate

$$I = \int_C z^2 dz,$$

where C is along x -axis from 0 to 1 and then along the line parallel to y -axis from 1 to $1 + 2i$.

- 1 It is possible to construct a domain D such that contour C lies **within the domain**.
- 2 Here $f(z) = z^2$ is **continuous**.

Application of the theorem

Example

Evaluate

$$I = \int_C z^2 dz,$$

where C is along x -axis from 0 to 1 and then along the line parallel to y -axis from 1 to $1 + 2i$.

- 1 It is possible to construct a domain D such that contour C lies **within the domain**.
- 2 Here $f(z) = z^2$ is **continuous**.
- 3 Also we can find $F(z) = z^3/3$ such that $F' = f$ on D .

Application of the theorem

Example

Evaluate

$$I = \int_C z^2 dz,$$

where C is along x -axis from 0 to 1 and then along the line parallel to y -axis from 1 to $1 + 2i$.

- 1 It is possible to construct a domain D such that contour C lies **within the domain**.
- 2 Here $f(z) = z^2$ is **continuous**.
- 3 Also we can find $F(z) = z^3/3$ such that $F' = f$ on D .
- 4 Hence, we **CAN** apply the result.

Example-contd

1

$$\int_C z^2 dz = \left[\frac{z^3}{3} \right]_0^{1+2i} = -\frac{11 + 2i}{3}.$$

Example-contd

1

$$\int_C z^2 dz = \left[\frac{z^3}{3} \right]_0^{1+2i} = -\frac{11 + 2i}{3}.$$

- 2 Now we have obtain the **SAME** answer as we had calculated along two smooth curves.

Example-contd

1

$$\int_C z^2 dz = \left[\frac{z^3}{3} \right]_0^{1+2i} = -\frac{11 + 2i}{3}.$$

- 2 Now we have obtain the **SAME** answer as we had calculated along two smooth curves.
- 3 Good news. We do not calculate line integration now.

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from i to $-i$?

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from i to $-i$?

- 1 $z = 0$ lies on the straight line.

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from i to $-i$?

- 1 $z = 0$ lies on the straight line.
- 2 Here $f(z) = 1/z$ is NOT continuous at $z = 0$.

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from i to $-i$?

- 1 $z = 0$ lies on the straight line.
- 2 Here $f(z) = 1/z$ is NOT continuous at $z = 0$.
- 3 We cannot apply the theorem.

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C |z|^2 dz$, where C is an arc of unit circle $|z| = 1$ traversed in the clockwise direction with initial point -1 and final point i ?

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C |z|^2 dz$, where C is an arc of unit circle $|z| = 1$ traversed in the clockwise direction with initial point -1 and final point i ?

- 1 Here $f(z) = |z|^2$ does not have any **anti-derivative**.

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C |z|^2 dz$, where C is an arc of unit circle $|z| = 1$ traversed in the clockwise direction with initial point -1 and final point i ?

- 1 Here $f(z) = |z|^2$ does not have any anti-derivative.
HOW????

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C |z|^2 dz$, where C is an arc of unit circle $|z| = 1$ traversed in the clockwise direction with initial point -1 and final point i ?

- 1 Here $f(z) = |z|^2$ does not have any **anti-derivative**.
HOW????
- 2 We **cannot** apply the theorem.

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from $-i$ to 1 and then from 1 to i ?

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from $-i$ to 1 and then from 1 to i ?

- 1 We construct a domain $D = \mathbb{C} \setminus \{x + iy : y = 0, x \leq 0\}$.

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from $-i$ to 1 and then from 1 to i ?

- 1 We construct a domain $D = \mathbb{C} \setminus \{x + iy : y = 0, x \leq 0\}$.
- 2 We know $\frac{d}{dz} \text{Log } z = \frac{1}{z}$ for all $z \in D$.

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from $-i$ to 1 and then from 1 to i ?

- 1 We construct a domain $D = \mathbb{C} \setminus \{x + iy : y = 0, x \leq 0\}$.
- 2 We know $\frac{d}{dz} \text{Log } z = \frac{1}{z}$ for all $z \in D$.
- 3 Obviously, C lies in D .

Application of the theorem

Example

Can you apply the theorem to evaluate $\int_C \frac{1}{z} dz$, where C is the straight line from $-i$ to 1 and then from 1 to i ?

- 1 We construct a domain $D = \mathbb{C} \setminus \{x + iy : y = 0, x \leq 0\}$.
- 2 We know $\frac{d}{dz} \text{Log } z = \frac{1}{z}$ for all $z \in D$.
- 3 Obviously, C lies in D .
- 4 Therefore,

$$\int_C \frac{1}{z} dz = [\text{Log } z]_{-i}^i = \text{Log}(i) - \text{Log}(-i) = i\frac{\pi}{2} - i\frac{-\pi}{2} = i\pi.$$

Arc length of a contour

Arc length of a contour

The arc length of a contour $C : \gamma(t) = x(t) + iy(t)$, $a \leq t \leq b$ is given by

Arc length of a contour

The arc length of a contour $C : \gamma(t) = x(t) + iy(t)$, $a \leq t \leq b$ is given by

$$\begin{aligned} L(\gamma) &= \int_a^b \sqrt{(x'(t))^2 + (y'(t))^2} dt \\ &= \int_a^b |\gamma'(t)| dt \end{aligned}$$

ML formula or ML inequality

ML formula or ML inequality

If L is the arc length of a contour $C : \gamma(t)$, $a \leq t \leq b$ and M is a positive number such that $|f(z)| \leq M$ for all $z \in C = \gamma$ ($f(z)$ is a piece-wise continuous function), then

ML formula or ML inequality

If L is the arc length of a contour $C : \gamma(t)$, $a \leq t \leq b$ and M is a positive number such that $|f(z)| \leq M$ for all $z \in C = \gamma$ ($f(z)$ is a piece-wise continuous function), then

$$\left| \int_C f(z) dz \right| \leq ML.$$

Proof

Proof

$$\begin{aligned}\left| \int_C f(z) dz \right| &= \left| \int_a^b f(\gamma(t)) \gamma'(t) dt \right| \\ &\leq \int_a^b |f(\gamma(t))| |\gamma'(t)| dt \\ &\leq \int_a^b M |\gamma'(t)| dt \\ &= M \int_a^b |\gamma'(t)| dt \\ &= ML\end{aligned}$$

Proof

$$\begin{aligned}\left| \int_C f(z) dz \right| &= \left| \int_a^b f(\gamma(t)) \gamma'(t) dt \right| \\ &\leq \int_a^b |f(\gamma(t))| |\gamma'(t)| dt \\ &\leq \int_a^b M |\gamma'(t)| dt \\ &= M \int_a^b |\gamma'(t)| dt \\ &= ML\end{aligned}$$

- 1 It give some **estimate** of an upper bound of the modulus of the integration.

Proof

$$\begin{aligned}\left| \int_C f(z) dz \right| &= \left| \int_a^b f(\gamma(t)) \gamma'(t) dt \right| \\ &\leq \int_a^b |f(\gamma(t))| |\gamma'(t)| dt \\ &\leq \int_a^b M |\gamma'(t)| dt \\ &= M \int_a^b |\gamma'(t)| dt \\ &= ML\end{aligned}$$

- 1 It give some **estimate** of an upper bound of the modulus of the integration.
- 2 We can prove some **important theorems** using this ML-inequality.

Important Integral Theorems

We present three important theorems

Important Integral Theorems

We present three important theorems

1 Cauchy's Integral Theorem

Important Integral Theorems

We present three important theorems

- 1 Cauchy's Integral Theorem
- 2 The Cauchy-Goursat Theorem

Important Integral Theorems

We present three important theorems

- 1 Cauchy's Integral Theorem
- 2 The Cauchy-Goursat Theorem
- 3 Cauchy's Integral Formula

Cauchy's Integral Theorem: by Cauchy in 1846

Cauchy's Integral Theorem: by Cauchy in 1846

Let $f(z)$ be a function such that

Cauchy's Integral Theorem: by Cauchy in 1846

Let $f(z)$ be a function such that

- 1 $f(z)$ is analytic **on and inside** a simple closed contour C , and

Cauchy's Integral Theorem: by Cauchy in 1846

Let $f(z)$ be a function such that

- 1 $f(z)$ is analytic on and inside a simple closed contour C , and
- 2 $f'(z)$ be continuous on and inside C .

Then

$$\int_C f(z) dz = 0.$$

Proof

Review of Green's theorem

- 1 Let C be a piece-wise, simple closed curve enclosing a region R .

Proof

Review of Green's theorem

- 1 Let C be a piece-wise, simple closed curve enclosing a region R .
- 2 Let $M(x, y)$ and $N(x, y)$ have continuous **first order partial derivatives** in an open set containing R .

Proof

Review of Green's theorem

- 1 Let C be a piece-wise, simple closed curve enclosing a region R .
- 2 Let $M(x, y)$ and $N(x, y)$ have continuous **first order partial derivatives** in an open set containing R .

Then

$$\int_C (M dx + N dy) = \iint_R (N_x - M_y) dxdy,$$

where the line integral is taken in **counterclockwise sense**.

Proof-Contd.

- 1 We know (alternative approach can be found in many books)

$$\int_C f(z) dz = \int_C (u dx - v dy) + i \int_C (v dx + u dy)$$

Proof-Contd.

- 1 We know (alternative approach can be found in many books)

$$\int_C f(z) dz = \int_C (u dx - v dy) + i \int_C (v dx + u dy)$$

- 2 In view of **Green's theorem**

$$\int_C f(z) dz = \iint_D (-v_x - u_y) dx dy + i \iint_D (u_x - v_y) dx dy$$

Proof-Contd.

- 1 We know (alternative approach can be found in many books)

$$\int_C f(z) dz = \int_C (udx - vdy) + i \int_C (vdx + udy)$$

- 2 In view of **Green's theorem**

$$\int_C f(z) dz = \iint_D (-v_x - u_y) dx dy + i \iint_D (u_x - v_y) dx dy$$

- 3 Because of **analyticity**, CR equations are satisfied on D .

- 4 Hence

$$\int_C f(z) dz = 0 + i0 = 0.$$

Cauchy-Goursat theorem

Let $f(z)$ be a function such that $f(z)$ is analytic on and inside a simple closed contour C . Then

$$\int_C f(z) dz = 0.$$

Information-Cauchy-Goursat theorem

Information-Cauchy-Goursat theorem

- 1 What are the **difference** between Cauchy's integral theorem and Cauchy-Goursat theorem.

Information-Cauchy-Goursat theorem

- 1 What are the **difference** between Cauchy's integral theorem and Cauchy-Goursat theorem.
- 2 Note that **continuity** of $f'(z)$ is not involved in Cauchy's integral theorem.

Information-Cauchy-Goursat theorem

- 1 What are the **difference** between Cauchy's integral theorem and Cauchy-Goursat theorem.
- 2 Note that **continuity** of $f'(z)$ is not involved in Cauchy's integral theorem.
- 3 Goursat (French mathematician in 1903) proved Cauchy's theorem without **continuity** of $f'(z)$.

Information-Cauchy-Goursat theorem

- 1 What are the **difference** between Cauchy's integral theorem and Cauchy-Goursat theorem.
- 2 Note that **continuity** of $f'(z)$ is not involved in Cauchy's integral theorem.
- 3 Goursat (French mathematician in 1903) proved Cauchy's theorem without **continuity** of $f'(z)$.
- 4 It is also called Goursat theorem.

Information-Cauchy-Goursat theorem

- 1 What are the **difference** between Cauchy's integral theorem and Cauchy-Goursat theorem.
- 2 Note that **continuity** of $f'(z)$ is not involved in Cauchy's integral theorem.
- 3 Goursat (French mathematician in 1903) proved Cauchy's theorem without **continuity** of $f'(z)$.
- 4 It is also called Goursat theorem.
- 5 Cauchy-Goursat theorem is a **stronger version** of Cauchy's integral theorem.

Information-Cauchy-Goursat theorem

- 1 What are the **difference** between Cauchy's integral theorem and Cauchy-Goursat theorem.
- 2 Note that **continuity** of $f'(z)$ is not involved in Cauchy's integral theorem.
- 3 Goursat (French mathematician in 1903) proved Cauchy's theorem without **continuity** of $f'(z)$.
- 4 It is also called Goursat theorem.
- 5 Cauchy-Goursat theorem is a **stronger version** of Cauchy's integral theorem.
- 6 We use this theorem to **solve problems**.

Alternative form of Cauchy-Goursat theorem

Alternative form of Cauchy-Goursat theorem

Let $f(z)$ be a function analytic throughout a **simply connected domain** D and C be a simple closed contour lying completely inside D . Then

$$\int_C f(z) dz = 0.$$

Alternative form of Cauchy-Goursat theorem

Let $f(z)$ be a function analytic throughout a **simply connected domain** D and C be a simple closed contour lying completely inside D . Then

$$\int_C f(z) dz = 0.$$




Figure:

Examples- Application of Cauchy-Goursat theorem

Examples- Application of Cauchy-Goursat theorem

Evaluate the following integrals where C denotes the circle of unit radius with center at zero.

1 $\int_C \sin z.$

Examples- Application of Cauchy-Goursat theorem

Evaluate the following integrals where C denotes the circle of unit radius with center at zero.

1 $\int_C \sin z.$

2 $\int_C e^{z^n}.$

Examples- Application of Cauchy-Goursat theorem

Evaluate the following integrals where C denotes the circle of unit radius with center at zero.

1 $\int_C \sin z.$

2 $\int_C e^{z^n}.$

3 $\int_C \frac{e^{iz^2}}{z^2 + 4} dz.$

Examples- Application of Cauchy-Goursat theorem

Examples- Application of Cauchy-Goursat theorem

- 1 Can you apply Cauchy-Goursat's theorem to evaluate $\int_C \operatorname{cosec}^2 z$, where C denotes the circle of unit radius with center at zero.

Examples- Application of Cauchy-Goursat theorem

- 1 Can you apply Cauchy-Goursat's theorem to evaluate $\int_C \operatorname{cosec}^2 z$, where C denotes the circle of **unit radius with center at zero**.
- 2 Determine the value of the integration.

Solution

Solution

- 1 $f(z)$ is not analytic inside the closed counter as $f(z)$ is **not** analytic at $z = 0$.

Solution

- 1 $f(z)$ is not analytic inside the closed counter as $f(z)$ is **not** analytic at $z = 0$.
- 2 So we **cannot** apply Cauchy-Goursat's theorem to evaluate $\int_C \operatorname{cosec}^2 z$.

Solution

- 1 $f(z)$ is not analytic inside the closed counter as $f(z)$ is **not** analytic at $z = 0$.
- 2 So we **cannot** apply Cauchy-Goursat's theorem to evaluate $\int_C \operatorname{cosec}^2 z$.
- 3 Note that $f(z)$ is analytic at $\mathbb{C} \setminus \{n\pi : n \in \mathbb{Z}\}$.

Solution

- 1 $f(z)$ is not analytic inside the closed counter as $f(z)$ is **not analytic at $z = 0$** .
- 2 So we **cannot** apply Cauchy-Goursat's theorem to evaluate $\int_C \operatorname{cosec}^2 z$.
- 3 Note that $f(z)$ is analytic at $\mathbb{C} \setminus \{n\pi : n \in \mathbb{Z}\}$.
- 4 Consider the annulus $D = \{z : 0 < |z| < \pi\}$. This is our **modified domain**.

Solution

Solution

- 1 $f(z)$ is continuous on the modified domain.

Solution

- 1 $f(z)$ is continuous on the modified domain.
- 2 The contour C lies entirely inside D .

Solution

- 1 $f(z)$ is continuous on the modified domain.
- 2 The contour C lies entirely inside D .
- 3 Also $F'(z) = f(z)$ on D , where $F(z) = -\cot z$.

Solution

- 1 $f(z)$ is continuous on the modified domain.
- 2 The contour C lies entirely inside D .
- 3 Also $F'(z) = f(z)$ on D , where $F(z) = -\cot z$.
- 4 Since C is a closed contour with starting and end points z_1 and z_2 . Then $z_1 = z_2$.
- 5 Hence $\int_C \operatorname{cosec}^2 z = F(z_2) - F(z_1) = 0$.

Complex Analysis

LECTURE 11

August 22, 2023

Deformation of simple closed counter

Deformation of simple closed counter

- 1 Let C_1 be a **simple closed contour** with **ugly shape**.

Deformation of simple closed counter

- 1 Let C_1 be a **simple closed contour** with **ugly shape**.
- 2 Then the ugly contour **can be deformed** into another closed contour C_2 (lies inside C_1) with a known shape such as triangle, rectangle, or circle, etc..

Deformation of simple closed counter

- 1 Let C_1 be a **simple closed contour** with **ugly shape**.
- 2 Then the ugly contour **can be deformed** into another closed contour C_2 (lies inside C_1) with a known shape such as triangle, rectangle, or circle, etc..
- 3 **Visualize** the statement.

Deformation of simple closed counter

- 1 Let C_1 be a **simple closed contour** with **ugly shape**.
- 2 Then the ugly contour **can be deformed** into another closed contour C_2 (lies inside C_1) with a known shape such as triangle, rectangle, or circle, etc..
- 3 **Visualize** the statement.
- 4 What is the **advantage/application** of such deformation?

Trouble with ugly contour

Trouble with ugly contour

- 1 Let C be a **simple closed contour** and with **ugly shape**.

Trouble with ugly contour

- 1 Let C be a **simple closed contour** and with **ugly shape**.
- 2 Let $f(z)$ be an analytic function on the contour and the domain enclosed by C .

Trouble with ugly contour

- 1 Let C be a **simple closed contour** and with **ugly shape**.
- 2 Let $f(z)$ be an analytic function on the contour and the domain enclosed by C .
- 3 Then by Cauchy-Goursat theorem we can obtain

$$\int_C f(z) dz = 0.$$

Trouble with ugly contour

- 1 Let C be a **simple closed contour** and with **ugly shape**.
- 2 Let $f(z)$ be an analytic function on the contour and the domain enclosed by C .
- 3 Then by Cauchy-Goursat theorem we can obtain

$$\int_C f(z) dz = 0.$$

- 4 We do not need to think about the **parametric representation of the ugly contour**.
- 5 Thus we have **NO trouble**.

Trouble with ugly contour

- 1 Let C be a **simple closed contour** and with **ugly shape**.
- 2 Let $f(z)$ be an analytic function on the contour and the domain enclosed by C .
- 3 Then by Cauchy-Goursat theorem we can obtain

$$\int_C f(z) dz = 0.$$

- 4 We do not need to think about the **parametric representation of the ugly contour**.
- 5 Thus we have **NO trouble**.
- 6 OK, think about the integration of a function which is **NOT analytic** on the domain enclosed by the ugly contour.

Trouble with ugly contour

- 1 Let C be a **simple closed contour** and with **ugly shape**.
- 2 Let $f(z)$ be an analytic function on the contour and the domain enclosed by C .
- 3 Then by Cauchy-Goursat theorem we can obtain

$$\int_C f(z) dz = 0.$$

- 4 We do not need to think about the **parametric representation of the ugly contour**.
- 5 Thus we have **NO trouble**.
- 6 OK, think about the integration of a function which is **NOT analytic** on the domain enclosed by the ugly contour.
- 7 Thus, we **CANNOT** apply Cauchy-Goursat theorem.

Trouble with ugly contour

Trouble with ugly contour

- 1 For simplicity, we consider the function $f(z) = 1/z$ and the contour encloses a domain **including origin**.

Trouble with ugly contour

- 1 For simplicity, we consider the function $f(z) = 1/z$ and the contour encloses a domain **including origin**.
- 2 Thus, we **CANNOT** apply Cauchy-Goursat theorem.

Trouble with ugly contour

- 1 For simplicity, we consider the function $f(z) = 1/z$ and the contour encloses a domain **including origin**.
- 2 Thus, we **CANNOT** apply Cauchy-Goursat theorem.
- 3 Thus, we need to calculate the integration **DIRECTLY** along the curve.

Trouble with ugly contour

- 1 For simplicity, we consider the function $f(z) = 1/z$ and the contour encloses a domain **including origin**.
- 2 Thus, we **CANNOT** apply Cauchy-Goursat theorem.
- 3 Thus, we need to calculate the integration **DIRECTLY** along the curve.
- 4 So **parametric representation of the ugly contour** is required.
- 5 This makes our **life terrible**.

Trouble with ugly contour

Trouble with ugly contour

- 1 It would be great if we can **replace the ugly contour** by a one with simple shape (such as circle) then the direct integration easier.

Trouble with ugly contour

- 1 It would be great if we can **replace the ugly contour** by a one with simple shape (such as circle) then the direct integration easier.
- 2 Now we learn the advantage of computing the direct integration over the contour with **a simple shape**.

Trouble with ugly contour

- 1 It would be great if we can **replace the ugly contour** by a one with simple shape (such as circle) then the direct integration easier.
- 2 Now we learn the advantage of computing the direct integration over the contour with **a simple shape**.
- 3 Follow the **next theorem**.

Theorem on deformation contour

Theorem on deformation contour

Theorem

Theorem on deformation contour

Theorem

Let C_1 and C_2 be two simple closed **positively** oriented contours such that C_2 lies **interior** to C_1 . If $f(z)$ is analytic in a domain D that contains both the contours and the region between them, then

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

Theorem on deformation contour

Theorem

Let C_1 and C_2 be two simple closed **positively** oriented contours such that C_2 lies **interior** to C_1 . If $f(z)$ is analytic in a domain D that contains both the contours and the region between them, then

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

- 1 Both the contours have **positive orientation**.

Theorem on deformation contour

Theorem

Let C_1 and C_2 be two simple closed **positively** oriented contours such that C_2 lies **interior** to C_1 . If $f(z)$ is analytic in a domain D that contains both the contours and the region between them, then

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

- 1 Both the contours have **positive orientation**.
- 2 It is not necessary that the function needs to be analytic inside the domain enclosed by C_2 .

Theorem on deformation contour

Theorem

Let C_1 and C_2 be two simple closed **positively** oriented contours such that C_2 lies **interior** to C_1 . If $f(z)$ is analytic in a domain D that contains both the contours and the region between them, then

$$\int_{C_1} f(z) dz = \int_{C_2} f(z) dz$$

- 1 Both the contours have **positive orientation**.
- 2 It is not necessary that the function needs to be analytic inside the domain enclosed by C_2 .
- 3 In practical situations (examples), $f(z)$ is **NOT** analytic inside the domain enclosed by C_2 .

Proof-outline

Proof-outline




Figure:

Proof-outline

Proof-outline




Figure: $f(z)$ is analytic on the curve and shaded region. It may not be analytic on the white domain enclosed by C_2 .

Proof-outline




Figure: Integration along $C_{1,0} + L_1 + C_{2,0} + L_2$ is 0 by Cauchy-Goursat theorem.

Proof-outline

$C_{1,1}$




Figure: Integration along $C_{1,1} - L_2 + C_{2,1} - L_1$ is 0 by Cauchy-Goursat theorem.

1

$$\int_{C_{1,0}+L_1+C_{2,0}+L_2} = 0$$

1

$$\int_{C_{1,0}+L_1+C_{2,0}+L_2} = 0$$

2

$$\int_{C_{1,1}-L_2+C_{2,1}-L_1} = 0$$

1

$$\int_{C_{1,0}+L_1+C_{2,0}+L_2} = 0$$

2

$$\int_{C_{1,1}-L_2+C_{2,1}-L_1} = 0$$

3 Adding together

$$\int_{\textcolor{red}{C_{1,0}+C_{1,1}}} + \int_{C_{2,0}+C_{2,1}} = 0$$

1

$$\int_{C_{1,0}+L_1+C_{2,0}+L_2} = 0$$

2

$$\int_{C_{1,1}-L_2+C_{2,1}-L_1} = 0$$

3 Adding together

$$\int_{\textcolor{red}{C_{1,0}} + \textcolor{red}{C_{1,1}} + C_{2,0} + C_{2,1}} = 0$$

4 Adding together

$$\int_{\textcolor{red}{C_1} - C_2} = 0$$

1 Therefore,

$$\int_{C_1} = \int_{C_2}$$

1 Therefore,

$$\int_{C_1} = \int_{C_2}$$

2 Thus we can obtain the same integration value along both the contours.

1 Therefore,

$$\int_{C_1} = \int_{C_2}$$

- 2 Thus we can obtain the same integration value along both the contours.
- 3 In practical situation we consider C_1 with simple form such as CIRCLE.

1 Therefore,

$$\int_{C_1} = \int_{C_2}$$

- 2 Thus we can obtain the same integration value along both the contours.
- 3 In practical situation we consider C_1 with simple form such as CIRCLE.
- 4 Note that we have applied Cauchy-Goursat theorem TWICE.

Example

Example

1 Compute

$$\int_{C_{\text{ugly}}} \frac{1}{z} dz, \text{ where the contour encloses origin.}$$

Example

- 1 Compute

$$\int_{C_{\text{ugly}}} \frac{1}{z} dz, \text{ where the contour encloses origin.}$$

- 2 We deform the ugly contour into a circle $|z| = r$ whose parametric equation is $z = re^{i\theta}$, $0 \leq \theta \leq 2\pi$.

Example

- 1 Compute

$$\int_{C_{\text{ugly}}} \frac{1}{z} dz, \text{ where the contour encloses origin.}$$

- 2 We deform the ugly contour into a circle $|z| = r$ whose parametric equation is $z = re^{i\theta}$, $0 \leq \theta \leq 2\pi$.
- 3 r will be chosen in such a way that the circle lies inside the ugly curve.

Example

- 1 Compute


$$\int_{C_{\text{ugly}}} \frac{1}{z} dz, \text{ where the contour encloses origin.}$$

- 2 We deform the ugly contour into a circle $|z| = r$ whose parametric equation is $z = re^{i\theta}$, $0 \leq \theta \leq 2\pi$.
- 3 r will be chosen in such a way that the circle lies inside the ugly curve.
- 4 Direct integration yield $2\pi i$ for any r .

ALTERNATIVE Proof-outline

ALTERNATIVE Proof-outline

We can prove the theorem by applying cauchy-Goursat theorem only **ONE** time.



ALTERNATIVE Proof-outline

ALTERNATIVE Proof-outline




Figure: One one simply connected domain (shaded portion). Integration along $C_1 + L_1 - C_2 + L_2$ is 0 by Cauchy-Goursat theorem.

- 1 We assume that L_1 and L_2 are parallel lines and close to each other.

- 1 We assume that L_1 and L_2 are parallel lines and close to each other.
- 2 Thus, $\int_{L_1} = - \int_{-L_2}$

- 1 We assume that L_1 and L_2 are parallel lines and close to each other.
- 2 Thus, $\int_{L_1} = - \int_{-L_2}$
- 3 Because of analyticity, integrations over the whole contours C_1 and C_2 do not change value if an infinitesimal arch is removed.
- 4 Hence,

$$\int_{C_1+L_1} - C_2+L_2 = 0$$

- 1 We assume that L_1 and L_2 are parallel lines and close to each other.
- 2 Thus, $\int_{L_1} = - \int_{-L_2}$
- 3 Because of analyticity, integrations over the whole contours C_1 and C_2 do not change value if an infinitesimal arch is removed.
- 4 Hence,

$$\int_{C_1+L_1} - C_2+L_2 = 0$$

- 5 Finally,

$$\int_{C_1} = \int_{C_2}$$

Cauchy-Goursat's theorem for multiply connected domain

Cauchy-Goursat's theorem for multiply connected domain

Theorem

Cauchy-Goursat's theorem for multiply connected domain

Theorem

Let C be a positively oriented simple closed contour and C_k , $k = 1, 2, \dots, n$ denote a finite number of positively oriented simple closed contours all lying wholly within C , but each C_k lies in the exterior of every other whose interior have no points in common. If a function f is analytic throughout the closed region D consisting of all points within and on C except for the points interior to each C_k .

Cauchy-Goursat's theorem for multiply connected domain

Theorem

Let C be a positively oriented simple closed contour and C_k , $k = 1, 2, \dots, n$ denote a finite number of positively oriented simple closed contours all lying wholly within C , but each C_k lies in the exterior of every other whose interior have no points in common. If a function f is analytic throughout the closed region D consisting of all points within and on C except for the points interior to each C_k . Then

$$\int_C f(z) dz = \int_{C_1} f(z) dz + \int_{C_2} f(z) dz + \cdots + \int_{C_n} f(z) dz.$$

Proof:- for doubly connected domain

Proof:- for doubly connected domain




Figure: Think about the proof with for doubly connected domain

Proof:- for doubly connected domain

Proof:- for doubly connected domain




Figure: Apply the concepts from earlier theorems

Example with multiply connected domain

Example with multiply connected domain

1 $\int_C \frac{1}{z^2 + 1} dz$ where $C := |z| = 3$

Example with multiply connected domain

- 1 $\int_C \frac{1}{z^2 + 1} dz$ where $C := |z| = 3$
- 2 The function is not analytic at $z = i$ and $z = -i$ which are inside C

Example with multiply connected domain

- 1 $\int_C \frac{1}{z^2 + 1} dz$ where $C := |z| = 3$
- 2 The function is not analytic at $z = i$ and $z = -i$ which are inside C
- 3 We make two circles $C_1 := |z - i| = 1/2$ and $C_2 := |z + i| = 1/2$.

Example with multiply connected domain

- 1 $\int_C \frac{1}{z^2 + 1} dz$ where $C := |z| = 3$
- 2 The function is not analytic at $z = i$ and $z = -i$ which are inside C
- 3 We make two circles $C_1 := |z - i| = 1/2$ and $C_2 := |z + i| = 1/2$.
- 4 Then by Cauchy-Goursat theorem for doubly connected domain

$$\int_C \frac{1}{z^2 + 1} dz = \int_{C_1} \frac{1}{z^2 + 1} dz + \int_{C_2} \frac{1}{z^2 + 1} dz$$

Example with multiply connected domain

Example with multiply connected domain

1

$$\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} \left[\int_{C_1} \frac{1}{z - i} dz - \int_{C_1} \frac{1}{z + i} dz \right]$$

Example with multiply connected domain

1

$$\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} \left[\int_{C_1} \frac{1}{z - i} dz - \int_{C_1} \frac{1}{z + i} dz \right]$$

2 Now $\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} [2\pi i - 0]$

Example with multiply connected domain

1

$$\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} \left[\int_{C_1} \frac{1}{z - i} dz - \int_{C_1} \frac{1}{z + i} dz \right]$$

2 Now $\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} [2\pi i - 0]$

3 $2\pi i$ is calculated by taking $z = (1/2)e^{i\theta}$ and second integral yield zero as the integrand is analytic on and inside C_1 .

Example with multiply connected domain

1

$$\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} \left[\int_{C_1} \frac{1}{z - i} dz - \int_{C_1} \frac{1}{z + i} dz \right]$$

2 Now $\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} [2\pi i - 0]$

3 $2\pi i$ is calculated by taking $z = (1/2)e^{i\theta}$ and second integral yield zero as the integrand is analytic on and inside C_1 .

4 Similarly, $\int_{C_2} \frac{1}{z^2 + 1} dz = \frac{1}{2i} [0 - 2\pi i]$

Example with multiply connected domain

1

$$\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} \left[\int_{C_1} \frac{1}{z - i} dz - \int_{C_1} \frac{1}{z + i} dz \right]$$

2 Now $\int_{C_1} \frac{1}{z^2 + 1} dz = \frac{1}{2i} [2\pi i - 0]$

3 $2\pi i$ is calculated by taking $z = (1/2)e^{i\theta}$ and second integral yield zero as the integrand is analytic on and inside C_1 .

4 Similarly, $\int_{C_2} \frac{1}{z^2 + 1} dz = \frac{1}{2i} [0 - 2\pi i]$

5 Finally, $\int_C \frac{1}{z^2 + 1} dz = \frac{1}{2i} [2\pi i - 0] + \frac{1}{2i} [0 - 2\pi i] = 0.$

Example with multiply connected domain

Example with multiply connected domain

- 1 OK, we are very much *successfully* to solve the previous problem.

Example with multiply connected domain

- 1 OK, we are very much **successfully** to solve the previous problem.
- 2 What about $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$?

Example with multiply connected domain

- 1 OK, we are very much **successfully** to solve the previous problem.
- 2 What about $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$?
- 3 Ok, the function is not analytic at $z = 2$ and $z = 3$ which are inside C

Example with multiply connected domain

- 1 OK, we are very much **successfully** to solve the previous problem.
- 2 What about $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$?
- 3 Ok, the function is not analytic at $z = 2$ and $z = 3$ which are inside C
- 4 We make two circles $C_1 := |z - 2| = 1/2$ and $C_2 := |z - 3| = 1/2$.

Example with multiply connected domain

- 1 OK, we are very much **successfully** to solve the previous problem.
- 2 What about $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$?
- 3 Ok, the function is not analytic at $z = 2$ and $z = 3$ which are inside C
- 4 We make two circles $C_1 := |z - 2| = 1/2$ and $C_2 := |z - 3| = 1/2$.
- 5 **Difficult** to calculate by applying parametric curve due the presence of $\sin(\pi z^2)$ and $\cos(\pi z^2)$.

Example with multiply connected domain

- 1 OK, we are very much **successfully** to solve the previous problem.
- 2 What about $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$?
- 3 Ok, the function is not analytic at $z = 2$ and $z = 3$ which are inside C
- 4 We make two circles $C_1 := |z - 2| = 1/2$ and $C_2 := |z - 3| = 1/2$.
- 5 **Difficult** to calculate by applying parametric curve due the presence of $\sin(\pi z^2)$ and $\cos(\pi z^2)$.
- 6 But we can find an **easy way** follows from the next theorem.

Cauchy's integral formula

Cauchy's integral formula

Statement

Cauchy's integral formula

Statement

Let the function $f(z)$ be analytic on and inside a positively oriented simple closed contour C and z_0 be **any point** inside C .

Cauchy's integral formula

Statement

Let the function $f(z)$ be analytic on and inside a positively oriented simple closed contour C and z_0 be **any point** inside C .

Then

$$f(z_0) = \frac{1}{2\pi i} \int_C \frac{f(z)}{z - z_0} dz. \quad (3)$$

Example

Example

- 1 Determine $\int_C \frac{\cos(e^z)}{z} dz$ where $C := |z| = 1$.

Example

- 1 Determine $\int_C \frac{\cos(e^z)}{z} dz$ where $C := |z| = 1$.
- 2 It is difficult to solve by making use of $z = e^{i\theta}$.

Example

- 1 Determine $\int_C \frac{\cos(e^z)}{z} dz$ where $C := |z| = 1$.
- 2 It is difficult to solve by making use of $z = e^{i\theta}$.
- 3 We apply Cauchy's integral formula

Example

- 1 Determine $\int_C \frac{\cos(e^z)}{z} dz$ where $C := |z| = 1$.
- 2 It is difficult to solve by making use of $z = e^{i\theta}$.
- 3 We apply Cauchy's integral formula
- 4 Solution:

$$\int_C \frac{\cos(e^z)}{z} dz = 2\pi i \cos(e^z)|_{z=0} = 2\pi i \cos(1).$$

Example

Example

- 1 Determine $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$.

Example

- 1 Determine $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$.
- 2 The function is not analytic at $z = 2$ and $z = 3$ which are inside C .

Example

- 1 Determine $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$.
- 2 The function is not analytic at $z = 2$ and $z = 3$ which are inside C .
- 3 We make two circles $C_1 := |z - 2| = 1/2$ and $C_2 := |z - 3| = 1/2$.

Example

- 1 Determine $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz$ where $C := |z| = 4$.
- 2 The function is not analytic at $z = 2$ and $z = 3$ which are inside C .
- 3 We make two circles $C_1 := |z - 2| = 1/2$ and $C_2 := |z - 3| = 1/2$.
- 4 Then by Cauchy-Goursat theorem for doubly connected domain

$$\begin{aligned}\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz &= \int_{C_1} \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz \\ &\quad + \int_{C_2} \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz\end{aligned}$$

Example

1 Now

$$\begin{aligned}\int_{C_1} \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z - 2)(z - 3)} dz &= \int_{C_1} \frac{[\sin(\pi z^2) + \cos(\pi z^2)]/(z - 3)}{(z - 2)} dz \\ &= 2\pi i [\sin(\pi z^2) + \cos(\pi z^2)/(z - 3)]|_{z=2} \\ &= -2\pi i [\sin(4\pi) + \cos(4\pi)] = -2\pi i\end{aligned}$$

Example

1 Now

$$\begin{aligned}\int_{C_1} \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z-2)(z-3)} dz &= \int_{C_1} \frac{[\sin(\pi z^2) + \cos(\pi z^2)]/(z-3)}{(z-2)} dz \\ &= 2\pi i [\sin(\pi z^2) + \cos(\pi z^2)]/(z-3)|_{z=2} \\ &= -2\pi i [\sin(4\pi) + \cos(4\pi)] = -2\pi i\end{aligned}$$

2 Similarly,

$$\begin{aligned}\int_{C_2} \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z-2)(z-3)} dz &= \int_{C_1} \frac{[\sin(\pi z^2) + \cos(\pi z^2)]/(z-2)}{(z-3)} dz \\ &= 2\pi i [\sin(\pi z^2) + \cos(\pi z^2)]/(z-2)|_{z=3} \\ &= 2\pi i [\sin(9\pi) + \cos(9\pi)] = -2\pi i\end{aligned}$$

Example

1 Now

$$\begin{aligned}\int_{C_1} \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z-2)(z-3)} dz &= \int_{C_1} \frac{[\sin(\pi z^2) + \cos(\pi z^2)]/(z-3)}{(z-2)} dz \\ &= 2\pi i [\sin(\pi z^2) + \cos(\pi z^2)]/(z-3)|_{z=2} \\ &= -2\pi i [\sin(4\pi) + \cos(4\pi)] = -2\pi i\end{aligned}$$

2 Similarly,

$$\begin{aligned}\int_{C_2} \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z-2)(z-3)} dz &= \int_{C_1} \frac{[\sin(\pi z^2) + \cos(\pi z^2)]/(z-2)}{(z-3)} dz \\ &= 2\pi i [\sin(\pi z^2) + \cos(\pi z^2)]/(z-2)|_{z=3} \\ &= 2\pi i [\sin(9\pi) + \cos(9\pi)] = -2\pi i\end{aligned}$$

3 Finally, $\int_C \frac{\sin(\pi z^2) + \cos(\pi z^2)}{(z-2)(z-3)} dz = -4\pi i.$

Derivative of contour integrals

Derivative of contour integrals

Statement

Derivative of contour integrals

Statement

If f is analytic on a simply connected domain D then f has derivatives of all orders in D (which are then analytic in D) and for any $z_0 \in D$, one has

$$f^n(z_0) = \frac{n!}{2\pi i} \int_C \frac{f(z)}{(z - z_0)^{n+1}} dz, \quad (4)$$

where C is a simple closed contour (oriented counterclockwise) around z_0 in D .

Example

Example

- 1 Determine $\int_C \frac{e^{z^2}}{z^2} dz$ where C is a simple closed curve enclosing origin.

Example

- 1 Determine $\int_C \frac{e^{z^2}}{z^2} dz$ where C is a simple closed curve enclosing origin.
- 2 Solution:

$$\int_C \frac{e^{z^2}}{z^2} dz = \frac{2\pi i}{1!} \frac{d}{dz}(e^{z^2})|_{z=0} = 0.$$

Converse of Cauchy-Goursat's Theorem

Converse of Cauchy-Goursat's Theorem

Problem statement:

Converse of Cauchy-Goursat's Theorem

Problem statement: Suppose $f(z)$ is a function with domain D such that

$$\int_C f(z) dz = 0$$

for every closed contour C lying inside D . Can we conclude from this that f is analytic in D ?

Converse of Cauchy-Goursat's Theorem

Problem statement: Suppose $f(z)$ is a function with domain D such that

$$\int_C f(z) dz = 0$$

for every closed contour C lying inside D . Can we conclude from this that f is analytic in D ?

1 Answer is

Converse of Cauchy-Goursat's Theorem

Problem statement: Suppose $f(z)$ is a function with domain D such that

$$\int_C f(z) dz = 0$$

for every closed contour C lying inside D . Can we conclude from this that f is analytic in D ?

- 1 Answer is NO.

Converse of Cauchy-Goursat's Theorem

Problem statement: Suppose $f(z)$ is a function with domain D such that

$$\int_C f(z) dz = 0$$

for every closed contour C lying inside D . Can we conclude from this that f is analytic in D ?

- 1 Answer is NO.
- 2 Consider the domain $D := \{z : |z| < 1\}$

Converse of Cauchy-Goursat's Theorem

Problem statement: Suppose $f(z)$ is a function with domain D such that

$$\int_C f(z) dz = 0$$

for every closed contour C lying inside D . Can we conclude from this that f is analytic in D ?

- 1 Answer is NO.
- 2 Consider the domain $D := \{z : |z| < 1\}$

$$3 f(z) = \begin{cases} z & \text{if } z \in D \setminus \{\frac{1}{2}\} \\ 1 & \text{if } z = \frac{1}{2} \end{cases}.$$

Converse of Cauchy-Goursat's Theorem

Problem statement: Suppose $f(z)$ is a function with domain D such that

$$\int_C f(z) dz = 0$$

for every closed contour C lying inside D . Can we conclude from this that f is analytic in D ?

- 1 Answer is NO.
- 2 Consider the domain $D := \{z : |z| < 1\}$
- 3 $f(z) = \begin{cases} z & \text{if } z \in D \setminus \{\frac{1}{2}\} \\ 1 & \text{if } z = \frac{1}{2} \end{cases}$.
- 4 For every closed contour, one can see:

$$\int_C f(z) dz = 0.$$

Converse of Cauchy-Goursat's Theorem

Problem statement: Suppose $f(z)$ is a function with domain D such that

$$\int_C f(z) dz = 0$$

for every closed contour C lying inside D . Can we conclude from this that f is analytic in D ?

- 1 Answer is NO.
- 2 Consider the domain $D := \{z : |z| < 1\}$
- 3 $f(z) = \begin{cases} z & \text{if } z \in D \setminus \{\frac{1}{2}\} \\ 1 & \text{if } z = \frac{1}{2} \end{cases}$.
- 4 For every closed contour, one can see:

$$\int_C f(z) dz = 0.$$

- 5 But, note that $f(z)$ is NOT analytic in D .

Morera's Theorem

Morera's Theorem

Statement:

Morera's Theorem

Statement: Suppose $f(z)$ is continuous inside a simply connected domain D and

$$\int_C f(z) dz = 0,$$

for any simple closed contour C lying inside D . Then $f(z)$ is analytic throughout D .

Morera's Theorem

Statement: Suppose $f(z)$ is continuous inside a simply connected domain D and

$$\int_C f(z) dz = 0,$$

for any simple closed contour C lying inside D . Then $f(z)$ is analytic throughout D .

- 1 We say that Morera's theorem is **partially** converse of Cauchy-Goursat's theorem.

Morera's Theorem

Statement: Suppose $f(z)$ is continuous inside a simply connected domain D and

$$\int_C f(z) dz = 0,$$

for any simple closed contour C lying inside D . Then $f(z)$ is analytic throughout D .

- 1 We say that Morera's theorem is **partially** converse of Cauchy-Goursat's theorem.
- 2 Why **partially**?

Morera's Theorem

Statement: Suppose $f(z)$ is continuous inside a simply connected domain D and

$$\int_C f(z) dz = 0,$$

for any simple closed contour C lying inside D . Then $f(z)$ is analytic throughout D .

- 1 We say that Morera's theorem is **partially** converse of Cauchy-Goursat's theorem.
- 2 Why **partially**?
- 3 Because: Instead of the function $f(z)$, we have imposed continuity property on $f(z)$ on the domain.

Consequences of Cauchy's Integral Formula

We discuss about three consequences:

1 Cauchy Estimates

Consequences of Cauchy's Integral Formula

We discuss about three consequences:

- 1 Cauchy Estimates
- 2 Liouville's Theorem

Consequences of Cauchy's Integral Formula

We discuss about three consequences:

- 1 Cauchy Estimates
- 2 Liouville's Theorem
- 3 Fundamental Theorem of Algebra

Cauchy Estimates

Let $f(z)$ be analytic on and inside the circle $C : |z - z_0| = r$. If $|f(z)| \leq M$ for all $z \in C$, then for all $n \geq 0$,

$$|f^{(n)}(z_0)| \leq \frac{n!M}{r^n}.$$

Proof

Liouville's Theorem

Liouville's Theorem: Bounded entire function must be a constant function.

Proof

Fundamental Theorem of Algebra

Before we present the **Fundamental Theorem of Algebra** we prove the following result.

Fundamental Theorem of Algebra

Before we present the **Fundamental Theorem of Algebra** we prove the following result.

Result: $|P(z)| \rightarrow \infty$ as $|z| \rightarrow \infty$ for every polynomial $P(z)$ of degree $n \geq 1$.

Proof

Proof

Fundamental Theorem of Algebra

Fundamental Theorem of Algebra: Every polynomial $P(z)$ of degree $n \geq 1$ has a root in \mathbb{C} .

Proof

Fundamental Theorem of Algebra

Corollary: Every polynomial $P(z)$ of degree $n \geq 1$ has **exactly** n (not necessarily distinct) roots in \mathbb{C} .

Proof

Thank You