

MA 204 Numerical Methods

Dr. Debopriya Mukherjee
Lecture-2

January 11, 2024

Contents

- Solution of a nonlinear equation, bisection and secant methods, Newton's method, rate of convergence, solution of a system of nonlinear equations.

Contents

- Solution of a nonlinear equation, bisection and secant methods, Newton's method, rate of convergence, solution of a system of nonlinear equations.
- Interpolation by polynomials, divided differences, error of the interpolating polynomial, piecewise linear and cubic spline interpolation.

Motivation

Nonlinearity

One of the most frequent problem in engineering and science is to find the root(s) of a non-linear equation

$$f(x) = 0. \quad (1)$$

Here,

- $f : [a, b] \rightarrow \mathbb{R}$ is a nonlinear function in x ;
- $f \in C^1[a, b]$;
- Roots are **isolated**.

Root of the equation

Definition 1

Given a nonlinear function $f : [a, b] \rightarrow \mathbb{R}$, find a value of r for which $f(r) = 0$. Such a solution value for r is called a **root** of the equation

$$f(x) = 0,$$

or **zero** of the function f .

Root of the equation

Definition 1

Given a nonlinear function $f : [a, b] \rightarrow \mathbb{R}$, find a value of r for which $f(r) = 0$. Such a solution value for r is called a **root** of the equation

$$f(x) = 0,$$

or **zero** of the function f .

Problem

Given $f : [a, b] \rightarrow \mathbb{R}$ continuous. Find $r \in \mathbb{R}$ such that $f(r) = 0$.

Root of the equation

Definition 1

Given a nonlinear function $f : [a, b] \rightarrow \mathbb{R}$, find a value of r for which $f(r) = 0$. Such a solution value for r is called a **root** of the equation

$$f(x) = 0,$$

or **zero** of the function f .

Problem

Given $f : [a, b] \rightarrow \mathbb{R}$ continuous. Find $r \in \mathbb{R}$ such that $f(r) = 0$.

Approximation to a root

A point $x^* \in \mathbb{R}$ such that

- $|r - x^*|$ is **very small**, and
- $f(x^*)$ is **very close** to 0.

Types of Iterative Methods

In this course, we find the root/root(s) upto some precision.

1 Open Domain Methods (Bracketing Methods)

Types of Iterative Methods

In this course, we find the root/root(s) upto some precision.

- 1 Open Domain Methods (Bracketing Methods)
 - Bisection method
 - Regular Falsi method (Method of False position)

Types of Iterative Methods

In this course, we find the root/root(s) upto some precision.

1 Open Domain Methods (Bracketing Methods)

- Bisection method
- Regular Falsi method (Method of False position)

Advantages: Always converges

Disadvantages: Locating the root initially

2 Closed Domain Methods (Non- Bracketing Methods)

Types of Iterative Methods

In this course, we find the root/root(s) upto some precision.

1 Open Domain Methods (Bracketing Methods)

- Bisection method
- Regular Falsi method (Method of False position)

Advantages: Always converges

Disadvantages: Locating the root initially

2 Closed Domain Methods (Non- Bracketing Methods)

- Secant method
- Fixed point theorem
- Newton's method (Newton Raphson method)

Types of Iterative Methods

In this course, we find the root/root(s) upto some precision.

1 Open Domain Methods (Bracketing Methods)

- Bisection method
- Regular Falsi method (Method of False position)

Advantages: Always converges

Disadvantages: Locating the root initially

2 Closed Domain Methods (Non- Bracketing Methods)

- Secant method
- Fixed point theorem
- Newton's method (Newton Raphson method)

Advantages: No need to locate the root initially

Disadvantages: May not converge

For each of the method, we study the following:

- Description of the method/ Basic idea

For each of the method, we study the following:

- Description of the method/ Basic idea
- Error analysis of the iteration and convergence

For each of the method, we study the following:

- Description of the method/ Basic idea
- Error analysis of the iteration and convergence
- Rate/Order of convergence

For each of the method, we study the following:

- Description of the method/ Basic idea
- Error analysis of the iteration and convergence
- Rate/Order of convergence
- Geometry of the method (iteration) process

For each of the method, we study the following:

- Description of the method/ Basic idea
- Error analysis of the iteration and convergence
- Rate/Order of convergence
- Geometry of the method (iteration) process
- Computational Scheme

For each of the method, we study the following:

- Description of the method/ Basic idea
- Error analysis of the iteration and convergence
- Rate/Order of convergence
- Geometry of the method (iteration) process
- Computational Scheme
- Application and example

Secant Method: Basic idea

This method is based on Mean Value Theorem.

- Assume that two initial guesses to α are known. Let these be x_0 and x_1 . They may occur on opposite side of α or on the same side of α .

Secant Method: Basic idea

This method is based on Mean Value Theorem.

- Assume that two initial guesses to α are known. Let these be x_0 and x_1 . They may occur on opposite side of α or on the same side of α .
- The two points $(x_0, f(x_0))$ and $(x_1, f(x_1))$ on the graph determine a straight line called a **secant line**.

Secant Method: Basic idea

This method is based on Mean Value Theorem.

- Assume that two initial guesses to α are known. Let these be x_0 and x_1 . They may occur on opposite side of α or on the same side of α .
- The two points $(x_0, f(x_0))$ and $(x_1, f(x_1))$ on the graph determine a straight line called a **secant line**.
- Equation of the secant line

$$y - f(x_1) = \frac{f(x_1) - f(x_0)}{x_1 - x_0}(x - x_1) \quad (2)$$

Secant Method: Basic idea

This method is based on Mean Value Theorem.

- The intersection of this line with x-axis is the next approximation to α . Let us denote it by x_2 .

Secant Method: Basic idea

This method is based on Mean Value Theorem.

- The intersection of this line with x-axis is the next approximation to α . Let us denote it by x_2 .
- The root x_2 or $(x_2, 0)$ lies on the line (2). So, we have

Secant Method: Basic idea

This method is based on Mean Value Theorem.

- The intersection of this line with x-axis is the next approximation to α . Let us denote it by x_2 .
- The root x_2 or $(x_2, 0)$ lies on the line (2). So, we have

$$\begin{aligned} 0 - f(x_1) &= \frac{f(x_1) - f(x_0)}{x_1 - x_0} (x_2 - x_1) \\ \Rightarrow \boxed{x_2 = x_1 - \frac{f(x_1) - f(x_0)}{x_1 - x_0} (x_2 - x_1)}. \end{aligned} \quad (3)$$

Secant Method: Basic idea

This method is based on Mean Value Theorem.

- The intersection of this line with x-axis is the next approximation to α . Let us denote it by x_2 .
- The root x_2 or $(x_2, 0)$ lies on the line (2). So, we have

$$\begin{aligned} 0 - f(x_1) &= \frac{f(x_1) - f(x_0)}{x_1 - x_0}(x_2 - x_1) \\ \Rightarrow \boxed{x_2 = x_1 - \frac{f(x_1) - f(x_0)}{x_1 - x_0}(x_2 - x_1)}. \end{aligned} \quad (3)$$

- Having found x_2 , we can drop x_0 and use x_1, x_2 as a new set of approximate value for α . This leads to an improved value x_3 ; and this process can be continued indefinitely.

Secant Method: General Iteration

- The general iteration formula for the secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} \quad n \geq 1. \quad (4)$$

Secant Method: General Iteration

- The general iteration formula for the secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} \quad n \geq 1. \quad (4)$$

- The sequence of iterates does not need to converge to root of the function. In fact it might also diverge. Then, **why does one use secant method instead of bisection method, which gives the security of convergence?**

Secant Method: General Iteration

- The general iteration formula for the secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} \quad n \geq 1. \quad (4)$$

- The sequence of iterates does not need to converge to root of the function. In fact it might also diverge. Then, **why does one use secant method instead of bisection method, which gives the security of convergence?**
- It is called a two-point method, since two approximation values are needed to obtain an improved value.

Secant Method: General Iteration

- The general iteration formula for the secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} \quad n \geq 1. \quad (4)$$

- The sequence of iterates does not need to converge to root of the function. In fact it might also diverge. Then, **why does one use secant method instead of bisection method, which gives the security of convergence?**
- It is called a two-point method, since two approximation values are needed to obtain an improved value.
- The Bisection method is also a two-point method, but the Secant method will almost always converge faster than bisection.

- Note that only one function evaluation is needed per step for the Secant method after x_2 has been determined. In contrast, each step of Newton's method requires an evaluation of both the function and its derivative.

- Note that only one function evaluation is needed per step for the Secant method after x_2 has been determined. In contrast, each step of Newton's method requires an evaluation of both the function and its derivative.
- Newton's method or the Secant method is often used to refine an answer obtained by another technique, such as the bisection method, since these methods requires good first approximation but generally give rapid convergence.

How fast is the convergence?

Definition 2

Let $\{x_n\}_{n \geq 1}$ be a sequence that converges to α . If positive constants λ and p exist with

$$\lim_{n \rightarrow \infty} \frac{|x_{n+1} - \alpha|}{|x_n - \alpha|^p} = \lambda,$$

then $\{x_n\}_{n \geq 1}$ is said to **converge to α of order p , with asymptotic error constant λ** . If $p = 1$, the method is called **linear**. If $p = 2$, the method is called **quadratic**.

How fast is the convergence?

- Can we find the exponent p such that

$$|x_{n+1} - \alpha| \approx |x_n - \alpha|^p? \quad (5)$$

- **Answer:** $p = \frac{1}{2}(1 + \sqrt{5}) \approx 1.62$. This is called **super linear convergence** ($1 < p < 2$).

Error analysis of the iteration and convergence

- The general iteration formula for the secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} \quad n \geq 1. \quad (6)$$

Error analysis of the iteration and convergence

- The general iteration formula for the secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} \quad n \geq 1. \quad (6)$$

- Let $\varepsilon_n = x_n - \alpha$ and $\varepsilon_{n+1} = x_{n+1} - \alpha$. So, ε_n and ε_{n+1} denote the errors in the root at the n th and $(n+1)$ th iterations.

Error analysis of the iteration and convergence

- The general iteration formula for the secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} \quad n \geq 1. \quad (6)$$

- Let $\varepsilon_n = x_n - \alpha$ and $\varepsilon_{n+1} = x_{n+1} - \alpha$. So, ε_n and ε_{n+1} denote the errors in the root at the n th and $(n+1)$ th iterations.
- (6) \implies

$$\varepsilon_{n+1} = \varepsilon_n - \frac{f(\alpha + \varepsilon_n)(\varepsilon_{n+1} - \varepsilon_n)}{f(\alpha + \varepsilon_n) - f(\alpha + \varepsilon_{n-1})} \quad (7)$$

Error analysis of the iteration and convergence

- Assume that f is twice differentiable and $f'(\alpha), f''(\alpha) \neq 0$. By Taylor's formula (with very small ε)

$$f(\alpha + \varepsilon) = f(\alpha) + \varepsilon f'(\alpha) + \frac{\varepsilon^2}{2} f''(\alpha) + R_2(\varepsilon).$$

Error analysis of the iteration and convergence

- Assume that f is twice differentiable and $f'(\alpha), f''(\alpha) \neq 0$. By Taylor's formula (with very small ε)

$$f(\alpha + \varepsilon) = f(\alpha) + \varepsilon f'(\alpha) + \frac{\varepsilon^2}{2} f''(\alpha) + R_2(\varepsilon).$$

Now, $f(\alpha) = 0$, and ε is very small, and $R_2(\varepsilon)$ is the remainder term. $R_2(\varepsilon)$ vanishes as $\varepsilon \rightarrow 0$ at a faster rate than ε^2 . Therefore,

$$f(\alpha + \varepsilon) \approx \varepsilon f'(\alpha) + \frac{\varepsilon^2}{2} f''(\alpha).$$

Error analysis of the iteration and convergence

- Let $M = \frac{f''(\alpha)}{2f'(\alpha)}$.



$$f(\alpha + \varepsilon_n) \approx \varepsilon_n f'(\alpha) \left(1 + \varepsilon_n M\right).$$

$$\varepsilon_{n+1} \approx \frac{f''(\alpha)}{2f'(\alpha)} \varepsilon_{n-1} \varepsilon_n \quad (8)$$

- (8) tells us that, as $n \rightarrow \infty$, the error tends to zero faster than a linear function but not quadratically!

Error Estimate

By Mean Value Theorem,

$$f(\alpha) - f(x_n) = f'(c_n)(\alpha - x_n) \quad (9)$$

where c_n lies between x_n and α . So, if $x_n \rightarrow \alpha$, then $c_n \approx x_n$ for large n , and we have

$$\begin{aligned} \alpha - x_n &\approx -\frac{f(x_n)}{f'(c_n)} \\ &\approx -\frac{f(x_n)}{f'(x_n)} \\ &\approx -f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})} \\ &\approx x_{n+1} - x_n. \end{aligned}$$

Thus, $\boxed{\alpha - x_n \approx x_{n+1} - x_n}$.