

MA 204 Numerical Methods

Dr. Debopriya Mukherjee
Lecture-3

January 11, 2024

Contents

- Solution of a nonlinear equation, bisection and secant methods, Newton's method, rate of convergence, solution of a system of nonlinear equations.

Contents

- Solution of a nonlinear equation, bisection and secant methods, Newton's method, rate of convergence, solution of a system of nonlinear equations.
- Interpolation by polynomials, divided differences, error of the interpolating polynomial, piecewise linear and cubic spline interpolation.

Newton Raphson Method

- Order of Convergence of Secant method: 1.62

Newton Raphson Method

- Order of Convergence of Secant method: 1.62
- Can this method be modified to have a **quadratic convergence**?

Newton Raphson Method

- Order of Convergence of Secant method: 1.62
- Can this method be modified to have a **quadratic convergence**? How?

Newton Raphson Method

- Order of Convergence of Secant method: 1.62
- Can this method be modified to have a **quadratic convergence**? How?
- In Secant method, if the iterative sequence converges, then

As n increases, x_{n-1} approaches x_n .

$$\Rightarrow f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}}, \quad \text{as } n \text{ increases}$$

Newton Raphson Method

- Order of Convergence of Secant method: 1.62
- Can this method be modified to have a **quadratic convergence**? How?
- In Secant method, if the iterative sequence converges, then

As n increases, x_{n-1} approaches x_n .

$$\Rightarrow f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}}, \quad \text{as } n \text{ increases}$$

- Recall the Secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})},$$

Newton Raphson Method

- Order of Convergence of Secant method: 1.62
- Can this method be modified to have a **quadratic convergence**? How?
- In Secant method, if the iterative sequence converges, then

As n increases, x_{n-1} approaches x_n .

$$\implies f'(x_n) \approx \frac{f(x_n) - f(x_{n-1})}{x_n - x_{n-1}}, \quad \text{as } n \text{ increases}$$

- Recall the Secant method is

$$x_{n+1} = x_n - f(x_n) \frac{x_n - x_{n-1}}{f(x_n) - f(x_{n-1})},$$

- The iterative procedure (given below) is called the **Newton Raphson's method**.

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}, \quad n = 0, 1, 2, \dots$$

History

- In numerical analysis, Newton's method, also known as the Newton–Raphson method, named after **Isaac Newton** and **Joseph Raphson**, is a root-finding algorithm which produces successively better approximations to the roots (or zeroes) of a real-valued function.

History

- In numerical analysis, Newton's method, also known as the Newton–Raphson method, named after **Isaac Newton** and **Joseph Raphson**, is a root-finding algorithm which produces successively better approximations to the roots (or zeroes) of a real-valued function.
- The most basic version starts with a real-valued function f , its derivative f' , and an initial guess x_0 for a root of f .

History

- In numerical analysis, Newton's method, also known as the Newton–Raphson method, named after **Isaac Newton** and **Joseph Raphson**, is a root-finding algorithm which produces successively better approximations to the roots (or zeroes) of a real-valued function.
- The most basic version starts with a real-valued function f , its derivative f' , and an initial guess x_0 for a root of f .
- This is one of the most powerful methods for solving a root-finding problem. There are many ways of introducing Newton's method.

Method Description

- Consider the sample graph of $y = f(x)$. Let the estimate of the root α be x_0 (initial guess).

Method Description

- Consider the sample graph of $y = f(x)$. Let the estimate of the root α be x_0 (initial guess).
- To improve on this estimate, consider the straight line that is tangent to the graph at $(x_0, f(x_0))$.

Method Description

- Consider the sample graph of $y = f(x)$. Let the estimate of the root α be x_0 (initial guess).
- To improve on this estimate, consider the straight line that is tangent to the graph at $(x_0, f(x_0))$.
- If x_0 is near α , the tangent line at x_0 cuts the x-axis at x_1 , which is near to α .

Method Description

- Consider the sample graph of $y = f(x)$. Let the estimate of the root α be x_0 (initial guess).
- To improve on this estimate, consider the straight line that is tangent to the graph at $(x_0, f(x_0))$.
- If x_0 is near α , the tangent line at x_0 cuts the x-axis at x_1 , which is near to α .
- To find a formula for x_1 , consider the equation of tangent to the graph of $y = f(x)$ at $(x_0, f(x_0))$. It is simply the graph of

$$y = f(x_0) + f'(x_0)(x - x_0).$$

Method Description

- Consider the sample graph of $y = f(x)$. Let the estimate of the root α be x_0 (initial guess).
- To improve on this estimate, consider the straight line that is tangent to the graph at $(x_0, f(x_0))$.
- If x_0 is near α , the tangent line at x_0 cuts the x-axis at x_1 , which is near to α .
- To find a formula for x_1 , consider the equation of tangent to the graph of $y = f(x)$ at $(x_0, f(x_0))$. It is simply the graph of

$$y = f(x_0) + f'(x_0)(x - x_0).$$

- $(x_1, 0)$ lies on this line:

$$\implies 0 = f(x_0) + f'(x_0)(x_1 - x_0)$$

$$\implies x_1 = x_0 - \frac{f(x_0)}{f'(x_0)}.$$

Method Description

- Since x_1 is expected to be an improvement over x_0 as an estimate of α , this entire procedure can be repeated with x_1 as the initial guess.

Method Description

- Since x_1 is expected to be an improvement over x_0 as an estimate of α , this entire procedure can be repeated with x_1 as the initial guess.
- This leads to the new estimate

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}.$$

Method Description

- Since x_1 is expected to be an improvement over x_0 as an estimate of α , this entire procedure can be repeated with x_1 as the initial guess.
- This leads to the new estimate

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}.$$

- Repeating this process, we obtain a sequence of numbers x_1, x_2, \dots that we hope will approach the root α . These numbers are called iterates, and they are defined recursively by the following general iteration formula:

Method Description

- Since x_1 is expected to be an improvement over x_0 as an estimate of α , this entire procedure can be repeated with x_1 as the initial guess.
- This leads to the new estimate

$$x_2 = x_1 - \frac{f(x_1)}{f'(x_1)}.$$

- Repeating this process, we obtain a sequence of numbers x_1, x_2, \dots that we hope will approach the root α . These numbers are called iterates, and they are defined recursively by the following general iteration formula:

$$x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}, \quad n = 0, 1, 2, \dots \quad (1)$$

This is **Newton's method** for solving $f(x) = 0$.

Convergence Analysis

- Assume that $f(x)$ has at least two continuous derivatives for all x in some interval about the root α .

Convergence Analysis

- Assume that $f(x)$ has atleast two continuous derivatives for all x in some interval about the root α . Further assume that

$$f'(\alpha) \neq 0 \quad (2)$$

Convergence Analysis

- Assume that $f(x)$ has atleast two continuous derivatives for all x in some interval about the root α . Further assume that

$$f'(\alpha) \neq 0 \tag{2}$$

- This says that the graph of $y = f(x)$ is not tangent to the x -axis when the graph intersects it $x = \alpha$.

Convergence Analysis

- Assume that $f(x)$ has atleast two continuous derivatives for all x in some interval about the root α . Further assume that

$$f'(\alpha) \neq 0 \quad (2)$$

- This says that the graph of $y = f(x)$ is not tangent to the x -axis when the graph intersects it $x = \alpha$. This case when $f'(\alpha) = 0$ will be discussed later.

Convergence Analysis

- Assume that $f(x)$ has atleast two continuous derivatives for all x in some interval about the root α . Further assume that

$$f'(\alpha) \neq 0 \quad (2)$$

- This says that the graph of $y = f(x)$ is not tangent to the x -axis when the graph intersects it $x = \alpha$. This case when $f'(\alpha) = 0$ will be discussed later.
- Combining (2) with the continuity of $f'(x)$ implies

$$f'(x) \neq 0 \quad \forall x \text{ near } \alpha.$$

Convergence Analysis

- By Taylor's theorem,

$$\begin{aligned} f(\alpha) &= f(x_n + \alpha - x_n) \\ &= f(x_n) + (\alpha - x_n)f'(x_n) + \frac{(\alpha - x_n)^2}{2}f''(c_n) \quad (3) \end{aligned}$$

where c_n is an unknown point between α and x_n .

Convergence Analysis

- Note that $f(\alpha) = 0$. Using $x_{n+1} = x_n - \frac{f(x_n)}{f'(x_n)}$ This implies

$$\begin{aligned} 0 &= f(x_n) + (\alpha - x_n)f'(x_n) + \frac{1}{2}(\alpha - x_n)^2 f''(c_n) \\ \implies 0 &= \frac{f(x_n)}{f'(x_n)} + (\alpha - x_n) + \frac{1}{2}(\alpha - x_n)^2 \frac{f''(c_n)}{f'(x_n)} \\ \implies 0 &= x_n - x_{n+1} + \alpha - x_n + (\alpha - x_n)^2 \frac{f''(c_n)}{f'(x_n)} \end{aligned} \quad (4)$$

- $$\alpha - x_{n+1} = (\alpha - x_n)^2 \left[-\frac{f''(c_n)}{2f'(x_n)} \right]$$

Error Analysis

- This formula says that the error in x_{n+1} is nearly proportional to the square of the error in x_n .

Error Analysis

- This formula says that the error in x_{n+1} is nearly proportional to the square of the error in x_n .
- When the initial error is sufficiently small, this shows that the error in the succeeding iterates will decrease very rapidly.

When to stop iterations?

- Noting that $f(\alpha) = 0$, by Mean Value Theorem

$$f(x_n) = f(x_n) - f(\alpha) = f'(\xi_n)(x_n - \alpha).$$

Thus, error $\varepsilon_n = \alpha - x_n = -\frac{f(x_n)}{f'(x_n)}$ provided that x_n is close to α that $f'(x_n) \approx f'(\xi_n)$. This implies $\boxed{\alpha - x_n \approx x_{n+1} - x_n}$.

When to stop iterations?

- Noting that $f(\alpha) = 0$, by Mean Value Theorem

$$f(x_n) = f(x_n) - f(\alpha) = f'(\xi_n)(x_n - \alpha).$$

Thus, error $\varepsilon_n = \alpha - x_n = -\frac{f(x_n)}{f'(x_n)}$ provided that x_n is close to α that $f'(x_n) \approx f'(\xi_n)$. This implies $\boxed{\alpha - x_n \approx x_{n+1} - x_n}$.

- This is the standard error estimation formula for Newton's method and it is usually fairly accurate. However, this formula is not valid if $f'(\alpha) = 0$.

Multiple Root

- The zero of the function f is said to be of multiplicity m if

$$f(x) = (x - \alpha)^m g(x)$$

for some continuous function g with $g(\alpha) \neq 0$, m is a positive integer.

Multiple Root

- The zero of the function f is said to be of multiplicity m if

$$f(x) = (x - \alpha)^m g(x)$$

for some continuous function g with $g(\alpha) \neq 0$, m is a positive integer.

- If we assume that f is sufficiently differentiable, an equivalent definition is that

$$f(\alpha) = f'(\alpha) = \dots = f^{(m-1)}(\alpha) = 0, f^{(m)}(\alpha) \neq 0.$$

Multiple Root

- The zero of the function f is said to be of multiplicity m if

$$f(x) = (x - \alpha)^m g(x)$$

for some continuous function g with $g(\alpha) \neq 0$, m is a positive integer.

- If we assume that f is sufficiently differentiable, an equivalent definition is that

$$f(\alpha) = f'(\alpha) = \dots = f^{(m-1)}(\alpha) = 0, f^{(m)}(\alpha) \neq 0.$$

- A zero of multiplicity 1 is called a simple root or a simple zero.

Remedy

- One method of handling the problem of multiple roots of a function f is to define

$$\mu(x) = \frac{f(x)}{f'(x)}.$$

Remedy

- One method of handling the problem of multiple roots of a function f is to define

$$\mu(x) = \frac{f(x)}{f'(x)}.$$

- If α is a zero of f of multiplicity m . Then, one can write

$$f(x) = (x - \alpha)^m g(x), \quad g(\alpha) \neq 0.$$

Remedy

- One method of handling the problem of multiple roots of a function f is to define

$$\mu(x) = \frac{f(x)}{f'(x)}.$$

- If α is a zero of f of multiplicity m . Then, one can write

$$f(x) = (x - \alpha)^m g(x), \quad g(\alpha) \neq 0.$$

- This implies

$$\begin{aligned}\mu(x) &= \frac{(x - \alpha)^m g(x)}{m(x - \alpha)^{m-1} g(x) + (x - \alpha)^m g'(x)} \\ &= (x - \alpha) \frac{g(x)}{mg(x) + (x - \alpha)g'(x)}\end{aligned}$$

also has a zero at α . However, $g(\alpha) \neq 0$.

Newton's method for multiple roots

- Thus, $\mu'(\alpha) = \frac{1}{m} \neq 0$, hence α is called a simple zero of μ .
- Newton's method can be applied to $\mu(x)$ to give

$$x_{n+1} = x_n - \frac{f(x_n)f'(x_n)}{[f'(x_n)]^2 - f(x_n)f''(x_n)}$$

- This is called Newton's modified method. This has quadratic convergence regardless of multiplicity of the zeros of f .
- For a simple zero, the original Newton's method requires significantly low computations.

Newton versus Secant

- Note that only one function evaluation is needed per step for the Secant method after x_2 has been determined. In contrast, each step of Newton's method requires an evaluation of both the function and its derivative.

Newton versus Secant

- Note that only one function evaluation is needed per step for the Secant method after x_2 has been determined. In contrast, each step of Newton's method requires an evaluation of both the function and its derivative.
- Newton's method or the Secant method is often used to refine an answer obtained by another technique, such as the bisection method, since these methods requires good first approximation but generally give rapid convergence.