

12.8.3.8

EE24BTECH11006 - Arnav Mahishi

Question:

Find the area of the smaller region bounded by the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$ and the line $\frac{x}{3} + \frac{y}{2} = 1$

input	Description	value
a	Length of semi major axis of ellipse	3
b	Length of semi minor axis of ellipse	2
v	Quadratic form of matrix	$\begin{pmatrix} b^2 & 0 \\ 0 & a^2 \end{pmatrix}$
u	Linear coefficient vector	0
f	Constant Term	$-(a^2 b^2)$
h	One of the points the line passes through	$\begin{pmatrix} a \\ 0 \end{pmatrix}$
m	Slope of line	$\begin{pmatrix} \frac{1}{b} \\ \frac{-1}{a} \end{pmatrix}$
n	number of subintervals we are taking	1000
x_0	semi-major axis of ellipse	3
x_n	semi-minor axis of ellipse	2

TABLE 0: Variables Used

Theoretical Solution:

The point of intersection of the line with the ellipse is $x_i = h + k_i m$, where, k_i is a constant and is calculated as follows:-

$$k_i = \frac{1}{m^T V m} \left(-m^T (V h + u) \pm \sqrt{[m^T (V h + u)]^2 - g(h) (m^T V m)} \right)$$

Substituting the input parameters in k_i ,

$$k_i = \frac{1}{\begin{pmatrix} \frac{1}{b} & \frac{-1}{a} \end{pmatrix} \begin{pmatrix} b^2 & 0 \\ 0 & a^2 \end{pmatrix} \begin{pmatrix} \frac{1}{b} \\ \frac{-1}{a} \end{pmatrix}} \left(-\begin{pmatrix} \frac{1}{b} & \frac{-1}{a} \end{pmatrix} \left(\begin{pmatrix} b^2 & 0 \\ 0 & a^2 \end{pmatrix} \begin{pmatrix} a \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) \pm \sqrt{\left[\begin{pmatrix} \frac{1}{b} & \frac{-1}{a} \end{pmatrix} \left(\begin{pmatrix} b^2 & 0 \\ 0 & a^2 \end{pmatrix} \begin{pmatrix} a \\ 0 \end{pmatrix} + \begin{pmatrix} 0 \\ 0 \end{pmatrix} \right) \right]^2 - g(h) \left(\begin{pmatrix} \frac{1}{b} & \frac{-1}{a} \end{pmatrix} \begin{pmatrix} b^2 & 0 \\ 0 & a^2 \end{pmatrix} \begin{pmatrix} \frac{1}{b} \\ \frac{-1}{a} \end{pmatrix} \right)} \right) \quad (0.1)$$

We get,

$$k_i = 0, -ab$$

Substituting k_i in $x_i = h + k_i m$ we get,

$$x_1 = \begin{pmatrix} a \\ 0 \end{pmatrix} + (0) \begin{pmatrix} \frac{1}{b} \\ \frac{-1}{a} \end{pmatrix} \quad (0.2)$$

$$\Rightarrow x_1 = \begin{pmatrix} a \\ 0 \end{pmatrix} \quad (0.3)$$

$$x_2 = \begin{pmatrix} a \\ 0 \end{pmatrix} + (-ab) \begin{pmatrix} \frac{1}{b} \\ \frac{-1}{a} \end{pmatrix} \quad (0.4)$$

$$\Rightarrow x_2 = \begin{pmatrix} a \\ 0 \end{pmatrix} + \begin{pmatrix} -a \\ b \end{pmatrix} \quad (0.5)$$

$$\Rightarrow x_2 = \begin{pmatrix} 0 \\ b \end{pmatrix} \quad (0.6)$$

The area of the smaller region bounded by the ellipse $\frac{x^2}{9} + \frac{y^2}{4} = 1$ and the line $\frac{x}{3} + \frac{y}{2} = 1$ is

$$= \int_0^a \frac{b}{a} \sqrt{a^2 - x^2} dx - \int_0^a \frac{b}{a} (a - x) dx \quad (0.7)$$

$$= \frac{b}{a} \left(\frac{x}{2} \sqrt{a^2 - x^2} + \frac{a^2}{2} \sin^{-1} \frac{x}{a} - ax + \frac{x^2}{2} \right)_0^a \quad (0.8)$$

$$= \frac{b}{a} \left(\frac{\pi a^2}{4} - \frac{a^2}{2} \right) = \frac{ab}{2} \left(\frac{\pi}{2} - 1 \right) \quad (0.9)$$

The given area is $\frac{ab}{2} \left(\frac{\pi}{2} - 1 \right)$ sq. units

\therefore Upon substituting $a = 3, b = 2$ the given area is $3 \left(\frac{\pi}{2} - 1 \right)$ sq. units ≈ 1.712 sq. units

Computational Solution:

Using the Trapezoidal rule which approximates the integral of a function $f(x)$ over an interval $[a, b]$ by dividing the interval into n subintervals and approximating the area under the curve as a series of trapezoids

$$\int_a^b f(x) dx \approx \frac{h}{2} \left[f(x_0) + 2 \sum_{i=1}^{n-1} f(x_i) + f(x_n) \right] \quad (0.10)$$

Where x_0 is semi-major axis of ellipse and x_n is semi-minor axis of the ellipse and h is the width of each subinterval.

$$x_i = x_0 + i \cdot h \quad (0.11)$$

$$h = \frac{b - a}{n} \quad (0.12)$$

In the case of our problem of the area between the line and ellipse the area is computed

by:

$$A = \int_{x_{start}}^{x_{end}} (f_{ellipse}(x) - f_{line}(x)) dx \quad (0.13)$$

$$f_{ellipse}(x) = \sqrt{4 \left(1 - \frac{x^2}{9}\right)} \quad (0.14)$$

$$f_{line}(x) = 2 - \frac{2x}{3} \quad (0.15)$$

Where $[x_{start}, x_{end}]$ are the intersection points. Using the difference equation which iteratively computes the integral by adding the contributions of each subinterval. Let A_k represent the area approximation after k subintervals:

$$A_k = A_{k-1} + \frac{h}{2} [f(x_k) + f(x_{k-1})] \quad (0.16)$$

$$x_k = x_{start} + k \cdot h \quad (0.17)$$

$$\Rightarrow A_k = A_{k-1} + \frac{h}{2} [(f_{ellipse}(x_k) - f_{line}(x_k)) + (f_{ellipse}(x_{k-1}) - f_{line}(x_{k-1}))] \quad (0.18)$$

Start with $A_0 = 0$ then iteratively calculate $f_{ellipse}(x_i)$ and $f_{line}(x_i)$ for each i from x_{start} to x_{end} and calculate each A_k from 0 to n . The value of A_n will be the approximate area of the ellipse. As n tends to infinity A_n will be the exact area of the ellipse.




Fig. 0.1: Plot of area enclosed between the line and ellipse