

Analysis 2 - Concise Notes

MATH50001

Total Content

Arnav Singh



Colour Code - **Definitions** are **green** in these notes, **Consequences** are **red** and **Causes** are **blue**

Content from MATH40002 assumed to be known.

Mathematics
Imperial College London
United Kingdom
March 15, 2022

Contents

1	Holomorphic Functions	2
1.1	Complex Numbers	2
1.2	Sets in Complex Plane	2
1.3	Complex Functions	3
1.4	Complex Derivative	3
1.5	Cauchy-Riemann equations	4
1.6	Cauchy-Riemann equations in polar	4
1.7	Power Series	4
1.8	Elementary functions	5
1.8.1	Exponential function	5
1.8.2	Trigonometric functions	5
1.8.3	Logarithmic Functions	6
1.8.4	Powers	6
2	Cauchy's Integral Formula	6
2.1	Parametrised Curve	6
2.2	Integration along Curves	7
2.3	Primitive Functions	7
2.4	Properties of Holomorphic functions	7
2.5	Local existence of primitives and Cauch-Goursat theorem in a disc	7
2.6	Homotopies and simply connected domains	8
2.7	Cauchy's Integral Formulae	8
3	Applications of Cauchy's integral formula	9
3.1	Taylor + Maclaurin Series	9
3.2	Sequences of holomorphic functions	9
3.3	Holomorphic functions defined in terms of integrals	9
3.4	Schwarz reflection principle	10
4	Meromorphic Functions	10
4.1	Complex Logarithm	10
4.2	Laurent Series	11
4.3	Poles of holomorphic functions	11
4.4	Residue Theory	11
4.5	The argument principle	12
4.6	Evaluation of definite integrals	12
5	Harmonic Functions	12
5.1	Harmonic functions	12
5.2	Properties of real + imaginary parts of holomorphic function	13
5.3	Preservation of angles	13
5.4	Möbius Transformations	14
5.5	Cross-ratios Mobius Transformations	14
5.6	Conformal mapping of half-plane to unit disc	14
5.7	Riemann mapping theorem	14

1 Holomorphic Functions

1.1 Complex Numbers

Definition 1.1. i

$$i = \sqrt{-1}, \quad i^2 = -1$$

Root of $x^2 + 1 = 0$

Basic properties

$$z = x + iy, \quad \operatorname{Re}(z) = x, \quad \operatorname{Im}(z) = y$$

The complex conjugate:

$$\bar{z} = x - iy$$

Polar Coordinates

$$z = x + iy$$

$$r = |z| = \sqrt{x^2 + y^2}$$

$$x = r \cos \theta, \quad y = r \sin \theta$$

$$z = r(\cos \theta + i \sin \theta)$$

De-Moivre's Formula

$$z^n = r^n(\cos(n\theta) + i \sin(n\theta)), \quad n \in \mathbb{Z}^+$$

Eulers's Formula

$$e^{i\theta} = (\cos \theta + i \sin \theta)$$

1.2 Sets in Complex Plane

Definition 1.2. Discs in \mathbb{C}

Open Disc : $D_r(z_0) = \{z \in \mathbb{C} : |z - z_0| < r\}$

Boundary of Disc : $C_r(z_0) = \{z \in \mathbb{C} : |z - z_0| = r\}$

Unit Disc : $\mathbb{D} = \{z \in \mathbb{C} : |z| < 1\}$

Definition 1.3. Interior Point

$\Omega \in \mathbb{C}, z_0$ an **interior point** of Ω if $\exists r > 0$ s.t $D_r(z_0) \subset \Omega$

Definition 1.4.

Set Ω **open** if $\forall \omega \in \Omega, \omega$ an interior point

Definition 1.5.

Set Ω **closed** if $\Omega^C = \mathbb{C} \setminus \Omega$ open

Closed \iff contains all its limit points.

Definition 1.6. Closure

Closure of $\Omega = \bar{\Omega} = \{\Omega \cup \text{limit points of } \Omega\}$

Definition 1.7. Boundary

$$\text{Boundary of } \Omega = \underbrace{\bar{\Omega}}_{\text{Closure}} \setminus \underbrace{\Omega}_{\text{interior}}$$

Definition 1.8. Bounded

Ω bounded if $\exists M > 0$ s.t $|\omega| < M \quad \forall \omega \in \Omega$

Definition 1.9. Diameter

$$\text{diam}(\Omega) = \sup_{z, w \in \Omega} |z - w|$$

Definition 1.10. Compact

Ω compact if closed and bounded

Theorem 1.1.

$$\begin{aligned} \Omega \text{ compact} &\iff \text{every sequence } \{z_n\} \subset \Omega \text{ has a subsequence convergent in } \Omega \\ &\iff \text{every open covering of } \Omega \text{ has a finite subcover} \end{aligned}$$

Theorem 1.2.

if $\Omega_1 \supset \Omega_2 \supset \dots \Omega_n \supset \dots$ a sequence of non-empty compact sets

s.t $\lim_{n \rightarrow \infty} \text{diam}(\Omega_n) \rightarrow 0$

$$\implies \exists! w \in \mathbb{C}, w \in \Omega_n \forall n$$

Definition 1.11. Connected

Open set Ω **connected** \iff any 2 points in Ω joined by curve γ entirely contained in Ω

1.3 Complex Functions

Definition 1.12.

$\Omega_1, \Omega_2 \subset \mathbb{C}$

$$f : \Omega_1 \rightarrow \Omega_2$$

a **mapping** $\Omega_1 \rightarrow \Omega_2$ if

$$\forall z = x + iy \in \Omega_1$$

$$\exists! w = u + iv \in \Omega_2, \text{ s.t } w = f(z)$$

We have $w = f(z) = u(x, y) + iv(x, y)$

$u, v : \mathbb{R}^2 \rightarrow \mathbb{R}$

Definition 1.13.

f defined on $\Omega_1 \subset \mathbb{C}$ **f continuous** at $z_0 \in \Omega$ if

$$\forall \epsilon > 0 \exists \delta > 0 \text{ s.t } |z - z_0| < \delta \implies |f(z) - f(z_0)| < \epsilon$$

f continuous if continuous $\forall z \in \Omega$

1.4 Complex Derivative

Definition 1.14. **Holomorphic**

$\Omega_1, \Omega_2 \subset \mathbb{C}$ open sets

$f : \Omega_1 \rightarrow \Omega_2$

Say f **differentiable/holomorphic** at z_0 if

$$\lim_{h \rightarrow 0} \frac{f(z_0 + h) - f(z_0)}{h} = f'(z_0) \text{ exists}$$

f holomorphic on open set Ω if holomorphic at every point of Ω

Lemma

f holomorphic at $z_0 \in \Omega \iff \exists a \in \mathbb{C}$ s.t

$$f(z_0 + h) - f(z_0) - ah = h\Psi(h)$$

For Ψ a function defined for all small h with $\lim_{h \rightarrow 0} \Psi(h) = 0$, $a = f'(z_0)$

Corollary

f holomorphic $\implies f$ continuous

Proposition

f, g holomorphic in $\Omega \implies$

- (i) $(f + g)' = f' + g'$
- (ii) $(fg)' = f'g + fg'$
- (iii) $g(z_0) \neq 0 \implies \left(\frac{f}{g}\right)' = \frac{f'g - fg'}{g^2}$
- (iv) $f : \Omega \rightarrow V, g : \Omega \rightarrow \mathbb{C}$ holomorphic
 $\implies [g \circ f(z)]' = g'(f(z))f'(z) \forall z \in \Omega$

1.5 Cauchy-Riemann equations

$$\begin{aligned} \frac{\partial u}{\partial x} &= \frac{\partial v}{\partial y} & \frac{\partial u}{\partial y} &= -\frac{\partial v}{\partial x} \\ u'_x &= v'_y & u'_y &= -v'_x \end{aligned}$$

Definition 1.15.

$$\frac{\partial}{\partial z} = \frac{1}{2} \left(\frac{\partial}{\partial x} + \frac{1}{i} \frac{\partial}{\partial y} \right) \quad \frac{\partial}{\partial \bar{z}} = \frac{1}{2} \left(\frac{\partial}{\partial x} - \frac{1}{i} \frac{\partial}{\partial y} \right)$$

Theorem 1.3.

$f(z) = u(x, y) + iv(x, y) \quad z = x + iy$
 f holomorphic at $z_0 \implies$

$$\frac{\partial f}{\partial \bar{z}}(z_0) = 0 \quad f'(z_0) = \frac{\partial f}{\partial z}(z_0) = 2 \frac{\partial u}{\partial z}(z_0)$$

Theorem 1.4.

$f = u + iv$ complex-valued function on open set Ω
 u, v continuously differentiable, satisfying Cauchy-Riemann equations $\implies f$ holomorphic on Ω with $f'(z) = \frac{\partial f}{\partial z}(z)$

1.6 Cauchy-Riemann equations in polar

For $f = u + iv$ we have

$$u'_r = \frac{1}{r} v'_\theta \quad v'_r = -\frac{1}{r} u'_\theta$$

1.7 Power Series

Definition 1.16. Power Series

Of the form

$$\sum_{n=0}^{\infty} a_n z^n \quad a_n \in \mathbb{C}$$

Series converge at z if $S_N(z) = \sum_{n=0}^N a_n z^n$ has limit $S(z) = \lim_{N \rightarrow \infty} S_N(z)$

Theorem 1.5.

Given power series $\sum_{n=0}^{\infty} a_n z^n, \exists 0 \leq R \leq \infty$ s.t

- (i) if $|z| < R \implies$ series converges absolutely
- (ii) $|z| > R \implies$ series diverges

$$\frac{1}{R} = \limsup_{n \rightarrow \infty} |a_n|^{1/n} \quad (\text{Radius of Convergence})$$

Theorem 1.6.

$$f(z) = \sum_{n=0}^{\infty} a_n z^n$$

Defines holomorphic function on its disc of convergence. With

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}$$

with same radius of convergence as f .

Power series infinitely differentiable in the disc of convergence, achieved through term-wise differentiation.

Definition 1.17. Entire

A function said to be **entire** if holomorphic $\forall z \in \mathbb{C}$

1.8 Elementary functions**1.8.1 Exponential function**

$$e^z = e^x \cos y + i e^x \sin y \quad z = x + iy \in \mathbb{C}$$

Properties

- (i) $y = 0 \implies e^z = e^x$
- (ii) e^z is entire
- (iii) $g(z)$ holomorphic
 $\implies \frac{\partial}{\partial z} e^{g(z)} = e^{g(z)} g'(z)$
- (iv) $z_1, z_2 \in \mathbb{C} \quad e^{z_1+z_2} = e^{z_1} e^{z_2}$
- (v) $|e^z| = |e^x| |e^{iy}| = e^x \sqrt{\cos^2 x + \sin^2(x)} = e^x$
- (vi) $(e^{iy})^n = e^{iny}$
- (vii) $\arg(z) = \arctan(y/x)$
 $\arg(e^z) = y + 2\pi k, \quad k \in \mathbb{Z}$

1.8.2 Trigonometric functions**Definition 1.18.**

$$\cos z = \frac{1}{2} (e^{iz} + e^{-iz}) \quad \sin z = \frac{1}{2i} (e^{iz} - e^{-iz})$$

Properties

- (i) $\sin z, \cos z$ are entire
- (ii) $\frac{\partial}{\partial z} \sin z = \cos z \quad \frac{\partial}{\partial z} \cos z = -\sin z$
- (iii) $\sin^2 z + \cos^2 z = 1$
- (iv) $\sin(z_1 \pm z_2) = \sin z_1 \cos z_2 \pm \cos z_1 \sin z_2$
 $\cos(z_1 \pm z_2) = \cos z_1 \cos z_2 \mp \sin z_1 \sin z_2$

1.8.3 Logarithmic Functions

Definition 1.19.

$$\log(z) = \ln|z| + i \arg(z) = \log(r) + i(\theta + 2\pi k) \quad z \neq 0, k \in \mathbb{Z}$$

$\log(z)$ a multi-valued function

Definition 1.20.

$\text{Log}(z) = \ln|z| + i \text{Arg}(z)$ for $\text{Arg}(z)$ principal value $\in [-\pi, \pi]$

Properties

- (i) $\log(z_1 z_2) = \log(z_1) + \log(z_2)$
- (ii) $\text{Log}(z)$ holomorphic in $\mathbb{C} \setminus \{(\infty, 0]\}$

1.8.4 Powers

Definition 1.21.

$\alpha \in \mathbb{C}$ define $z^\alpha := e^{\alpha \log(z)}$ as a multi-valued function

Definition 1.22.

Principal value of z^α , $\alpha \in \mathbb{C}$ as $z^\alpha = e^{\alpha \text{Log}(z)}$ **Properties**

- (i) $z^{a_1} z^{a_2} = z^{a_1 + a_2}$

2 Cauchy's Integral Formula

2.1 Parametrised Curve

Definition 2.1.

Parametrised curve a function $z(t) : [a, b] \rightarrow \mathbb{C}$

Smooth if $z'(t)$ exists and is continuous on $[a, b]$ with $z'(t) \neq 0 \forall t \in [a, b]$ Taking one-sided limits for $z'(a), z'(b)$.

Piecewise-smooth if z continuous on $[a, b]$ and if \exists finitely many points $a = a_0 < a_1 < \dots < a_n = b$ s.t $z(t)$ smooth on $[a_k, a_{k+1}]$

$$z : [a, b] \rightarrow \mathbb{C} \quad \tilde{z} : [c, d] \rightarrow \mathbb{C}$$

equivalent if \exists continuously differentiable bijection $s \rightarrow t(s)$ from $[c, d]$ to $[a, b]$ s.t $t'(s) > 0$ and $\tilde{z}(s) = z(t(s)) =$

Definition 2.2. Path Integrals

Path integral given smooth $\gamma \subset \mathbb{C}$ parametrised by $z : [a, b] \rightarrow \mathbb{C}$.

f continuous function on γ

$$\int_{\gamma} f(z) dz = \int_a^b f(z(t)) z'(t) dt$$

independent of choice of parametrization.

If γ piece-wise smooth

$$\int_{\gamma} f(z) dz = \sum_{k=0}^{n-1} \int_{a_k}^{a_{k+1}} f(z(t)) z'(t) dt$$

Definition 2.3.

Define curve γ^- obtained by reversing orientation of γ

Can take $z^- : [a, b] \rightarrow \mathbb{C}$ s.t $z^-(t) = z(b + a - t)$

Definition 2.4. Closed Curve

Smooth/piece-wise smooth curve closed if $z(a) = z(b)$ for any parametrisation.

Definition 2.5. Simple Curve

Smooth/piece-wise smooth curve simple if not **self-intersecting**

$$z(t) \neq z(s) \text{ unless } s = t \in [a, b]$$

2.2 Integration along Curves

Definition 2.6. Length of smooth curve

$$\text{Length}(\gamma) = \int_a^b |z'(t)| dt = \int_a^b \sqrt{x'(t)^2 + y'(t)^2} dt$$

Theorem 2.1. Properties of Integration

- (i) $\int_{\gamma} af(z) + bg(z)dz = a \int_{\gamma} f(z)dz + b \int_{\gamma} g(z)dz$
- (ii) γ^- reverse orientation of γ

$$\implies \int_{\gamma} f(z)dz = - \int_{\gamma^-} f(z)dz$$

- (iii) **M-L inequality**

$$\left| \int_{\gamma} f(z)dz \right| \leq \sup_{z \in \gamma} |f(z)| \cdot \text{length}(\gamma) = \int_a^b \sqrt{x'(t)^2 + y'(t)^2} dt$$

2.3 Primitive Functions

Definition 2.7. Primitive

A **Primitive** for f on $\Omega \subset \mathbb{C}$ a function F holomorphic on Ω s.t $F'(z) = f(z) \forall z \in \Omega$

Theorem 2.2.

Continuous function f with primitive F in open set Ω and curve γ in Ω from $w_1 \rightarrow w_2$

$$\int_{\gamma} f(z)dz = F(w_2) - F(w_1)$$

Corollary

γ closed curve in open set Ω f continuous and has primitive in $\Omega \implies$

$$\oint_{\gamma} f(z)dz = 0$$

Corollary

Ω with $f' = 0 \implies f$ constant

2.4 Properties of Holomorphic functions

Theorem 2.3.

Let $\Omega \subset \mathbb{C}$ open set

$T \subset \Omega$ a triangle whose interior contained in Ω

$$\implies \oint_T f(z)dz = 0$$

for f holomorphic in Ω

Corollary

f holomorphic on open set Ω containing rectangle R in its interior

$$\implies \oint_R f(z)dz = 0$$

2.5 Local existence of primitives and Cauchy-Goursat theorem in a disc

Theorem 2.4.

Holomorphic functions in open disc have a primitive in that disc

Corollary - (Cauchy-Goursat Theorem for a disc)

f holomorphic in disc $\implies \oint_{\gamma} f(z)dz = 0$

for any closed curve γ in that disc

Corollary

Suppose f holomorphic in open set containing circle C and its interior

$$\implies \oint_C f(z)dz = 0$$

2.6 Homotopies and simply connected domains

Definition 2.8. Homotopic

γ_0, γ_1 **homotopic** in Ω if $\forall s \in [0, 1], \exists$ curve $\gamma \subset \Omega$ with $\gamma_s(t)$ s.t

$$\gamma_s(a) = \alpha \quad \gamma_s(b) = \beta$$

$$\forall t \in [a, b] : \gamma_s(t)|_{s=0} = \gamma_0(t) \quad \gamma_s(t)|_{s=1} = \gamma_1(t)$$

With $\gamma_s(t)$ jointly continuous in $s \in [0, 1]$ and $t \in [a, b]$

Theorem 2.5.

γ_0, γ_1 homotopic, f holomorphic

$$\int_{\gamma_0} f(z)dz = \int_{\gamma_1} f(z)dz$$

Definition 2.9.

Open set $\Omega \subset \mathbb{C}$ **simply connected** if any 2 pair of curves in Ω with shared end-points homotopic.

Theorem 2.6.

Any holomorphic function in simply connected domain has a primitive.

Corollary - (Cauchy-Goursat Theorem)

f holomorphic in simply connected open set Ω

$$\implies \oint_{\gamma} f(z)dz = 0$$

for any closed piecewise-smooth curve $\gamma \subset \Omega$

Theorem 2.7. (Deformation Theorem)

γ_1 and γ_2 , 2 simple closed piecewise-smooth curves with γ_2 lying wholly inside γ_1

f holomorphic in domain containing region between γ_1, γ_2

$$\implies \oint_{\gamma_1} f(z)dz = \oint_{\gamma_2} f(z)dz$$

2.7 Cauchy's Integral Formulae

Theorem 2.8. (Cauchy's Integral Formula)

f holomorphic inside and on simple closed piecewise-smooth curve γ

$\forall z_0$ interior to γ

$$f(z_0) = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(z)}{z - z_0} dz$$

Theorem 2.9. (Generalised Cauchy's integral formula)

f holomorphic in open set Ω .

γ simple, closed piecewise-smooth Ω

$\forall z$ interior to γ

$$\implies \frac{d^n f(z)}{dz^n} = \frac{n!}{2\pi i} \oint_{\gamma} \frac{f(t)}{(t - z)^{n+1}} dt$$

Corollary

f holomorphic \implies all its derivatives are too.

3 Applications of Cauchy's integral formula

Corollary - (Liouville's theorem)

if an entire function bounded $\implies f$ constant

Theorem 3.1. (Fundamental theorem of algebra)

Every polynomial of degree > 0 with complex coefficients has at least one zero.

Corollary Every polynomial $P(z) = a_n z^n + \dots + a_0$ of degree $n \geq 1$ has precisely n roots in \mathbb{C}

Theorem 3.2. (Morera's theorem)

Suppose f continuous in open disc D s.t \forall triangle $T \subset D$

$$\int_T f(z)dz = 0 \implies f \text{ holomorphic}$$

3.1 Taylor + Maclaurin Series

Theorem 3.3. (Taylor's expansion theorem)

f holomorphic in Ω , $z_0 \in \Omega$

$$\implies f(z) = f(z_0) + f'(z_0)(z - z_0) + \frac{f''(z_0)}{2!}(z - z_0)^2 + \dots$$

Valid in all circles $\{z : |z - z_0| < r\} \subset \Omega$

Definition 3.1. (Taylor Series)

$$f(z) = f(z_0) + f'(z_0)(z - z_0) + \frac{f''(z_0)}{2!}(z - z_0)^2 + \dots = \sum_{i=0}^{\infty} \frac{f^{(i)}(z_0)}{i!}(z - z_0)^i$$

Definition 3.2. (Maclaurin Series)

Taylor series for $z_0 = 0$

$$f(z) = \sum_{n=0}^{\infty} \frac{f^n(0)}{n!} z^n$$

3.2 Sequences of holomorphic functions

Theorem 3.4.

if $\{f_n\}_{n=1}^{\infty}$ a sequence of holomorphic functions converging uniformly to f in every compact subset of $\Omega \implies f$ holomorphic in Ω

Corollary

$$F(z) = \sum_{n=1}^{\infty} f_n(z)$$

f_n holomorphic in $\Omega \subset \mathbb{C}$

Given series converges uniformly in compact subsets of $\Omega \implies F(z)$ holomorphic

Theorem 3.5.

Sequence $\{f_n\}_{n=1}^{\infty} \xrightarrow{\text{unif}} f$ in every compact subset of $\Omega \implies$ sequence $\{f'_n\}_{n=1}^{\infty} \xrightarrow{\text{unif}} f'$ in every compact subset of Ω

3.3 Holomorphic functions defined in terms of integrals

Theorem 3.6.

Let $F(z, s)$ defined for $(z, s) \in \Omega \times [0, 1]$

$\Omega \subset \mathbb{C}$ open set. Given F satisfies

(i) $F(z, s)$ holomorphic in $\Omega \forall s$

(ii) F continuous on $\Omega \times [0, 1]$

$\implies f(z) := \int_0^1 F(z, s)ds$ holomorphic

3.4 Schwarz reflection principle

Definition 3.3.

$\Omega \subset \mathbb{C}$ open and **symmetric** w.r.t real line

$$z \in \Omega \iff \bar{z} \in \Omega$$

Definition 3.4.

$$\Omega^+ = \{z \in \Omega : \text{Im}(z) > 0\} \quad \Omega^- = \{z \in \Omega : \text{Im}(z) < 0\} \quad I = \{z \in \Omega : \text{Im}(z) = 0\}$$

Theorem 3.7. (Symmetry Principle)

f^+, f^- holomorphic in Ω^+, Ω^- respectively.

Extend continuously to I s.t $f^+(x) = f^-(x) \quad \forall x \in I$

$$f(z) := \begin{cases} f^+(z), & z \in \Omega^+ \\ f^+(z) = f^-(z), & z \in I \\ f^-(z), & z \in \Omega^- \end{cases} \quad \text{holomorphic}$$

Theorem 3.8. (Schwarz reflection principle)

f holomorphic in Ω^+ extend continuously to I s.t f real-valued on I

$\implies \exists F$ holomorphic in Ω s.t $F|_{\Omega^+} = f$

4 Meromorphic Functions

4.1 Complex Logarithm

Theorem 4.1.

Ω simply connected, $1 \in \Omega, 0 \notin \Omega$

\implies in Ω there is a branch of logarithm

$$F(z) = \log_{\Omega}(z)$$

Satisfying

(i) F holomorphic in Ω

(ii) $e^{F(z)} = z \quad \forall z \in \Omega$

(iii) $F(r) = \log(r), \quad r \in \mathbb{R} \text{ close to } 1$

Theorem 4.2.

Holomorphic f has 0 of order m at z_0

\iff can be written in form

$$f(z) = (z - z_0)^m g(z)$$

g holomorphic at $z_0, g(z_0) \neq 0$

Corollary

0s of non-constant holomorphic function are isolated.

Every zero has neighbourhood, inside of which it is the only 0

4.2 Laurent Series

Definition 4.1.

Laurent Series for f at z_0 , where series converge

$$f(z) = \sum_{-\infty}^{\infty} a_n(z - z_0)^n = \cdots + a_{-2}(z - z_0)^{-2} + a_{-1}(z - z_0)^{-1} + a_0 + a_1(z - z_0)^1 + a_2(z - z_0)^2 + \cdots$$

Theorem 4.3. (Laurent Expansion theorem)

f holomorphic in annulus $D = \{z : r < |z - z_0| < R\}$

$\implies f(z)$ expressed in form $f(z) = \sum_{-\infty}^{\infty} a_n(z - z_0)^n$

$$a_n = \frac{1}{2\pi i} \oint_{\gamma} \frac{f(\eta)}{(\eta - z_0)^{n+1}} d\eta$$

γ simple, closed piecewise smooth curve in D with z_0 in its interior.

4.3 Poles of holomorphic functions

Definition 4.2.

z_0 a **singularity** of complex function f

if f not holomorphic at z_0 , but every neighbourhood of z_0 has at least 1 holomorphic point.

Definition 4.3.

Singularity z_0 is **isolated** if \exists neighbourhood of z_0 , where it is the only singularity.

Definition 4.4.

f holomorphic with isolated singularity z_0

Considering Laurent expansion valid in some annulus

$$f(z) = \sum_{-\infty}^{\infty} a_n(z - z_0)^n$$

\implies

- $a_n = 0 \ \forall n < 0 \implies z_0$ a **removable singularity**
- $a_n = 0 \ \forall n < -m, m \in \mathbb{Z}^+, a_{-m} \neq 0 \implies z_0$ pole of order m
- $a_n \neq 0$ for infinitely many negative $n \implies z_0$ a **essential singularity**

Theorem 4.4.

f has pole of order m at $z_0 \iff$ written in form

$$f(z) = \frac{g(z)}{(z - z_0)^m}$$

g holomorphic at $z_0, g(z_0) \neq 0$

4.4 Residue Theory

Definition 4.5.

Let $f(z) = \sum_{-\infty}^{\infty} a_n(z - z_0)^n$ for $0 < |z - z_0| < R$ the Laurent series for f at z_0

Residue of f at z_0 is

$$\implies \text{Res}[f, z_0] = a_{-1}$$

Theorem 4.5.

$\gamma \subset \{z : 0 < |z - z_0| < R\}$ simple closed piecewise-smooth curve containing z_0

$$\implies \text{Res}[f, z_0] = \frac{1}{2\pi i} \oint_{\gamma} f(z) dz$$

Theorem 4.6.

f holomorphic function inside and on simple closed piecewise-smooth curve γ except at the singularities z_1, \dots, z_n in its interior

$$\implies \oint_{\gamma} f(z) dz = 2\pi i \sum_{j=1}^n \text{Res}[f, z_j]$$

4.5 The argument principle

Theorem 4.7. (Principle of argument)

f holomorphic in open Ω , except for finitely many poles.

γ simple closed piecewise-smooth curve in Ω not passing through poles or zeroes of f

$$\implies \oint_{\gamma} \frac{f'(z)}{f(z)} dz = 2\pi i(N - P)$$

$$N = \sum \text{order}(\text{zeroes}) \quad P = \sum \text{order}(\text{poles})$$

Theorem 4.8. (Rouche's Theorem)

f, g holomorphic in open Ω

$\gamma \subset \Omega$ simple closed piecewise-smooth curve with interior containing only points of Ω

if $|g(z)| < |f(z)|, z \in \gamma$

$$\implies \sum_{\text{0s of } f+g \text{ in } \gamma} \text{order}(\text{zeros}) = \sum_{\text{0s of } f \text{ in } \gamma} \text{order}(\text{zeros})$$

Definition 4.6.

Mapping **open** if maps open sets \mapsto open sets

Theorem 4.9. (Open mapping theorem)

if f holomorphic and non-constant in open $\Omega \subset \mathbb{C}$

$$\implies f \text{ open}$$

Remark

f open $\implies |f|$ open

Theorem 4.10. (Max modulus principle)

f non-constant holomorphic in open $\Omega \subset \mathbb{C}$

$\implies f$ cannot attain maximum in Ω

Corollary

Ω open with closure $\bar{\Omega}$ compact

f holomorphic on Ω and continuous on $\bar{\Omega}$

$$\sup_{z \in \Omega} |f(z)| \leq \sup_{z \in \text{Omega} \setminus \Omega} |f(z)|$$

4.6 Evaluation of definite integrals

5 Harmonic Functions

5.1 Harmonic functions

Definition 5.1.

$\varphi = \varphi(x, y) : \mathbb{R}^2 \rightarrow \mathbb{R}, x, y \in \mathbb{R}$

φ **harmonic** in open $\Omega \subset \mathbb{R}^2$ if

$$\begin{aligned} \underbrace{\Delta \varphi(x, y)}_{\text{laplace operator}} &:= \frac{\partial^2 \varphi}{\partial x^2}(x, y) + \frac{\partial^2 \varphi}{\partial y^2}(x, y) \\ &:= \varphi''_{xx}(x, y) + \varphi''_{yy}(x, y) \\ &:= 0 \end{aligned}$$

Theorem 5.1.

$f(z) = u(x, y) + iv(x, y)$ holomorphic in open $\Omega \subset \mathbb{C}$

$\implies u, v$ harmonic

Theorem 5.2. (Harmonic conjugate)

u harmonic in open disc $D \subset \mathbb{C}$

$\implies \exists$ harmonic v s.t $f = u + iv$ holomorphic in D

v the **harmonic conjugate** to u

Remark

In simply connected domain $\Omega \subset \mathbb{R}^2$ every harmonic function u has harmonic conjugate v s.t

$$v(x, y) = \int_{\gamma} \left(-\frac{\partial u}{\partial y} dx + \frac{\partial u}{\partial x} dy \right)$$

Integral independent of path, by Green's theorem as u harmonic and Ω simply connected.

5.2 Properties of real + imaginary parts of holomorphic function

Theorem 5.3.

Assume $f = u + iv$ holomorphic on open connected $\Omega \subset \mathbb{C}$

$$u(x, y) = C \quad (1)$$

$$v(x, y) = K \quad (2)$$

$$C, K \in \mathbb{R} \quad (3)$$

If (1) and (2) have same solution (x_0, y_0) and $f'(x_0 + iy_0) \neq 0$

\implies curve defined by (1) orthogonal to curve defined by (2)

5.3 Preservation of angles

Definition 5.2.

Consider smooth curve $\gamma \subset \mathbb{C}$

$$z(t) = x(t) + iy(t) \quad t \in [a, b]$$

$\forall t_0 \in [a, b]$ we have direction vector

$$\begin{aligned} L_{t_0} &= \{z(t_0) + tz'(t_0) : t \in \mathbb{R}\} \\ &= \{x(t_0) + tx'(t_0) + i(y(t_0) + ty'(t_0)) : t \in \mathbb{R}\} \end{aligned}$$

For γ_1, γ_2 curves parameterised by functions $z_1(t), z_2(t), t \in [0, 1]$ s.t $z_1(0) = z_2(0)$

Define angle between γ_1, γ_2 as angle between tangents

$$\arg z_2'(0) - \arg z_1'(0)$$

Theorem 5.4. (Angle preservation theorem)

f holomorphic in open $\Omega \subset \mathbb{C}$

Given γ_1, γ_2 inside Ω , parameterised by $z_1(t), z_2(t)$

Take $z_0 = z_1(0) = z_2(0)$ with $z_1'(0), z_2'(0), f'(z_0) \neq 0$

$$\underbrace{\arg z_2'(t) - \arg z_1'(t)}_{\text{angle between } z_1(0), z_2(0)} \Big|_{t=0} = \underbrace{\arg f(z_2'(t)) - \arg f(z_1'(t))}_{\text{angle between } f(z_1(0)), f(z_2(0))} \Big|_{t=0} \pmod{2\pi}$$

Definition 5.3.

Ω open $\subset \mathbb{C}$

$f : \Omega \rightarrow \mathbb{C}$ **conformal** if holomorphic in Ω and if $f'(z) \neq 0 \forall z \in \Omega$

Conformal mappings preserve angles.

Definition 5.4.

Holomorphic function a **local injection** on open $\Omega \subset \mathbb{C}$

if

$$\forall z_0 \in \Omega, \exists D = \{z : |z - z_0| < r\} \subset \Omega \text{ s.t } f : D \rightarrow f(D) \text{ an injection}$$

Theorem 5.5.

$f : \Omega \rightarrow \mathbb{C}$ local injection and holomorphic

$$\implies f'(z) \neq 0 \quad \forall z \in \Omega$$

Inverse of f defined on its range holomorphic

\implies inverse of conformal mapping also holomorphic

5.4 Möbius Transformations

Definition 5.5.

Möbius Transformation/ Bilinear transformation a map

$$f(z) = \frac{az + b}{cz + d} \quad a, b, c, d \in \mathbb{C}, ad - bc \neq 0$$

Remark

Möbius Transformations holomorphic except for simple pole $z = -\frac{d}{c}$ with derivative

$$f'(z) = \frac{ad - bc}{(cz + d)^2}$$

\implies mapping conformal for $\mathbb{C} \setminus \{-\frac{d}{c}\}$

Theorem 5.6.

- (i) Inverse of Möbius transformation a Möbius transformation
- (ii) Composition of Möbius transformations a Möbius transformation

Corresponding to matrix multiplication and inverses

Definition 5.6. (Special/Simple Möbius transformations)

(M1) $f(z) = az$ Scaling and rotation by a

(M2) $f(z) = z + b$ Translation by b

(M3) $f(z) = \frac{1}{z}$ Inverse and reflection w.r.t real axis

Theorem 5.7.

Every Möbius transformation a composition of $M1, M2, M3$

Corollary

Möbius transformations:

circles \mapsto circles
interior points \mapsto interior points

Straight lines, considered to be circles of infinite radius

5.5 Cross-ratios Möbius Transformations

Theorem 5.8.

$w = f(z)$ a Möbius Transformation

s.t distinct $(z_1, z_2, z_3) \mapsto (w_1, w_2, w_3)$

$$\implies \left(\frac{z - z_1}{z - z_3} \right) \left(\frac{z_2 - z_3}{z_2 - z_1} \right) = \left(\frac{w - w_1}{w - w_3} \right) \left(\frac{w_2 - w_3}{w_2 - w_1} \right) \quad \forall z$$

5.6 Conformal mapping of half-plane to unit disc

Theorem 5.9.

$$\begin{aligned} \mathbb{D} &= \{z : |z| < 1\} & \mathbb{H} &= \{z = x + iy : \operatorname{Im}(z) = y > 0\} \\ w = f(z) &= \frac{i - z}{i + z} & g(w) &= \frac{1 - w}{1 + w} \end{aligned}$$

5.7 Riemann mapping theorem

Definition 5.7.

$\Omega \subset \mathbb{C}$ **proper** if non-empty and $\Omega \neq \mathbb{C}$

Theorem 5.10.

Ω proper and simply connected

if $z_0 \in \Omega \implies \exists!$ conformal $f : \Omega \rightarrow \mathbb{D}$ s.t $f(z_0) = 0$ and $f'(z_0) > 0$

Corollary

Any 2 simply connected open subsets in \mathbb{C} conformally equivalent.