

1. You drive down a road whose speed limit is 60 miles per hour. An observer sees you at 12pm, and a second observer 35 miles away sees you at 12:30pm. Assuming they've watched their analysis lectures, how can they prove you were speeding?

*Solution.* The mean value theorem guarantees that there is some time strictly between 12:00 and 12:30 when your velocity was

$$\frac{35 \text{ miles}}{\frac{1}{2} \text{ hour}} = 70 \text{ miles per hour.}$$

2. Prove using l'Hôpital's rule that  $\lim_{x \rightarrow \infty} \left(1 + \frac{r}{x}\right)^x = e^r$ . (Hint: take logs first.)

*Solution.* We write the limiting term as

$$\left(1 + \frac{r}{x}\right)^x = e^{x \log\left(1 + \frac{r}{x}\right)},$$

so by the continuity of  $f(x) = e^x$  it will suffice to compute

$$\lim_{x \rightarrow \infty} \frac{\log\left(1 + \frac{r}{x}\right)}{1/x} = \lim_{y \downarrow 0} \frac{\log(1 + ry)}{y}.$$

The derivative of  $\log(1 + ry)$  is  $\frac{r}{1+ry}$ , so we apply l'Hôpital's rule to get

$$\lim_{y \downarrow 0} \frac{\log(1 + ry)}{y} = \lim_{y \downarrow 0} \frac{r/(1 + ry)}{1} = r,$$

and so  $\lim_{x \rightarrow \infty} x \log\left(1 + \frac{r}{x}\right) = r$  and it follows that  $\lim_{x \rightarrow \infty} \left(1 + \frac{r}{x}\right)^x = e^r$ .

3. Let  $H_n$  denote the harmonic sum  $\frac{1}{1} + \frac{1}{2} + \cdots + \frac{1}{n}$ .
- (a) Show using the mean value theorem that  $\frac{1}{n+1} < \log(n+1) - \log(n) < \frac{1}{n}$  for all  $n \in \mathbb{N}$ .
  - (b) Prove that  $H_n - 1 < \log(n) < H_{n-1}$  for all  $n \geq 2$ , where  $H_k = \frac{1}{1} + \frac{1}{2} + \cdots + \frac{1}{k}$ , and deduce that  $\log(n+1) < H_n < \log(n) + 1$ .
  - (c) Prove that the sequence  $(H_n - \log(n))$  is decreasing, and that  $\lim_{n \rightarrow \infty} (H_n - \log(n))$  exists. (This limit is called the *Euler–Mascheroni constant*  $\gamma \approx 0.577 \dots$ )

*Solution.* (a) Since  $\log(x)$  has derivative  $\frac{1}{x}$ , there is some  $z \in (n, n+1)$  such that

$$\frac{\log(n+1) - \log(n)}{(n+1) - n} = \frac{1}{z} \in \left(\frac{1}{n+1}, \frac{1}{n}\right),$$

or equivalently  $\frac{1}{n+1} < \log(n+1) - \log(n) < \frac{1}{n}$ .

- (b) We sum each side of the inequality  $\frac{1}{k+1} < \log(k+1) - \log(k) < \frac{1}{k}$  from  $k = 1$  to  $n - 1$  to get

$$\sum_{k=1}^{n-1} \frac{1}{k+1} < \log(n) - \log(1) < \sum_{k=1}^{n-1} \frac{1}{k},$$

noticing that lots of cancellation occurs in the middle. Since  $\log(1) = 0$ , this is equivalent to  $H_n - 1 < \log(n) < H_{n-1}$ . The left half of this gives us  $H_n < \log(n) + 1$ , and when we replace  $n$  with  $n + 1$  the right half gives us  $\log(n + 1) < H_n$ , so we combine these to get  $\log(n + 1) < H_n < \log(n) + 1$ .

- (c) From part (b) we have

$$0 < \log(n + 1) - \log(n) < H_n - \log(n) < 1,$$

so the sequence  $a_n = H_n - \log(n)$  is bounded, and thus if it is monotone decreasing then it converges. We compute

$$a_{n+1} - a_n = \frac{1}{n+1} - \log(n+1) + \log(n),$$

so if we let  $f(x) = \frac{1}{x+1} - \log(x+1) + \log(x)$ , then we want to show that  $f(n) < 0$  for all integers  $n \geq 1$ . We have

$$\begin{aligned} f'(x) &= -\frac{1}{(x+1)^2} - \frac{1}{x+1} + \frac{1}{x} \\ &= -\frac{1}{(x+1)^2} + \frac{1}{x(x+1)} = \frac{1}{x(x+1)^2} \end{aligned}$$

and so  $f'(x) > 0$  for all  $x > 0$ , meaning that  $f(x)$  is strictly increasing. But

$$\lim_{x \rightarrow \infty} f(x) = \lim_{x \rightarrow \infty} \left( \frac{1}{x+1} - \log \left( 1 + \frac{1}{x} \right) \right) = 0.$$

Since  $f(x)$  is strictly increasing, this implies that  $f(x) < 0$  for all  $x > 0$ . (Proof: assuming otherwise, let  $\epsilon = f(y)$  be a positive value of  $f$ . Then there is no  $N > 0$  such that if  $x \geq N$  then  $|f(x) - 0| < \epsilon$ , because as soon as  $x > \max(y, N)$  we have  $f(x) > f(y) = \epsilon$ . This contradicts  $f(x) \rightarrow 0$ .) So in particular, for all integers  $n \geq 1$  we have  $a_{n+1} - a_n = f(n) < 0$ , hence  $a_{n+1} < a_n$ .

4. (\*) Let  $f : \mathbb{R} \rightarrow \mathbb{R}$  be differentiable, and suppose there is a constant  $C < 1$  such that  $|f'(x)| \leq C$  for all  $x \in \mathbb{R}$ . We will prove that  $f$  has exactly one fixed point, meaning there is a unique  $y \in \mathbb{R}$  such that  $f(y) = y$ . Pick some  $x_0 \in \mathbb{R}$  and let

$$x_{n+1} = f(x_n) \text{ for all } n \geq 0.$$

- (a) Prove that  $|x_{n+2} - x_{n+1}| \leq C|x_{n+1} - x_n|$  for all  $n$ .
- (b) Prove that the sequence  $(x_n)$  converges, and that if its limit is  $y$  then  $f(y) = y$ .
- (c) Prove that  $f$  cannot have two different fixed points.

*Solution.* (a) If  $x_{n+1} = x_n$  then  $x_{n+2} = f(x_{n+1}) = f(x_n) = x_{n+1}$ , and so both sides of the desired inequality are zero. Otherwise, the mean value theorem tells us that there is some  $t$  between  $x_n$  and  $x_{n+1}$  such that

$$\frac{f(x_{n+1}) - f(x_n)}{x_{n+1} - x_n} = f'(t) \Rightarrow \left| \frac{x_{n+2} - x_{n+1}}{x_{n+1} - x_n} \right| = |f'(t)| \leq C.$$

Thus  $|x_{n+2} - x_{n+1}| \leq C|x_{n+1} - x_n|$  as desired.

(b) Write  $d = |x_1 - x_0|$ . Then  $|x_{n+1} - x_n| \leq C^n d$  by induction and part (a). The triangle inequality says that for any integers  $m \geq n$ ,

$$\begin{aligned} |x_m - x_n| &\leq |x_m - x_{m-1}| + \cdots + |x_{n+2} - x_{n+1}| + |x_{n+1} - x_n| \\ &\leq C^{m-1}d + \cdots + C^{n+1}d + C^n d \\ &< \sum_{i=n}^{\infty} C^i d = \frac{C^n d}{1 - C}. \end{aligned}$$

For any  $\epsilon > 0$  we can find  $N \geq 0$  such that  $\frac{C^N d}{1 - C} < \epsilon$ , since the left side approaches 0 as  $N \rightarrow \infty$ . Then given  $m, n \geq N$  we have shown that

$$|x_m - x_n| < \frac{C^N d}{1 - C} < \epsilon,$$

which proves that the sequence  $(x_n)$  is Cauchy and hence convergent, say with limit  $y$ . Since  $x_n \rightarrow y$  and  $f$  is continuous, we have  $f(x_n) \rightarrow f(y)$ . But  $f(x_n) = x_{n+1} \rightarrow y$ , so it must be the case that  $f(y) = y$ .

(c) Suppose that  $y$  and  $z$  are distinct fixed points of  $f$ . By the mean value theorem, there is some  $t$  between  $y$  and  $z$  such that

$$\frac{f(y) - f(z)}{y - z} = f'(t) \Rightarrow \frac{y - z}{y - z} = f'(t) \Rightarrow f'(t) = 1.$$

But this contradicts the assumption that  $|f'(x)| \leq C < 1$  for all  $x \in \mathbb{R}$ .

5. (a) Compute the Taylor series  $P(x)$  of  $f(x) = \log(1 + x)$  centered at  $x = 0$ , and prove that it converges absolutely on  $(-1, 1)$ .
- (b) Prove using Taylor's theorem that  $f(x) = P(x)$  on some open neighborhood of 0, by showing that the sequence of  $n$ th order Taylor polynomials  $P_n(x)$  converges uniformly to  $f(x)$ . Show that the same is true at  $x = 1$ , and so  $\log(2) = \frac{1}{1} - \frac{1}{2} + \frac{1}{3} - \frac{1}{4} + \frac{1}{5} - \cdots$ .

*Solution.* (a) We have  $f'(0) = \frac{1}{1+x}$ , and we claim by induction that

$$f^{(n)}(x) = (-1)^{n-1}(n-1)!(1+x)^{-n}$$

for all  $n \geq 1$ : if it's true for  $n = k$  then we have

$$f^{(k+1)}(x) = (-1)^{k-1}(k-1)! \cdot (-k)(1+x)^{-k-1} = (-1)^k k!(1+x)^{-(k+1)}$$

as desired. Then  $f^{(n)}(0) = (-1)^{n-1}(n-1)!$  for  $n \geq 1$ , and  $f(0) = \log(1) = 0$ , so  $f(x)$  has Taylor series

$$P(x) = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}(n-1)!x^n}{n!} = \sum_{n=1}^{\infty} \frac{(-1)^{n-1}x^n}{n},$$

which has the form  $\frac{x}{1} - \frac{x^2}{2} + \frac{x^3}{3} - \frac{x^4}{4} + \dots$ . Absolute convergence follows from the comparison test, since  $|\frac{(-1)^{n-1}x^n}{n}| \leq |x^n|$  and  $\sum_{n=1}^{\infty} x^n$  is a geometric series which converges absolutely on  $(-1, 1)$ .

(b) By Taylor's theorem, if  $x > -1$  is nonzero then we have

$$f(x) = P_n(x) + \frac{f^{(n+1)}(t)}{(n+1)!}x^{n+1}$$

for some  $t$  between 0 and  $x$ . The same computation as in part (a) says that  $f^{(n+1)}(t) = (-1)^n n!(1+t)^{-(n+1)}$ , so

$$|f(x) - P_n(x)| = \left| \frac{(-1)^n n!(1+t)^{-(n+1)}}{(n+1)!} x^{n+1} \right| = \frac{1}{n+1} \left| \frac{x}{1+t} \right|^{n+1}.$$

Now if  $0 < x \leq 1$  then we have  $0 < t < x$ , so  $1+t > 1 \geq x$  and hence  $\left| \frac{x}{1+t} \right| < 1$ . If instead  $-\frac{1}{2} \leq x < 0$  then we have  $1+t > 1+x \geq \frac{1}{2} > |x|$ , so again  $\left| \frac{x}{1+t} \right| < 1$ . Thus for any nonzero  $x \in [-\frac{1}{2}, 1]$  we have

$$|f(x) - P_n(x)| \leq \frac{1}{n+1},$$

and so  $P_n$  converges uniformly to  $f(x) = \log(1+x)$  on this interval.

Remark: In fact  $f(x) = P(x)$  on all of  $(-1, 1)$ , but we need better control over  $t$  to prove this on the interval  $(-1, \frac{1}{2})$ .

6. Suppose that  $f : \mathbb{R} \rightarrow \mathbb{R}$  has at least six continuous derivatives, and that  $f^{(i)}(0) = 0$  for  $i = 1, 2, 3, 4, 5$  but  $f^{(6)}(0) = 1$ . Prove that  $f(x)$  has a local minimum at  $x = 0$ .

*Solution.* We apply Taylor's theorem to see that if  $x \in (-\delta, \delta)$  is nonzero, then there is some  $t$  between 0 and  $x$  such that

$$f(x) = \sum_{i=0}^5 \frac{f^{(i)}(0)x^i}{i!} + \frac{f^{(6)}(t)x^6}{6!} = f(0) + \frac{f^{(6)}(t)x^6}{6!}.$$

Since  $f^{(6)}(x)$  is continuous, there is some  $\delta > 0$  such that

$$|y - 0| < \delta \Rightarrow |f^{(6)}(y) - f^{(6)}(0)| < 1,$$

hence  $f^{(6)}(y) > 0$  for all  $y \in (-\delta, \delta)$ . If we take  $x \in (-\delta, \delta)$  above then  $t \in (-\delta, \delta)$  as well, so  $f^{(6)}(t) > 0$ , and then since  $\frac{x^6}{6!} \geq 0$  we conclude that  $f(x) \geq f(0)$  for all  $x \in (-\delta, \delta)$ .

7. (a) Prove that  $f(x) = e^x$  is convex on all of  $\mathbb{R}$ .  
 (b) Let  $a, b > 0$ . Prove the *arithmetic mean–geometric mean inequality*

$$\frac{a+b}{2} \geq \sqrt{ab}$$

by using the convexity of  $e^x$ . (Hint: think about  $\alpha = \log(a)$  and  $\beta = \log(b)$ .)

- (c) Prove for any  $a, b > 0$  and  $s \in [0, 1]$  that  $sa + (1-s)b \geq a^s b^{1-s}$ .  
 (d) Prove *Young's inequality*: for any  $x, y \geq 0$  and  $p, q > 0$  with  $\frac{1}{p} + \frac{1}{q} = 1$ ,

$$\frac{x^p}{p} + \frac{y^q}{q} \geq xy.$$

*Solution.* (a) It suffices to check that  $f''(x) \geq 0$  for all  $x$ , and this is certainly true since  $f''(x) = e^x$ .

- (b) Assuming  $\alpha < \beta$  without loss of generality, the convexity of  $e^x$  implies for  $\alpha < \frac{\alpha+\beta}{2} < \beta$  that

$$\frac{e^\alpha + e^\beta}{2} \geq e^{(\alpha+\beta)/2} = \sqrt{e^\alpha \cdot e^\beta}$$

which is equivalent to  $\frac{a+b}{2} \geq \sqrt{ab}$ .

- (c) Since  $e^x$  is convex, we know that

$$se^\alpha + (1-s)e^\beta \geq e^{s\alpha + (1-s)\beta},$$

and the left side is  $sa + (1-s)b$  while the right side is  $(e^\alpha)^s (e^\beta)^{1-s} = a^s b^{1-s}$ .

- (d) We may assume that  $x, y > 0$ , since otherwise the inequality reduces to  $\frac{x^p}{p} + \frac{y^q}{q} = 0$ , which is true. We now use part (c), setting  $s = \frac{1}{p}$  (so  $1-s = \frac{1}{q}$ ) and  $(a, b) = (x^p, y^q)$ , to get

$$\frac{x^p}{p} + \frac{y^q}{q} \geq (x^p)^{1/p} (y^q)^{1/q} = xy.$$

8. Define  $f : \mathbb{R} \rightarrow \mathbb{R}$  by  $f(x) = \begin{cases} e^{-1/x^2}, & x \neq 0 \\ 0, & x = 0. \end{cases}$

- (a) Prove that for all integers  $n \geq 0$ , there is a polynomial  $p_n(x)$  such that

$$f^{(n)}(x) = \frac{p_n(x)}{x^{3n}} e^{-1/x^2} \text{ for all } x \neq 0.$$

- (b) Prove that  $f^{(n)}(0) = 0$  for all  $n$ , and hence that  $f(x)$  does not equal its Taylor series (centered at  $a = 0$ ) at any nonzero  $x$ .

- (c) Define  $g : \mathbb{R} \rightarrow \mathbb{R}$  by  $g(x) = \begin{cases} 0, & x \leq 0 \\ e^{-1/x^2}, & x > 0. \end{cases}$  Prove that  $g^{(n)}(x)$  exists for all  $n \geq 0$  and all  $x \in \mathbb{R}$ , and that  $g^{(n)}(0) = 0$  for all  $n$ .

- (d) Define  $h : \mathbb{R} \rightarrow \mathbb{R}$  by  $h(x) = g(x)g(1-x)$ . Prove that  $h^{(n)}(x)$  exists for all  $n \geq 0$  and all  $x \in \mathbb{R}$ , and that  $h(x) \neq 0$  if and only if  $0 < x < 1$ .

The function  $h$  is called a *bump function*: it is infinitely differentiable, and it is zero outside a compact set (namely  $[0, 1]$ ) but also takes positive values.

*Solution.* (a) When  $n = 0$  we take  $p_0(x) = 1$ . If this holds for  $n = k$ , we compute

$$\begin{aligned} f^{(k+1)}(x) &= \frac{d}{dx} \left( \frac{p_k(x)}{x^{3k}} e^{-1/x^2} \right) \\ &= \frac{p'_k(x)x^{3k} - 3kx^{3k-1}p_k(x)}{x^{6k}} e^{-1/x^2} + \frac{p_k(x)}{x^{3k}} \left( \frac{2}{x^3} e^{-1/x^2} \right) \\ &= \left( \frac{p'_k(x)x^3 - 3kx^2p_k(x) + 2p_k(x)}{x^{3k+3}} \right) e^{-1/x^2}, \end{aligned}$$

so we can take  $p_{k+1} = x^3p'_k - (3kx^2 - 2)p_k$  and the proof follows by induction.

- (b) When  $n = 0$  it is true by definition. If we have proved it for all  $n \leq k$ , then for  $n = k + 1$  we have

$$f^{(k+1)}(0) = \lim_{x \rightarrow 0} \frac{f^{(k)}(x) - f^{(k)}(0)}{x - 0} = \lim_{x \rightarrow 0} \frac{f^{(k)}(x)}{x},$$

assuming the limit exists, and so using part (a) we wish to prove that

$$\lim_{x \rightarrow 0} \frac{p_k(x)}{x^{3k+1}e^{1/x^2}} = 0.$$

Since  $p_k(x)$  is continuous, it will suffice to prove that  $\lim_{x \rightarrow 0} |x^{3k+1}e^{1/x^2}| = \infty$ . By the substitution  $y = \frac{1}{x}$  we have

$$\lim_{x \rightarrow 0} |x^{3k+1}e^{1/x^2}| = \lim_{|y| \rightarrow \infty} \left| \frac{e^{y^2}}{y^{3k+1}} \right| = \lim_{y \rightarrow \infty} \frac{e^{y^2}}{y^{3k+1}}.$$

But since  $y^2 \geq 0$ , every term in the power series  $e^{y^2} = \sum_{i=0}^{\infty} \frac{(y^2)^i}{i!}$  is nonnegative, and so if we single out the  $i = 2k + 1$  term then

$$e^{y^2} \geq \frac{(y^2)^{2k+1}}{(2k+1)!} \Rightarrow \frac{e^{y^2}}{y^{3k+1}} \geq \frac{y^{4k+2}/(2k+1)!}{y^{3k+1}} = \frac{y^{k+1}}{(2k+1)!}.$$

The right side certainly goes to  $\infty$  as  $y \rightarrow \infty$ , hence  $\lim_{x \rightarrow 0} |x^{3k+1}e^{1/x^2}| = \infty$  and this proves that  $f^{(k+1)}(0) = 0$ . The proof follows for all  $n$  by induction.

The Taylor series of  $f$  at  $a = 0$  is  $\sum_{n=0}^{\infty} \frac{f^{(n)}(0)}{n!} x^n = 0$ , but clearly for all  $x \neq 0$  this is not equal to  $f(x) = e^{-1/x^2} > 0$ .

(c) For all  $n$ , we have

$$g^{(n)}(x) = \begin{cases} 0, & x < 0 \\ f^{(n)}(x), & x > 0. \end{cases},$$

so we only need to check that  $g^{(n)}(0)$  exists and is zero for all  $n$ . Again, we induct: it is true when  $n = 0$ , and if it is true for  $n = k$  then

$$\frac{g^{(k)}(x) - g^{(k)}(0)}{x - 0} = \frac{g^{(k)}(x)}{x} = \begin{cases} 0, & x < 0 \\ f^{(k)}(x), & x > 0. \end{cases}$$

Thus  $\lim_{x \uparrow 0} \frac{g^{(k)}(x) - g^{(k)}(0)}{x - 0} = 0$  by inspection, and


$$\lim_{x \downarrow 0} \frac{g^{(k)}(x) - g^{(k)}(0)}{x - 0} = \lim_{x \downarrow 0} \frac{f^{(k)}(x)}{x} = f^{(k+1)}(0) = 0$$

by part (b), so  $g^{(k+1)}(0) = \lim_{x \rightarrow 0} \frac{g^{(k)}(x) - g^{(k)}(0)}{x - 0}$  exists and is zero as well.

(d) Since  $g(x)$  and  $g(1-x)$  are infinitely differentiable, repeated application of the product rule says that  $h(x) = g(x)g(1-x)$  has  $n$  derivatives for all  $n$  as well. Moreover, we have  $g(x) = 0$  for  $x \leq 0$  and  $g(1-x) = 0$  for  $x \geq 1$ , so  $h(x) = 0$  for all  $x \notin (0, 1)$ ; and if  $0 < x < 1$  then

$$h(x) = g(x)g(1-x) = e^{-1/x^2} \cdot e^{-1/(1-x)^2} > 0.$$

Here is a graph of  $h(x)$ :



Note that  $h(x)$  is very small on the interval  $(0, 1)$  – the maximum value is  $h(\frac{1}{2}) = e^{-8} \approx 0.000335 \dots$  – and it decays to zero so quickly that it's hard to see from the graph that  $h(x) > 0$  for most  $x$  on this interval, but  $h(x)$  is indeed positive there.

9. Define functions  $f_n : \mathbb{R} \rightarrow \mathbb{R}$  by  $f_n(x) = \sqrt{x^2 + \frac{1}{n^2}}$  for all  $n \geq 1$ .

- (a) Prove that  $f_n$  is continuously differentiable, and that  $|x| \leq f_n(x) \leq |x| + \frac{1}{n}$ .
- (b) Prove that  $(f_n)$  converges uniformly to a continuous function  $f$ .

- (c) Prove that  $(f'_n)$  doesn't converge uniformly on  $[-1, 1]$ , so the theorem from lecture about limits of differentiable functions doesn't apply to tell us that  $f$  should be differentiable on  $[-1, 1]$ . (Is  $f$  differentiable there?)

*Solution.* (a) Since  $\sqrt{x} = x^{1/2}$  is differentiable at  $x^2 + \frac{1}{n^2} > 0$ , we can apply the chain rule:

$$f'_n(x) = \frac{d}{dx} \left( \left( x^2 + \frac{1}{n^2} \right)^{1/2} \right) = \frac{1}{2} \left( x^2 + \frac{1}{n^2} \right)^{-1/2} (2x) = \frac{x}{\sqrt{x^2 + \frac{1}{n^2}}}.$$

We also use the fact that  $\sqrt{x}$  is monotone increasing to check that

$$|x| = \sqrt{|x|^2} < \sqrt{|x|^2 + \frac{1}{n^2}} \leq \sqrt{|x|^2 + \frac{2|x|}{n} + \frac{1}{n^2}} = |x| + \frac{1}{n}.$$

The middle term is  $f_n(x)$  since  $|x|^2 = x^2$ , so we have  $|x| < f_n(x) \leq |x| + \frac{1}{n}$ .

- (b) The continuous function is  $f(x) = |x|$ , since for any  $\epsilon > 0$ , part (a) says that

$$n > \frac{1}{\epsilon} \implies |f_n(x) - f(x)| \leq \frac{1}{n} < \epsilon$$

for all  $x \in \mathbb{R}$ .

- (c) If  $f'_n$  converges uniformly on  $[-1, 1]$ , then its pointwise limit should be continuous. But a quick computation shows that

$$\lim_{n \rightarrow \infty} f'_n(x) = \begin{cases} 1, & x > 0 \\ 0, & x = 0 \\ -1, & x < 0 \end{cases}$$

is discontinuous at  $x = 0$ , which is exactly where  $f(x) = |x|$  is not differentiable.

10. In an upcoming lecture, we'll need to know that  $\lim_{x \rightarrow \infty} xs^{x-1} = 0$  for all  $s \in (0, 1)$ .

- (a) Prove that for all  $c > 0$ , there exists  $N > 0$  such that  $\log(x) < cx$  for all  $x \geq N$ .  
 (b) Prove for  $s \in (0, 1)$  that  $\lim_{x \rightarrow \infty} xs^x = 0$ , and that this implies the above claim.

*Solution.* (a) It's enough to prove that  $\lim_{x \rightarrow \infty} \frac{\log(x)}{x} = 0$ , since then there's an  $N > 0$  such that  $0 < \frac{\log(x)}{x} < c$  for all  $x \geq N$ . This limit exists by l'Hôpital's rule, which says that it is equal to  $\lim_{x \rightarrow \infty} \frac{1/x}{1} = 0$ .

- (b) For any  $c > 0$ , part (a) says that  $0 < xs^x < e^{cx}s^x = (e^c s)^x$  for all large enough  $x$ . Since  $0 < s < 1$ , we can choose a positive  $c < \log(1/s)$  so that  $0 < e^c s < 1$ , and then

$$\lim_{x \rightarrow \infty} (e^c s)^x = 0.$$



Thus the squeeze theorem says that  $\lim_{x \rightarrow \infty} xs^x = 0$  as well. We conclude that

$$\lim_{x \rightarrow \infty} xs^{x-1} = \frac{1}{s} \left( \lim_{x \rightarrow \infty} xs^x \right) = 0.$$