Session 6: Protein Structure

KB8029 — Bioinformatics & Deep Learning

Arne Elofsson

SciLifeLab

SciLifeLab

Log in to the lecture quiz

Log in to the quiz

Scan the code, or go to quiz.bioinfo.se/live

Log in with your university account

Answering counts as being at the lecture: up to 3 bonus points on the exam

Right or wrong is never graded, only whether you answered. Results are shown without names.

quiz.bioinfo.se/live

SciLifeLab

Warm-up: what do you know already?

3 quick questions

On the topics of today, one at a time, about 30 seconds each

Answer on your own, and guess if you do not know

It shows me where to start; right or wrong is never graded

SciLifeLab

An α-helix is held together mainly by …

A Disulphide bonds

B Backbone hydrogen bonds, residue i to i + 4

C Salt bridges between side chains

D Hydrogen bonds between neighbouring strands

Answer now
Scan the code, or go to
quiz.bioinfo.se/live

SciLifeLab

What does a protein's contact map show?

A Which residues are close to each other in 3D

B Which proteins interact in the cell

C The hydrophobicity along the sequence

D Where in the cell the protein is found

Answer now

Scan the code, or go to

quiz.bioinfo.se/live

SciLifeLab

During evolution, which usually changes more slowly?

A The sequence

B Both change equally fast

C The structure

D Neither changes

Answer now

Scan the code, or go to

quiz.bioinfo.se/live

SciLifeLab

What the room knows already

Where we start

Which topics most of you know, and which we take slowly today.

Show the results ▸ quiz.bioinfo.se/live/report

Your questions before today

What you asked

The questions you sent in on Canvas before the lecture

The most common ones now; the others in the block where they belong

SciLifeLab

In up to three words: what do you want to learn today?

Four levels

sequence to 3D, contact maps

Comparing in 3D

RMSD, TM-score, DALI, Foldseek

Domains and states

CATH, T and R, prediction

Answer now

Scan the code, or go to
quiz.bioinfo.se/live

SciLifeLab


What you want to learn today

What the room wants to learn

Show the results ▸ quiz.bioinfo.se/live/report

SciLifeLab


What does a sequence encode in 3D?



- **Today the structure is the data**
- Coordinates → distances, domains, interfaces
- Sessions 7-12: can a computer predict it?

See course.bioinfo.se/session06.html, 'Learning goals'.

Three blocks: about 30 min talk and 15 min discussion



Block 1: the four levels of structure and how to represent a structure. Block 2: comparing structures (alignment, RMSD, TM-score, DALI, Foldseek). Block 3: domains, conformational states, and structure as a prediction target.

Times are approximate: a 10 min start (warm-up quiz and your questions), two 15 min breaks and a 5 min close. Each block ends with a quiz question: guess, run the notebook and work out why it behaves as it does, discuss, answer again.

SciLifeLab

BLOCK 1 · ABOUT 30 MIN

Sequence becomes structure

SciLifeLab

Four levels of structure, all in haemoglobin

primary

the sequence

VHLTPEEKSA
VTALWGKVNV
DEVGGEALGR
LLVVYPWTQR
FFESFGDLST
PDAVMGNPKV
…


secondary

one α-helix




tertiary

one chain's fold



quaternary

four chains



Primary: the amino-acid sequence. Secondary: local backbone shapes (α-helix, β-strand, coil). Tertiary: the 3D fold of one chain. Quaternary: several chains packed together. Rendered from PDB 2HHB with PyMOL.

SciLifeLab

Each level comes with its own computational task

level	what it is	computational task	where
primary	sequence	alignment, BLAST, profiles	Sessions 3-5
secondary	helix, strand, coil	prediction from sequence	Sessions 7-11
tertiary	one chain's fold	comparison, prediction	today; Session 12
quaternary	several chains	interfaces, complexes	today; Session 12

The course follows the levels in order: sequences (Sessions 3-5), secondary structure (Sessions 7-11), folds and complexes (today and Session 12).

See course.bioinfo.se/session06.html, Table 6.1.

Helix, strand, coil

Ubiquitin (PDB 1UBQ)




helix strand coil

- **Helix: H-bonds between residues i and i+4**
- **Strand: H-bonded edge to edge in a sheet**
- Coil: everything else
- DSSP assigns them from coordinates

Ubiquitin (PDB 1UBQ), rendered with PyMOL.

The fold hides its hydrophobic side chains


β-globin: hydrophobic side chains (orange)



- **Hydrophobic effect: non-polar side chains away from water**
- Plus H-bonds, electrostatics, van der Waals
- Many proteins work in complexes

β-globin (2HHB chain B); Ala, Val, Ile, Leu, Met, Phe, Trp, Cys as spheres.

Haemoglobin: two α and two β chains




- **α: 141 residues; β (HBB): 146**
- One haem per chain (orange: iron)
- **One β chain: 80.8% helix, 0% strand**

PDB 2HHB, rendered with PyMOL; numbers from session06-protein-structure-in-code.ipynb.

SciLifeLab

Does the sequence determine the structure?

Anfinsen, 1960s: ribonuclease refolds on its own




- **Often yes**
- That makes prediction possible
- Not a law for every protein

But not always: chaperones · ligands and partners · alternative states · disorder

See course.bioinfo.se/session06.html, 'Does the sequence determine the structure?'.

SciLifeLab


Local is easy, long-range is hard



- **Local: secondary structure**
- **Long-range: the tertiary fold**
- A window of sequence sees only local

See course.bioinfo.se/session06.html, 'Does the sequence determine the structure?'.

A structure as a matrix: distance and contact maps




β-globin as a structure, a distance map and a contact map (Cβ within 8 Å). Diagonal band: each helix touching itself (local). Off-diagonal stripes: helices packed together (long-range). The maps do not change when the protein is rotated.

tools/builders/make_session06_figures.py; notebook.

SciLifeLab

One protein, many representations





Each representation makes different patterns easy to see. Later in the course it also decides which patterns a model can easily learn.


See course.bioinfo.se/session06.html, callout 'Representations'.

SciLifeLab

PyMOL: three ways to draw the same chain







Cartoon shows the secondary structure, sticks every bond (for a side chain or a ligand), spheres every atom (for packing). The lab uses PyMOL hands-on; the notebook drives it from Python.


β-globin (2HHB chain B), haem in red.

SciLifeLab

What does dss find for myoglobin?

Haemoglobin β: 80.8% helix, 0% strand, but it works only as four chains. Myoglobin: the same fold, one chain.





A About 80% helix, 0% strand: the same

B About 50% helix: some helices unfold

C About 40% helix, 30% strand

D Hardly any helix or strand: mostly loop

Guess on your own

Scan the code, or go to
quiz.bioinfo.se/live

SciLifeLab

What does dss find for myoglobin?

How did the room answer?

Halt the round when most have answered, then show the results.

Show the results ▸ quiz.bioinfo.se/live/report

SciLifeLab

What does dss find for myoglobin?

Fix it, run it, discuss

Open the notebook: scan the code

Fix the line marked FIX, then run the cells

Did it come out as you guessed? Discuss with your neighbour

The notebook: https://colab.research.google.com/github/ElofssonLab/kb8029-book/blob/main/notebooks/session06-discussion-1.ipynb

SciLifeLab

What does dss find for myoglobin?

Answer again

Same question, after the discussion.

quiz.bioinfo.se/live

SciLifeLab

What does dss find for myoglobin?

Results, and who explains

Both rounds side by side.


Draw two to explain: one of you says how you reasoned.

Show the results ▸ quiz.bioinfo.se/live/report

To check it with code afterwards: discussion notebook 1 https://colab.research.google.com/github/ElofssonLab/kb8029-book/blob/main/notebooks/session06-discussion-1.ipynb

SciLifeLab

The answer: the same fold, alone or in a tetramer



Myoglobin is 79.7% helix and 0% strand, haemoglobin's β chain 80.8% and 0%. Secondary and tertiary structure are properties of one folded chain; quaternary structure is a separate level on top. Haemoglobin needs four chains for cooperative oxygen binding, not for its fold. And a result of exactly zero is worth a second look.

SciLifeLab

BLOCK 1 DONE

Break


SciLifeLab

BLOCK 2 • ABOUT 30 MIN

Comparing proteins in 3D

SciLifeLab


A structure alignment pairs residues by their place in 3D



- **Sequence alignment: from substitution scores**
- **Structure alignment: from coordinates**
- Both are hypotheses about correspondence

Superposition with PyMOL; TM-score from the notebook (TM-align, tmtools).

Sequence identity fades, the fold stays




- **Five proteins vs β-globin**
- Identity 43% → 6%
- TM-score stays above 0.5 for all globins

Notebook: Smith-Waterman/BLOSUM62 and TM-align (tmtools). TM-scores normalised by the length of β-globin.

SciLifeLab

When sequence signal is gone, structure still finds the relative




- **Truncated Hb: no better than shuffled**
- 0% of its pairs right
- **Structure is more conserved than sequence**

Notebook: best Smith-Waterman score against the best of 200 alignments to shuffled sequences.

SciLifeLab

RMSD and TM-score weight a bad pair differently




RMSD: the root-mean-square distance between corresponding atoms after superposition; it grows with length, one misplaced loop inflates it, and it means little without the number of aligned residues. TM-score: weights each pair by how close it is and normalises by length, from 0 to 1.

See course.bioinfo.se/session06.html, 'Measuring structural similarity: RMSD and TM-score'.

SciLifeLab

Reading a TM-score




Unrelated proteins typically score 0.2-0.3; above about 0.5 usually means the same fold. A rule of thumb, not a law: scores in between need other evidence.

TM-scores from the notebook.

SciLifeLab

DALI: comparing patterns of internal distances



β-globin and myoglobin (25% identical) give almost the same distance map; GFP's β-barrel does not. Sequence alignment compares residues; DALI compares residue-residue geometry.

Holm & Sander (1993). Figure: tools/builders/make_session06_figures.py.

SciLifeLab

Searching by sequence, searching by structure

Sequence (Session 5)

BLAST, MMseqs2

- Search with the amino-acid sequence
- Billions of sequences

Structure (today)

Foldseek

- Search with a 'structural sequence'
- Hundreds of millions of structures in minutes

van Kempen, Steinegger et al. (2024), Nature Biotechnology 42:243-246.

SciLifeLab

Foldseek turns a structure into a string



the 20 states were learned by a variational autoencoder, not designed by hand

Each residue's local 3D environment becomes one of 20 '3Di' letters. The structure is then searched with fast sequence-alignment machinery instead of expensive superposition.

See course.bioinfo.se/session06.html, 'Foldseek' and the callout 'Machine-learning preview'.

SciLifeLab

A worked comparison with Foldseek

- **Same protein, two states: T (2HHB) vs R (1HHO)**
- α vs β globin (43% identical): TM 0.90-0.91
- **2HHB vs GFP: no hits, even with a loose threshold**

0.98
TM-score
T vs R


0.7 Å
RMSD
T vs R

Foldseek run locally on 2HHB, 1HHO and 1EMA (course.bioinfo.se/session06.html).

SciLifeLab

What does dss find for GFP?

The globins: about 80% helix, 0% strand. GFP (1EMA): a barrel of β-strands around one central helix.



A About 80% helix, 0% strand, like the globins

B About 50% helix, 50% strand

C About 10% helix, 54% strand

D About 0% helix, 100% strand

Guess on your own

Scan the code, or go to

quiz.bioinfo.se/live

SciLifeLab

What does dss find for GFP?

How did the room answer?

Halt the round when most have answered, then show the results.

Show the results ▸ quiz.bioinfo.se/live/report

SciLifeLab

What does dss find for GFP?

Fix it, run it, discuss

Open the notebook: scan the code

Fix the line marked FIX, then run the cells

Did it come out as you guessed? Discuss with your neighbour

The notebook: https://colab.research.google.com/github/ElofssonLab/kb8029-book/blob/main/notebooks/session06-discussion-2.ipynb

SciLifeLab

What does dss find for GFP?

Answer again

Same question, after the discussion.

quiz.bioinfo.se/live

SciLifeLab

What does dss find for GFP?

Results, and who explains

Both rounds side by side.


Draw two to explain: one of you says how you reasoned.

Show the results ▸ quiz.bioinfo.se/live/report

To check it with code afterwards: discussion notebook 2 https://colab.research.google.com/github/ElofssonLab/kb8029-book/blob/main/notebooks/session06-discussion-2.ipynb

SciLifeLab

The answer: GFP is mostly strand



GFP is 9.8% helix and 53.8% strand, almost the mirror image of the globins. The globins are an all-α family, not a typical protein, so secondary-structure prediction cannot rely on one fixed rule: it has to learn position-specific patterns (Session 7). A copied block of code that still names the old object is a common mistake; a function called twice avoids it.

SciLifeLab

BLOCK 2 DONE


Break

SciLifeLab

BLOCK 3 · ABOUT 30 MIN

Domains, states, and structure as a prediction target

CATH classifies domains in four levels




- **Domain: a compact, reusable unit**
- Grouped by 3D shape
- Myoglobin, neuroglobin, truncated Hb: same code

Orengo et al. (1997); live PDBe CATH lookups. GFP: class 2 (Mainly Beta), Beta Barrel.

SciLifeLab

Structure sees one superfamily, sequence sees two families



CATH puts the truncated haemoglobin in the same superfamily as β-globin. Pfam, working from sequence, needs a separate family (PF01152, 'Bacterial-like globin') and joins it to PF00042 'Globin' only in a clan.

Live PDBe CATH and InterPro lookups (course.bioinfo.se/session06.html).

SciLifeLab

Other classifications, other philosophies

SCOP

1995

- Expert manual curation

ECOD

2014

- Organised by evolutionary relationship

TED

2024


- Parsed automatically from the whole AlphaFold Database

Murzin et al. (1995); Cheng et al. (2014); Lau, Bordin et al. (2024).

SciLifeLab

Interfaces and ligands: where function happens

Deoxyhaemoglobin: interface atoms of one α chain (blue) and the β chain it touches (orange) as spheres; haem in red




- **Interface: where two chains touch**
- More hydrophobic, more conserved
- Ligands: haem, ions, small molecules
- Predicting them: Session 12

Deoxyhaemoglobin (2HHB): atoms within 5 Å of the other chain as spheres, rendered with PyMOL.

SciLifeLab

One protein, two structures: haemoglobin T and R

Superimposed on one α/β pair: the other pair has rotated



- **Each chain hardly changes (TM 0.98)**
- The α/β pairs rotate ≈13-15°
- β-β Fe-Fe: 39.5 → 34.6 Å
- Cooperative oxygen binding

Baldwin & Chothia (1979); notebook (Fe-Fe distances); 1HHO biological assembly.

SciLifeLab

A PDB entry is one snapshot, not 'the' structure

- **States depend on ligands, partners, conditions**
- Matters whenever a model is compared with 'the' experiment
- **AlphaFold usually predicts one state (Session 12)**

4.9 Å

β-β haem Fe-Fe contraction, T → R

0.98


TM-score of each chain, T vs R

Notebook: session06-protein-structure-in-code.ipynb.

SciLifeLab

Secondary structure as a prediction target

input: a window of sequence (or profile) around each residue



target: H, E or C for every residue, from solved structures (DSSP); β-globin residues 51-76

Input: the sequence (or its profile), one window per residue. Target: H, E or C for every residue, with labels from solved structures. What could help? The residue itself, its neighbours, and how the position varies among homologs (Session 5).

β-globin residues 51-76 with the PDB entry's secondary structure; window 57-69 around His63 (as in Session 7).

SciLifeLab

Four ways to predict a whole fold

physics	comparative	contact prediction	deep learning
lowest-energy chain; Rosetta fragments	a related known structure (S4-5)	coevolving pairs in alignments (S5)	AlphaFold2, CASP14 (S12)

CASP: blind tests on unpublished structures since 1994

Physics is rigorous but expensive; comparative modelling needs a template; coevolution fills in a contact map; deep learning (AlphaFold2, 2020) combined all of it. 2024 Nobel Prize in Chemistry: Baker, Hassabis, Jumper.

SciLifeLab

Residues in contact mutate together

A	K	L	E	D	G	E
A	R	L	E	E	G	D
S	K	V	E	D	A	E
A	E	L	Q	K	G	K
S	E	V	Q	K	A	R

→ these two residues are probably in contact

illustrative: K/R in one column goes with E/D in the other, and vice versa

- **Large alignments reveal coevolving pairs**
- Coevolving pairs → a contact map
- Session 12: AlphaFold

Illustrative alignment, not real sequences. See course.bioinfo.se/session06.html, 'Predicting the whole fold'.

One structure becomes many

ensembles	inside the cell	Nobel Prize 2024
AF-Cluster: AlphaFold2 samples both folds of the fold-switching protein KaiB	cryo-electron tomography: RNA polymerase and ribosome coupled, inside Mycoplasma cells	Baker: computational design Hassabis & Jumper: structure prediction
Wayment-Steele et al. 2024	*O'Reilly et al. 2020*	*Chemistry*

haemoglobin has (at least) two: T and R. Can one predicted structure ever be enough?

- Proteins move: ensembles, not snapshots
- Structures inside cells, by cryo-ET
- Nobel Prize in Chemistry 2024
- **Can one predicted structure ever be enough?**

Wayment-Steele et al. (2024), Nature 625:832; O'Reilly et al. (2020), Science 369:554; nobelprize.org (2024).

SciLifeLab

The course's path from sequence to structure




Today: structures as data, compared by alignment, TM-score, distance maps and Foldseek, and the reminder that one entry is one state. Next: models that learn the steps from sequence to structure.

See course.bioinfo.se/session06.html, 'Predicting the whole fold'.

SciLifeLab

Which iron–iron distance changes the least?

T (2HHB) to R (1HHO): each chain hardly changes shape, but the β1–β2 iron–iron distance shrinks by about 5 Å.



Fe–Fe distances, T (2HHB) → R (1HHO)

β1–β2: about 5 Å shorter

each chain: Cα RMSD below 1 Å

Which pair changes the least?

A α1–β1, the "fixed" interface

B α1–β2, the "sliding" interface

C α1–α2

D None: all shrink by about 5 Å

Guess on your own

Scan the code, or go to
quiz.bioinfo.se/live

SciLifeLab

Which iron–iron distance changes the least?

How did the room answer?

Halt the round when most have answered, then show the results.

Show the results ▸ quiz.bioinfo.se/live/report

SciLifeLab

Which iron–iron distance changes the least?

Fix it, run it, discuss

Open the notebook: scan the code

Fix the line marked FIX, then run the cells

Did it come out as you guessed? Discuss with your neighbour

The notebook: https://colab.research.google.com/github/ElofssonLab/kb8029-book/blob/main/notebooks/session06-discussion-3.ipynb

SciLifeLab

Which iron–iron distance changes the least?

Answer again

Same question, after the discussion.



quiz.bioinfo.se/live

SciLifeLab

Which iron–iron distance changes the least?

Results, and who explains

Both rounds side by side.


Draw two to explain: one of you says how you reasoned.

Show the results ▸ quiz.bioinfo.se/live/report

To check it with code afterwards: discussion notebook 3 https://colab.research.google.com/github/ElofssonLab/kb8029-book/blob/main/notebooks/session06-discussion-3.ipynb

SciLifeLab

The answer: α1–α2 barely moves; one distance is not the change



the two α/β pairs rotate about 13° as rigid bodies

From T to R, β1–β2 shrinks by 4.9 Å but α1–α2 changes by only 0.6 Å. The chains hardly change shape: the two α/β pairs rotate about 13° against each other as rigid bodies, and a rotation moves each atom by an amount that depends on its distance from the axis. Describing a conformational change needs the rigid units and their motion, not one distance.

SciLifeLab

In up to three words: what was the hardest part of today?

Four levels

sequence to 3D, contact maps

Comparing in 3D

RMSD, TM-score, DALI, Foldseek

Domains and states

CATH, T and R, prediction

Answer now

Scan the code, or go to
quiz.bioinfo.se/live

SciLifeLab

The hardest part of today

What the room found hardest

Show the results ▸ quiz.bioinfo.se/live/report

Session 7 next

Session 7: the PSSM's weighted sum becomes a perceptron with learned weights
First target: the secondary structure you saw today