

MA2001 Linear Algebra I Notes

Contents

1	Linear Systems & Gaussian Elimination	17
1.1	Linear Systems & Their Solutions	17
1.1.1	Linear Equations	17
1.1.1.1	Zero Equation	17
1.1.1.2	Nonzero Equation	17
1.1.1.3	Inconsistent Equation	17
1.1.2	Solutions of a Linear Equation	17
1.1.2.1	Solution Set	18
1.1.2.2	General Solution	18
1.1.3	Solution Set of Zero Equation	18
1.1.4	Solution Set of Inconsistent Linear Equation	18
1.1.5	General Solution of Consistent Nonzero Equation	18
1.1.6	Geometrical Interpretation of Linear Equations	18
1.1.6.1	In Two Variables	18
1.1.6.2	In Three Variables	19
1.1.7	Linear Systems	19
1.1.7.1	Zero System	19
1.1.7.2	Nonzero System	19
1.1.8	Solution of Linear Systems	19
1.1.8.1	Solution Set	19
1.1.8.2	General Solution	19
1.1.9	Consistency of Linear Systems	20
1.1.10	Number of Solutions to Linear Systems	20
1.1.11	Geometric Interpretation of Linear Systems	20
1.1.11.1	In Two Variables of Two Equations	20
1.1.11.2	In Three Variables of Two Equations	20
1.2	Elementary Row Operations	21
1.2.1	Augmented Matrix Representation of a Linear System	21
1.2.2	Elementary Row Operations on Augmented Matrices	21
1.2.2.1	Correspondence to Operations on Equations in Linear System	21
1.2.2.2	Interchanging two rows can be decomposed further	22
1.2.2.3	Inverse Elementary Row Operations	22

1.2.3	Row Equivalent Matrices	22
1.2.4	Row Equivalence as an Equivalence Relation	22
1.2.5	Row Equivalence Implies Same Solution Set	22
1.3	Row-Echelon Form	23
1.3.1	Leading Entry	23
1.3.2	Row-Echelon Form (REF)	23
1.3.3	Pivot Points, Pivot Columns, and Non-pivot Columns	23
1.3.3.1	Pivot Point	23
1.3.3.2	Pivot and Non-pivot Columns	23
1.3.3.3	Each Pivot Column Contains Exactly One Pivot Point	23
1.3.4	Reduced Row-Echelon Form (RREF)	23
1.3.5	General Solution From Row-echelon Form	24
1.3.6	General Solution From Reduced Row-echelon Form	24
1.4	Gaussian Elimination	24
1.4.1	REF/RREF of Augmented Matrices	24
1.4.2	Gaussian Elimination: Finding REF of Augmented Matrices	24
1.4.3	Gauss-Jordan Elimination: Finding RREF of Augmented Matrices	25
1.4.4	Infinitely Many REFs	25
1.4.5	Uniqueness of RREF	25
1.4.6	Deducing Consistency of Linear Systems from its REF	25
1.4.6.1	Inconsistent With No Solutions	25
1.4.6.2	Consistent With One Solution	26
1.4.6.3	Consistent With Infinitely Many Solutions & Arbitrary Parameters	26
1.4.7	Geometrical Interpretation of Linear Systems with RREF	26
1.4.7.1	Linear System in Three Variables of Three Equations	26
1.5	Homogeneous Linear Systems	26
1.5.1	Homogeneous Linear Equation	26
1.5.1.1	Remarks	27
1.5.2	Geometrical Interpretation of Homogeneous Linear Equations	27
1.5.2.1	In Two Variables	27
1.5.2.2	In Three Variables	27
1.5.3	Homogeneous Linear System	27
1.5.4	Trivial Solution of Homogeneous Linear Systems	27
1.5.5	Geometric Interpretation of Homogeneous Linear Systems	28
1.5.5.1	In Two Variables of Two Equations	28
1.5.5.2	In Three Variables of Two Equations	28
2	Matrices	29
2.1	Introduction to Matrices	29
2.1.1	Definition of Matrix	29
2.1.1.1	Remarks	29

2.1.2	Notation of Matrices	30
2.1.3	Row Matrix	30
2.1.4	Column Matrix	30
2.1.5	Square Matrix	30
2.1.5.1	Order of Square Matrix	30
2.1.5.2	Diagonal / Diagonal Entries of Square Matrix	30
2.1.6	Diagonal Matrix	31
2.1.7	Scalar Matrix	31
2.1.8	Identity Matrix	31
2.1.8.1	Notation for Identity Matrix	31
2.1.9	Zero Matrix	31
2.1.9.1	Notation for Zero Matrix	31
2.1.10	Symmetric Square Matrix	31
2.1.11	Upper Triangular Square Matrix	32
2.1.12	Lower Triangular Square Matrix	32
2.1.13	Triangular Square Matrix	32
2.1.14	Diagonal and Triangular Square Matrices	32
2.2	Matrix Operations	32
2.2.1	Identical Matrices	32
2.2.2	Matrix Addition, Subtraction & Scalar Multiplication	33
2.2.3	Properties of Matrix Addition, Subtraction & Scalar Multiplication	33
2.2.4	Matrix Multiplication	34
2.2.4.1	Matrix Multiplication Explained in Words	34
2.2.5	Noncommutativity of Matrix Multiplication	34
2.2.6	Properties of Matrix Multiplication	34
2.2.7	Nonnegative Integer Powers of Square Matrices	35
2.2.7.1	Multiplying a Matrix With Itself	35
2.2.7.2	Nonnegative Integer Powers of Square Matrices	35
2.2.8	Properties of Nonnegative Integer Powers of Square Matrices	35
2.2.9	Matrix Representation	35
2.2.9.1	Matrix as a Column of Row Vectors	36
2.2.9.2	Matrix as a Row of Column Vectors	36
2.2.10	Decomposing Matrix Multiplication	36
2.2.10.1	Decomposing Into Entries	37
2.2.10.2	Decomposing Into Rows	37
2.2.10.3	Decomposing Into Columns	37
2.2.10.4	Decomposing Into Blocks	37
2.2.11	Matrix Representation of Linear Equation	37
2.2.12	Matrix Representation of Linear System	38
2.2.12.1	Solution Vector to Linear System	38

2.2.12.2	Alternative Matrix Representation of Linear System	39
2.2.13	Transpose of Matrix	39
2.2.14	Properties of Transpose of Matrix	39
2.3	Inverse of Square Matrices	39
2.3.1	Additive Inverse of Matrices	39
2.3.2	Inverse of Square Matrices	40
2.3.3	Uniqueness of Inverse of Invertible Matrices	40
2.3.4	Cancellation Law for Invertible Matrices	40
2.3.5	Inverse of Square Matrix of Order Two	40
2.3.6	Properties of Invertible Matrices	41
2.3.7	Negative Integer Powers of Square Matrices	41
2.3.8	Properties of Integer Powers of Square Matrices	41
2.4	Elementary Matrices	41
2.4.1	Definition of Elementary Matrices	41
2.4.2	Connection of Elementary Matrices to Pre-Matrix Multiplication	41
2.4.3	Elementary Matrices are Invertible	42
2.4.4	Row Equivalent Matrices are Linked by Elementary Matrices	42
2.4.5	Main Theorem for Invertible Matrices	42
2.4.6	Checking Whether a Square Matrix is Invertible	43
2.4.7	Finding Inverse of an Invertible Matrix	43
2.4.8	Weaker Requirement for Invertible Matrices	43
2.4.9	Invertibility of Product of Matrices	43
2.4.10	Elementary Column Operations	43
2.4.11	Performing a Single Elementary Column Operation on Identity Matrix Gives an Elementary Matrix	44
2.4.12	Connection of Elementary Matrices to Matrix Post-Multiplication	44
2.5	Determinant	44
2.5.1	(i, j) -cofactor	44
2.5.2	Definition of Determinant of Square Matrix	44
2.5.3	Determinant of 2×2 Matrix	45
2.5.4	Broken Diagonal Formula for Determinant of 3×3 Matrix	45
2.5.5	Properties of Determinant	45
2.5.5.1	Properties of Determinant	45
2.5.5.2	Change in Determinant by Elementary Row Operations	45
2.5.5.3	Change in Determinant by Elementary Matrices	46
2.5.5.4	Change in Determinant Between Row Equivalent Matrices	46
2.5.5.5	Determinant of Square Matrix With Zero Row	46
2.5.5.6	Invertibility and Determinant	46
2.5.5.7	Determinant of Triangular Matrix	46
2.5.6	Cofactor Expansion	47

2.5.7	Finding Determinant Efficiently	47
2.5.8	Adjoint Matrix	47
2.5.8.1	Definition of Adjoint Matrix	47
2.5.8.2	Properties of Adjoint Matrix	48
2.5.9	Cramer's Rule	48
3	Vector Spaces	49
3.1	Euclidean n -Spaces	49
3.1.1	Introduction to n -vectors	49
3.1.1.1	Definition of n -vector	49
3.1.1.2	Zero Vector	49
3.1.2	n -vector Operations	49
3.1.2.1	Equal n -vectors	49
3.1.2.2	n -vector Addition, Subtraction & Scalar Multiplication	49
3.1.3	Properties of n -vectors	50
3.1.4	n -vectors as Matrices	50
3.1.5	Vectors in xy -Plane	50
3.1.5.1	Vectors as Points on the xy -Plane	50
3.1.5.2	Vectors as Change from Initial Point to End Point	51
3.1.5.3	Length of Vectors	51
3.1.5.4	Properties of Length of Vectors	51
3.1.5.5	Geometrical Interpretation of Scalar Multiplication	51
3.1.5.6	Geometrical Interpretation of Vector Addition	51
3.1.5.7	Geometrical Interpretation of Vector Subtraction	51
3.1.6	Vectors in xyz -Space	52
3.1.6.1	Vectors as Points on the xyz -Space	52
3.1.7	Euclidean Spaces	52
3.1.8	Implicit and Explicit Forms of Linear Systems	52
3.1.8.1	Definition of Implicit and Explicit Forms	52
3.1.8.2	Lines in \mathbb{R}^2	53
3.1.8.2.1	Vector Equation for line in \mathbb{R}^2	53
3.1.8.2.2	Implicit to Explicit Form	53
3.1.8.2.3	Explicit to Implicit Form	53
3.1.8.3	Planes in \mathbb{R}^3	53
3.1.8.3.1	Vector Equation for plane in \mathbb{R}^3	53
3.1.8.3.2	Implicit to Explicit Form	53
3.1.8.3.3	Explicit to Implicit Form	53
3.1.8.4	Lines in \mathbb{R}^3	54
3.1.8.4.1	Vector Equation for line in \mathbb{R}^3	54
3.1.8.4.2	Implicit to Explicit Form	54
3.1.8.4.3	Explicit to Implicit Form	54

3.2	Linear Combinations and Linear Spans	54
3.2.1	Definition of Linear Combination	54
3.2.2	Definition of Linear Span	55
3.2.3	Criterion for $\text{span}(S) = \mathbb{R}^n$	55
3.2.3.1	When $k \geq n$	55
3.2.3.2	When $k < n$	55
3.2.4	Properties of Linear Spans	55
3.2.4.1	Linear Spans Always Contains Zero Vector	55
3.2.4.2	Linear Spans Are Closed Under Linear Combination	56
3.2.4.3	When a Span is a Subset of Another Span	56
3.2.4.4	Linear Spans with Redundant Vectors	56
3.2.5	Criterion for Vector Belongs to Span	56
3.3	Subspaces	57
3.3.1	Definition of Subspaces	57
3.3.2	Zero Space	57
3.3.3	Euclidean n-Space is a Subspace	57
3.3.4	Showing That a Subset is a Subspace	57
3.3.5	Showing That a Subset is Not a Subspace	57
3.3.6	Subspaces of $\mathbb{R}^1, \mathbb{R}^2, \mathbb{R}^3$	58
3.3.6.1	Subspaces of $\mathbb{R}^1 = \mathbb{R}$	58
3.3.6.2	Subspaces of \mathbb{R}^2	58
3.3.6.3	Subspaces of \mathbb{R}^3	58
3.3.7	Solution Set of Homogeneous Linear System is a Subspace of \mathbb{R}^n	58
3.3.8	Solution Space	58
3.3.9	A Subspace of \mathbb{R}^n is Always The Solution Space of a Homogeneous Linear System	59
3.4	Vector Spaces	59
3.4.1	Definition of Vector Space	59
3.4.2	Expanded Definition of Subspace	59
3.5	Linear Independence	59
3.5.1	Definition of Linear Independence	59
3.5.2	Linear Independence of Subset/Superset	59
3.5.2.1	Superset Preserves Linear Dependence	59
3.5.2.2	Subset Preserves Linear Independence	59
3.5.3	Linear Independence of Sets Containing Zero Vector	60
3.5.4	Linear Independence of Sets Containing Only One Vector	60
3.5.5	Linear Independence and Linear Combinations	60
3.5.6	Cardinality Upper Bound for Linear Independence	60
3.5.7	Adding Vectors to a Linearly Independent Set	60
3.6	Bases	60

3.6.1	Definition of Bases	60
3.6.1.1	Remarks	60
3.6.2	Basis and Unique Linear Combination	61
3.6.3	Coordinate Vector	61
3.6.3.1	Remark	61
3.6.4	Standard Basis	61
3.6.5	Properties of Coordinate Vector	61
3.6.5.1	Zero Coordinate Vector	61
3.6.5.2	Homogeneity of Degree One / Scalar Multiplication Preserving	61
3.6.5.3	Additivity / Addition Preserving	62
3.6.5.4	Equal Coordinate Vectors	62
3.6.5.5	Linear Combination Preserving	62
3.6.5.6	Linear Independence Preserving	62
3.6.5.7	Span Preserving	62
3.7	Dimensions	62
3.7.1	Dimension Theorem for Vector Spaces	62
3.7.1.1	Corollary	62
3.7.2	Definition of Dimension	62
3.7.2.1	Examples	63
3.7.3	Dimension of Solution Space	63
3.7.4	Easier Criterion for Basis	63
3.7.5	Dimension for Subspaces	63
3.7.5.1	Corollary	63
3.7.6	Invertibility and Rows and Columns as Basis	64
3.8	Transition Matrices	64
3.8.1	Coordinate Vector As a Column Vector	64
3.8.2	Definition of Transition Matrix	64
3.8.3	Pre-Multiplication of Transition Matrix	64
3.8.4	Inverse of Transition Matrices	64
3.8.4.1	Transition Matrices Are Invertible	64
3.8.4.2	Inverse is a Transition Matrix	64
4	Vector Spaces Associated With Matrices	65
4.1	Row Spaces and Column Spaces	65
4.1.1	Definition of Row Spaces and Column Spaces	65
4.1.1.1	Row Space	65
4.1.1.2	Column Space	65
4.1.2	Row Equivalent Matrices And Their Rows	65
4.1.2.1	Row Equivalence Implies Same Row Space	65
4.1.2.1.1	Remark	66
4.1.2.2	Nonzero Rows of REF are Linearly Independent	66

4.1.2.3	Finding Basis/Dimension For Row Space	66
4.1.3	Row Equivalent Matrices And Their Columns	66
4.1.3.1	Row Equivalence Preserve Linear Relations Between Columns .	66
4.1.3.2	Pivot Columns of REF is a Basis For Column Space	66
4.1.3.3	Finding Basis/Dimension For Column Space	66
4.1.4	Finding Basis for Vector Space Spanned By a Set of Vectors	66
4.1.4.1	Method 1: From New Vectors Using Row Space	67
4.1.4.2	Method 2: Select From Original Vectors Using Column Space .	67
4.1.5	Consistency of Linear Systems And Column Space of Coefficient Matrix .	67
4.2	Rank	67
4.2.1	The Dimension of Row Space And Column Space Are The Same	67
4.2.2	Definition of Rank	67
4.2.3	Properties of Rank	67
4.2.4	Rank & Consistency of Linear System	68
4.2.4.1	Remark	68
4.2.5	Row/Column Spaces and Matrix Multiplication	68
4.2.5.1	Ranks and Matrix Multiplication	68
4.3	Nullspaces and Nullities	69
4.3.1	Definition of Nullspace	69
4.3.2	Definition of Nullity	69
4.3.3	Finding Nullity of a Matrix	69
4.3.4	Dimension Theorem	69
4.3.5	Solution to Inhomogeneous Linear System and Nullspace	69
4.3.5.1	Condition for Linear System to have Unique Solution	69
4.3.5.2	Condition for the Solution Set of a Linear System to be a Vector Space	70
5	Orthogonality	71
5.1	The Dot Product	71
5.1.1	Pythagoras' Theorem	71
5.1.2	Cosine Rule	71
5.1.3	Definitions	71
5.1.3.1	Dot Product (Inner Product)	71
5.1.3.2	Norm (Length)	71
5.1.3.3	Unit Vector	71
5.1.3.4	Distance	72
5.1.3.5	Angle	72
5.1.4	Dot Product and Matrix Multiplication	72
5.1.4.1	Dot Product As Matrix Multiplication	72
5.1.4.2	Matrix Multiplication As Dot Products	72
5.1.5	Properties of Dot Product and Norm	72

5.1.5.1	Dot Product is Commutative	72
5.1.5.2	Dot Product is Distributive Over Vector Addition	72
5.1.5.3	Dot Product and Scalar Multiplication	72
5.1.5.4	Absolute Homogeneity of Norm	73
5.1.5.5	Dot Product of a Vector with Itself	73
5.1.5.6	Norm and Dot Product	73
5.1.5.7	Nonnegativity of Norm and When Norm is Zero	73
5.1.6	Inequalities of Norm and Distance	73
5.1.6.1	Cauchy-Schwarz Inequality	73
5.1.6.2	Triangle Inequality (Norm Version)	73
5.1.6.3	Triangle Inequality (Distance Version)	73
5.1.7	Results for Multiplying a Matrix with its Transpose	73
5.1.7.1	When Multiplying a Matrix with its Transpose Gives the Zero Matrix	73
5.1.7.2	When the Trace of Multiplying a Matrix with its Transpose is Zero	73
5.2	Orthogonality, Orthogonal and Orthonormal Sets	74
5.2.1	Definition of Orthogonal for Two Vectors	74
5.2.1.1	Zero Vector is Orthogonal to Every Vector in \mathbb{R}^n	74
5.2.2	Definition of Orthogonal / Orthonormal Sets	74
5.2.2.1	Definition of Orthogonal for a Subset of \mathbb{R}^n	74
5.2.2.2	Definition of Orthonormal for a Subset of \mathbb{R}^n	74
5.2.3	Properties of Orthogonal / Orthonormal Sets	74
5.2.4	Normalizing: Converting an Orthogonal Set to an Orthonormal Set	74
5.2.5	Using Matrix Multiplication to Check for Orthogonal / Orthonormal Set	75
5.2.5.1	Standard Basis is An Orthonormal Set	75
5.2.5.2	An Orthonormal Subset of \mathbb{R}^n with n Vectors is a Basis for \mathbb{R}^n	75
5.2.6	Orthogonal Nonzero Sets are Linearly Independent	75
5.2.6.1	Orthonormal Sets are Linearly Independent	75
5.3	Orthogonal and Orthonormal Bases	75
5.3.1	Orthogonal / Orthonormal Bases	75
5.3.2	Criterion for Orthogonal / Orthonormal Set to be Basis	76
5.3.3	Simpler Formula for Coordinate Vectors Relative to Orthogonal / Orthonormal Bases	76
5.3.3.1	Orthogonal Bases	76
5.3.3.2	Orthonormal Bases	76
5.3.4	Orthogonality between a Vector and a Subspace	76
5.3.4.1	Example: Normal Vector of a Plane in \mathbb{R}^3	76
5.3.5	Easier Criterion for Vector to be Orthogonal to a Vector Space	76
5.3.6	The Set of All Vectors Orthogonal to a Subspace is a Subspace	76
5.3.7	Projection of a Vector Onto a Vector Space	77

5.3.7.1	With Orthonormal Basis	77
5.3.7.2	With Orthogonal Basis	77
5.3.8	Gram-Schmidt Process: Generating Orthogonal / Orthonormal Basis . .	77
5.3.9	Solving a Linear System whose Coefficient Matrix's Columns are Linearly Independent	77
5.3.9.1	Decomposition	77
5.3.9.2	Algorithm	78
5.3.9.3	Remark	78
5.4	Best Approximations	78
5.4.1	The Projection of a Vector on a Subspace is its Best Approximation in that Subspace	78
5.4.2	Least Squares Solution	78
5.4.2.1	Definition of Least Squares Solution	78
5.4.2.2	Least Squares Solution and Projection	78
5.4.3	Methodology: Finding Least Squares Solution	79
5.4.3.1	Tedious Method	79
5.4.3.2	Easy Method	79
5.5	Orthogonal Matrices	79
5.5.1	Definition of Orthogonal Matrices	79
5.5.1.1	Example: Identity Matrix is an Orthogonal Matrix	79
5.5.2	Properties of Orthogonal Matrices	79
5.5.2.1	Rows and Columns of Orthogonal Matrix Form an Orthonormal Basis for \mathbb{R}^n	79
5.5.2.2	The Transpose or Inverse of an Orthogonal Matrix is Orthogonal	80
5.5.2.3	The Product of Two Orthogonal Matrices is Orthogonal	80
5.5.2.4	Condition for Rows/Columns of a Matrix to Form an Orthonor- mal Set	80
5.5.2.4.1	Condition for Columns of a Matrix to Form an Or- thonormal Set	80
5.5.2.4.2	Condition for Rows of a Matrix to Form an Orthonormal Set	80
5.5.2.5	Pre-Multiplication of an Orthogonal Matrix to an Orthonormal Set	80
5.5.3	Transition Matrices Between Orthonormal Bases	80
5.5.3.1	Formula for Transition Matrices Between Orthonormal Bases .	80
5.5.3.2	Transition Matrices Between Orthonormal Bases are Orthogonal	80
5.5.4	Classification of Orthogonal Matrices	80
5.5.4.1	Table of Orthogonal Matrices By Order	80
5.5.5	Geometric Representation of Orthogonal Matrix	81
5.5.5.1	Rotation Matrix	81

5.5.5.2	Composition of Rotation Matrices	81
5.5.5.3	Deriving Sum Laws for Sine and Cosine	81

6 Diagonalization 83

6.1	Eigenvalues and Eigenvectors	83
6.1.1	Definition of Eigenvalues and Eigenvectors	83
6.1.2	Characteristic Polynomial	83
6.1.2.1	Definition of Characteristic Polynomial	83
6.1.2.2	Characteristic Polynomial is a Monic Polynomial of Degree n	83
6.1.3	Definition of Characteristic Equation	83
6.1.4	Eigenvalues are the Roots to the Characteristic Equation	83
6.1.5	Eigenvalue and Invertibility	84
6.1.6	Eigenvalues of Triangular Matrices are its Diagonal Entries	84
6.1.7	Eigenspace	84
6.1.7.1	Definition of Eigenspace	84
6.1.7.2	Properties of Eigenspace	84
6.1.7.2.1	Vectors in Eigenspace	84
6.1.7.2.2	Dimension of Eigenspace	84
6.1.7.2.3	Eigenspace Associated To Eigenvalue Zero	84
6.2	Diagonalization	84
6.2.1	Definition of Diagonalizable Matrix	84
6.2.2	Criterion for Diagonalizability	84
6.2.3	Eigenvectors associated to Distinct Eigenvalues are Linearly Independent	85
6.2.4	Algorithm of Diagonalization	85
6.2.5	Matrix with n Distinct Eigenvalues is Diagonalizable	85
6.2.6	Application of Diagonalization	86
6.2.6.1	Matrix Exponentiation	86
6.2.6.2	Example: Fibonacci Numbers	86
6.3	Orthogonal Diagonalization	86
6.3.1	Definition of Orthogonally Diagonalizable Matrix	86
6.3.1.1	Remarks	86
6.3.2	Only Symmetric Matrices are Orthogonally Diagonalizable	87
6.3.3	Algorithm of Orthogonal Diagonalization	87
6.4	Quadratic Forms and Conic Sections	87
6.4.1	Quadratic Forms	87
6.4.1.1	Definition of Quadratic Form	87
6.4.1.2	Matrix Representation of Quadratic Forms	87
6.4.1.3	Simplification of Quadratic Forms	88
6.4.2	Quadratic Equation	88
6.4.2.1	Quadratic Equation in One Variable	88
6.4.2.2	Quadratic Equation in Two Variables	88

6.4.2.2.1	Matrix Representation	88
6.4.2.2.2	Quadratic Form Associated With Quadratic Equation	88
6.4.2.3	Graph of a Quadratic Equation	88
6.4.3	Classification of Conic Sections	89
6.4.3.1	Table of Conics	89
6.4.3.2	Algorithm to Classify Conic Sections	89
6.4.3.2.1	Determinant of Symmetric Matrix and Type of Conic Sections	89
6.4.3.2.2	Rotating and Reflecting Conic Sections	89
6.4.3.2.3	Can Always Orthogonally Diagonalize by Matrix with Determinant One	90
7	Linear Transformation	91
7.1	Linear Transformations from \mathbb{R}^n to \mathbb{R}^m	91
7.1.1	Definition of Linear Transformation	91
7.1.1.1	Linear Operator	91
7.1.2	Linear Transformation As Matrix Form	91
7.1.3	Examples of Linear Transformation	91
7.1.3.1	Identity Transformation	91
7.1.3.2	Zero Transformation	92
7.1.4	Standard Matrix of a Linear Transformation is Unique	92
7.1.5	To Prove that a Mapping is a Linear Transformation	92
7.1.6	Linearity of Linear Transformations	92
7.1.7	To Prove that a Mapping is Not a Linear Transformation	92
7.1.8	Representation of Linear Transformations	93
7.1.8.1	Linear Transformations are Completely Determined by a Basis of \mathbb{R}^n and its Images	93
7.1.8.2	Columns of Standard Matrix are Images of Standard Basis of \mathbb{R}^n	93
7.1.9	Linear Combination Preserving Implies Linear Transformation	93
7.1.9.1	With k vectors	93
7.1.9.2	With Two Vectors	93
7.1.10	General Definition of Linear Transformations	93
7.1.11	Change of Bases	94
7.1.11.1	Change Between the Images of Two Bases	94
7.1.11.2	Change Basis for Linear Operator	94
7.1.12	Composition	94
7.1.12.1	Definition of Function Composition	94
7.1.12.2	Function Composition is not Commutative in General	94
7.1.12.3	Linear Transformation Composition	94
7.1.12.4	Linear Transformation Composition is not Commutative in General	94

7.1.12.5	Composition of Linear Transformations is a Linear Transformation and Formula for its Standard Matrix	95
7.1.12.6	Properties of Linear Transformation Composition	95
7.2	Ranges and Kernels	95
7.2.1	Range	95
7.2.1.1	Range of Function	95
7.2.1.2	Range of Linear Transformation	95
7.2.1.3	Representation of Range	96
7.2.1.3.1	Range as Subspace Spanned by the Images of Any Basis for \mathbb{R}^n	96
7.2.1.3.2	Range as Column Space of Standard Matrix	96
7.2.2	Rank of Linear Transformation	96
7.2.3	Kernel	96
7.2.3.1	Definition of Kernel of Linear Transformation	96
7.2.3.2	Representation of Kernel	96
7.2.3.2.1	Kernel as Nullspace of Standard Matrix and Subspace of \mathbb{R}^n	96
7.2.3.2.2	Finding General Solution for Kernel	96
7.2.4	Nullity of Linear Transformation	97
7.2.5	Dimension Theorem for Linear Transformations	97
7.2.6	Linear Transformation Between Vector Spaces	97
7.2.6.1	Range	97
7.2.6.2	Rank	97
7.2.6.3	Kernel	97
7.2.6.4	Nullity	97
7.2.6.5	Dimension Theorem	97
7.3	Geometric Linear Transformations	97
7.3.1	Geometric Interpretation Completely Determined by Effect on a Basis for its Domain	97
7.3.2	Scaling	98
7.3.2.1	Scaling in \mathbb{R}^2	98
7.3.2.1.1	Dilation & Contraction	98
7.3.2.2	Scaling in other Axes in \mathbb{R}^2	98
7.3.2.3	Scaling in \mathbb{R}^3	98
7.3.2.3.1	Dilation & Contraction	99
7.3.2.4	Scaling in other Axes in \mathbb{R}^3	99
7.3.3	Reflection	99
7.3.3.1	Reflection in \mathbb{R}^2	99
7.3.3.1.1	Reflection w.r.t. x -axis	99
7.3.3.1.2	Reflection w.r.t. y -axis	100

7.3.3.1.3	Reflection w.r.t. the line $y = x$	100
7.3.3.1.4	Reflection w.r.t. a Line Through the Origin	100
7.3.3.2	Reflections in \mathbb{R}^3	101
7.3.3.2.1	Reflections w.r.t. Planes Formed by Coordinate Axes .	101
7.3.3.2.2	Reflections w.r.t. Any Plane	101
7.3.4	Rotation	101
7.3.4.1	Rotation in \mathbb{R}^2	101
7.3.4.1.1	Orthogonal Matrices of Determinant 1	101
7.3.4.2	Rotation in \mathbb{R}^3	101
7.3.5	Reflections and Rotations in \mathbb{R}^2	102
7.3.5.1	Determinant Determines Rotation vs Reflection	102
7.3.5.2	Reflection about a Line can be Decomposed into Reflection and Rotation	102
7.3.6	Shears	103
7.3.6.1	Shears in \mathbb{R}^2	103
7.3.6.1.1	Shear in x -direction	103
7.3.6.1.2	Shear in y -direction	103
7.3.6.2	Shears in \mathbb{R}^3	103
7.3.7	Translations	103
7.3.8	2D Computer Graphic System	104
7.3.8.1	Representation of 2D Figures	104
7.3.8.2	Scaling, Reflection, Rotation, Shearing of 2D Figures	104
7.3.8.3	Translating 2D Figure with Homogeneous Coordinate System .	104

Chapter 1

Linear Systems & Gaussian Elimination

1.1 Linear Systems & Their Solutions

1.1.1 Linear Equations

Definition. A **linear equation** in n **variables (unknowns)** x_1, x_2, \dots, x_n is an equation of the form

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = b$$

where a_1, a_2, \dots, a_n and b are real constants.

1.1.1.1 Zero Equation

Definition. A linear equation is a **zero equation** if

$$a_1 = a_2 = \dots = a_n = b = 0$$

1.1.1.2 Nonzero Equation

Definition. A linear equation is a **nonzero equation** if it is not a zero equation

1.1.1.3 Inconsistent Equation

Definition. A linear equation is **inconsistent** if

$$a_1 = a_2 = \dots = a_n = 0 \text{ but } b \neq 0$$

1.1.2 Solutions of a Linear Equation

Let $a_1x_1 + a_2x_2 + \dots + a_nx_n = b$ be a linear equation in n variables x_1, x_2, \dots, x_n .

Definition. For all $s_1, s_2, \dots, s_n \in \mathbb{R}$, if

$$a_1s_1 + a_2s_2 + \dots + a_ns_n = b$$

then $\mathbf{x}_1 = \mathbf{s}_1, \mathbf{x}_2 = \mathbf{s}_2, \dots, \mathbf{x}_n = \mathbf{s}_n$ is a **solution** to the given linear equation.

1.1.2.1 Solution Set

Definition. The set of all solutions is called the **solution set**.

1.1.2.2 General Solution

Definition. An expression that gives the entire solution set is a **general solution**.

1.1.3 Solution Set of Zero Equation

The zero equation in n variables x_1, x_2, \dots, x_n is satisfied by any values of x_1, x_2, \dots, x_n .

1.1.4 Solution Set of Inconsistent Linear Equation

An inconsistent linear equation in n variables x_1, x_2, \dots, x_n is not satisfied by any values of x_1, x_2, \dots, x_n .

1.1.5 General Solution of Consistent Nonzero Equation

The general solution of a consistent nonzero equation $a_1x_1 + a_2x_2 + \dots + a_nx_n = b$ in n variables x_1, x_2, \dots, x_n has $n - 1$ arbitrary parameters and is of the following form: For any integer $i \in [1, n]$,

$$\left\{ \begin{array}{ll} x_1 = & s_1, \\ x_2 = & s_2, \\ \vdots & \\ x_{i-1} = & s_{i-1}, \\ x_i = \frac{1}{a_i}(b - a_1s_1 - a_2s_2 - \dots - a_{i-1}s_{i-1} - a_{i+1}s_{i+1} - \dots - a_ns_n), \\ x_{i+1} = & s_{i+1}, \\ \vdots & \\ x_n = & s_n \end{array} \right.$$

where $s_1, s_2, \dots, s_{i-1}, s_{i+1}, \dots, s_n$ are arbitrary parameters.

1.1.6 Geometrical Interpretation of Linear Equations

1.1.6.1 In Two Variables

The solution set of the linear equation

$$ax + by = c \text{ (in } x, y\text{)}$$

where a and b are not both zero represents a **straight line** in the xy -plane.

The solution set of the linear equation

where a, b, c are not all zero represents a **plane** in the xyz -space.

Definition. A **linear system** (**system of linear equations**) of m linear equations in n variables x_1, x_2, \dots, x_n is

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1, \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2, \\ \qquad \qquad \qquad \vdots \qquad \qquad \qquad \qquad \qquad \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m, \end{cases}$$

where a_{ij} and b_i are real constants.

- a_{ij} is the **coefficient** of x_j in the i th equation,
- b_i is the **constant term** of the i th equation.

Definition. If all a_{ij} and b_i are zero, the linear system is a **zero system**.

Definition. If some a_{ij} or b_i is nonzero, the linear system is a **nonzero system**.

Given a linear system in n variables x_1, x_2, \dots, x_n , if $x_1 = s_1, x_2 = s_2, \dots, x_n = s_n$ is a solution to **every equation** of the linear system, then it is a **solution** to the system.

Definition. The set of all solutions to the linear system is called the **solution set**.

Definition. An expression that gives the entire solution set of the linear system is a **general solution**.

1.1.9 Consistency of Linear Systems

Definition. A linear system is

- **consistent** if it has at least one solution;
- **inconsistent** if it has no solution.

1.1.10 Number of Solutions to Linear Systems

A linear system has either

- no solution, or
- exactly one solution, or
- infinitely many solutions

1.1.11 Geometric Interpretation of Linear Systems

1.1.11.1 In Two Variables of Two Equations

Given a linear system in variables x, y of two equations

$$\begin{cases} a_1x + b_1y = c_1, & (L_1) \\ a_2x + b_2y = c_2, & (L_2) \end{cases}$$

where a_1, b_1 are not both zero and a_2, b_2 are not both zero. The system has

- no solution $\iff L_1$ and L_2 are parallel but distinct;
- exactly one solution $\iff L_1$ and L_2 are not parallel;
- infinitely many solutions $\iff L_1$ and L_2 are the same line.

1.1.11.2 In Three Variables of Two Equations

Given a linear system in variables x, y, z of two equations

$$\begin{cases} a_1x + b_1y + c_1z = d_1, & (P_1) \\ a_2x + b_2y + c_2z = d_2, & (P_2) \end{cases}$$

where a_1, b_1, c_1 are not all zero and a_2, b_2, c_2 are not all zero. The system has

- no solution $\iff P_1$ and P_2 are parallel but distinct
- $\iff \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2};$
- infinitely many solutions $\iff P_1$ and P_2 intersect at a straight line

$$\iff \frac{a_1}{a_2}, \frac{b_1}{b_2}, \frac{c_1}{c_2} \text{ are not all the same;}$$

- infinitely many solutions $\iff P_1$ and P_2 are the same plane

$$\iff \frac{a_1}{a_2} = \frac{b_1}{b_2} = \frac{c_1}{c_2} = \frac{d_1}{d_2}.$$

1.2 Elementary Row Operations

1.2.1 Augmented Matrix Representation of a Linear System

Definition. Given a linear system in variables x_1, x_2, \dots, x_n :

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1, \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2, \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m, \end{cases}$$

The rectangular array of constants

$$\left(\begin{array}{cccc|c} a_{11} & a_{12} & \dots & a_{1n} & b_1 \\ a_{21} & a_{22} & \dots & a_{2n} & b_2 \\ \vdots & \vdots & & \vdots & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} & b_m \end{array} \right)$$

is called the **augmented matrix** of the linear system.

1.2.2 Elementary Row Operations on Augmented Matrices

Definition. The **elementary row operations** are the following operations on rows of an augmented matrix:

Description of Operation	Notation
Multiply the i th row by a nonzero constant k	kR_i
Interchange the i th and j th rows	$R_i \leftrightarrow R_j$
Add k times the i th row to the j th row	$R_j + kR_i$

1.2.2.1 Correspondence to Operations on Equations in Linear System

Each elementary row operation corresponds to operations on the equations of the linear system as follows:

Elementary Row Operation	Equation Operation
kR_i	Multiply the i th equation by a nonzero constant k
$R_i \leftrightarrow R_j$	Interchange the i th and j th equations
$R_j + kR_i$	Add k times the i th equation to the j th equation

1.2.2.2 Interchanging two rows can be decomposed further

Interchanging two rows can be obtained by using the other two operations.

$$\begin{aligned} \begin{pmatrix} a \\ b \end{pmatrix} &\xrightarrow{R_1 + R_2} \begin{pmatrix} a + b \\ b \end{pmatrix} \xrightarrow{R_2 - R_1} \begin{pmatrix} a + b \\ -a \end{pmatrix} \\ &\xrightarrow{R_1 + R_2} \begin{pmatrix} b \\ -a \end{pmatrix} \xrightarrow{-R_2} \begin{pmatrix} b \\ a \end{pmatrix} \end{aligned}$$

1.2.2.3 Inverse Elementary Row Operations

Each **elementary row operation** has an inverse operation, which undoes the said **elementary row operation** and is also an **elementary row operation**, as follows:

Elementary Row Operation	Its Inverse
$A \xrightarrow{kR_i} B$	$B \xrightarrow{\frac{1}{k}R_i} A$
$A \xrightarrow{R_i \leftrightarrow R_j} B$	$B \xrightarrow{R_i \leftrightarrow R_j} A$
$A \xrightarrow{R_j + kR_i} B$	$B \xrightarrow{R_j - kR_i} A$

1.2.3 Row Equivalent Matrices

Definition. Two **augmented matrices** are **row equivalent** if one can be obtained from the other by a **series of elementary row operations**.

1.2.4 Row Equivalence as an Equivalence Relation

Theorem. Let A, B, C be any **augmented matrices**.

- A is row equivalent to A (reflexive);
- A is row equivalent to $B \iff B$ is row equivalent to A (symmetric);
- A is row equivalent to B and B is row equivalent to $C \implies A$ is row equivalent to C (transitive).

Therefore **row equivalence** is an **equivalence relation**.

1.2.5 Row Equivalence Implies Same Solution Set

Theorem. Let A and B be **augmented matrices** of two **linear systems**. Suppose A and B are **row equivalent**.

- Then the corresponding linear systems have the same solution set.

1.3 Row-Echelon Form

1.3.1 Leading Entry

Definition. The **leading entry** for any **nonzero row** of any **augmented matrix** is the first nonzero number of that row, from leftmost to rightmost column.

1.3.2 Row-Echelon Form (REF)

Definition. An **augmented matrix** is in **row-echelon form (REF)** if the following properties are satisfied:

- The **zero rows** are grouped together at the bottom;
- For any two successive **nonzero rows**, the **leading entry** in the lower row appears to the right of the **leading entry** in the higher row.

1.3.3 Pivot Points, Pivot Columns, and Non-pivot Columns

Definition. Suppose an **augmented matrix** is in **row-echelon form**.

1.3.3.1 Pivot Point

Then the **leading entry** of a **nonzero row** is a **pivot point**.

1.3.3.2 Pivot and Non-pivot Columns

A column of the **augmented matrix** is called a

- **pivot column** if it contains a **pivot point**;
- **non-pivot column** if it contains no **pivot point**.

1.3.3.3 Each Pivot Column Contains Exactly One Pivot Point

By the second property of the **row-echelon form**, every **pivot column** contains exactly one **pivot point**.

1.3.4 Reduced Row-Echelon Form (RREF)

Definition. Suppose an **augmented matrix** is in **row-echelon form**. It is in **reduced row-echelon form (RREF)** if

- The **leading entry**, or equivalently the **pivot point**, of every **nonzero row** is 1;
- In each **pivot column**, all entries except the **pivot point** are 0.

1.3.5 General Solution From Row-echelon Form

Suppose that the **augmented matrix** corresponding to a **linear system** is in **row-echelon form**. Then the **general solution** of the **linear system** can be obtained by the following algorithm:

1. **Set** the variables corresponding to **non-pivot columns** to be arbitrary parameters.
2. **Solve** the variables corresponding to **pivot columns** by **back substitution** (from the bottom equation to the top).

1.3.6 General Solution From Reduced Row-echelon Form

Suppose that the **augmented matrix** corresponding to a **linear system** is in **reduced row-echelon form**. Then the **general solution** of the **linear system** can be obtained by the following algorithm:

1. **Set** the variables corresponding to **non-pivot columns** to be arbitrary parameters.
2. **Solve** the variables corresponding to **pivot columns** in any order.

1.4 Gaussian Elimination

1.4.1 REF/RREF of Augmented Matrices

Definition. Let A and R be **augmented matrices**. Suppose that A is **row equivalent** to R .

- If R is in **row-echelon form**,
 1. then R is a **row-echelon form** of A ;
- If R is in **reduced row-echelon form**,
 1. then R is a **reduced row-echelon form** of A .

Therefore, by section 1.16 and 1.21, solving a **linear system** with **augmented matrix** $A \iff$ solving a **linear system** with **augmented matrix** that is the **REF/RREF** of A .

1.4.2 Gaussian Elimination: Finding REF of Augmented Matrices

Given an **augmented matrix**, we can find its **row-echelon form** by an algorithm called **Gaussian Elimination**. The algorithm is as follows:

1. Find the **leftmost column** which is not entirely zero.
2. Check the **top entry** of such column. If it is 0,

- replace it by a nonzero number by interchanging the top row with another row below.
3. For **each row below** the top row,
 - add a suitable multiple of the **top row** to it so that its **leading entry** becomes 0.
 4. If the entire matrix is not in **row-echelon form**,
 - then cover the top row and repeat steps 1-3 to the remaining matrix.

1.4.3 Gauss-Jordan Elimination: Finding RREF of Augmented Matrices

Given an **augmented matrix**, we can find its **reduced row-echelon form** by an algorithm called **Gauss-Jordan Elimination**. The algorithm is as follows:

1. Use **Gaussian Elimination** to get a **row-echelon form**.
2. For **each nonzero row**, multiply a suitable constant so that the **pivot point** becomes 1.
3. For each row, starting from the last nonzero row and working backwards,
 - Add a suitable multiple of the current row to each of the rows above to introduce 0 above the **pivot point** of the current row.

1.4.4 Infinitely Many REFs

Every nonzero matrix has infinitely many non-reduced **row-echelon forms**.

1.4.5 Uniqueness of RREF

Every matrix has a unique **reduced row-echelon form**.

1.4.6 Deducing Consistency of Linear Systems from its REF

Suppose that \mathbf{A} is the **augmented matrix** of a **linear system**, and \mathbf{R} is a **row-echelon form** of \mathbf{A} .

1.4.6.1 Inconsistent With No Solutions

The **linear system** is **inconsistent**, i.e. **no solution** if

- the last column is a **pivot column** \iff the last nonzero row has a **pivot point** in the last column.

1.4.6.2 Consistent With One Solution

The **linear system** is **consistent** with exactly one solution if

- the last column is a **non-pivot column**, and
- all other columns are **pivot columns**.

1.4.6.3 Consistent With Infinitely Many Solutions & Arbitrary Parameters

The **linear system** is **consistent** with infinitely many solution if

- the last column is a **non-pivot column**, and
- some other column(s) is/are **non-pivot column(s)**.

The number of arbitrary parameters = the number of **non-pivot columns** excluding the last column.

1.4.7 Geometrical Interpretation of Linear Systems with RREF

1.4.7.1 Linear System in Three Variables of Three Equations

Given a linear system in variables x, y, z of three equations:

$$\begin{cases} a_{11}x + a_{12}y + a_{13}z = b_1 & (P_1), \\ a_{21}x + a_{22}y + a_{23}z = b_2 & (P_2), \\ a_{31}x + a_{32}y + a_{33}z = b_3 & (P_3) \end{cases}$$

where a_{i1}, a_{i2}, a_{i3} are not all zero for $i = 1, 2, 3$. The **reduced row-echelon form** \mathbf{R} has three rows and four columns. The following table summarises the possible solutions sets:

Consistent	Last Column	Other Pivot Columns	Other Non-pivot Columns / Arbitrary Parameters	Intersection
No	Pivot	0-2	1-3	null set
Yes	Non-pivot	1	2	plane (three planes coincide)
Yes	Non-pivot	2	1	line
Yes	Non-pivot	3	0	point

1.5 Homogeneous Linear Systems

1.5.1 Homogeneous Linear Equation

Definition. A **linear equation** in variables x_1, x_2, \dots, x_n is **homogeneous** if it is of the form

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = 0$$

1.5.1.1 Remarks

A **linear equation** is homogeneous $\iff x_1 = 0, x_2 = 0, \dots, x_n = 0$ is a solution.

1.5.2 Geometrical Interpretation of Homogeneous Linear Equations

1.5.2.1 In Two Variables

The solution set of the linear equation

$$ax + by = 0 \text{ (in } x, y\text{)}$$

where a and b are not both zero represents a **straight line**, in the xy -plane, that passes through the origin $O(0, 0)$.

1.5.2.2 In Three Variables

The solution set of the linear equation

$$ax + by + cz = 0 \text{ (in } x, y, z\text{)}$$

where a, b, c are not all zero represents a **plane**, in the xyz -space, that contains the origin $O(0, 0, 0)$.

1.5.3 Homogeneous Linear System

Definition. A **linear system** is **homogeneous** if every linear equation of the system is homogeneous. That is the linear system is of the form

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = 0, \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = 0, \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = 0, \end{cases}$$

1.5.4 Trivial Solution of Homogeneous Linear Systems

A **linear system** in x_1, x_2, \dots, x_n is **homogeneous** $\iff x_1 = 0, x_2 = 0, \dots, x_n = 0$ is a solution.

- This is the **trivial solution** of a **homogeneous linear system**
- Other solutions are called **non-trivial solutions**.

1.5.5 Geometric Interpretation of Homogeneous Linear Systems

1.5.5.1 In Two Variables of Two Equations

Given a **homogeneous linear system** in variables x, y of two equations

$$\begin{cases} a_1x + b_1y = 0, & (L_1) \\ a_2x + b_2y = 0, & (L_2) \end{cases}$$

where a_1, b_1 are not both zero and a_2, b_2 are not both zero. The system has

- only the **trivial solution** $\iff L_1$ and L_2 are different;
- **non-trivial solutions** $\iff L_1$ and L_2 are the same.

1.5.5.2 In Three Variables of Two Equations

Given a linear system in variables x, y, z of two equations

$$\begin{cases} a_1x + b_1y + c_1z = 0, & (P_1) \\ a_2x + b_2y + c_2z = 0, & (P_2) \end{cases}$$

where a_1, b_1, c_1 are not all zero and a_2, b_2, c_2 are not all zero. The system has infinitely many solutions with only two cases

- The two planes are the same, or
- The two planes intersect at a straight line passing through $O(0, 0, 0)$.

Chapter 2

Matrices

2.1 Introduction to Matrices

2.1.1 Definition of Matrix

Definition. A **matrix** (plural **matrices**) is a rectangular array of numbers

$$\begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

where

- m is the number of **rows** in the matrix.
- n is the number of **columns** in the matrix.
- The **size** of the matrix is given by $m \times n$.
- The **(i, j) -entry** is the entry in the i th row and j th column.
 - In the given matrix, the (i, j) -entry is a_{ij} .

2.1.1.1 Remarks

- Some books use $[\dots]$ instead of (\dots) .
- A 1×1 matrix is usually treated as a real number in computation.

2.1.2 Notation of Matrices

A **matrix** is usually denoted by capital letters $\mathbf{A}, \mathbf{B}, \mathbf{C}, \dots$. Given a

$$m \times n \text{ matrix } \mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix},$$

- a_{ij} is the (i, j) -entry of \mathbf{A} .
- The matrix is denoted by $\mathbf{A} = (a_{ij})_{m \times n}$
- If the size of \mathbf{A} is known (or not important)
 - The matrix can also be denoted by $\mathbf{A} = (a_{ij})$.

2.1.3 Row Matrix

Definition. A **row matrix** (**row vector**) is a **matrix** with only one **row**.

2.1.4 Column Matrix

Definition. A **column matrix** (**column vector**) is a **matrix** with only one **column**.

2.1.5 Square Matrix

Definition. A **square matrix** is a **matrix** with the same number of **rows** and **columns**.

2.1.5.1 Order of Square Matrix

Definition. An $n \times n$ matrix is a **square matrix** of **order** n .

2.1.5.2 Diagonal / Diagonal Entries of Square Matrix

Definition. Let $\mathbf{A} = (a_{ij})$ be a **square matrix** of order n .

- The **diagonal/principle diagonal/major diagonal** of \mathbf{A} is the **sequence** of entries
 - $a_{11}, a_{22}, \dots, a_{nn}$
- The entries a_{ii} , for $i = 1, \dots, n$ are the **diagonal entries**
- The entries a_{ij} , $i \neq j$ are the **non-diagonal entries**
- The **anti-diagonal/minor diagonal** of \mathbf{A} is the **sequence** of entries from the right top to the left bottom
 - $a_{1n}, a_{2,n-1}, \dots, a_{n1}$

2.1.6 Diagonal Matrix

Definition. A square matrix is a **diagonal matrix** if all its **non-diagonal entries** are zero.

- $A = (a_{ij})_{n \times n}$ is **diagonal** $\iff a_{ij} = 0$ for all $i \neq j$

2.1.7 Scalar Matrix

Definition. A diagonal matrix is a **scalar matrix** if all its **diagonal entries** are the same.

- $A = (a_{ij})_{n \times n}$ is **scalar** $\iff a_{ij} = \begin{cases} c & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$

2.1.8 Identity Matrix

Definition. A scalar matrix is an **identity matrix** if all its **diagonal entries** are 1.

- $A = (a_{ij})_{n \times n}$ is **identity** $\iff a_{ij} = \begin{cases} 1 & \text{if } i = j \\ 0 & \text{if } i \neq j \end{cases}$

Note: There is exactly one identity matrix of order n .

2.1.8.1 Notation for Identity Matrix

- The **identity matrix** of **order** n is denoted by I_n .
- If no confusion in order, we may write I instead of I_n .

2.1.9 Zero Matrix

Definition. A matrix with all entries equal to zero is a **zero matrix**.

- $A = (a_{ij})_{m \times n}$ is **zero** $\iff a_{ij} = 0$ for all i, j

Note: There is exactly one zero matrix of size $m \times n$.

2.1.9.1 Notation for Zero Matrix

- The **zero matrix** of **size** $m \times n$ is denoted by $\mathbf{0}_{m \times n}$.
- If no confusion in size, we may write $\mathbf{0}$ instead of $\mathbf{0}_{m \times n}$.

2.1.10 Symmetric Square Matrix

Definition. A square matrix is **symmetric** if it is symmetric with respect to the diagonal.

- $A = (a_{ij})_{n \times n}$ is **symmetric** $\iff a_{ij} = a_{ji}$ for all i, j
 - There is no restriction to the **diagonal entries**.

2.1.11 Upper Triangular Square Matrix

Definition. A square matrix is **upper triangular** if all the entries **below** the diagonal are zero.

- $A = (a_{ij})_{n \times n}$ is **upper triangular** $\iff a_{ij} = 0$ if $i > j$
 - There is no restriction to the **diagonal entries**.

2.1.12 Lower Triangular Square Matrix

Definition. A square matrix is **lower triangular** if all the entries **above** the diagonal are zero.

- $A = (a_{ij})_{n \times n}$ is **lower triangular** $\iff a_{ij} = 0$ if $i < j$
 - There is no restriction to the **diagonal entries**.

2.1.13 Triangular Square Matrix

Definition. Both upper triangular matrices and lower triangular matrices are **triangular matrices**.

2.1.14 Diagonal and Triangular Square Matrices

A **square matrix** is both upper and lower triangular \iff it is diagonal.

2.2 Matrix Operations

2.2.1 Identical Matrices

- A matrix is completely determined by its **size** and **entries**.
- **Definition.** Two matrices are **equal** if
 - they have the same **size** (same number of rows and same number of columns), and
 - all the corresponding entries are the same.

Let $A = (a_{ij})_{m \times n}$ and $B = (b_{ij})_{p \times q}$. Then

- $A = B \iff m = p \ \& \ n = q \ \& \ a_{ij} = b_{ij} \text{ for all } i, j$

2.2.2 Matrix Addition, Subtraction & Scalar Multiplication

Definition. Let $\mathbf{A} = (a_{ij})_{m \times n}$ and $\mathbf{B} = (b_{ij})_{m \times n}$ be matrices, and c a constant. The following operations are defined:

- **Addition:** $\mathbf{A} + \mathbf{B} = (a_{ij} + b_{ij})_{m \times n}$
- **Subtraction:** $\mathbf{A} - \mathbf{B} = (a_{ij} - b_{ij})_{m \times n}$
- **Scalar Multiplication:** $c\mathbf{A} = (ca_{ij})_{m \times n}$

Remarks.

- $(-1)\mathbf{A}$ is usually denoted by $-\mathbf{A}$.
- It can be proved that $\mathbf{A} - \mathbf{B} = \mathbf{A} + (-\mathbf{B})$.
 - In the discussion we usually only consider addition and scalar multiplication.

2.2.3 Properties of Matrix Addition, Subtraction & Scalar Multiplication

Theorem. Let $\mathbf{A}, \mathbf{B}, \mathbf{C}$ be matrices of the same size, and c, d be constants. Then

- $\mathbf{A} - \mathbf{B} = \mathbf{A} + (-\mathbf{B})$
- Commutative Law for Matrix Addition:
 - $\mathbf{A} + \mathbf{B} = \mathbf{B} + \mathbf{A}$
- Associative Law for Matrix Addition:
 - $(\mathbf{A} + \mathbf{B}) + \mathbf{C} = \mathbf{A} + (\mathbf{B} + \mathbf{C})$
- Let $\mathbf{0}$ be the zero matrix of the same size as \mathbf{A} . Then
 - $\mathbf{0} + \mathbf{A} = \mathbf{A}; \quad \mathbf{A} - \mathbf{A} = \mathbf{0}; \quad \mathbf{0}\mathbf{A} = \mathbf{0}; \quad c\mathbf{0} = \mathbf{0}.$
- Distributive Law for Scalar Multiplication over Addition:
 - $c(\mathbf{A} + \mathbf{B}) = c\mathbf{A} + c\mathbf{B}$
 - $(c + d)\mathbf{A} = c\mathbf{A} + d\mathbf{A}$
- $c(d\mathbf{A}) = (cd)\mathbf{A}, \quad 1\mathbf{A} = \mathbf{A}$

2.2.4 Matrix Multiplication

Definition. Let $\mathbf{A} = (a_{ij})_{m \times p}$ and $\mathbf{B} = (b_{ij})_{p \times n}$. Then \mathbf{AB} is the $m \times n$ matrix such that its (i, j) -entry is

$$a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{ip}b_{pj} = \sum_{k=1}^p a_{ik}b_{kj}$$

Note: No. of columns of \mathbf{A} = the no. of rows of \mathbf{B} .

2.2.4.1 Matrix Multiplication Explained in Words

In order to get the (i, j) -entry of the **product** matrix:

1. Find the i th **row** of the first matrix;
2. Find the j th **column** of the second matrix;
3. Multiply the corresponding entries;
4. Add the products together.

2.2.5 Noncommutativity of Matrix Multiplication

Matrix Multiplication is **not commutative** in general. For example, let $\mathbf{A} = \begin{pmatrix} 0 & 1 \\ 0 & 0 \end{pmatrix}$ and

$$\mathbf{B} = \begin{pmatrix} 0 & 0 \\ 1 & 0 \end{pmatrix}. \quad \mathbf{AB} = \begin{pmatrix} 1 & 0 \\ 0 & 0 \end{pmatrix} \neq \begin{pmatrix} 0 & 0 \\ 0 & 1 \end{pmatrix} = \mathbf{BA}.$$

- \mathbf{AB} is the **pre-multiplication** of \mathbf{A} to \mathbf{B} (to \mathbf{B} by \mathbf{A}).
- \mathbf{BA} is the **post-multiplication** of \mathbf{A} to \mathbf{B} (to \mathbf{B} by \mathbf{A}).

2.2.6 Properties of Matrix Multiplication

Theorem.

- Let $\mathbf{A}, \mathbf{B}, \mathbf{C}$ be $m \times p, p \times q, q \times n$ matrices, resp. Then
 - Associative Law: $\mathbf{A}(\mathbf{BC}) = (\mathbf{AB})\mathbf{C}$.
- Let \mathbf{A} be a $m \times p$ matrix, $\mathbf{B}_1, \mathbf{B}_2$ be $p \times n$ matrices. Then
 - Distributive Law: $\mathbf{A}(\mathbf{B}_1 + \mathbf{B}_2) = \mathbf{AB}_1 + \mathbf{AB}_2$.
- Let $\mathbf{A}_1, \mathbf{A}_2$ be $m \times p$ matrices, \mathbf{B} be a $p \times n$ matrix. Then
 - Distributive Law: $(\mathbf{A}_1 + \mathbf{A}_2)\mathbf{B} = \mathbf{A}_1\mathbf{B} + \mathbf{A}_2\mathbf{B}$.
- Let \mathbf{A}, \mathbf{B} be $m \times p, p \times n$ matrices resp. and c be a constant. Then

- $c(\mathbf{AB}) = (c\mathbf{A})\mathbf{B} = \mathbf{A}(c\mathbf{B})$.
- Let \mathbf{A} be a $m \times n$ matrix. Then
 - $\mathbf{A}\mathbf{0}_{n \times p} = \mathbf{0}_{m \times p}$; $\mathbf{0}_{p \times m}\mathbf{A} = \mathbf{0}_{p \times n}$.
 - $\mathbf{A}\mathbf{I}_n = \mathbf{A}$; $\mathbf{I}_m\mathbf{A} = \mathbf{A}$.

2.2.7 Nonnegative Integer Powers of Square Matrices

2.2.7.1 Multiplying a Matrix With Itself

Let \mathbf{A} be a $m \times n$ matrix. Then

- \mathbf{AA} is well-defined $\iff m = n \iff \mathbf{A}$ is a **square matrix**.

2.2.7.2 Nonnegative Integer Powers of Square Matrices

Definition. Let \mathbf{A} be a **square matrix** of order n . For nonnegative integers k , the **powers** of \mathbf{A} are defined as

$$\mathbf{A}^k = \begin{cases} \mathbf{I}_n & \text{if } k = 0 \\ \underbrace{\mathbf{AA} \dots \mathbf{A}}_{k \text{ times}} & \text{if } k \geq 1 \end{cases}$$

2.2.8 Properties of Nonnegative Integer Powers of Square Matrices

Let \mathbf{A}, \mathbf{B} be square matrices of the same size, and m, n nonnegative integers. Then

- $\mathbf{A}^m \mathbf{A}^n = \mathbf{A}^{m+n}$.
- In general, $(\mathbf{AB})^n \neq \mathbf{A}^n \mathbf{B}^n$ for $n = 2, 3, \dots$
 - Since matrix multiplication is not commutative,

$$\begin{aligned} (\mathbf{AB})^n &= \underbrace{(\mathbf{AB})(\mathbf{AB}) \dots (\mathbf{AB})}_{n \text{ times}} \\ &\neq \underbrace{\mathbf{AA} \dots \mathbf{A}}_{n \text{ times}} \underbrace{\mathbf{BB} \dots \mathbf{B}}_{n \text{ times}} \\ &= \mathbf{A}^n \mathbf{B}^n \end{aligned}$$

- However, suppose that $\mathbf{AB} = \mathbf{BA}$. Then
 - $(\mathbf{AB})^n = \mathbf{A}^n \mathbf{B}^n$ for all nonnegative integers n .

2.2.9 Matrix Representation

$$\text{Let } \mathbf{A} = (a_{ij})_{m \times n} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$$

2.2.9.1 Matrix as a Column of Row Vectors

- Let \mathbf{a}_i denote the i th row of \mathbf{A} , for $i = 1, 2, \dots, m$.

$$\begin{aligned}\mathbf{a}_1 &= \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \end{pmatrix} \\ \mathbf{a}_2 &= \begin{pmatrix} a_{21} & a_{22} & \dots & a_{2n} \end{pmatrix} \\ &\vdots \\ \mathbf{a}_m &= \begin{pmatrix} a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}\end{aligned}$$

- Then each \mathbf{a}_i is a $1 \times n$ matrix (row vector).
- Then \mathbf{A} can be represented as

$$\mathbf{A} = \begin{pmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \vdots \\ \mathbf{a}_m \end{pmatrix}$$

2.2.9.2 Matrix as a Row of Column Vectors

- Let \mathbf{b}_j denote the j th column of \mathbf{A} , for $j = 1, 2, \dots, n$.

$$\mathbf{b}_1 = \begin{pmatrix} a_{11} \\ a_{21} \\ \vdots \\ a_{m1} \end{pmatrix}, \mathbf{b}_2 = \begin{pmatrix} a_{12} \\ a_{22} \\ \vdots \\ a_{m2} \end{pmatrix}, \dots, \mathbf{b}_n = \begin{pmatrix} a_{1n} \\ a_{2n} \\ \vdots \\ a_{mn} \end{pmatrix}$$

- Then each \mathbf{b}_j is a $m \times 1$ matrix (column vector).
- Then \mathbf{A} can be represented as

$$\mathbf{A} = \begin{pmatrix} \mathbf{b}_1 & \mathbf{b}_2 & \dots & \mathbf{b}_n \end{pmatrix}$$

2.2.10 Decomposing Matrix Multiplication

Suppose $\mathbf{A} = (a_{ij})_{m \times p}$.

- Let $\mathbf{a}_i = \begin{pmatrix} a_{i1} & a_{i2} & \dots & a_{ip} \end{pmatrix}$ be the i th row of \mathbf{A} .

Suppose $\mathbf{B} = (b_{ij})_{p \times n}$.

- Let $\mathbf{b}_j = \begin{pmatrix} b_{1j} \\ b_{2j} \\ \vdots \\ b_{pj} \end{pmatrix}$ be the j th column of \mathbf{B} .

Then

$$\mathbf{AB} = \begin{pmatrix} \mathbf{a}_1\mathbf{b}_1 & \mathbf{a}_1\mathbf{b}_2 & \dots & \mathbf{a}_1\mathbf{b}_n \\ \mathbf{a}_2\mathbf{b}_1 & \mathbf{a}_2\mathbf{b}_2 & \dots & \mathbf{a}_2\mathbf{b}_n \\ \vdots & \vdots & & \vdots \\ \mathbf{a}_m\mathbf{b}_1 & \mathbf{a}_m\mathbf{b}_2 & \dots & \mathbf{a}_m\mathbf{b}_n \end{pmatrix}$$

2.2.10.1 Decomposing Into Entries

$$\begin{aligned} (i, j)\text{-entry of } \mathbf{AB} &= a_{i1}b_{1j} + a_{i2}b_{2j} + \dots + a_{ip}b_{pj} \\ &= \mathbf{a}_i\mathbf{b}_j \end{aligned}$$

2.2.10.2 Decomposing Into Rows

$$\begin{aligned} i\text{th row of } \mathbf{AB} &= (\mathbf{a}_i\mathbf{b}_1 \quad \mathbf{a}_i\mathbf{b}_2 \quad \dots \quad \mathbf{a}_i\mathbf{b}_n) \\ &= \mathbf{a}_i(\mathbf{b}_1 \quad \mathbf{b}_2 \quad \dots \quad \mathbf{b}_n) \\ &= \mathbf{a}_i\mathbf{B} \end{aligned}$$

$$\text{Thus, } \mathbf{AB} = \begin{pmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \vdots \\ \mathbf{a}_m \end{pmatrix} \mathbf{B} = \begin{pmatrix} \mathbf{a}_1\mathbf{B} \\ \mathbf{a}_2\mathbf{B} \\ \vdots \\ \mathbf{a}_m\mathbf{B} \end{pmatrix}$$

2.2.10.3 Decomposing Into Columns

$$\begin{aligned} j\text{th column of } \mathbf{AB} &= \begin{pmatrix} \mathbf{a}_1\mathbf{b}_j \\ \mathbf{a}_2\mathbf{b}_j \\ \vdots \\ \mathbf{a}_m\mathbf{b}_j \end{pmatrix} = \begin{pmatrix} \mathbf{a}_1 \\ \mathbf{a}_2 \\ \vdots \\ \mathbf{a}_m \end{pmatrix} \mathbf{b}_j \\ &= \mathbf{A}\mathbf{b}_j \end{aligned}$$

$$\text{Thus, } \mathbf{AB} = \mathbf{A}(\mathbf{b}_1 \quad \mathbf{b}_2 \quad \dots \quad \mathbf{b}_n) = (\mathbf{A}\mathbf{b}_1 \quad \mathbf{A}\mathbf{b}_2 \quad \dots \quad \mathbf{A}\mathbf{b}_n)$$

2.2.10.4 Decomposing Into Blocks

Matrices can be multiplied in blocks (provided that the sizes are matched).

2.2.11 Matrix Representation of Linear Equation

Given a **linear equation** in n variables x_1, \dots, x_n

$$a_1x_1 + a_2x_2 + \dots + a_nx_n = b$$

The corresponding matrix representation is

$$\begin{pmatrix} a_1 & a_2 & \dots & a_n \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} = b$$

2.2.12 Matrix Representation of Linear System

Given a **linear system** of m equations in n variables x_1, \dots, x_n

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1 \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2 \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m \end{cases}$$

Then the corresponding matrix representation is as follows:

- The **coefficient matrix** is $\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & \dots & a_{1n} \\ a_{21} & a_{22} & \dots & a_{2n} \\ \vdots & \vdots & & \vdots \\ a_{m1} & a_{m2} & \dots & a_{mn} \end{pmatrix}$

- The **variable matrix** is $\mathbf{x} = \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix}$

- The **constant matrix** is $\mathbf{b} = \begin{pmatrix} b_1 \\ \vdots \\ b_n \end{pmatrix}$

- Then $\mathbf{Ax} = \mathbf{b}$

2.2.12.1 Solution Vector to Linear System

Let $\mathbf{u} = \begin{pmatrix} u_1 \\ \vdots \\ u_n \end{pmatrix}$. Then

$x_1 = u_1, \dots, x_n = u_n$ is a solution to the system

$$\iff \mathbf{Au} = \mathbf{b}$$

$$\iff \mathbf{u} \text{ is a solution to } \mathbf{Ax} = \mathbf{b}$$

2.2.12.2 Alternative Matrix Representation of Linear System

Let \mathbf{a}_j denote the j th column of \mathbf{A} . Then

$$\begin{aligned}\mathbf{b} = \mathbf{Ax} &= \begin{pmatrix} a_{11} & \dots & a_{1n} \\ \vdots & & \vdots \\ a_{m1} & \dots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ \vdots \\ x_n \end{pmatrix} = \begin{pmatrix} a_{11}x_1 + \dots + a_{1n}x_n \\ \vdots \\ a_{m1}x_1 + \dots + a_{mn}x_n \end{pmatrix} \\ &= x_1 \begin{pmatrix} a_{11} \\ \vdots \\ a_{m1} \end{pmatrix} + \dots + x_n \begin{pmatrix} a_{1n} \\ \vdots \\ a_{mn} \end{pmatrix} = x_1 \mathbf{a}_1 + \dots + x_n \mathbf{a}_n \\ &= \sum_{j=1}^n x_j \mathbf{a}_j\end{aligned}$$

2.2.13 Transpose of Matrix

Definition. Let $\mathbf{A} = (a_{ij})_{m \times n}$ be a **matrix**. The **transpose** of \mathbf{A} is the $n \times m$ matrix \mathbf{A}^T (or \mathbf{A}^t)

- whose (i, j) -entry is a_{ji} .

Remarks.

- The i th row of \mathbf{A}^T is the i th column of \mathbf{A} .
- The j th column of \mathbf{A}^T is the j th row of \mathbf{A} .

2.2.14 Properties of Transpose of Matrix

Theorem. Let \mathbf{A} be an $m \times n$ matrix. Then

- $(\mathbf{A}^T)^T = \mathbf{A}$.
- \mathbf{A} is symmetric $\iff \mathbf{A} = \mathbf{A}^T$.
- Let c be a scalar. Then $(c\mathbf{A})^T = c\mathbf{A}^T$.
- Let \mathbf{B} be $m \times n$. Then $(\mathbf{A} + \mathbf{B})^T = \mathbf{A}^T + \mathbf{B}^T$.
- Let \mathbf{B} be $n \times p$. Then $(\mathbf{AB})^T = \mathbf{B}^T \mathbf{A}^T$.
 - **Note:** In general, $(\mathbf{AB})^T \neq \mathbf{A}^T \mathbf{B}^T$.

2.3 Inverse of Square Matrices

2.3.1 Additive Inverse of Matrices

Let \mathbf{A} be a **matrix**. Then $-\mathbf{A}$ is the **additive inverse** of \mathbf{A} .

2.3.2 Inverse of Square Matrices

Definition. Let \mathbf{A} be a **square matrix** of order n .

- If there exists a **square matrix** \mathbf{B} of order n so that

$$\circ \mathbf{AB} = \mathbf{I}_n \text{ and } \mathbf{BA} = \mathbf{I}_n,$$

then \mathbf{A} is **invertible**, and \mathbf{B} is an **inverse** of \mathbf{A} .

- If \mathbf{A} is not **invertible**, then \mathbf{A} is **singular**.

Note: Non-square matrix is neither **invertible** nor **singular**.

2.3.3 Uniqueness of Inverse of Invertible Matrices

Theorem. Let \mathbf{A} be a **square matrix**. If \mathbf{A} is **invertible**

- then its **inverse** is unique.
- **Notation.** The unique inverse of \mathbf{A} , is denoted by \mathbf{A}^{-1}

$$\circ \mathbf{AA}^{-1} = \mathbf{A}^{-1}\mathbf{A} = \mathbf{I}.$$

2.3.4 Cancellation Law for Invertible Matrices

Theorem. Let \mathbf{A} be an **invertible matrix**. Then

- $\mathbf{AB}_1 = \mathbf{AB}_2 \implies \mathbf{B}_1 = \mathbf{B}_2.$
- $\mathbf{C}_1\mathbf{A} = \mathbf{C}_2\mathbf{A} \implies \mathbf{C}_1 = \mathbf{C}_2.$

Remark. The cancellation law fails if \mathbf{A} is **singular**.

2.3.5 Inverse of Square Matrix of Order Two

Theorem. Let $\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Then

- \mathbf{A} is **invertible** $\iff ad - bc \neq 0$.
- If \mathbf{A} is **invertible**, then $\mathbf{A}^{-1} = \frac{1}{ad - bc} \begin{pmatrix} d & -b \\ -c & a \end{pmatrix}.$

2.3.6 Properties of Invertible Matrices

Theorem. Let \mathbf{A}, \mathbf{B} be invertible matrices of same size. Then

- Let $c \neq 0$. $c\mathbf{A}$ is invertible, and $(c\mathbf{A})^{-1} = \frac{1}{c}\mathbf{A}^{-1}$.
- \mathbf{A}^T is invertible, and $(\mathbf{A}^T)^{-1} = (\mathbf{A}^{-1})^T$.
- \mathbf{A}^{-1} is invertible, and $(\mathbf{A}^{-1})^{-1} = \mathbf{A}$.
- \mathbf{AB} is invertible, and $(\mathbf{AB})^{-1} = \mathbf{B}^{-1}\mathbf{A}^{-1}$.

Let $\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_k$ be invertible matrices of the same size. Then

- $(\mathbf{A}_1\mathbf{A}_2 \dots \mathbf{A}_k)^{-1} = \mathbf{A}_k^{-1} \dots \mathbf{A}_2^{-1}\mathbf{A}_1^{-1}$.

In particular, $(\underbrace{\mathbf{AA} \dots \mathbf{A}}_{k \text{ times}})^{-1} = \underbrace{\mathbf{A}^{-1} \dots \mathbf{A}^{-1}}_{k \text{ times}}\mathbf{A}^{-1}$. Thus,

- $(\mathbf{A}^k)^{-1} = (\mathbf{A}^{-1})^k$.

2.3.7 Negative Integer Powers of Square Matrices

Definition. Let \mathbf{A} be an invertible matrix. For any positive integer k , $\mathbf{A}^{-k} = (\mathbf{A}^{-1})^k$.

2.3.8 Properties of Integer Powers of Square Matrices

Theorem. Let \mathbf{A} be an invertible matrix. For any integers m and n ,

- $\mathbf{A}^{m+n} = \mathbf{A}^m\mathbf{A}^n$ and $(\mathbf{A}^m)^n = \mathbf{A}^{mn}$.

Note: If \mathbf{A} is singular, then \mathbf{A}^{-1} is undefined.

2.4 Elementary Matrices

2.4.1 Definition of Elementary Matrices

Definition. A square matrix is called an **elementary matrix** if it can be obtained from the identity matrix by performing a single **elementary row operation**.

2.4.2 Connection of Elementary Matrices to Pre-Matrix Multiplication

Theorem. Let \mathbf{E} be an elementary matrix obtained by performing an elementary row operation on \mathbf{I}_m . Then for any $m \times n$ matrix \mathbf{A} , \mathbf{EA} can be obtained by performing the same elementary row operation to \mathbf{A} .

- Let \mathbf{A} be an $m \times n$ matrix.

- $\mathbf{I}_m \xrightarrow{cR_i} \mathbf{E} \implies \mathbf{A} \xrightarrow{cR_i} \mathbf{EA}.$
- $\mathbf{I}_m \xrightarrow{R_i \leftrightarrow R_j} \mathbf{E} \implies \mathbf{A} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{EA}.$
- $\mathbf{I}_m \xrightarrow{R_i + cR_j} \mathbf{E} \implies \mathbf{A} \xrightarrow{R_i + cR_j} \mathbf{EA}.$

2.4.3 Elementary Matrices are Invertible

Theorem.

- Every **elementary matrix** is **invertible**.
- The **inverse** of an **elementary matrix** is **elementary**.

Let \mathbf{E} be an **elementary matrix**.

- $\mathbf{I} \xrightarrow{cR_i} \mathbf{E} \implies \mathbf{I} \xrightarrow{\frac{1}{c}R_i} \mathbf{E}^{-1}.$
- $\mathbf{I} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{E} \implies \mathbf{I} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{E}^{-1}. \text{ (So } \mathbf{E} = \mathbf{E}^{-1}).$
- $\mathbf{I} \xrightarrow{R_i + cR_j} \mathbf{E} \implies \mathbf{I} \xrightarrow{R_i - cR_j} \mathbf{E}^{-1}.$

2.4.4 Row Equivalent Matrices are Linked by Elementary Matrices

Theorem. Two matrices \mathbf{A} and \mathbf{B} are **row equivalent** \iff there exist **elementary matrices** $\mathbf{E}_1, \mathbf{E}_2, \dots, \mathbf{E}_k$ such that $\mathbf{B} = \mathbf{E}_k \mathbf{E}_{k-1} \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A}.$

Remarks. Suppose that $\mathbf{B} = \mathbf{E}_k \mathbf{E}_{k-1} \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A}.$ Then $\mathbf{A} = \mathbf{E}_1^{-1} \mathbf{E}_2^{-1} \dots \mathbf{E}_{k-1}^{-1} \mathbf{E}_k^{-1} \mathbf{B}$

2.4.5 Main Theorem for Invertible Matrices

Theorem. Let \mathbf{A} be a **square matrix**. Then the following are equivalent:

1. \mathbf{A} is an **invertible matrix**.
2. Linear system $\mathbf{Ax} = \mathbf{b}$ has a unique solution.
3. Linear system $\mathbf{Ax} = \mathbf{0}$ has only the trivial solution.
4. The **reduced row-echelon form** of \mathbf{A} is $\mathbf{I}.$
5. \mathbf{A} is the product of **elementary matrices**.

2.4.6 Checking Whether a Square Matrix is Invertible

- A square matrix is **invertible**

- \iff Its reduced row-echelon form is \mathbf{I}
- \iff All the columns in its row-echelon form are pivot.
- \iff All the rows in its row-echelon form are nonzero.

- A square matrix is **singular**

- \iff Its reduced row-echelon form is not \mathbf{I}
- \iff Some columns in its row-echelon form are non-pivot.
- \iff Some rows in its row-echelon form are zero.

2.4.7 Finding Inverse of an Invertible Matrix

Theorem. Let \mathbf{A} be an invertible matrix. The reduced row-echelon form of $\left(\mathbf{A} \mid \mathbf{I} \right)$ is $\left(\mathbf{I} \mid \mathbf{A}^{-1} \right)$.

2.4.8 Weaker Requirement for Invertible Matrices

Theorem. Let \mathbf{A} and \mathbf{B} be square matrices of the same size. If $\mathbf{AB} = \mathbf{I}$, then \mathbf{A} and \mathbf{B} are invertible, and $\mathbf{A}^{-1} = \mathbf{B}$, $\mathbf{B}^{-1} = \mathbf{A}$.

2.4.9 Invertibility of Product of Matrices

Corollary. Let $\mathbf{A}_1, \mathbf{A}_2, \dots, \mathbf{A}_k$ be square matrices of the same size. Then

- $\mathbf{A}_1\mathbf{A}_2 \dots \mathbf{A}_k$ is invertible \iff all \mathbf{A}_i are invertible.
- $\mathbf{A}_1\mathbf{A}_2 \dots \mathbf{A}_k$ is singular \iff some \mathbf{A}_i are singular.

2.4.10 Elementary Column Operations

Definition. The elementary column operations are the following operations on columns of a matrix:

Description of Operation	Notation
Multiply the i th column by a nonzero constant k	kC_i
Interchange the i th and j th columns	$C_i \leftrightarrow C_j$
Add k times the j th column to the i th column	$C_i + kC_j$

2.4.11 Performing a Single Elementary Column Operation on Identity Matrix Gives an Elementary Matrix

Let \mathbf{E} be the **matrix** obtained from \mathbf{I} by a single **elementary column operation**. Then \mathbf{E} is an **elementary matrix** (i.e. \mathbf{E} can be obtained from \mathbf{I} by a single **elementary row operation**).

- $\mathbf{I} \xrightarrow{kC_i} \mathbf{E} \iff \mathbf{I} \xrightarrow{kR_i} \mathbf{E}.$
- $\mathbf{I} \xrightarrow{C_i \leftrightarrow C_j} \mathbf{E} \iff \mathbf{I} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{E}.$
- $\mathbf{I} \xrightarrow{C_i + kC_j} \mathbf{E} \iff \mathbf{I} \xrightarrow{R_j + kR_i} \mathbf{E}.$

2.4.12 Connection of Elementary Matrices to Matrix Post-Multiplication

Theorem. Let \mathbf{E} be an **elementary matrix** obtained by performing an **elementary column operation** on \mathbf{I}_n . Then for any $m \times n$ matrix \mathbf{A} , \mathbf{AE} can be obtained by performing the same **elementary column operation** to \mathbf{A} .

- $\mathbf{I} \xrightarrow{kC_i} \mathbf{E} \implies \mathbf{A} \xrightarrow{kC_i} \mathbf{AE}.$
- $\mathbf{I} \xrightarrow{C_i \leftrightarrow C_j} \mathbf{E} \implies \mathbf{A} \xrightarrow{C_i \leftrightarrow C_j} \mathbf{AE}.$
- $\mathbf{I} \xrightarrow{C_i + kC_j} \mathbf{E} \implies \mathbf{A} \xrightarrow{C_i + kC_j} \mathbf{AE}.$

2.5 Determinant

2.5.1 (i, j) -cofactor

Definition. Let $\mathbf{A} = (a_{ij})_{n \times n}$. Let \mathbf{M}_{ij} be the **submatrix** obtained from \mathbf{A} by deleting the i th row and j th column. Then for $1 \leq i, j \leq n$, the **(i, j) -cofactor** of \mathbf{A} is

$$A_{ij} = (-1)^{i+j} \det(\mathbf{M}_{ij})$$

2.5.2 Definition of Determinant of Square Matrix

Definition. Let $\mathbf{A} = (a_{ij})_{n \times n}$. Its **determinant** is:

- If $n = 1$, define $\det(\mathbf{A}) = a_{11}$.
- If $n > 1$, let A_{ij} be its (i, j) -cofactor, define

$$\circ \det(\mathbf{A}) = a_{11}A_{11} + a_{12}A_{12} + \dots + a_{1n}A_{1n}$$

2.5.3 Determinant of 2×2 Matrix

Let $\mathbf{A} = \begin{pmatrix} a & b \\ c & d \end{pmatrix}$. Then $\det(\mathbf{A}) = ad - bc$.

2.5.4 Broken Diagonal Formula for Determinant of 3×3 Matrix

Let $\mathbf{A} = \begin{pmatrix} a_{11} & a_{12} & a_{13} \\ a_{21} & a_{22} & a_{23} \\ a_{31} & a_{32} & a_{33} \end{pmatrix}$. Then

$$\det(\mathbf{A}) = (a_{11}a_{22}a_{33} + a_{12}a_{23}a_{31} + a_{13}a_{21}a_{32}) - (a_{11}a_{23}a_{32} + a_{12}a_{21}a_{33} + a_{13}a_{22}a_{31})$$

- The positive terms come from the
 - 3 (broken) diagonals from the top left to the bottom right.
- The negative terms come from the
 - 3 (broken) diagonals from the top right to the bottom left.

Warning: The "diagonal expansion" of $\det(\mathbf{A})$ for 2×2 or 3×3 matrices is not valid if the order ≥ 4 .

2.5.5 Properties of Determinant

2.5.5.1 Properties of Determinant

Theorem. For any **square matrices** \mathbf{A} and \mathbf{B} of the same size and any scalar c ,

- For $n \in \mathbb{N}$, $\det(\mathbf{I}_n) = 1$.
- $\det(\mathbf{A}) = \det(\mathbf{A}^T)$.
- $\det(\mathbf{AB}) = \det(\mathbf{A}) \det(\mathbf{B})$.
- $\det(c\mathbf{A}) = c^n \det(\mathbf{A})$, where \mathbf{A} is $n \times n$.
- $\det(\mathbf{A}^{-1}) = \det(\mathbf{A})^{-1}$ if \mathbf{A} is **invertible**.

2.5.5.2 Change in Determinant by Elementary Row Operations

Theorem. For any square matrices \mathbf{A} and \mathbf{B} ,

- $\mathbf{A} \xrightarrow{cR_i} \mathbf{B} \implies \det(\mathbf{B}) = c \det(\mathbf{A})$.
 - In particular, $\mathbf{I} \xrightarrow{cR_i} \mathbf{E} \implies \det(\mathbf{E}) = c$.
- $\mathbf{A} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{B} \implies \det(\mathbf{B}) = -\det(\mathbf{A})$.

- In particular, $\mathbf{I} \xrightarrow{R_i \leftrightarrow R_j} \mathbf{E} \implies \det(\mathbf{E}) = -1$.
- $\mathbf{A} \xrightarrow{R_i + cR_j} \mathbf{B} \implies \det(\mathbf{B}) = \det(\mathbf{A})$.
- In particular, $\mathbf{I} \xrightarrow{R_i + cR_j} \mathbf{E} \implies \det(\mathbf{E}) = 1$.

2.5.5.3 Change in Determinant by Elementary Matrices

Theorem. Let \mathbf{A} be a **square matrix**. For any **elementary matrix** of the same order,

$$\det(\mathbf{EA}) = \det(\mathbf{E}) \det(\mathbf{A})$$

2.5.5.4 Change in Determinant Between Row Equivalent Matrices

Theorem. Suppose **square matrices** \mathbf{A} and \mathbf{B} are **row equivalent**. Then there exist **elementary matrices** $\mathbf{E}_1, \mathbf{E}_2, \dots, \mathbf{E}_k$ such that

$$\mathbf{B} = \mathbf{E}_k \dots \mathbf{E}_2 \mathbf{E}_1 \mathbf{A}$$

Then

$$\det(\mathbf{B}) = \det(\mathbf{E}_k) \dots \det(\mathbf{E}_2) \det(\mathbf{E}_1) \det(\mathbf{A})$$

Since $\det(\mathbf{E}) \neq 0$ for every **elementary matrix** \mathbf{E} ,

- $\det(\mathbf{A}) = 0 \iff \det(\mathbf{B}) = 0$;
- Equivalently, $\det(\mathbf{A}) \neq 0 \iff \det(\mathbf{B}) \neq 0$.

2.5.5.5 Determinant of Square Matrix With Zero Row

Theorem. Suppose a **square matrix** \mathbf{A} has a **zero row**. Then $\det(\mathbf{A}) = 0$.

2.5.5.6 Invertibility and Determinant

Theorem. For any **square matrix** \mathbf{A}

- $\det(\mathbf{A}) = 0 \iff \mathbf{A}$ is **singular**;
- Equivalently, $\det(\mathbf{A}) \neq 0 \iff \mathbf{A}$ is **invertible**,

2.5.5.7 Determinant of Triangular Matrix

Theorem. Suppose $\mathbf{A} = (a_{ij})_{n \times n}$ is **triangular**. Then

$$\det(\mathbf{A}) = a_{11}a_{22} \dots a_{nn}$$

Remarks.

- Note that a **row-echelon form** of a **square matrix** is always **upper triangular**.
 - To find the **determinant** using **elementary row operation**, it suffices to use **Gaussian elimination** to get a **row-echelon form**.

2.5.6 Cofactor Expansion

Theorem. Let $\mathbf{A} = (a_{ij})_{n \times n}$ and A_{ij} denote the (i, j) -cofactor of \mathbf{A} . Then for $1 \leq i \leq n$,

- $\det(\mathbf{A}) = a_{i1}A_{i1} + a_{i2}A_{i2} + \dots + a_{in}A_{in}$
- This is called the **cofactor expansion along the i th row**.

and for $1 \leq j \leq n$,

- $\det(\mathbf{A}) = a_{1j}A_{1j} + a_{2j}A_{2j} + \dots + a_{nj}A_{nj}$
- This is called the **cofactor expansion along the j th column**.

Remarks.

- In evaluating the **determinant** using **cofactor expansion**,
 - expand along the row or column with the most zeros.

2.5.7 Finding Determinant Efficiently

Given $\mathbf{A} = (a_{ij})_{n \times n}$, $\det(\mathbf{A})$ may be found as follows:

- \mathbf{A} has a zero row/column $\implies \det(\mathbf{A}) = 0$.
- \mathbf{A} is **triangular** $\implies \det(\mathbf{A}) = a_{11}a_{22} \dots a_{nn}$.
- Suppose that \mathbf{A} is not **triangular**.
 - $n = 2 \implies \det(\mathbf{A}) = a_{11}a_{22} - a_{12}a_{21}$
 - If a row/column has many 0, use **cofactor expansion**.
 - Otherwise, use **elementary row operations** to get **row-echelon form**.
 - * $\det(\mathbf{EA}) = \det(\mathbf{E})\det(\mathbf{A})$.

2.5.8 Adjoint Matrix

2.5.8.1 Definition of Adjoint Matrix

Definition. Let \mathbf{A} be a square matrix of order n . The **(classical) adjoint** (or **adjugate**, or **adjunct**) of \mathbf{A} is

$$\text{adj}(\mathbf{A}) = (A_{ji})_{n \times n}$$

where A_{ij} is the (i, j) -cofactor of \mathbf{A} .

2.5.8.2 Properties of Adjoint Matrix

Theorem. Let \mathbf{A} and \mathbf{B} be square matrices of order n . Then

- $\mathbf{A}[\text{adj}(\mathbf{A})] = \det(\mathbf{A})\mathbf{I}$.
- $[\text{adj}(\mathbf{A})]\mathbf{A} = \det(\mathbf{A})\mathbf{I}$.
- If \mathbf{A} is invertible, then
 - $\mathbf{A}^{-1} = \frac{1}{\det(\mathbf{A})} \text{adj}(\mathbf{A})$
 - $[\text{adj}(\mathbf{A})]^{-1} = \frac{1}{\det(\mathbf{A})} \mathbf{A}$
 - $\text{adj}(\mathbf{A}^{-1}) = [\text{adj}(\mathbf{A})]^{-1}$
 - $\det(\text{adj}(\mathbf{A})) = [\det(\mathbf{A})]^{n-1}$
 - $\text{adj}(\mathbf{AB}) = \text{adj}(\mathbf{B}) \text{adj}(\mathbf{A})$

2.5.9 Cramer's Rule

Let \mathbf{A} be an invertible matrix of order n . Then for every column matrix \mathbf{b} of size $n \times 1$, the linear system $\mathbf{Ax} = \mathbf{b}$ has a unique solution:

$$\mathbf{x} = \frac{1}{\det(\mathbf{A})} \begin{pmatrix} \det(\mathbf{A}_1) \\ \det(\mathbf{A}_2) \\ \vdots \\ \det(\mathbf{A}_n) \end{pmatrix}$$

where \mathbf{A}_j is obtained from \mathbf{A} by replacing its j th column by \mathbf{b} . Therefore, for $j = 1, 2, \dots, n$,

$$x_j = \frac{\det(\mathbf{A}_j)}{\det(\mathbf{A})}$$

Chapter 3

Vector Spaces

3.1 Euclidean n -Spaces

3.1.1 Introduction to n -vectors

3.1.1.1 Definition of n -vector

Definition. An n -vector or ordered n -tuple of real numbers is

$$\mathbf{v} = (v_1, v_2, \dots, v_n)$$

where $v_i \in \mathbb{R}$ is the i th component or i th coordinate of \mathbf{v} .

3.1.1.2 Zero Vector

Definition. The n -vector $\mathbf{0} = (0, 0, \dots, 0)$ is the **zero vector**.

3.1.2 n -vector Operations

3.1.2.1 Equal n -vectors

Definition. Let $\mathbf{u} = (u_1, u_2, \dots, u_n)$ and $\mathbf{v} = (v_1, v_2, \dots, v_n)$. Then \mathbf{u} and \mathbf{v} are **equal** if $u_i = v_i$ for all $i = 1, \dots, n$.

3.1.2.2 n -vector Addition, Subtraction & Scalar Multiplication

Definition. Let $\mathbf{u} = (u_1, u_2, \dots, u_n)$, $\mathbf{v} = (v_1, v_2, \dots, v_n)$, and $c \in \mathbb{R}$. The following operations are defined:

- **Scalar Multiplication:** $c\mathbf{v} = (cv_1, cv_2, \dots, cv_n)$
- **Negative:** $-\mathbf{v} = (-1)\mathbf{v}$
- **Addition:** $\mathbf{u} + \mathbf{v} = (u_1 + v_1, u_2 + v_2, \dots, u_n + v_n)$
- **Subtraction:** $\mathbf{u} - \mathbf{v} = (u_1 - v_1, u_2 - v_2, \dots, u_n - v_n)$

3.1.3 Properties of n -vectors

Theorem. Let $\mathbf{u}, \mathbf{v}, \mathbf{w}$ be n -vectors and $c, d \in \mathbb{R}$. Then

- $\mathbf{u} - \mathbf{v} = \mathbf{u} + (-\mathbf{v})$
- Commutative Law for Vector Addition:
 - $\mathbf{u} + \mathbf{v} = \mathbf{v} + \mathbf{u}$
- Associative Law for Vector Addition:
 - $(\mathbf{u} + \mathbf{v}) + \mathbf{w} = \mathbf{u} + (\mathbf{v} + \mathbf{w})$
- Additive Identity and Additive Inverse:
 - $\mathbf{v} + \mathbf{0} = \mathbf{v}$ and $\mathbf{v} + (-\mathbf{v}) = \mathbf{0}$
- Distributive Law for Scalar Multiplication over Addition:
 - $c(\mathbf{u} + \mathbf{v}) = c\mathbf{u} + c\mathbf{v}$
 - $(c + d)\mathbf{v} = c\mathbf{v} + d\mathbf{v}$
- $c(d\mathbf{v}) = (cd)\mathbf{v} = d(c\mathbf{v})$
 - In particular, $-c\mathbf{v} = (-c)\mathbf{v} = c(-\mathbf{v})$ and $-(-\mathbf{v}) = \mathbf{v}$
- $1\mathbf{v} = \mathbf{v}$

3.1.4 n -vectors as Matrices

An n -vector (v_1, v_2, \dots, v_n) can be viewed as

- a **row matrix** (**row vector**) $\begin{pmatrix} v_1 & v_2 & \dots & v_n \end{pmatrix}$, or
- a **column matrix** (**column vector**) $\begin{pmatrix} v_1 \\ v_2 \\ \vdots \\ v_n \end{pmatrix}$

3.1.5 Vectors in xy -Plane

3.1.5.1 Vectors as Points on the xy -Plane

Every point P on the xy -plane is represented by a vector $\mathbf{v} = (a, b)$ where

- a is the x -coordinate and b is the y -coordinate; and
- \mathbf{v} is the arrow from the origin O to the point P , denoted by \overrightarrow{OP}

3.1.5.2 Vectors as Change from Initial Point to End Point

A vector (a, b) represents the change from the **initial point** (x_1, y_1) to the **end point** (x_2, y_2) where $a = x_2 - x_1, b = y_2 - y_1$.

3.1.5.3 Length of Vectors

Definition. Let $\mathbf{v} = (v_1, v_2)$ be a vector in xy -plane. Its **length** is

$$\|\mathbf{v}\| = \sqrt{v_1^2 + v_2^2}$$

3.1.5.4 Properties of Length of Vectors

Theorem.

- $\|c\mathbf{v}\| = |c|\|\mathbf{v}\|$
- $\mathbf{v} = \mathbf{0} \iff \|\mathbf{v}\| = 0$

3.1.5.5 Geometrical Interpretation of Scalar Multiplication

Let $\mathbf{v} = (v_1, v_2)$ and $c \in \mathbb{R}$. Then

- $c\mathbf{v}$ is a vector **parallel** to \mathbf{v} such that
 - its **length** is $|c|$ times the **length** of \mathbf{v}
- 1. $c = 0 \implies c\mathbf{v} = 0\mathbf{v} = \mathbf{0}$ is the **zero vector**.
- 2. $c > 0 \implies c\mathbf{v}$ has the same direction as \mathbf{v}
- 3. $c < 0 \implies c\mathbf{v}$ has the opposite direction of \mathbf{v}

3.1.5.6 Geometrical Interpretation of Vector Addition

Let $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$. Then $\mathbf{u} + \mathbf{v}$ is the vector obtained as follows:

1. Parallel shift \mathbf{v} so that its initial point is the same as the end of \mathbf{u} .
2. Then $\mathbf{u} + \mathbf{v}$ is the vector from the initial point of \mathbf{u} to the end point of \mathbf{v} .

3.1.5.7 Geometrical Interpretation of Vector Subtraction

Let $\mathbf{u} = (u_1, u_2)$ and $\mathbf{v} = (v_1, v_2)$. Then $\mathbf{u} - \mathbf{v}$ is the vector obtained as follows:

1. Parallel shift \mathbf{v} so that \mathbf{u} and \mathbf{v} have the same initial point.
2. Then $\mathbf{u} - \mathbf{v}$ is the vector from the end point of \mathbf{v} to the end point of \mathbf{u} .

3.1.6 Vectors in xyz -Space

3.1.6.1 Vectors as Points on the xyz -Space

Every point P on the xyz -space is represented by a vector $\mathbf{v} = (a, b, c)$ where

- a is the x -coordinate, b is the y -coordinate, and c is the z -coordinate; and
- \mathbf{v} is the arrow from the origin O to the point P , denoted by \overrightarrow{OP}

3.1.7 Euclidean Spaces

Definition. The **Euclidean n -space** (or simply **n -space**) is the set of all n -vectors of real numbers, denoted by \mathbb{R}^n where

$$\mathbb{R}^n = \{(v_1, v_2, \dots, v_n) \mid v_1, v_2, \dots, v_n \in \mathbb{R}\}$$

$\mathbf{v} \in \mathbb{R}^n \iff \mathbf{v}$ is of the form $\mathbf{v} = (v_1, v_2, \dots, v_n)$ for real numbers v_1, v_2, \dots, v_n

- In particular,
 - $n = 1 \implies \mathbb{R} = \mathbb{R}^1$ is the real line.
 - $n = 2 \implies \mathbb{R}^2$ is the xy -plane.
 - $n = 3 \implies \mathbb{R}^3$ is the xyz -space.
- Given a **linear system** $\mathbf{Ax} = \mathbf{b}$ in m equations and n variables
 - \mathbf{x} as be viewed as an n -vector, i.e. $\mathbf{x} \in \mathbb{R}^n$
 - Then the **solution set** of $\mathbf{Ax} = \mathbf{b}$ is a subset of \mathbb{R}^n

3.1.8 Implicit and Explicit Forms of Linear Systems

3.1.8.1 Definition of Implicit and Explicit Forms

Definition.

- A **linear system** is given in the **implicit form** as follows:

$$\begin{cases} a_{11}x_1 + a_{12}x_2 + \dots + a_{1n}x_n = b_1, \\ a_{21}x_1 + a_{22}x_2 + \dots + a_{2n}x_n = b_2, \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \dots + a_{mn}x_n = b_m, \end{cases}$$

- Its general solution is in the **explicit form**.

3.1.8.2 Lines in \mathbb{R}^2 **3.1.8.2.1 Vector Equation for line in \mathbb{R}^2**

A **straight line** in \mathbb{R}^2 is determined by a point (x_0, y_0) on the line, and its direction vector $(a, b) \neq \mathbf{0}$.

- A point on the line is of the form $(x_0, y_0) + t(a, b)$, where $t \in \mathbb{R}$

3.1.8.2.2 Implicit to Explicit Form

Refer to Chapter 1.1.5 on how to convert implicit form to explicit form.

3.1.8.2.3 Explicit to Implicit Form

Given an explicit form of a line in \mathbb{R}^2 :

$$\{(x_0 + at, y_0 + bt) \mid t \in \mathbb{R}\}$$

where x_0, y_0, a, b are real constants. An implicit form may be obtained as follows:

1. Let $x = x_0 + at$, and $y = y_0 + bt$
2. Then $t = \frac{x - x_0}{a}$ and $t = \frac{y - y_0}{b} \implies \frac{x - x_0}{a} = \frac{y - y_0}{b}$
3. Cross multiply to get $bx - bx_0 = ay - ay_0 \implies bx - ay = bx_0 - ay_0$
4. Hence, the implicit form is $\{(x, y) \mid bx - ay = bx_0 - ay_0\}$

3.1.8.3 Planes in \mathbb{R}^3 **3.1.8.3.1 Vector Equation for plane in \mathbb{R}^3**

A **plane** in \mathbb{R}^3 is determined by a point (x_0, y_0, z_0) on the plane, and two non-parallel vectors parallel to the plane (a_1, b_1, c_1) and (a_2, b_2, c_2) .

- A point on the plane is of the form $(x_0, y_0, z_0) + s(a_1, b_1, c_1) + t(a_2, b_2, c_2)$, where $s, t \in \mathbb{R}$

3.1.8.3.2 Implicit to Explicit Form

Refer to Chapter 1.1.5 on how to convert implicit form to explicit form.

3.1.8.3.3 Explicit to Implicit Form

Given an explicit form of a plane in \mathbb{R}^3 :

$$\{(x_0 + a_1s + a_2t, y_0 + b_1s + b_2t, z_0 + c_1s + c_2t) \mid t \in \mathbb{R}\}$$

where $x_0, y_0, z_0, a_i, b_i, c_i$ are real constants. An implicit form may be obtained as follows:

1. Let $x = x_0 + a_1s + a_2t$, $y = y_0 + b_1s + b_2t$, and $z = z_0 + c_1s + c_2t$

2. Then we obtain the following **linear system** in s, t :

$$\begin{cases} a_1s + a_2t = x - x_0 \\ b_1s + b_2t = y - y_0 \\ c_1s + c_2t = z - z_0 \end{cases}$$

3. Perform **Gaussian Elimination** to its corresponding **augmented matrix**.

4. The $(3, 3)$ -entry of its **row-echelon form** is a function f in variables x, y, z

5. Since the system is **consistent**, $f(x, y, z) = 0$.

6. Hence, the implicit form is $\{(x, y, z) \mid f(x, y, z) = 0\}$

3.1.8.4 Lines in \mathbb{R}^3

3.1.8.4.1 Vector Equation for line in \mathbb{R}^3

A **straight line** in \mathbb{R}^3 is determined by a point (x_0, y_0, z_0) on the line, and its direction vector $(a, b, c) \neq \mathbf{0}$.

- A point on the line is of the form $(x_0, y_0, z_0) + t(a, b, c)$, where $t \in \mathbb{R}$

3.1.8.4.2 Implicit to Explicit Form

Solve the **linear system** of two equations (each representing a plane) in three variables to convert implicit form to explicit form.

3.1.8.4.3 Explicit to Implicit Form

Given an explicit form of a line in \mathbb{R}^3 :

$$\{(x_0 + at, y_0 + bt, z_0 + ct) \mid t \in \mathbb{R}\}$$

where x_0, y_0, z_0, a, b, c are real constants. An implicit form may be obtained as follows:

1. Let $x = x_0 + at$, $y = y_0 + bt$, and $z = z_0 + ct$
2. Find the relation between x and y , say $f(x, y) = 0$
3. Find the relation between x and z , say $g(x, z) = 0$
4. Then, the implicit form is $\{(x, y, z) \mid f(x, y) = 0 \text{ \& } g(x, z) = 0\}$

3.2 Linear Combinations and Linear Spans

3.2.1 Definition of Linear Combination

Definition. Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ be vectors in \mathbb{R}^n . A **linear combination** of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ has the form

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k$$

where $c_1, c_2, \dots, c_k \in \mathbb{R}$

Remarks. In particular, $\mathbf{0}$ is a **linear combination** of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$:

$$\mathbf{0} = 0\mathbf{v}_1 + 0\mathbf{v}_2 + \dots + 0\mathbf{v}_k$$

3.2.2 Definition of Linear Span

Definition. Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ be a subset of \mathbb{R}^n . The **set of all linear combinations** of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ is

$$\{c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k \mid c_1, c_2, \dots, c_k \in \mathbb{R}\}$$

and is called the **linear span** of S (or $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$), denoted by $\text{span}(S)$ or $\text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ respectively.

Remarks. \mathbf{v} is a linear combination of $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \iff \mathbf{v} \in \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$

3.2.3 Criterion for $\text{span}(S) = \mathbb{R}^n$

3.2.3.1 When $k \geq n$

Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\} \subseteq \mathbb{R}^n$. The following steps allow us to check whether $\text{span}(S) = \mathbb{R}^n$:

1. View each \mathbf{v}_j as a column vector
2. Let $\mathbf{A} = \begin{pmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_k \end{pmatrix}$
3. Find a **row-echelon form** \mathbf{R} of \mathbf{A}
 - If \mathbf{R} has a zero row, then $\text{span}(S) \neq \mathbb{R}^n$
 - If \mathbf{R} has no zero row, then $\text{span}(S) = \mathbb{R}^n$

3.2.3.2 When $k < n$

Theorem. Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\} \subseteq \mathbb{R}^n$. $k < n \implies \text{span}(S) \neq \mathbb{R}^n$

3.2.4 Properties of Linear Spans

3.2.4.1 Linear Spans Always Contains Zero Vector

Theorem. Let $S = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\} \subseteq \mathbb{R}^n$. Then

- $\mathbf{0} \in \text{span}(S)$, where $\mathbf{0}$ is the zero vector in \mathbb{R}^n
 - Hence, $\text{span}(S) \neq \emptyset$

3.2.4.2 Linear Spans Are Closed Under Linear Combination

Theorem. Let $S = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\} \subseteq \mathbb{R}^n$, $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_r \in \text{span}(S)$, and $c_1, c_2, \dots, c_r \in \mathbb{R}$. Then

$$c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_r\mathbf{v}_r \in \text{span}(S)$$

In particular,

- $\text{span}(S)$ is **closed** under scalar multiplication.
 - $\mathbf{v} \in \text{span}(S)$ and $c \in \mathbb{R} \implies c\mathbf{v} \in \text{span}(S)$
- $\text{span}(S)$ is **closed** under addition.
 - $\mathbf{u} \in \text{span}(S)$ and $\mathbf{v} \in \text{span}(S) \implies \mathbf{u} + \mathbf{v} \in \text{span}(S)$

3.2.4.3 When a Span is a Subset of Another Span

Theorem. Given two subsets of \mathbb{R}^n :

$$S_1 = \{\mathbf{u}_1, \mathbf{u}_2, \dots, \mathbf{u}_k\}, S_2 = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m\}$$

Then

$$\text{span}(S_1) \subseteq \text{span}(S_2) \iff \text{Every } \mathbf{u}_i \text{ is a linear combination of } \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_m$$

3.2.4.4 Linear Spans with Redundant Vectors

Theorem. Let $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{k-1}, \mathbf{v}_k \in \mathbb{R}^n$. Then

$$\mathbf{v}_k \text{ is a linear combination of } \mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{k-1} \implies \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{k-1}\} = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_{k-1}, \mathbf{v}_k\}$$

3.2.5 Criterion for Vector Belongs to Span

Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\} \subseteq \mathbb{R}^n$. The following steps allow us to check whether a vector $\mathbf{v} \in \text{span}(S)$:

1. View each \mathbf{v}_j as a **column vector**.
2. Let $\mathbf{A} = \begin{pmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \dots & \mathbf{v}_k \end{pmatrix}$
3. Check if the **linear system** $\mathbf{A}\mathbf{x} = \mathbf{v}$ is consistent.
 - If $\mathbf{A}\mathbf{x} = \mathbf{v}$ is consistent, then $\mathbf{v} \in \text{span}(S)$
 - If $\mathbf{A}\mathbf{x} = \mathbf{v}$ is inconsistent, then $\mathbf{v} \notin \text{span}(S)$

3.3 Subspaces

3.3.1 Definition of Subspaces

Definition. Let V be a subset of \mathbb{R}^n . Then V is called a **subspace** of \mathbb{R}^n if there exist $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$ such that:

$$V = \text{span}\{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$$

More precisely,

- V is the **subspace spanned** by $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$;
- $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\}$ **spans** the subspace V .

3.3.2 Zero Space

Let $\mathbf{0} \in \mathbb{R}^n$ be the zero vector. Then

$$\{\mathbf{0}\} = \text{span}\{\mathbf{0}\} \text{ is a subspace of } \mathbb{R}^n \text{ called the } \mathbf{zero\ space}$$

3.3.3 Euclidean n-Space is a Subspace

Let \mathbf{e}_i denote the n -vector whose i th coordinate is 1 and elsewhere 0. Then

$$\mathbb{R}^n = \text{span}\{\mathbf{e}_1, \mathbf{e}_2, \dots, \mathbf{e}_n\} \text{ is a subspace of } \mathbb{R}^n$$

3.3.4 Showing That a Subset is a Subspace

To show that a subset V of \mathbb{R}^n is a subspace:

1. Find $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k \in \mathbb{R}^n$ such that
 - V is the set containing all vectors of the form:

$$\mathbf{v} = c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k$$

where $c_1, c_2, \dots, c_k \in \mathbb{R}$

3.3.5 Showing That a Subset is Not a Subspace

A subset V of \mathbb{R}^n is not a subspace if any of the following fails:

- $\mathbf{0} \in V$
- $c \in \mathbb{R} \ \& \ \mathbf{v} \in V \implies c\mathbf{v} \in V$
- $\mathbf{u} \in V \ \& \ \mathbf{v} \in V \implies \mathbf{u} + \mathbf{v} \in V$

3.3.6 Subspaces of $\mathbb{R}^1, \mathbb{R}^2, \mathbb{R}^3$

3.3.6.1 Subspaces of $\mathbb{R}^1 = \mathbb{R}$

Let the nonzero vector $\mathbf{v} = v \in \mathbb{R}$. The following are the **subspaces** of \mathbb{R}^1 :

- $\{0\}$,
- $\mathbb{R} = \text{span}\{\mathbf{v}\}$.

3.3.6.2 Subspaces of \mathbb{R}^2

Let the nonzero vectors $\mathbf{u} = (u_1, u_2) \in \mathbb{R}^2, \mathbf{v} = (v_1, v_2) \in \mathbb{R}^2$. The following are the **subspaces** of \mathbb{R}^2 :

- $\{\mathbf{0}\} = \{(0, 0)\}$,
- $\text{span}\{\mathbf{v}\}$ (straight line passing through the origin),
- $\mathbb{R}^2 = \text{span}\{\mathbf{u}, \mathbf{v}\}$ if \mathbf{u} and \mathbf{v} are not parallel.

3.3.6.3 Subspaces of \mathbb{R}^3

Let the nonzero vectors $\mathbf{u} = (u_1, u_2, u_3) \in \mathbb{R}^3, \mathbf{v} = (v_1, v_2, v_3) \in \mathbb{R}^3$. The following are the **subspaces** of \mathbb{R}^3 :

- $\{\mathbf{0}\} = \{(0, 0, 0)\}$,
- $\text{span}\{\mathbf{v}\}$ (straight line passing through the origin or intersection of two planes containing the origin),
- $\text{span}\{\mathbf{u}, \mathbf{v}\}$ if \mathbf{u} and \mathbf{v} are not parallel (a plane containing the origin),
- \mathbb{R}^3

3.3.7 Solution Set of Homogeneous Linear System is a Subspace of \mathbb{R}^n

Theorem. The solution set of a homogeneous linear system of n variables is a **subspace** of \mathbb{R}^n .

3.3.8 Solution Space

Definition. The solution set of a homogeneous linear system is called the **solution space** of the system.

3.3.9 A Subspace of \mathbb{R}^n is Always The Solution Space of a Homogeneous Linear System

Theorem. The subspace of \mathbb{R}^n is always the solution space of a homogeneous linear system.

3.4 Vector Spaces

3.4.1 Definition of Vector Space

Definition. A set V is a vector space if V is a subspace of \mathbb{R}^n for some $n \in \mathbb{N}$.

3.4.2 Expanded Definition of Subspace

Definition. If W and V are vector spaces such that $W \subseteq V$, then W is a subspace of V .

3.5 Linear Independence

3.5.1 Definition of Linear Independence

Definition. Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k\} \subseteq \mathbb{R}^n$. If the equation $c_1\mathbf{v}_1 + c_2\mathbf{v}_2 + \dots + c_k\mathbf{v}_k = \mathbf{0}$ has a non-trivial solution, then

- S is a linearly dependent set,
- $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly dependent.

If the equation has only the trivial solution, then

- S is a linearly independent set,
- $\mathbf{v}_1, \mathbf{v}_2, \dots, \mathbf{v}_k$ are linearly independent.

3.5.2 Linear Independence of Subset/Superset

Theorem. Let S_1, S_2 be finite subsets of \mathbb{R}^n such that $S_1 \subseteq S_2$. Then

3.5.2.1 Superset Preserves Linear Dependence

S_1 linearly dependent $\implies S_2$ linearly dependent.

3.5.2.2 Subset Preserves Linear Independence

S_2 linearly independent $\implies S_1$ linearly independent.

3.5.3 Linear Independence of Sets Containing Zero Vector

Theorem. $0 \in S \subseteq \mathbb{R}^n \implies S$ is linearly dependent.

3.5.4 Linear Independence of Sets Containing Only One Vector

Theorem. Let $v \in \mathbb{R}^n$. Then $\{v\}$ is linearly independent $\iff v \neq 0$.

3.5.5 Linear Independence and Linear Combinations

Theorem. Let $S = \{v_1, v_2, \dots, v_k\} \subseteq \mathbb{R}^n, k \geq 2$. Then S is linearly dependent \iff there exists v_i such that $v_i \in \text{span}\{v_1, \dots, v_{i-1}, v_{i+1}, \dots, v_k\}$.

3.5.6 Cardinality Upper Bound for Linear Independence

Theorem. Let $S \subseteq \mathbb{R}^n$. $|S| > n \implies S$ is linearly dependent.

3.5.7 Adding Vectors to a Linearly Independent Set

Theorem. Suppose $S \subseteq \mathbb{R}^n$ is linearly independent, $v \in \mathbb{R}^n$, and $v \notin \text{span}(S)$. Then $S \cup \{v\}$ is linearly independent.

3.6 Bases

3.6.1 Definition of Bases

Definition. Let S be a finite subset of a vector space V . Then S is a **basis** (plural **bases**) for V if

- S is linearly independent, and
- $\text{span}(S) = V$.

3.6.1.1 Remarks

- A **basis** for a vector space V contains
 - the smallest possible number of vectors that spans V , and
 - the largest possible number of vectors that are linearly independent
- For convenience, the **basis** for $\{0\}$ is defined to be \emptyset .
- Other than $\{0\}$, any vector space has infinitely many different **bases**.

3.6.2 Basis and Unique Linear Combination

Theorem. Let S be a finite subset of a **vector space** V . Then the following are equivalent

- S is a **basis** for V .
- Every $\mathbf{v} \in V$ can be **uniquely** expressed (there is exactly one $c_1, \dots, c_k \in \mathbb{R}$) as

$$\circ \mathbf{v} = c_1 \mathbf{v}_1 + \dots + c_k \mathbf{v}_k$$

3.6.3 Coordinate Vector

Definition. Let S be a finite subset of a **vector space** V .

- For every $\mathbf{v} \in V$, there exist unique $c_1, \dots, c_k \in \mathbb{R}$ such that

$$\circ \mathbf{v} = c_1 \mathbf{v}_1 + \dots + c_k \mathbf{v}_k$$

- c_1, \dots, c_k are the **coordinates** of \mathbf{v} **relative** to S .
- (c_1, \dots, c_k) is the **coordinate vector** of \mathbf{v} **relative** to the basis S , denoted by $(\mathbf{v})_S$.

3.6.3.1 Remark

The order of $\mathbf{v}_1, \dots, \mathbf{v}_k$ is fixed.

3.6.4 Standard Basis

Definition. Let $E = \{\mathbf{e}_1, \dots, \mathbf{e}_n\} \subseteq \mathbb{R}^n$ where

- $\mathbf{e}_1 = (1, 0, \dots, 0), \mathbf{e}_2 = (0, 1, \dots, 0), \dots, \mathbf{e}_n = (0, 0, \dots, 1)$

E is the **standard basis** for \mathbb{R}^n .

- For any $\mathbf{v} = (v_1, \dots, v_n) \in \mathbb{R}^n$,

$$\circ (\mathbf{v})_E = (v_1, \dots, v_n) = \mathbf{v}$$

3.6.5 Properties of Coordinate Vector

Theorem. Let S be a **basis** for a **vector space** V . Suppose $|S| = k$. Let $\mathbf{v}_1, \dots, \mathbf{v}_r \in V$

3.6.5.1 Zero Coordinate Vector

$$(\mathbf{v})_S = \mathbf{0} \Leftrightarrow \mathbf{v} = \mathbf{0}$$

3.6.5.2 Homogeneity of Degree One / Scalar Multiplication Preserving

For any $c \in \mathbb{R}$ and $\mathbf{v} \in V$, $(c\mathbf{v})_S = c(\mathbf{v})_S$.

3.6.5.3 Additivity / Addition Preserving

For any $\mathbf{u}, \mathbf{v} \in V$, $(\mathbf{u} + \mathbf{v})_S = (\mathbf{u})_S + (\mathbf{v})_S$.

3.6.5.4 Equal Coordinate Vectors

For any $\mathbf{u}, \mathbf{v} \in V$, $\mathbf{u} = \mathbf{v} \Leftrightarrow (\mathbf{u})_S = (\mathbf{v})_S$.

3.6.5.5 Linear Combination Preserving

For any $c_1, \dots, c_r \in \mathbb{R}$, $(c_1\mathbf{v}_1 + \dots + c_r\mathbf{v}_r)_S = c_1(\mathbf{v}_1)_S + \dots + c_r(\mathbf{v}_r)_S$.

3.6.5.6 Linear Independence Preserving

$\mathbf{v}_1, \dots, \mathbf{v}_r$ are **linearly independent** $\Leftrightarrow (\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S$ are **linearly independent**.

3.6.5.7 Span Preserving

$\text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_r\} = V \Leftrightarrow \text{span}\{(\mathbf{v}_1)_S, \dots, (\mathbf{v}_r)_S\} = \mathbb{R}^k$.

3.7 Dimensions

3.7.1 Dimension Theorem for Vector Spaces

Theorem. Let V be a **vector space** having a **basis** with k vectors. Then

- Any subset of V of $> k$ vectors is **linearly dependent**.
- Any subset of V of $< k$ vectors cannot span V .

3.7.1.1 Corollary

All **bases** of a **vector space** have the same cardinality.

- To be more precise, if S_1 and S_2 are two **bases** of a **vector space** V ,
 - then $|S_1| = |S_2|$

3.7.2 Definition of Dimension

Definition. Let V be a **vector space** and S a **basis** for V . Then

- the **dimension** of V , denoted by $\dim(V)$, is $|S|$

3.7.2.1 Examples

- $\dim(\{\mathbf{0}\}) = 0$
- $\dim(\mathbb{R}^n) = n$
- **Straight Line Through the Origin in \mathbb{R}^2 and \mathbb{R}^3**
 - $\dim(\text{span}(\mathbf{v})) = 1$ with $\mathbf{v} \neq \mathbf{0}$.
- **Plane Containing Origin in \mathbb{R}^3**
 - $\dim(\text{span}(\mathbf{u}, \mathbf{v})) = 2$ where \mathbf{u} and \mathbf{v} are **linearly independent**.

3.7.3 Dimension of Solution Space

Let $\mathbf{A}\mathbf{x} = \mathbf{0}$ be a **homogeneous linear system** and V be its **solution space**. Let \mathbf{R} be a **row-echelon form** of \mathbf{A} . Then

$$\begin{aligned} \text{no. of non-pivot cols of } \mathbf{R} &= \text{no. of arbitrary parameters in soln} \\ &= \dim(V) \end{aligned}$$

3.7.4 Easier Criterion for Basis

Theorem. Let S be a subset of a **vector space** V . The following are equivalent:

1. S is a **basis** for V .
2. S is **linearly independent**, and $|S| = \dim(V)$.
3. $\text{span}(S) = V$ and $|S| = \dim(V)$.

3.7.5 Dimension for Subspaces

Let U be a **subspace** of a **vector space** V . Then

- $U = V \Leftrightarrow \dim(U) = \dim(V)$.

3.7.5.1 Corollary

Let U be a **subspace** of a **vector space** V . Then

- $U \neq V \Leftrightarrow \dim(U) < \dim(V)$.

3.7.6 Invertibility and Rows and Columns as Basis

Theorem. Let \mathbf{A} be a **square matrix** of **order** n . Then the following are equivalent:

1. \mathbf{A} is **invertible**.
2. The rows of \mathbf{A} form a **basis** for \mathbb{R}^n .
3. The columns of \mathbf{A} form a **basis** for \mathbb{R}^n .

3.8 Transition Matrices

3.8.1 Coordinate Vector As a Column Vector

Definition. Let S be a **basis** for a **vector space** V and $\mathbf{v} \in V$ such that $(\mathbf{v})_S = (c_1, \dots, c_k)$.

Then the **column vector** $[\mathbf{v}]_S = \begin{pmatrix} c_1 \\ \vdots \\ c_k \end{pmatrix}$ is also called the **coordinate vector** of \mathbf{v} relative to S .

3.8.2 Definition of Transition Matrix

Definition. Let V be a **vector space**, and $S = \{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ and T be **bases** for V . Then the **transition matrix** from S to T is

$$\mathbf{P} = \begin{pmatrix} [\mathbf{u}_1]_T & \cdots & [\mathbf{u}_k]_T \end{pmatrix}$$

3.8.3 Pre-Multiplication of Transition Matrix

Theorem. Let S and T be **bases** for a **vector space** V , and \mathbf{P} be the **transition matrix** from S to T . Then $\forall \mathbf{w} \in V \quad \mathbf{P}[\mathbf{w}]_S = [\mathbf{w}]_T$.

3.8.4 Inverse of Transition Matrices

Theorem. Let S and T be **bases** for a **vector space** V , and \mathbf{P} be the **transition matrix** from S to T . Then

3.8.4.1 Transition Matrices Are Invertible

\mathbf{P} is **invertible**.

3.8.4.2 Inverse is a Transition Matrix

\mathbf{P}^{-1} is the **transition matrix** from T to S .

Chapter 4

Vector Spaces Associated With Matrices

4.1 Row Spaces and Column Spaces

4.1.1 Definition of Row Spaces and Column Spaces

Definition. Let $\mathbf{A} = (a_{ij})_{m \times n}$.

4.1.1.1 Row Space

For $1 \leq i \leq m$, let

$$\begin{aligned}\mathbf{r}_i &= i\text{th row of } \mathbf{A} \\ &= (a_{i1} \quad \cdots \quad a_{in}) \in \mathbb{R}^n\end{aligned}$$

Then the **row space** of $\mathbf{A} = \text{span}\{\mathbf{r}_1, \dots, \mathbf{r}_m\}$ is a **subspace** of \mathbb{R}^n .

4.1.1.2 Column Space

For $1 \leq j \leq n$, let

$$\begin{aligned}\mathbf{c}_j &= j\text{th column of } \mathbf{A} \\ &= \begin{pmatrix} a_{1j} \\ \vdots \\ a_{mj} \end{pmatrix} \in \mathbb{R}^m\end{aligned}$$

Then the **column space** of $\mathbf{A} = \text{span}\{\mathbf{c}_1, \dots, \mathbf{c}_n\}$ is a **subspace** of \mathbb{R}^m .

4.1.2 Row Equivalent Matrices And Their Rows

4.1.2.1 Row Equivalence Implies Same Row Space

Theorem. Suppose \mathbf{A} and \mathbf{B} are row equivalent matrices. Then row space of $\mathbf{A} =$ row space of \mathbf{B} .

4.1.2.1.1 Remark

Let \mathbf{R} be a row-echelon form of \mathbf{A} . Then the row space of $\mathbf{A} = \text{row space of } \mathbf{R}$.

4.1.2.2 Nonzero Rows of REF are Linearly Independent

Theorem. Let \mathbf{R} be a row-echelon form of a matrix \mathbf{A} . Then the nonzero rows of \mathbf{R} are linearly independent.

4.1.2.3 Finding Basis/Dimension For Row Space

Theorem. Let \mathbf{R} be a row-echelon form of a matrix \mathbf{A} . Then

- the nonzero rows of \mathbf{R} form a **basis** for the row space of \mathbf{A} , and
- $\dim(\text{row space of } \mathbf{A}) = \text{no. of nonzero rows of } \mathbf{R}$.

4.1.3 Row Equivalent Matrices And Their Columns

4.1.3.1 Row Equivalence Preserve Linear Relations Between Columns

Theorem. Let \mathbf{A} and \mathbf{B} be row equivalent matrices. Then

- There is a linear relation among a given set of columns of $\mathbf{A} \Leftrightarrow$ the same linear relation exists among the corresponding set of columns of \mathbf{B} .
- A given set of columns of \mathbf{A} is **linearly independent** \Leftrightarrow the corresponding set of columns of \mathbf{B} is **linearly independent**.
- A given set of columns of \mathbf{A} is a **basis** for the column space of $\mathbf{A} \Leftrightarrow$ the corresponding set of columns of \mathbf{B} is a **basis** for the column space of \mathbf{B} .

4.1.3.2 Pivot Columns of REF is a Basis For Column Space

Theorem. Let \mathbf{R} be a row-echelon form of a matrix \mathbf{A} . Then the pivot columns of \mathbf{R} is a basis for the column space of \mathbf{R} .

4.1.3.3 Finding Basis/Dimension For Column Space

Theorem. Let \mathbf{R} be a row-echelon form of a matrix \mathbf{A} . The pivot columns of \mathbf{R} is a basis for the column space of \mathbf{R} . Then

- the corresponding columns of \mathbf{A} is a **basis** for the column space of \mathbf{A} , and
- $\dim(\text{column space of } \mathbf{A}) = \text{no. of pivot columns of } \mathbf{R}$.

4.1.4 Finding Basis for Vector Space Spanned By a Set of Vectors

To find a **basis** for a vector space $V = \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_m\}$, choose either one of the methods below:

4.1.4.1 Method 1: From New Vectors Using Row Space

1. Let each $\mathbf{v}_1, \dots, \mathbf{v}_m$ be a **row vector**

2. Let matrix $\mathbf{A} = \begin{pmatrix} \mathbf{v}_1 \\ \vdots \\ \mathbf{v}_m \end{pmatrix}$

3. The problem is now equivalent to finding the **basis** for the **row space** of \mathbf{A} (refer to section 1.1.2.3).

4.1.4.2 Method 2: Select From Original Vectors Using Column Space

1. Let each $\mathbf{v}_1, \dots, \mathbf{v}_m$ be a **column vector**

2. Let matrix $\mathbf{A} = \begin{pmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_m \end{pmatrix}$

3. The problem is now equivalent to finding the **basis** for the **column space** of \mathbf{A} (refer to section 1.1.3.3).

4.1.5 Consistency of Linear Systems And Column Space of Coefficient Matrix

Theorem. Let \mathbf{A} be a $m \times n$ matrix. The linear system $\mathbf{A}\mathbf{x} = \mathbf{b}$ is consistent $\Leftrightarrow \mathbf{b} \in \{\mathbf{A}\mathbf{v} \mid \mathbf{v} \in \mathbb{R}^n\} = \text{column space of } \mathbf{A}$.

4.2 Rank

4.2.1 The Dimension of Row Space And Column Space Are The Same

Theorem. Let \mathbf{A} be a matrix. Then

$$\dim(\text{row space of } \mathbf{A}) = \dim(\text{column space of } \mathbf{A})$$

4.2.2 Definition of Rank

Definition. Let \mathbf{A} be a matrix. The **rank** of \mathbf{A} , denoted by $\text{rank}(\mathbf{A})$ is

$$\text{rank}(\mathbf{A}) = \dim(\text{row space of } \mathbf{A}) = \dim(\text{column space of } \mathbf{A})$$

4.2.3 Properties of Rank

Theorem. Let \mathbf{A} be a $m \times n$ matrix. Then

- $\text{rank}(\mathbf{A}) = \text{rank}(\mathbf{A}^T)$

- $\text{rank}(\mathbf{A}) = 0 \Leftrightarrow \mathbf{A} = \mathbf{0}$
- $\text{rank}(\mathbf{A}) \leq m$ and $\text{rank}(\mathbf{A}) \leq n$
 - $\text{rank}(\mathbf{A}) \leq \min\{m, n\}$
 - \mathbf{A} is **full rank** if $\text{rank}(\mathbf{A}) = \min\{m, n\}$
- A square matrix \mathbf{A} is **full rank** $\Leftrightarrow \mathbf{A}$ is invertible.

4.2.4 Rank & Consistency of Linear System

Theorem. Let $\mathbf{Ax} = \mathbf{b}$ be a linear system. Let $\left(\mathbf{R} \mid \mathbf{b}' \right)$ be the row-echelon form of $\left(\mathbf{A} \mid \mathbf{b} \right)$. Then

$$\begin{aligned} \mathbf{Ax} = \mathbf{b} \text{ is consistent} &\Leftrightarrow \text{rank}(\mathbf{A}) = \text{rank} \left(\mathbf{A} \mid \mathbf{b} \right) \\ &\Leftrightarrow \text{rank}(\mathbf{R}) = \text{rank} \left(\mathbf{R} \mid \mathbf{b}' \right) \end{aligned}$$

4.2.4.1 Remark

In general, $\text{rank}(\mathbf{A}) \leq \text{rank} \left(\mathbf{A} \mid \mathbf{b} \right) \leq \text{rank}(\mathbf{A}) + 1$

4.2.5 Row/Column Spaces and Matrix Multiplication

Theorem. Let \mathbf{A} be a $m \times n$ matrix and \mathbf{B} be a $n \times p$ matrix. Then

- column space of $\mathbf{AB} \subseteq$ column space of \mathbf{A} ;
- row space of $\mathbf{AB} \subseteq$ row space of \mathbf{B} .

4.2.5.1 Ranks and Matrix Multiplication

In particular,

- $\text{rank}(\mathbf{AB}) \leq \text{rank}(\mathbf{A})$;
- $\text{rank}(\mathbf{AB}) \leq \text{rank}(\mathbf{B})$.

That is, $\text{rank}(\mathbf{AB}) \leq \min\{\text{rank}(\mathbf{A}), \text{rank}(\mathbf{B})\}$.

- \mathbf{A} is invertible $\implies \text{rank}(\mathbf{AB}) = \text{rank}(\mathbf{B})$
- \mathbf{B} is invertible $\implies \text{rank}(\mathbf{AB}) = \text{rank}(\mathbf{A})$

4.3 Nullspaces and Nullities

4.3.1 Definition of Nullspace

Definition. Let \mathbf{A} be a $m \times n$ matrix. The **nullspace** of \mathbf{A} is the **solution space** of $\mathbf{Ax} = \mathbf{0}$, which is

$$\{\mathbf{v} \in \mathbb{R}^n \mid \mathbf{Av} = \mathbf{0}\}.$$

4.3.2 Definition of Nullity

Definition. The **dimension** of the **nullspace** of a **matrix** \mathbf{A} is the **nullity** of \mathbf{A} , denoted by $\text{nullity}(\mathbf{A})$.

4.3.3 Finding Nullity of a Matrix

Let \mathbf{R} be a **row-echelon form** of \mathbf{A} . Then

- $\mathbf{Ax} = \mathbf{0} \Leftrightarrow \mathbf{Rx} = \mathbf{0}$
- **nullspace** of $\mathbf{A} = \text{nullspace of } \mathbf{R}$
- $\text{nullity}(\mathbf{A}) = \text{nullity}(\mathbf{R}) = \text{no. of non-pivot columns of } \mathbf{R}$.

4.3.4 Dimension Theorem

Theorem. Let \mathbf{A} be a $m \times n$ matrix. Then

$$\text{rank}(\mathbf{A}) + \text{nullity}(\mathbf{A}) = n$$

4.3.5 Solution to Inhomogeneous Linear System and Nullspace

Theorem. Suppose $\mathbf{Ax} = \mathbf{b}$ has a **solution** \mathbf{v} . Let $W = \text{nullspace of } \mathbf{A}$. Then the **solution set** of $\mathbf{Ax} = \mathbf{b}$ is

$$\mathbf{v} + W = \{\mathbf{v} + \mathbf{w} \mid \mathbf{w} \in W\}$$

Hence

(A general solution of $\mathbf{Ax} = \mathbf{b}$) = (A particular solution of $\mathbf{Ax} = \mathbf{b}$) + (A general solution of $\mathbf{Ax} = \mathbf{0}$)

4.3.5.1 Condition for Linear System to have Unique Solution

Theorem. Suppose that a **linear system** $\mathbf{Ax} = \mathbf{b}$ is **consistent**. Then

$\mathbf{Ax} = \mathbf{b}$ has a **unique solution** $\Leftrightarrow \mathbf{Ax} = \mathbf{0}$ has only the **trivial solution**

$$\Leftrightarrow \text{nullspace of } \mathbf{A} \text{ is } \{\mathbf{0}\}$$

$$\Leftrightarrow \text{nullity}(\mathbf{A}) = 0$$

$$\Leftrightarrow \text{rank}(\mathbf{A}) = \text{no. of columns of } \mathbf{A}$$

4.3.5.2 Condition for the Solution Set of a Linear System to be a Vector Space

Theorem. The solution set of a linear system $A\mathbf{x} = \mathbf{b}$ is a vector space $\Leftrightarrow \mathbf{b} = \mathbf{0}$.

Chapter 5

Orthogonality

5.1 The Dot Product

5.1.1 Pythagoras' Theorem

Theorem. In a right-angled triangle, let c be the length of the **hypotenuse**, and a and b be the lengths of the other two sides. Then

$$a^2 + b^2 = c^2$$

5.1.2 Cosine Rule

Theorem. Given any triangle with sides of length a , b , and c . Let θ be the angle contained between the sides of lengths a and b . Then

$$c^2 = a^2 + b^2 - 2ab \cos \theta$$

5.1.3 Definitions

Definition. Let $\mathbf{u} = (u_1, \dots, u_n)$, $\mathbf{v} = (v_1, \dots, v_n) \in \mathbb{R}^n$.

5.1.3.1 Dot Product (Inner Product)

The **dot product** (**inner product**) of \mathbf{u} and \mathbf{v} is

$$\mathbf{u} \cdot \mathbf{v} = u_1 v_1 + \dots + u_n v_n$$

5.1.3.2 Norm (Length)

The **norm** (**length**) of \mathbf{v} is

$$\|\mathbf{v}\| = \sqrt{v_1^2 + \dots + v_n^2}$$

5.1.3.3 Unit Vector

Definition. \mathbf{v} is a **unit vector** if $\|\mathbf{v}\| = 1$.

5.1.3.4 Distance

The **distance** between \mathbf{u} and \mathbf{v} is

$$d(\mathbf{u}, \mathbf{v}) = \|\mathbf{u} - \mathbf{v}\| = \sqrt{\sum_{i=1}^n (u_i - v_i)^2}$$

5.1.3.5 Angle

The **angle** between \mathbf{u} and \mathbf{v} ($\mathbf{u} \neq \mathbf{0}$ and $\mathbf{v} \neq \mathbf{0}$) is

$$\theta = \cos^{-1} \left(\frac{\mathbf{u} \cdot \mathbf{v}}{\|\mathbf{u}\| \|\mathbf{v}\|} \right), \quad 0 \leq \theta \leq \pi$$

5.1.4 Dot Product and Matrix Multiplication

5.1.4.1 Dot Product As Matrix Multiplication

Theorem. Let $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$.

- If \mathbf{u} and \mathbf{v} are viewed as **row vectors**, then $\mathbf{u} \cdot \mathbf{v} = \mathbf{u}\mathbf{v}^T$.
- If \mathbf{u} and \mathbf{v} are viewed as **column vectors**, then $\mathbf{u} \cdot \mathbf{v} = \mathbf{u}^T \mathbf{v}$.

5.1.4.2 Matrix Multiplication As Dot Products

Theorem. Let \mathbf{A} be a $m \times n$ **matrix** and \mathbf{B} be a $n \times p$ **matrix**. Let the **row vector** $\mathbf{a}_i \in \mathbb{R}^n$ be the i th row of \mathbf{A} and the **column vector** $\mathbf{b}_j \in \mathbb{R}^n$ be the j th column of \mathbf{B} . Then

$$(i, j)\text{-entry of } \mathbf{AB} = \mathbf{a}_i \mathbf{b}_j = \mathbf{a}_i \cdot \mathbf{b}_j$$

5.1.5 Properties of Dot Product and Norm

Theorem. Let $\mathbf{u}, \mathbf{v} = (v_1, \dots, v_n), \mathbf{w} \in \mathbb{R}^n$ and $c \in \mathbb{R}$.

5.1.5.1 Dot Product is Commutative

- $\mathbf{u} \cdot \mathbf{v} = \mathbf{v} \cdot \mathbf{u}$

5.1.5.2 Dot Product is Distributive Over Vector Addition

- $(\mathbf{u} + \mathbf{v}) \cdot \mathbf{w} = \mathbf{u} \cdot \mathbf{w} + \mathbf{v} \cdot \mathbf{w}$
- $\mathbf{w} \cdot (\mathbf{u} + \mathbf{v}) = \mathbf{w} \cdot \mathbf{u} + \mathbf{w} \cdot \mathbf{v}$

5.1.5.3 Dot Product and Scalar Multiplication

- $(c\mathbf{u}) \cdot \mathbf{v} = \mathbf{u} \cdot (c\mathbf{v}) = c(\mathbf{u} \cdot \mathbf{v})$

5.1.5.4 Absolute Homogeneity of Norm

- $\|c\mathbf{v}\| = |c|\|\mathbf{v}\|$

5.1.5.5 Dot Product of a Vector with Itself

- $\mathbf{v} \cdot \mathbf{v} \geq 0$
- $\mathbf{v} \cdot \mathbf{v} = 0 \Leftrightarrow \mathbf{v} = \mathbf{0}$

5.1.5.6 Norm and Dot Product

- $\|\mathbf{v}\| = \sqrt{v_1^2 + \cdots + v_n^2} = \sqrt{\mathbf{v} \cdot \mathbf{v}}$

5.1.5.7 Nonnegativity of Norm and When Norm is Zero

- $\|\mathbf{v}\| \geq 0$
- $\|\mathbf{v}\| = 0 \Leftrightarrow \mathbf{v} = \mathbf{0}$

5.1.6 Inequalities of Norm and Distance

Theorem. Let $\mathbf{u}, \mathbf{v}, \mathbf{w} \in \mathbb{R}^n$

5.1.6.1 Cauchy-Schwarz Inequality

- $|\mathbf{u} \cdot \mathbf{v}| \leq \|\mathbf{u}\|\|\mathbf{v}\|$

5.1.6.2 Triangle Inequality (Norm Version)

- $\|\mathbf{u} + \mathbf{v}\| \leq \|\mathbf{u}\| + \|\mathbf{v}\|$

5.1.6.3 Triangle Inequality (Distance Version)

- $d(\mathbf{u}, \mathbf{w}) \leq d(\mathbf{u}, \mathbf{v}) + d(\mathbf{v}, \mathbf{w})$

5.1.7 Results for Multiplying a Matrix with its Transpose

5.1.7.1 When Multiplying a Matrix with its Transpose Gives the Zero Matrix

Theorem. Let \mathbf{A} be a $m \times n$ matrix.

$$\mathbf{A}\mathbf{A}^T = \mathbf{0}_{m \times m} \implies \mathbf{A} = \mathbf{0}_{m \times n}$$

5.1.7.2 When the Trace of Multiplying a Matrix with its Transpose is Zero

Theorem. Let \mathbf{A} be a $m \times n$ matrix. Then

$$\text{tr}(\mathbf{A}\mathbf{A}^T) = 0 \Leftrightarrow \mathbf{A} = \mathbf{0}_{m \times n}$$

5.2 Orthogonality, Orthogonal and Orthonormal Sets

5.2.1 Definition of Orthogonal for Two Vectors

Definition. Let $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$. They are **orthogonal**, denoted by $\mathbf{u} \perp \mathbf{v}$, if $\mathbf{u} \cdot \mathbf{v} = 0$.

5.2.1.1 Zero Vector is Orthogonal to Every Vector in \mathbb{R}^n

Theorem. Let $\mathbf{0} \in \mathbb{R}^n$. Then $\forall \mathbf{v} \in \mathbb{R}^n \quad \mathbf{0} \perp \mathbf{v}$

5.2.2 Definition of Orthogonal / Orthonormal Sets

5.2.2.1 Definition of Orthogonal for a Subset of \mathbb{R}^n

Definition. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\} \subseteq \mathbb{R}^n$. S is **orthogonal** if $\mathbf{v}_i \perp \mathbf{v}_j$ for all $i \neq j$, that is

$$\mathbf{v}_i \cdot \mathbf{v}_j = 0 \quad \text{for all } i \neq j$$

5.2.2.2 Definition of Orthonormal for a Subset of \mathbb{R}^n

Definition. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\} \subseteq \mathbb{R}^n$. S is **orthonormal** if S is **orthogonal** and $\forall \mathbf{v}_i \in S \quad \|\mathbf{v}_i\| = 1$, that is

$$\mathbf{v}_i \cdot \mathbf{v}_j = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases}$$

5.2.3 Properties of Orthogonal / Orthonormal Sets

- S is **orthonormal** $\implies S$ is **orthogonal**.
- S is **orthogonal** \implies a subset of S is **orthogonal**.
- S is **orthonormal** \implies a subset of S is **orthonormal**.
- S is **orthogonal** $\implies S \cup \{\mathbf{0}\}$ is **orthogonal**.
- S is **orthonormal** $\implies \mathbf{0} \notin S$.

5.2.4 Normalizing: Converting an Orthogonal Set to an Orthonormal Set

The following process of converting an **orthogonal** set of nonzero vectors to an **orthonormal** set of vectors, is called **normalizing**:

1. Let $S = \{\mathbf{u}_1, \dots, \mathbf{u}_k\} \subseteq \mathbb{R}^n$ be an **orthogonal** set of **nonzero vectors**
2. For all $\mathbf{u}_i \in S$, set $\mathbf{v}_i = \frac{\mathbf{u}_i}{\|\mathbf{u}_i\|}$.
3. Then $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ is an **orthonormal** set.

5.2.5 Using Matrix Multiplication to Check for Orthogonal / Orthonormal Set

Theorem. To check whether a set $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\} \subseteq \mathbb{R}^n$ is **orthogonal** / **orthonormal**, let $\mathbf{A} = \begin{pmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_k \end{pmatrix}$. Then

- S is **orthogonal** $\Leftrightarrow \mathbf{v}_i \cdot \mathbf{v}_j = 0$ for all $i \neq j \Leftrightarrow \mathbf{A}^T \mathbf{A}$ is **diagonal**
- S is **orthonormal** $\Leftrightarrow \mathbf{v}_i \cdot \mathbf{v}_j = \begin{cases} 0 & \text{if } i \neq j \\ 1 & \text{if } i = j \end{cases} \Leftrightarrow \mathbf{A}^T \mathbf{A} = \mathbf{I}_k$.

5.2.5.1 Standard Basis is An Orthonormal Set

Theorem. The standard basis $E = \{\mathbf{e}_1, \dots, \mathbf{e}_n\} \subseteq \mathbb{R}^n$ is an **orthonormal** set.

5.2.5.2 An Orthonormal Subset of \mathbb{R}^n with n Vectors is a Basis for \mathbb{R}^n

Theorem. Let $\{\mathbf{u}_1, \dots, \mathbf{u}_n\} \subseteq \mathbb{R}^n$. Let $\mathbf{A} = \begin{pmatrix} \mathbf{u}_1 & \cdots & \mathbf{u}_n \end{pmatrix}$. Then

$$\begin{aligned} \mathbf{A}^T \mathbf{A} = \mathbf{I}_n &\implies \mathbf{A} \text{ is invertible} \\ &\implies \{\mathbf{u}_1, \dots, \mathbf{u}_n\} \text{ is a } \mathbf{basis} \text{ for } \mathbb{R}^n \quad (\text{refer to section 3.7.6}) \end{aligned}$$

5.2.6 Orthogonal Nonzero Sets are Linearly Independent

Theorem. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be an **orthogonal** set of **nonzero** vectors in \mathbb{R}^n . Then S is **linearly independent**.

5.2.6.1 Orthonormal Sets are Linearly Independent

Corollary. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be an **orthonormal** set in \mathbb{R}^n . Then S is **linearly independent**.

5.3 Orthogonal and Orthonormal Bases

5.3.1 Orthogonal / Orthonormal Bases

Definition. Let S be a **basis** for a **vector space**.

- S is an **orthogonal basis** if it is **orthogonal**.
- S is an **orthonormal basis** if it is **orthonormal**.

5.3.2 Criterion for Orthogonal / Orthonormal Set to be Basis

Theorem. Suppose S is a subset of **vector space** V , where S is an **orthogonal/orthonormal set** and $\mathbf{0} \notin S$, then

$|S| = \dim(V)$ or $\text{span}(S) = V \implies S$ is an **orthogonal/orthonormal** (respectively) **basis** for V

5.3.3 Simpler Formula for Coordinate Vectors Relative to Orthogonal / Orthonormal Bases

5.3.3.1 Orthogonal Bases

Theorem. Let $S = \{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ be an **orthogonal basis** for a **vector space** V . Then

$$\forall \mathbf{w} \in V \quad (\mathbf{w})_S = \left(\frac{\mathbf{w} \cdot \mathbf{u}_1}{\mathbf{u}_1 \cdot \mathbf{u}_1}, \dots, \frac{\mathbf{w} \cdot \mathbf{u}_k}{\mathbf{u}_k \cdot \mathbf{u}_k} \right)$$

5.3.3.2 Orthonormal Bases

Theorem. Let $S = \{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ be an **orthonormal basis** for a **vector space** V . Then

$$\forall \mathbf{w} \in V \quad (\mathbf{w})_S = (\mathbf{w} \cdot \mathbf{u}_1, \dots, \mathbf{w} \cdot \mathbf{u}_k)$$

5.3.4 Orthogonality between a Vector and a Subspace

Definition. Let V be a **subspace** of \mathbb{R}^n . $\mathbf{u} \in \mathbb{R}^n$ is **orthogonal** (perpendicular) to V if $\forall \mathbf{v} \in V \quad \mathbf{u} \perp \mathbf{v}$.

5.3.4.1 Example: Normal Vector of a Plane in \mathbb{R}^3

Let $V = \{(x, y, z) \mid ax + by + cz = 0\}$ be a plane in \mathbb{R}^3 containing the origin. Then $\mathbf{n} = (a, b, c)$ is **orthogonal** to V and is a **normal vector** of the plane V .

5.3.5 Easier Criterion for Vector to be Orthogonal to a Vector Space

Theorem. Let $V = \text{span}\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be a **vector space**. Then

$$\mathbf{w} \text{ is orthogonal to } V \Leftrightarrow \mathbf{w} \perp \mathbf{v}_i \text{ for all } \mathbf{v}_i \in V$$

5.3.6 The Set of All Vectors Orthogonal to a Subspace is a Subspace

Theorem. Let W be a **subspace** of \mathbb{R}^n . Then $W^\perp = \{\mathbf{v} \in \mathbb{R}^n \mid \mathbf{v} \text{ is orthogonal to } W\}$ is a **subspace** of \mathbb{R}^n .

5.3.7 Projection of a Vector Onto a Vector Space

5.3.7.1 With Orthonormal Basis

Theorem. Let $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be an **orthonormal basis** for a **vector space** V . Then the **projection** of \mathbf{w} onto V is

$$(\mathbf{w} \cdot \mathbf{v}_1)\mathbf{v}_1 + \dots + (\mathbf{w} \cdot \mathbf{v}_k)\mathbf{v}_k$$

5.3.7.2 With Orthogonal Basis

Theorem. Let $\{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ be an **orthogonal basis** for a **vector space** V . Then the **projection** of \mathbf{w} onto V is

$$\left(\frac{\mathbf{w} \cdot \mathbf{u}_1}{\mathbf{u}_1 \cdot \mathbf{u}_1} \right) \mathbf{u}_1 + \dots + \left(\frac{\mathbf{w} \cdot \mathbf{u}_k}{\mathbf{u}_k \cdot \mathbf{u}_k} \right) \mathbf{u}_k$$

Remarks. It is the sum of projections of \mathbf{w} onto $\mathbf{u}_1, \dots, \mathbf{u}_k$.

5.3.8 Gram-Schmidt Process: Generating Orthogonal / Orthonormal Basis

Let $\{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ be a **basis** for a **vector space** V . Define

$$\mathbf{v}_1 = \mathbf{u}_1$$

$$\mathbf{v}_2 = \mathbf{u}_2 - \frac{\mathbf{u}_2 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1$$

$$\mathbf{v}_3 = \mathbf{u}_3 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{u}_3 \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2$$

$$\vdots \quad \vdots$$

$$\mathbf{v}_k = \mathbf{u}_k - \frac{\mathbf{u}_k \cdot \mathbf{v}_1}{\mathbf{v}_1 \cdot \mathbf{v}_1} \mathbf{v}_1 - \frac{\mathbf{u}_k \cdot \mathbf{v}_2}{\mathbf{v}_2 \cdot \mathbf{v}_2} \mathbf{v}_2 - \dots - \frac{\mathbf{u}_k \cdot \mathbf{v}_{k-1}}{\mathbf{v}_{k-1} \cdot \mathbf{v}_{k-1}} \mathbf{v}_{k-1}$$

- Then $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ is an **orthogonal basis** for V .
- $\{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ can be **normalized** to obtain an **orthonormal basis** for V (refer to section 5.2.4).

5.3.9 Solving a Linear System whose Coefficient Matrix's Columns are Linearly Independent

5.3.9.1 Decomposition

Theorem. Let \mathbf{A} be a $m \times n$ matrix whose columns are **linearly independent**. Then there exists

- A $m \times n$ **matrix** \mathbf{Q} whose columns form an **orthonormal set**, and

- An invertible $n \times n$ upper triangular matrix R

such that $A = QR$.

5.3.9.2 Algorithm

To solve a linear system $Ax = b$ where the columns of A are linearly independent.

1. $(QR)x = b$
2. $Rx = IRx = Q^T QRx = Q^T b$
3. Solve x by back-substitution.

5.3.9.3 Remark

There exists a R such that the diagonal entries are all positive.

5.4 Best Approximations

5.4.1 The Projection of a Vector on a Subspace is its Best Approximation in that Subspace

Theorem. Let V be a subspace of \mathbb{R}^n . $\forall u \in \mathbb{R}^n$, let p be the projection of u onto V . Then

- p is the best approximation of u in V , i.e.
 - $\forall v \in V \quad d(u, p) \leq d(u, v)$
- Moreover, p is the only best approximation of u in V .
 - $d(u, p) = d(u, v) \Leftrightarrow v = p$

5.4.2 Least Squares Solution

5.4.2.1 Definition of Least Squares Solution

Definition. Let A be a $m \times n$ matrix and $b \in \mathbb{R}^m$. $u \in \mathbb{R}^n$ is a least squares solution to the linear system $Ax = b$ if

$$\forall v \in \mathbb{R}^n \quad \|b - Au\| \leq \|b - Av\|$$

5.4.2.2 Least Squares Solution and Projection

Theorem. Let A be a $m \times n$ matrix and $b \in \mathbb{R}^m$. Let p be the projection of b onto the column space of A . Then

- $\forall v \in \mathbb{R}^n \quad \|b - p\| \leq \|b - Av\|$
- u is a least squares solution to $Ax = b \Leftrightarrow u$ is a solution to $Ax = p$.

5.4.3 Methodology: Finding Least Squares Solution

5.4.3.1 Tedious Method

To find a **least squares solution** to $\mathbf{Ax} = \mathbf{b}$, proceed as follows:

1. Find an **orthogonal (orthonormal) basis** for $V = \text{column space of } \mathbf{A}$.
2. Find the **projection** \mathbf{p} of \mathbf{b} onto V .
3. Solve the **linear system** $\mathbf{Ax} = \mathbf{p}$.
 - A solution to $\mathbf{Ax} = \mathbf{p}$ is a **least squares solution** to $\mathbf{Ax} = \mathbf{b}$.

Remarks.

- The **linear system** $\mathbf{Ax} = \mathbf{p}$ is always **consistent** since \mathbf{p} lies in the **column space** of \mathbf{A} .
- If $\mathbf{Ax} = \mathbf{b}$ is already **consistent**, then
 - $\mathbf{b} = \mathbf{p} \in V$
 - (solution to $\mathbf{Ax} = \mathbf{b}$) = (least squares solution to $\mathbf{Ax} = \mathbf{b}$).

5.4.3.2 Easy Method

Theorem. \mathbf{u} is a least squares solution to $\mathbf{Ax} = \mathbf{b} \Leftrightarrow \mathbf{u}$ is a solution to $\mathbf{A}^T \mathbf{Ax} = \mathbf{A}^T \mathbf{b}$.

5.5 Orthogonal Matrices

5.5.1 Definition of Orthogonal Matrices

Definition. Let \mathbf{A} be a square matrix. Then

$$\begin{aligned}
 \mathbf{A} \text{ is an orthogonal matrix} &\Leftrightarrow \mathbf{A}^T \mathbf{A} = \mathbf{I} \\
 &\Leftrightarrow \mathbf{A}^{-1} = \mathbf{A}^T \\
 &\Leftrightarrow \mathbf{A} \mathbf{A}^T = \mathbf{I}
 \end{aligned}$$

5.5.1.1 Example: Identity Matrix is an Orthogonal Matrix

Theorem. The identity matrix \mathbf{I}_n is an orthogonal matrix.

5.5.2 Properties of Orthogonal Matrices

5.5.2.1 Rows and Columns of Orthogonal Matrix Form an Orthonormal Basis for \mathbb{R}^n

Theorem. Let \mathbf{A} be a square matrix of order n . Then

$$\mathbf{A} \text{ is an orthogonal matrix} \Leftrightarrow \text{columns of } \mathbf{A} \text{ form an orthonormal basis for } \mathbb{R}^n$$

\Leftrightarrow rows of \mathbf{A} form an **orthonormal basis** for \mathbb{R}^n

5.5.2.2 The Transpose or Inverse of an Orthogonal Matrix is Orthogonal

Theorem. If \mathbf{A} is an **orthogonal matrix**, then $\mathbf{A}^T = \mathbf{A}^{-1}$ is an **orthogonal matrix**.

5.5.2.3 The Product of Two Orthogonal Matrices is Orthogonal

Theorem. If \mathbf{A} and \mathbf{B} are **orthogonal matrices** of the same size, then \mathbf{AB} is an **orthogonal matrix**.

5.5.2.4 Condition for Rows/Columns of a Matrix to Form an Orthonormal Set

5.5.2.4.1 Condition for Columns of a Matrix to Form an Orthonormal Set

Refer to section 5.2.5.

5.5.2.4.2 Condition for Rows of a Matrix to Form an Orthonormal Set

Theorem. For any $m \times n$ matrix \mathbf{A} , $\mathbf{AA}^T = \mathbf{I}_m \Leftrightarrow$ the rows of \mathbf{A} form an **orthonormal set**.

5.5.2.5 Pre-Multiplication of an Orthogonal Matrix to an Orthonormal Set

Theorem. Let $S = \{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ be an **orthonormal subset** of \mathbb{R}^n and \mathbf{P} be a $n \times n$ **orthogonal matrix**. Then $\{\mathbf{Pu}_1, \dots, \mathbf{Pu}_k\}$ is an **orthonormal set**.

5.5.3 Transition Matrices Between Orthonormal Bases

5.5.3.1 Formula for Transition Matrices Between Orthonormal Bases

Theorem. Let $S = \{\mathbf{u}_1, \dots, \mathbf{u}_k\}$ and $T = \{\mathbf{v}_1, \dots, \mathbf{v}_k\}$ be **orthonormal bases** for a vector space V . Let $\mathbf{A} = \begin{pmatrix} \mathbf{u}_1 & \cdots & \mathbf{u}_k \end{pmatrix}$ and $\mathbf{B} = \begin{pmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_k \end{pmatrix}$. Then

- $\mathbf{P} = \mathbf{B}^T \mathbf{A}$ is the **transition matrix** from S to T .
- $\mathbf{Q} = \mathbf{A}^T \mathbf{B}$ is the **transition matrix** from T to S .

5.5.3.2 Transition Matrices Between Orthonormal Bases are Orthogonal

Theorem. Let S and T be **orthonormal bases** for a vector space V and \mathbf{P} be the **transition matrix** from S to T . Then \mathbf{P} is an **orthogonal matrix**.

5.5.4 Classification of Orthogonal Matrices

5.5.4.1 Table of Orthogonal Matrices By Order

Theorem. The following table lists all the **orthogonal matrices** in increasing **order**:

Order	Determinant	Formulae
1	1	$\begin{pmatrix} 1 \end{pmatrix}$
	-1	$\begin{pmatrix} -1 \end{pmatrix}$
2	1	$\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$
	-1	$\begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix}$

5.5.5 Geometric Representation of Orthogonal Matrix

5.5.5.1 Rotation Matrix

- Let $\mathbf{u}_1 = \begin{pmatrix} \cos \theta \\ \sin \theta \end{pmatrix}, \mathbf{u}_2 = \begin{pmatrix} -\sin \theta \\ \cos \theta \end{pmatrix}$.
 - Then $\{\mathbf{u}_1, \mathbf{u}_2\}$ is an **orthonormal basis** for \mathbb{R}^2 .
- Let $\mathbf{P}_\theta = \begin{pmatrix} \mathbf{u}_1 & \mathbf{u}_2 \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$. Then
 - \mathbf{P}_θ is **orthogonal**, and
 - \mathbf{P}_θ is the **transition matrix** from $\{\mathbf{u}_1, \mathbf{u}_2\}$ to $\{\mathbf{e}_1, \mathbf{e}_2\}$.
- $\forall \mathbf{u} \in \mathbb{R}^2$ $\mathbf{P}_\theta \mathbf{u}$ is the **rotation** of \mathbf{u} about the **origin** by θ anticlockwise.

5.5.5.2 Composition of Rotation Matrices

- Moreover, $\mathbf{P}_\beta \mathbf{P}_\alpha = \mathbf{P}_{\alpha+\beta}$

5.5.5.3 Deriving Sum Laws for Sine and Cosine

- $\cos(\alpha + \beta) = \cos \alpha \cos \beta - \sin \alpha \sin \beta$
- $\sin(\alpha + \beta) = \sin \alpha \cos \beta + \cos \alpha \sin \beta$

Chapter 6

Diagonalization

6.1 Eigenvalues and Eigenvectors

6.1.1 Definition of Eigenvalues and Eigenvectors

Definition. Let A be a square matrix of order n . Suppose that for some $\lambda \in \mathbb{R}$ and **nonzero** $v \in \mathbb{R}^n$, $Av = \lambda v$. Then

- λ is an **eigenvalue** of A .
- v is an **eigenvector** of A associated with λ .

6.1.2 Characteristic Polynomial

6.1.2.1 Definition of Characteristic Polynomial

Definition. Let A be a square matrix. $\det(\lambda I - A)$ is the **characteristic polynomial** of A .

6.1.2.2 Characteristic Polynomial is a Monic Polynomial of Degree n

Theorem. Let A be a square matrix. Then the **characteristic polynomial** of A is a **monic polynomial** in λ of **degree n** :

$$\lambda^n + c_{n-1}\lambda^{n-1} + \cdots + c_1\lambda + c_0$$

6.1.3 Definition of Characteristic Equation

Definition. Let A be a square matrix. $\det(\lambda I - A) = 0$ is the **characteristic equation** of A .

6.1.4 Eigenvalues are the Roots to the Characteristic Equation

Theorem. Let A be a square matrix. Then the **eigenvalues** of A are all the **roots** to the **characteristic equation** of A .

6.1.5 Eigenvalue and Invertibility

Theorem. Let \mathbf{A} be a square matrix of order n . Then

$$0 \text{ is not an eigenvalue of } \mathbf{A} \Leftrightarrow \det(\mathbf{A}) \neq 0 \Leftrightarrow \mathbf{A} \text{ is invertible}$$

6.1.6 Eigenvalues of Triangular Matrices are its Diagonal Entries

Theorem. Let \mathbf{A} be a triangular matrix. Then its eigenvalues are all the diagonal entries of \mathbf{A} .

6.1.7 Eigenspace

6.1.7.1 Definition of Eigenspace

Definition. Let \mathbf{A} be a square matrix and λ an eigenvalue of \mathbf{A} . Then the eigenspace of \mathbf{A} associated to λ , denoted by E_λ (or $E_{\mathbf{A},\lambda}$), is the nullspace of $\lambda\mathbf{I} - \mathbf{A}$.

6.1.7.2 Properties of Eigenspace

6.1.7.2.1 Vectors in Eigenspace

Theorem. $E_{\mathbf{A},\lambda}$ consists of all the eigenvectors of \mathbf{A} associated to λ and the zero vector $\mathbf{0}$.

6.1.7.2.2 Dimension of Eigenspace

Theorem. Since $\lambda\mathbf{I} - \mathbf{A}$ is singular, $\dim(E_\lambda) \geq 1$.

6.1.7.2.3 Eigenspace Associated To Eigenvalue Zero

Theorem. If \mathbf{A} is singular, then $E_0 = \text{nullspace of } \mathbf{A}$.

6.2 Diagonalization

6.2.1 Definition of Diagonalizable Matrix

Definition. Let \mathbf{A} be a square matrix. \mathbf{A} is diagonalizable if \exists an invertible matrix \mathbf{P} such that $\mathbf{P}^{-1}\mathbf{A}\mathbf{P}$ is a diagonal matrix.

6.2.2 Criterion for Diagonalizability

Theorem. Let \mathbf{A} be a square matrix of order n . Then

$$\mathbf{A} \text{ is diagonalizable} \Leftrightarrow \mathbf{A} \text{ has } n \text{ linearly independent eigenvectors}$$

6.2.3 Eigenvectors associated to Distinct Eigenvalues are Linearly Independent

Theorem. Let $\lambda_1, \dots, \lambda_k$ be distinct **eigenvalues** of \mathbf{A} and \mathbf{v}_i be an **eigenvector** of \mathbf{A} associated to λ_i . Then $\mathbf{v}_1, \dots, \mathbf{v}_k$ are **linearly independent**.

6.2.4 Algorithm of Diagonalization

Let \mathbf{A} be a **square matrix** of **order** n .

1. Find the **characteristic polynomial** $\det(\lambda \mathbf{I} - \mathbf{A})$.
2. **Factorise** $\det(\lambda \mathbf{I} - \mathbf{A})$ over \mathbb{R} (not \mathbb{C}) to find **eigenvalues** of \mathbf{A} .
 - If $\det(\lambda \mathbf{I} - \mathbf{A})$ cannot be completely factorised, then \mathbf{A} is not **diagonalizable**.
 - If $\det(\lambda \mathbf{I} - \mathbf{A})$ can be completely factorised, say $\det(\lambda \mathbf{I} - \mathbf{A}) = \prod_{i=1}^k (\lambda - \lambda_i)^{r_i}$ where $\lambda_1, \dots, \lambda_k$ are all distinct. Then
 - **algebraic multiplicity** of λ_i , denoted by $a(\lambda_i)$, is r_i
 - **geometric multiplicity** of λ_i , denoted by $g(\lambda_i)$, is $\dim(E_{\lambda_i})$
 - Moreover, $1 \leq g(\lambda_i) \leq a(\lambda_i)$, and $\sum_{i=1}^k a(\lambda_i) = n$.
3. For each **eigenvalue** λ_i of \mathbf{A} , find a **basis** S_i for the **eigenspace** E_{λ_i} .
 - $\exists i \quad g(\lambda_i) < a(\lambda_i) \Rightarrow \sum_{i=1}^k \dim(E_{\lambda_i}) < n \Rightarrow \mathbf{A}$ is not **diagonalizable**.
4. $\forall i \quad g(\lambda_i) = a(\lambda_i) \Rightarrow \sum_{i=1}^k \dim(E_{\lambda_i}) = n \Rightarrow \mathbf{A}$ is **diagonalizable**, then
 - $\bigcup_{i=1}^k S_i = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a **basis** for \mathbb{R}^n .
 - $\mathbf{P} = \begin{pmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_n \end{pmatrix}$ **diagonalizes** \mathbf{A} , i.e. $\mathbf{P}^{-1} \mathbf{A} \mathbf{P} = \mathbf{D}$ is **diagonal**.
 - \mathbf{v}_i is an **eigenvector** of \mathbf{A} associated to the i th **diagonal entry** (eigenvalue) of \mathbf{D} .

6.2.5 Matrix with n Distinct Eigenvalues is Diagonalizable

Theorem. Let \mathbf{A} be a **square matrix** of **order** n . \mathbf{A} has n distinct **eigenvalues** $\Rightarrow \mathbf{A}$ is **diagonalizable**.

6.2.6 Application of Diagonalization

6.2.6.1 Matrix Exponentiation

Theorem. Suppose that \mathbf{P} diagonalizes \mathbf{A} such that $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix}$. Then for

$$\begin{cases} m \in \mathbb{Z}, m \geq 0, & \mathbf{A} \text{ is singular} \\ m \in \mathbb{Z}, & \mathbf{A} \text{ is invertible} \end{cases} \quad \mathbf{A}^m = \mathbf{P} \begin{pmatrix} \lambda_1^m & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n^m \end{pmatrix} \mathbf{P}^{-1}$$

6.2.6.2 Example: Fibonacci Numbers

- The **fibonacci numbers** are defined as $a_n = \begin{cases} n, & n = 0 \text{ or } n = 1 \\ a_{n-1} + a_{n-2}, & n \geq 2 \end{cases}$
- Note that $a_{n+1} = a_{n-1} + a_n$ for $n \geq 1$
- Let $\mathbf{x}_n = \begin{pmatrix} a_n \\ a_{n+1} \end{pmatrix} = \begin{pmatrix} a_n \\ a_{n-1} + a_n \end{pmatrix} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix} \begin{pmatrix} a_{n-1} \\ a_n \end{pmatrix}$ and $\mathbf{A} = \begin{pmatrix} 0 & 1 \\ 1 & 1 \end{pmatrix}$. Then
- $\mathbf{x}_n = \mathbf{A}\mathbf{x}_{n-1} = \mathbf{A}^2\mathbf{x}_{n-2} = \cdots = \mathbf{A}^n\mathbf{x}_0$, where $\mathbf{x}_0 = \begin{pmatrix} 0 \\ 1 \end{pmatrix}$
- To find \mathbf{A}^n for large n , it will be easier to **diagonalize** \mathbf{A} .

6.3 Orthogonal Diagonalization

6.3.1 Definition of Orthogonally Diagonalizable Matrix

Definition. A square matrix \mathbf{A} is **orthogonally diagonalizable** if it can be **diagonalized** by an **orthogonal matrix**, i.e.,

- \exists an **orthogonal matrix** \mathbf{P} such that $\mathbf{P}^T\mathbf{A}\mathbf{P} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$ is a **diagonal matrix**.
- \mathbf{P} is said to **orthogonally diagonalize** \mathbf{A} .

6.3.1.1 Remarks

- For any **eigenvalue** λ of \mathbf{A} , we can always choose an **orthonormal basis** for E_λ .
- Suppose further that \mathbf{A} is **orthogonally diagonalizable**. Then
 - for distinct **eigenvalues** $\lambda \neq \mu$, every **eigenvector** of λ is **orthogonal** to that of μ .

6.3.2 Only Symmetric Matrices are Orthogonally Diagonalizable

Theorem. A square matrix is orthogonally diagonalizable \Leftrightarrow it is a symmetric matrix.

6.3.3 Algorithm of Orthogonal Diagonalization

Let \mathbf{A} be a symmetric matrix of order n .

1. Find the **characteristic polynomial** $\det(\lambda \mathbf{I} - \mathbf{A})$.
2. **Factorise** $\det(\lambda \mathbf{I} - \mathbf{A})$ over \mathbb{R} to find **eigenvalues** of \mathbf{A} .
 - $\det(\lambda \mathbf{I} - \mathbf{A})$ can definitely be completely factorised, say $\det(\lambda \mathbf{I} - \mathbf{A}) = \prod_{i=1}^k (\lambda - \lambda_i)^{r_i}$ where $\lambda_1, \dots, \lambda_k$ are all distinct.
 - $\sum_{i=1}^k a(\lambda_i) = n$.
3. For each **eigenvalue** λ_i of \mathbf{A} , find an **orthonormal basis** T_{λ_i} for E_{λ_i} .
 - (a) Find a basis S_{λ_i} for E_{λ_i} .
 - $\forall i \quad 1 \leq g(\lambda_i) = a(\lambda_i) \Rightarrow \sum_{i=1}^k \dim(E_{\lambda_i}) = n$
 - (b) Use Gram-Schmidt process to transfer S_{λ_i} to an **orthonormal basis** T_{λ_i} for E_{λ_i} .
4. $\forall i \quad g(\lambda_i) = a(\lambda_i) \Rightarrow \sum_{i=1}^k \dim(E_{\lambda_i}) = n \Rightarrow \mathbf{A}$ is **diagonalizable**, then
5. $\bigcup_{i=1}^k T_{\lambda_i} = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is an **orthonormal basis** for \mathbb{R}^n .
 - $\mathbf{P} = \begin{pmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_n \end{pmatrix}$ orthogonally diagonalizes \mathbf{A}

6.4 Quadratic Forms and Conic Sections

6.4.1 Quadratic Forms

6.4.1.1 Definition of Quadratic Form

Definition. A quadratic form/homogeneous polynomial in degree 2 in n variables x_1, \dots, x_n is

$$Q(x_1, \dots, x_n) = \sum_{i=1}^n q_{ii}x_i^2 + \sum_{i < j} q_{ij}x_i x_j$$

6.4.1.2 Matrix Representation of Quadratic Forms

Theorem. Let $Q(x_1, \dots, x_n) = \sum_{i=1}^n q_{ii}x_i^2 + \sum_{i < j} q_{ij}x_i x_j$ be a **quadratic form**. Let $\mathbf{x} = (x_1, \dots, x_n)^T$ and $\mathbf{A} = (a_{ij})_{n \times n}$ where $a_{ii} = q_{ii}$ and $a_{ij} = a_{ji} = \frac{1}{2}q_{ij}$ for $i < j$. Then $Q(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$, $\mathbf{x} \in \mathbb{R}^n$.

6.4.1.3 Simplification of Quadratic Forms

Theorem. Suppose $Q(\mathbf{x}) = \mathbf{x}^T \mathbf{A} \mathbf{x}$ is a **quadratic form** where $\mathbf{x} = (x_1, \dots, x_n)^T \in \mathbb{R}^n$ and \mathbf{A} is a **symmetrix matrix** of order n . Then

1. \exists an **orthogonal matrix** \mathbf{P} such that $\mathbf{P}^T \mathbf{A} \mathbf{P} = \begin{pmatrix} \lambda_1 & \cdots & 0 \\ \vdots & \ddots & \vdots \\ 0 & \cdots & \lambda_n \end{pmatrix}$
2. Let $\mathbf{y} = \mathbf{P}^T \mathbf{x} = (y_1, \dots, y_n)^T \in \mathbb{R}^n$. Then
3. $Q(\mathbf{x}) = \sum_{i=1}^n \lambda_i y_i^2$.

6.4.2 Quadratic Equation

6.4.2.1 Quadratic Equation in One Variable

Definition. A **quadratic equation** in variable x is of the form $ax^2 + bx = c$.

6.4.2.2 Quadratic Equation in Two Variables

Definition. A **quadratic equation** in variables x and y is

$$ax^2 + bxy + cy^2 + dx + ey = f$$

6.4.2.2.1 Matrix Representation

Let $\mathbf{x} = \begin{pmatrix} x \\ y \end{pmatrix}$, $\mathbf{A} = \begin{pmatrix} a & \frac{1}{2}b \\ \frac{1}{2}b & c \end{pmatrix}$, and $\mathbf{b} = \begin{pmatrix} d \\ e \end{pmatrix}$. Then $\mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} = f$.

6.4.2.2.2 Quadratic Form Associated With Quadratic Equation

Definition. $ax^2 + bxy + cy^2 = \mathbf{x}^T \mathbf{A} \mathbf{x}$ is the quadratic form **associated** with the quadratic equation.

6.4.2.3 Graph of a Quadratic Equation

Theorem. The graph of a **quadratic equation** is a **conic section**.

6.4.3 Classification of Conic Sections

6.4.3.1 Table of Conics

Degeneracy	Name	Equation / Standard Form
Degenerated	The whole plane \mathbb{R}^2	$0 = 0$
	Empty Set	$x^2 + y^2 = -1$
	A point	$x^2 + y^2 = 0$
	A line	$x = 0$ or $x^2 = 0$
	A pair of distinct lines	$x^2 - y^2 = 0$
Non-degenerated	Circle	$\frac{x^2}{\alpha^2} + \frac{y^2}{\alpha^2} = 1, \quad \alpha > 0$
	Ellipse	$\frac{x^2}{\alpha^2} + \frac{y^2}{\beta^2} = 1, \quad \alpha > 0, \beta > 0$
	Hyperbola	$\pm \frac{x^2}{\alpha^2} \mp \frac{y^2}{\beta^2} = 1, \quad \alpha > 0, \beta > 0$
	Parabola	$x^2 = \alpha y$ or $y^2 = \alpha x$

6.4.3.2 Algorithm to Classify Conic Sections

Given a quadratic equation $\mathbf{x}^T \mathbf{A} \mathbf{x} + \mathbf{b}^T \mathbf{x} = f, \mathbf{x} \in \mathbb{R}^2$.

1. Orthogonally diagonalize \mathbf{A} .

- $\mathbf{P}^T \mathbf{A} \mathbf{P} = \begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix}, \mathbf{P}$ an orthogonal matrix.

2. Let $\mathbf{y} = \mathbf{P}^T \mathbf{x}$. Then

- $\mathbf{y}^T \begin{pmatrix} \lambda & 0 \\ 0 & \mu \end{pmatrix} \mathbf{y} + \mathbf{b}^T \mathbf{P} \mathbf{y} = f$

3. Complete the squares.

6.4.3.2.1 Determinant of Symmetric Matrix and Type of Conic Sections

Suppose the conic section is **non-degenerate**. Since $\lambda\mu = \det(\mathbf{A})$, then

- $\det(\mathbf{A}) > 0 \Leftrightarrow$ ellipse (or circle).
- $\det(\mathbf{A}) < 0 \Leftrightarrow$ hyperbola.
- $\det(\mathbf{A}) = 0 \Leftrightarrow$ parabola.

6.4.3.2.2 Rotating and Reflecting Conic Sections

Let \mathbf{P} be **orthogonal** of order 2. Then $\det(\mathbf{P}) = \pm 1$.

- $\det(\mathbf{P}) = 1 \Rightarrow \mathbf{P} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$. Let $\mathbf{y} = \mathbf{P}^T \mathbf{x}$. Then

- the new axes are obtained by rotating the original axes about origin anticlockwise by θ .
- $\det(\mathbf{P}) = -1 \Rightarrow \mathbf{P} = \begin{pmatrix} \cos \theta & \sin \theta \\ \sin \theta & -\cos \theta \end{pmatrix} = \begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$. Let $\mathbf{y} = \mathbf{P}^T \mathbf{x}$.

Then

- the new axes are obtained by first rotating the original axes about origin anticlockwise by θ , then reflecting w.r.t. the x' -axis.

6.4.3.2.3 Can Always Orthogonally Diagonalize by Matrix with Determinant One

By multiplying the 2nd column of \mathbf{P} by -1 if necessary, we can always **diagonalize** a **symmetric matrix** \mathbf{A} by an **orthogonal matrix** with **determinant 1**.

Chapter 7

Linear Transformation

7.1 Linear Transformations from \mathbb{R}^n to \mathbb{R}^m

7.1.1 Definition of Linear Transformation

Definition. The mapping $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ defined by

$$T \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \right) = \begin{pmatrix} a_{11}x_1 + a_{12}x_2 + \cdots + a_{1n}x_n \\ a_{21}x_1 + a_{22}x_2 + \cdots + a_{2n}x_n \\ \vdots \\ a_{m1}x_1 + a_{m2}x_2 + \cdots + a_{mn}x_n \end{pmatrix}$$

a **linear transformation** from \mathbb{R}^n to \mathbb{R}^m .

7.1.1.1 Linear Operator

T is a **linear operator** on \mathbb{R}^n if $m = n$.

7.1.2 Linear Transformation As Matrix Form

A **linear transformation** is viewed as the **matrix form**:

$$T \left(\begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix} \right) = \begin{pmatrix} a_{11} & a_{12} & \cdots & a_{1n} \\ a_{21} & a_{22} & \cdots & a_{2n} \\ \vdots & \vdots & \ddots & \vdots \\ a_{m1} & a_{m2} & \cdots & a_{mn} \end{pmatrix} \begin{pmatrix} x_1 \\ x_2 \\ \vdots \\ x_n \end{pmatrix}$$

$T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ such that $\forall \mathbf{x} \in \mathbb{R}^n \quad T(\mathbf{x}) = \mathbf{Ax}$

- $\mathbf{A} = (a_{ij})_{m \times n}$ is the **standard matrix** for T .

7.1.3 Examples of Linear Transformation

7.1.3.1 Identity Transformation

Definition. Let $I : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be the **linear transformation** $\forall \mathbf{x} \in \mathbb{R}^n \quad I(\mathbf{x}) = \mathbf{x}$.

- It is the
 - **identity transformation**; or
 - **identity operator** on \mathbb{R}^n .
- $I(\mathbf{x}) = \mathbf{x} = \mathbf{I}_n \mathbf{x} \Rightarrow \mathbf{I}_n$ is the **standard matrix** for I .

7.1.3.2 Zero Transformation

Definition. Let $O : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be the **linear transformation** $\forall \mathbf{x} \in \mathbb{R}^n \quad O(\mathbf{x}) = \mathbf{0}$.

- It is the **zero transformation**.
- $O(\mathbf{x}) = \mathbf{0} = \mathbf{0}_{m \times n} \mathbf{x} \Rightarrow \mathbf{0}_{m \times n}$ is the **standard matrix** for O .

7.1.4 Standard Matrix of a Linear Transformation is Unique

Theorem. The **standard matrix** of a **linear transformation** is unique.

7.1.5 To Prove that a Mapping is a Linear Transformation

To show that $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation**, it suffices to find a **matrix** \mathbf{A} so that $\forall \mathbf{x} \in \mathbb{R}^n \quad T(\mathbf{x}) = \mathbf{A}\mathbf{x}$.

7.1.6 Linearity of Linear Transformations

Theorem. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation**. Then

- $T(\mathbf{0}) = \mathbf{0}$
- $\forall c \in \mathbb{R}, \forall \mathbf{v} \in \mathbb{R}^n, \quad T(c\mathbf{v}) = cT(\mathbf{v})$
- $\forall \mathbf{u}, \mathbf{v} \in \mathbb{R}^n, \quad T(\mathbf{u} + \mathbf{v}) = T(\mathbf{u}) + T(\mathbf{v})$
- $\forall c_1, \dots, c_k \in \mathbb{R}, \forall \mathbf{v}_1, \dots, \mathbf{v}_k \in \mathbb{R}^n, \quad T(c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k) = c_1T(\mathbf{v}_1) + \dots + c_kT(\mathbf{v}_k)$

7.1.7 To Prove that a Mapping is Not a Linear Transformation

To show that a mapping $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is **not** a **linear transformation**,

- Show that $T(\mathbf{0}) \neq \mathbf{0}$; or
- Find $c \in \mathbb{R}, \mathbf{v} \in \mathbb{R}^n$ such that $T(c\mathbf{v}) \neq cT(\mathbf{v})$; or
- Find $\mathbf{u}, \mathbf{v} \in \mathbb{R}^n$ such that $T(\mathbf{u} + \mathbf{v}) \neq T(\mathbf{u}) + T(\mathbf{v})$.

7.1.8 Representation of Linear Transformations

7.1.8.1 Linear Transformations are Completely Determined by a Basis of \mathbb{R}^n and its Images

Suppose that $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation** and $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ is a **basis** for \mathbb{R}^n . Then $\forall \mathbf{v} \in \mathbb{R}^n$ $(\mathbf{v})_S = (c_1, \dots, c_n)$, and

$$\begin{aligned} T(\mathbf{v}) &= T(c_1\mathbf{v}_1 + \dots + c_n\mathbf{v}_n) \\ &= c_1T(\mathbf{v}_1) + \dots + c_nT(\mathbf{v}_n) \end{aligned}$$

$T(\mathbf{v})$ is completely determined by $T(\mathbf{v}_1), \dots, T(\mathbf{v}_n)$. Let

- \mathbf{A} be the **standard matrix** for T ,
- $\mathbf{B} = \begin{pmatrix} T(\mathbf{v}_1) & \dots & T(\mathbf{v}_n) \end{pmatrix}$, and
- $\mathbf{P} = \begin{pmatrix} \mathbf{v}_1 & \dots & \mathbf{v}_n \end{pmatrix}$

Then $\mathbf{A} = \mathbf{BP}^{-1}$.

7.1.8.2 Columns of Standard Matrix are Images of Standard Basis of \mathbb{R}^n

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation** and \mathbf{A} be the **standard matrix** for T . Then $\mathbf{A} = \begin{pmatrix} T(\mathbf{e}_1) & \dots & T(\mathbf{e}_n) \end{pmatrix}$.

7.1.9 Linear Combination Preserving Implies Linear Transformation

7.1.9.1 With k vectors

Theorem. A mapping $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation**, i.e., $T(\mathbf{x}) = \mathbf{Ax} \Leftrightarrow \forall c_1, \dots, c_k \in \mathbb{R}, \forall \mathbf{v}_1, \dots, \mathbf{v}_k \in \mathbb{R}^n, \quad T(c_1\mathbf{v}_1 + \dots + c_k\mathbf{v}_k) = c_1T(\mathbf{v}_1) + \dots + c_kT(\mathbf{v}_k)$

7.1.9.2 With Two Vectors

Theorem. A mapping $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ is a **linear transformation**, i.e., $T(\mathbf{x}) = \mathbf{Ax} \Leftrightarrow \forall c, d \in \mathbb{R}, \forall \mathbf{u}, \mathbf{v} \in \mathbb{R}^n, \quad T(c\mathbf{u} + d\mathbf{v}) = cT(\mathbf{u}) + dT(\mathbf{v})$.

7.1.10 General Definition of Linear Transformations

Definition. Let V and W be **vector spaces**. A mapping $T : V \rightarrow W$ is a **linear transformation** if

$$\forall \mathbf{u}, \mathbf{v} \in \mathbb{R}^n, \forall c, d \in \mathbb{R}, \quad T(c\mathbf{u} + d\mathbf{v}) = cT(\mathbf{u}) + dT(\mathbf{v})$$

7.1.11 Change of Bases

7.1.11.1 Change Between the Images of Two Bases

Theorem. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation**. Let

- $S = \{\mathbf{u}_1, \dots, \mathbf{u}_n\}$ and $R = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be **bases** for \mathbb{R}^n ,
- $\mathbf{B} = \begin{pmatrix} T(\mathbf{u}_1) & \cdots & T(\mathbf{u}_n) \end{pmatrix}$ and $\mathbf{C} = \begin{pmatrix} T(\mathbf{v}_1) & \cdots & T(\mathbf{v}_n) \end{pmatrix}$, and
- \mathbf{P} be the **transition matrix** from S to R and $\mathbf{Q} = \mathbf{P}^{-1}$

Then $\mathbf{B} = \mathbf{C}\mathbf{P}$ and $\mathbf{C} = \mathbf{B}\mathbf{Q}$.

7.1.11.2 Change Basis for Linear Operator

Theorem. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^n$ be a **linear operation** on \mathbb{R}^n and \mathbf{A} be the **standard matrix**. Let $S = \{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be a **basis** for \mathbb{R}^n and $\mathbf{P} = \begin{pmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_n \end{pmatrix}$. Then

- $[T(\mathbf{v})]_S = \mathbf{B}[\mathbf{v}]_S$, where $\mathbf{B} = \mathbf{P}^{-1}\mathbf{A}\mathbf{P}$
- T can be represented by $[\mathbf{v}]_S \mapsto \mathbf{B}[\mathbf{v}]_S$
- \mathbf{A} and \mathbf{B} are **similar**.

7.1.12 Composition

7.1.12.1 Definition of Function Composition

Definition. Let $f : X \rightarrow Y$ and $g : Y \rightarrow Z$ be two functions. The **composition** of g with f , denoted by $g \circ f$, is the function $X \rightarrow Z$ such that $\forall x \in X \quad (g \circ f)(x) = g(f(x))$

7.1.12.2 Function Composition is not Commutative in General

Note: In general, $g \circ f \neq f \circ g$.

7.1.12.3 Linear Transformation Composition

Definition. Let $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $T : \mathbb{R}^m \rightarrow \mathbb{R}^k$ be **linear transformations**. The **composition** of T with S , denoted by $T \circ S$, is the mapping $\mathbb{R}^n \rightarrow \mathbb{R}^k$ such that $\forall \mathbf{u} \in \mathbb{R}^n, \quad (T \circ S)(\mathbf{u}) = T(S(\mathbf{u}))$

7.1.12.4 Linear Transformation Composition is not Commutative in General

Note: In general, $T \circ S \neq S \circ T$.

7.1.12.5 Composition of Linear Transformations is a Linear Transformation and Formula for its Standard Matrix

Theorem. Let $S : \mathbb{R}^n \rightarrow \mathbb{R}^m$ and $T : \mathbb{R}^m \rightarrow \mathbb{R}^k$ be **linear transformations**. Then

- $T \circ S : \mathbb{R}^n \rightarrow \mathbb{R}^k$ is a **linear transformation**

Let \mathbf{A} and \mathbf{B} be the **standard matrices** for S and T respectively. Then

- the **standard matrix** for $T \circ S$ is \mathbf{BA} .

7.1.12.6 Properties of Linear Transformation Composition

Theorem. Let $S, T, T_1, T_2, S_1, S_2, U$ be **linear transformations** and $c \in \mathbb{R}$. Then

- **Identity & Zero Composition:** $I \circ S = S \circ I = S$; $O \circ S = S \circ O = O$
- **Constant Multiple:** $c(T \circ S) = (cT) \circ S = T \circ (cS)$
- **Associativity:** $U \circ (T \circ S) = (U \circ T) \circ S$
- **Distributive Over Addition:**

$$\circ (T_1 + T_2) \circ S = T_1 \circ S + T_2 \circ S$$

$$\circ T \circ (S_1 + S_2) = T \circ S_1 + T \circ S_2$$

7.2 Ranges and Kernels

7.2.1 Range

7.2.1.1 Range of Function

Definition. Let $f : X \rightarrow Y$ be a **function**. The **range** of f , denoted by $R(f)$, is the set of all **images** of f :

$$R(f) = \{f(x) \mid x \in X\} \subseteq Y$$

7.2.1.2 Range of Linear Transformation

Definition. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation**. The **range** of T , denoted by $R(T)$, is the set of all **images** of T :

$$R(T) = \{T(\mathbf{v}) \mid \mathbf{v} \in \mathbb{R}^n\} \subseteq \mathbb{R}^m$$

7.2.1.3 Representation of Range

7.2.1.3.1 Range as Subspace Spanned by the Images of Any Basis for \mathbb{R}^n

Theorem. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation** and $\{\mathbf{v}_1, \dots, \mathbf{v}_n\}$ be any basis for \mathbb{R}^n . Then

- $R(T) = \text{span}\{T(\mathbf{v}_1), \dots, T(\mathbf{v}_n)\}$.
- $R(T)$ is a **subspace** of \mathbb{R}^m .

7.2.1.3.2 Range as Column Space of Standard Matrix

Theorem. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation** and \mathbf{A} be the **standard matrix** for T . Then $R(T) = \text{column space of } \mathbf{A}$.

7.2.2 Rank of Linear Transformation

Definition. Let T be a **linear transformation**. The **rank** of T , denoted by $\text{rank}(T)$, is defined as

$$\text{rank}(T) = \dim(R(T))$$

Let \mathbf{A} be the **standard matrix** for T . Then

$$\text{rank}(T) = \dim(R(T)) = \dim(\text{column space of } \mathbf{A}) = \text{rank}(\mathbf{A})$$

7.2.3 Kernel

7.2.3.1 Definition of Kernel of Linear Transformation

Definition. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation**. The **kernel** of T , denoted by $\text{Ker}(T)$, is

$$\text{Ker}(T) = \{\mathbf{v} \in \mathbb{R}^n \mid T(\mathbf{v}) = \mathbf{0}\} \subseteq \mathbb{R}^n$$

Recall that $T(\mathbf{0}) = \mathbf{0} \Rightarrow \mathbf{0} \in \text{Ker}(T) \subseteq \mathbb{R}^n$.

7.2.3.2 Representation of Kernel

7.2.3.2.1 Kernel as Nullspace of Standard Matrix and Subspace of \mathbb{R}^n

Theorem. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation** and \mathbf{A} be the **standard matrix** for T . Then $\text{Ker}(T) = (\text{nullspace of } \mathbf{A})$ which is a **subspace** of \mathbb{R}^n .

7.2.3.2.2 Finding General Solution for Kernel

Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation** and \mathbf{A} be the **standard matrix** for T . To find a general solution for $\text{Ker}(T)$, solve the linear system $\mathbf{A}\mathbf{x} = \mathbf{0}$.

7.2.4 Nullity of Linear Transformation

Definition. Let T be a **linear transformation**. The **nullity** of T , denoted by $\text{nullity}(T)$, is defined as

$$\text{nullity}(T) = \dim(\text{Ker}(T))$$

Let \mathbf{A} be the **standard matrix** for T . Then

$$\text{nullity}(T) = \dim(\text{Ker}(T)) = \dim(\text{nullspace of } \mathbf{A}) = \text{nullity}(\mathbf{A})$$

7.2.5 Dimension Theorem for Linear Transformations

Theorem. Let $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$ be a **linear transformation**. Then $\text{rank}(T) + \text{nullity}(T) = n$.

7.2.6 Linear Transformation Between Vector Spaces

Let $T : V \rightarrow W$ be a **linear transformation** between **vector spaces**.

7.2.6.1 Range

Definition. $R(T) = \{T(\mathbf{v}) \mid \mathbf{v} \in V\}$ is a **subspace** of W .

7.2.6.2 Rank

Definition. $\text{rank}(T) = \dim(R(T))$.

7.2.6.3 Kernel

Definition. $\text{Ker}(T) = \{\mathbf{v} \in V \mid T(\mathbf{v}) = \mathbf{0}\}$ is a **subspace** of V .

7.2.6.4 Nullity

Definition. $\text{nullity}(T) = \dim(\text{Ker}(T))$.

7.2.6.5 Dimension Theorem

Definition. $\text{rank}(T) + \text{nullity}(T) = \dim(V)$.

7.3 Geometric Linear Transformations

7.3.1 Geometric Interpretation Completely Determined by Effect on a Basis for its Domain

Since the images of any **basis** for \mathbb{R}^n completely determines a **linear transformation** $T : \mathbb{R}^n \rightarrow \mathbb{R}^m$, to study the geometric interpretation of a **linear transformation**, it suffices to check the effect of the **linear transformation** on a **basis** for its domain.

7.3.2 Scaling

7.3.2.1 Scaling in \mathbb{R}^2

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a **linear transformation** and $\begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$ be the **standard matrix** for T , where $\lambda_1, \lambda_2 > 0$.

$$T \left(\begin{pmatrix} x \\ y \end{pmatrix} \right) = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} = \begin{pmatrix} \lambda_1 x \\ \lambda_2 y \end{pmatrix}$$

Then T is a **scaling** in \mathbb{R}^2 ,

- along the x -axis by a factor of λ_1 , and
- along the y -axis by a factor of λ_2 .

7.3.2.1.1 Dilation & Contraction

Suppose further that $\lambda_1 = \lambda_2$. Let $\lambda = \lambda_1 = \lambda_2$. Then

- T is a **dilation** if $\lambda > 1$.
- T is a **contraction** if $0 < \lambda < 1$.

7.3.2.2 Scaling in other Axes in \mathbb{R}^2

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a **linear transformation** and \mathbf{A} be the **standard matrix**.

- Suppose \mathbf{A} is **diagonalizable**, say $\exists \mathbf{P}$ such that $\mathbf{P}^{-1} \mathbf{A} \mathbf{P} = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix}$, $\lambda_1, \lambda_2 > 0$
- Suppose $\mathbf{P} = (\mathbf{v}_1 \ \mathbf{v}_2)$. Let $S = \{\mathbf{v}_1, \mathbf{v}_2\}$. Then S is a **basis** for \mathbb{R}^2 .
- $\forall \mathbf{v} \in \mathbb{R}^2$, $[T(\mathbf{v})]_S = \begin{pmatrix} \lambda_1 & 0 \\ 0 & \lambda_2 \end{pmatrix} [\mathbf{v}]_S$ (from section 7.1.11.2)
- Then T can be viewed as a scaling
 - along the direction of \mathbf{v}_1 by factor λ_1 ,
 - along the direction of \mathbf{v}_2 by factor λ_2 ,

7.3.2.3 Scaling in \mathbb{R}^3

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a **linear transformation** with **standard matrix** $\begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$, where

$\lambda_1, \lambda_2, \lambda_3 > 0$.

$$T \left(\begin{pmatrix} x \\ y \\ z \end{pmatrix} \right) = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix} \begin{pmatrix} x \\ y \\ z \end{pmatrix} = \begin{pmatrix} \lambda_1 x \\ \lambda_2 y \\ \lambda_3 z \end{pmatrix}$$

Then T is a **scaling**,

- along the x -axis by a factor of λ_1 ,
- along the y -axis by a factor of λ_2 ,
- along the z -axis by a factor of λ_3 .

7.3.2.3.1 Dilation & Contraction

Suppose further that $\lambda_1 = \lambda_2 = \lambda_3 = \lambda$. Then

- T is a **dilation** if $\lambda > 1$.
- T is a **contraction** if $0 < \lambda < 1$.

7.3.2.4 Scaling in other Axes in \mathbb{R}^3

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a **linear transformation** with **standard matrix** \mathbf{A} .

- Suppose \mathbf{A} is **diagonalizable**, say $\exists \mathbf{P}$ such that $\mathbf{P}^{-1}\mathbf{A}\mathbf{P} = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix}$, $\lambda_1, \lambda_2, \lambda_3 > 0$
- Suppose $\mathbf{P} = (\mathbf{v}_1 \ \mathbf{v}_2 \ \mathbf{v}_3)$. Let $S = \{\mathbf{v}_1, \mathbf{v}_2, \mathbf{v}_3\}$. Then S is a **basis** for \mathbb{R}^3 .
- $\forall \mathbf{v} \in \mathbb{R}^3$, $[T(\mathbf{v})]_S = \begin{pmatrix} \lambda_1 & 0 & 0 \\ 0 & \lambda_2 & 0 \\ 0 & 0 & \lambda_3 \end{pmatrix} [\mathbf{v}]_S$ (from section 7.1.11.2)
- Then T can be viewed as a scaling
 - along the direction of \mathbf{v}_1 by factor λ_1 ,
 - along the direction of \mathbf{v}_2 by factor λ_2 ,
 - along the direction of \mathbf{v}_3 by factor λ_3 .

7.3.3 Reflection

7.3.3.1 Reflection in \mathbb{R}^2

7.3.3.1.1 Reflection w.r.t. x -axis

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a **linear transformation** with **standard matrix** $\begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$

$$T\left(\begin{pmatrix} x \\ y \end{pmatrix}\right) = \begin{pmatrix} x \\ -y \end{pmatrix}$$

T is the **reflection** w.r.t. the x -axis.

7.3.3.1.2 Reflection w.r.t. y -axis

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a **linear transformation** with **standard matrix** $\begin{pmatrix} -1 & 0 \\ 0 & 1 \end{pmatrix}$

$$T \left(\begin{pmatrix} x \\ y \end{pmatrix} \right) = \begin{pmatrix} -x \\ y \end{pmatrix}$$

T is the **reflection** w.r.t. the y -axis.

7.3.3.1.3 Reflection w.r.t. the line $y = x$

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a **linear transformation** with **standard matrix** $\begin{pmatrix} 0 & 1 \\ 1 & 0 \end{pmatrix}$

$$T \left(\begin{pmatrix} x \\ y \end{pmatrix} \right) = \begin{pmatrix} y \\ x \end{pmatrix}$$

T is the **reflection** w.r.t. the line $y = x$.

7.3.3.1.4 Reflection w.r.t. a Line Through the Origin

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be a **linear transformation** with **standard matrix** $\begin{pmatrix} \cos(2\theta) & \sin(2\theta) \\ \sin(2\theta) & -\cos(2\theta) \end{pmatrix}$

T is the **reflection** w.r.t. the line ℓ passing through the origin where θ is the angle between ℓ and the x -axis.

Orthogonal Matrices of Determinant -1

Every **orthogonal matrix** of $\det = -1$ is in this form.

Representation with Unit Vector Parallel to Line

- Let $\mathbf{n} = (\cos \theta, \sin \theta)^T$ be a unit vector on ℓ .
- \mathbf{p} is the projection of \mathbf{v} onto $\text{span}\{\mathbf{n}\}$
 - $\mathbf{p} = (\mathbf{v} \cdot \mathbf{n})\mathbf{n}$
- \mathbf{p} is the midpoint of \mathbf{v} and $T(\mathbf{v})$
 - $T(\mathbf{v}) = 2\mathbf{p} - \mathbf{v} = 2(\mathbf{v} \cdot \mathbf{n})\mathbf{n} - \mathbf{v}$

Representation with Unit Vector Orthogonal to Line

- Let $\mathbf{n} = (\sin \theta, -\cos \theta)^T$ be a unit vector orthogonal to ℓ .
- \mathbf{p} is the projection of \mathbf{v} onto $\text{span}\{\mathbf{n}\}$
 - $\mathbf{p} = (\mathbf{v} \cdot \mathbf{n})\mathbf{n}$
- Note that $T(\mathbf{v}) + 2\mathbf{p} = \mathbf{v}$
 - $T(\mathbf{v}) = \mathbf{v} - 2\mathbf{p} = \mathbf{v} - 2(\mathbf{v} \cdot \mathbf{n})\mathbf{n}$

7.3.3.2 Reflections in \mathbb{R}^3

7.3.3.2.1 Reflections w.r.t. Planes Formed by Coordinate Axes

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be a **linear transformation**.

- If the **standard matrix** is $\begin{pmatrix} 1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & -1 \end{pmatrix}$,
 - then T is the reflection w.r.t. the xy -plane.

- If the **standard matrix** is $\begin{pmatrix} 1 & 0 & 0 \\ 0 & -1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$,
 - then T is the reflection w.r.t. the xz -plane.

- If the **standard matrix** is $\begin{pmatrix} -1 & 0 & 0 \\ 0 & 1 & 0 \\ 0 & 0 & 1 \end{pmatrix}$,
 - then T is the reflection w.r.t. the yz -plane.

7.3.3.2.2 Reflections w.r.t. Any Plane

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the reflection w.r.t. the plane $ax + by + cz = 0$, where a, b, c not all zero. Then $\mathbf{n} = (a, b, c)^T$ and $\forall \mathbf{v} \in \mathbb{R}^3$, $T(\mathbf{v}) = \mathbf{v} - \left(2 \frac{\mathbf{v} \cdot \mathbf{n}}{\|\mathbf{n}\|^2}\right) \mathbf{n}$.

7.3.4 Rotation

7.3.4.1 Rotation in \mathbb{R}^2

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be the **rotation** about the origin by θ .

- Then T is a **linear transformation**.
- The **standard matrix** for T is $\begin{pmatrix} \cos \theta & -\sin \theta \\ \sin \theta & \cos \theta \end{pmatrix}$

7.3.4.1.1 Orthogonal Matrices of Determinant 1

Every **orthogonal matrix** of $\det = 1$ is in this form.

7.3.4.2 Rotation in \mathbb{R}^3

- Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the **rotation** about the z -axis anticlockwise by angle θ .
 - The z -coordinate does not change
 - It is the rotation about the origin on the plane $z = z_0$ anticlockwise by θ .

- **Standard Matrix** $\begin{pmatrix} \cos \theta & -\sin \theta & 0 \\ \sin \theta & \cos \theta & 0 \\ 0 & 0 & 1 \end{pmatrix}$

- Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the **rotation** about the x -axis anticlockwise by angle θ .

- The x -coordinate does not change

- It is the rotation about the origin on the plane $x = x_0$ anticlockwise by θ .

- **Standard Matrix** $\begin{pmatrix} 1 & 0 & 0 \\ 0 & \cos \theta & -\sin \theta \\ 0 & \sin \theta & \cos \theta \end{pmatrix}$

- Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be the **rotation** about the y -axis anticlockwise by angle θ .

- The y -coordinate does not change

- It is the rotation about the origin on the plane $y = y_0$ anticlockwise by θ .

- **Standard Matrix** $\begin{pmatrix} \cos \theta & 0 & -\sin \theta \\ 0 & 1 & 0 \\ \sin \theta & 0 & \cos \theta \end{pmatrix}$

7.3.5 Reflections and Rotations in \mathbb{R}^2

7.3.5.1 Determinant Determines Rotation vs Reflection

Suppose **standard matrix** A for $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ is **orthogonal**.

- If $\det(A) = 1$, T represents a **rotation** about the origin.
- If $\det(A) = -1$, T represents the **reflection** w.r.t. a line passing through the origin.

7.3.5.2 Reflection about a Line can be Decomposed into Reflection and Rotation

Since the **standard matrix** for **reflection** about a line ℓ passing through the origin where θ is the angle between ℓ and the x -axis is

$$\begin{pmatrix} \cos(2\theta) & \sin(2\theta) \\ \sin(2\theta) & -\cos(2\theta) \end{pmatrix} = \begin{pmatrix} \cos(2\theta) & -\sin(2\theta) \\ \sin(2\theta) & \cos(2\theta) \end{pmatrix} \begin{pmatrix} 1 & 0 \\ 0 & -1 \end{pmatrix}$$

Reflection w.r.t. ℓ can be decomposed into

1. Reflection w.r.t. x -axis
2. Rotation about the origin anticlockwise by 2θ .

7.3.6 Shears

7.3.6.1 Shears in \mathbb{R}^2

7.3.6.1.1 Shear in x -direction

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by

$$T \left(\begin{pmatrix} x \\ y \end{pmatrix} \right) = \begin{pmatrix} x + ky \\ y \end{pmatrix}$$

Then T is a **shear** in the x -direction by a factor k .

7.3.6.1.2 Shear in y -direction

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by

$$T \left(\begin{pmatrix} x \\ y \end{pmatrix} \right) = \begin{pmatrix} x \\ kx + y \end{pmatrix}$$

Then T is a **shear** in the y -direction by a factor k .

7.3.6.2 Shears in \mathbb{R}^3

Let $T : \mathbb{R}^3 \rightarrow \mathbb{R}^3$ be defined by

$$T \left(\begin{pmatrix} x \\ y \\ z \end{pmatrix} \right) = \begin{pmatrix} x + k_1 z \\ y + k_2 z \\ z \end{pmatrix}$$

Then T is a **shear** in the x -direction by a factor k_1 , and in the y -direction by factor k_2 .

- On yz -plane $x = 0$, it is a shear in y -direction by k_2 .
- On xz -plane $y = 0$, it is a shear in x -direction by k_1 .
- On the plane $z = 1$,

$$\circ T \left(\begin{pmatrix} x \\ y \\ 1 \end{pmatrix} \right) = \begin{pmatrix} x + k_1 \\ y + k_2 \\ 1 \end{pmatrix}.$$

7.3.7 Translations

Let $T : \mathbb{R}^2 \rightarrow \mathbb{R}^2$ be defined by

$$T \left(\begin{pmatrix} x \\ y \end{pmatrix} \right) = \begin{pmatrix} x + a \\ y + b \end{pmatrix}, a, b \in \mathbb{R}$$

- T is a **translation** by $(a, b)^T$.
- T is **not** a **linear transformation** unless $a = b = 0$.

7.3.8 2D Computer Graphic System

7.3.8.1 Representation of 2D Figures

- In 2D computer graphic, a figure is drawn by connecting points $(a_1, b_1), (a_2, b_2), \dots, (a_n, b_n)$.
- It can be written as an $2 \times n$ matrix:

$$\circ \mathbf{M} = \begin{pmatrix} a_1 & a_2 & \cdots & a_n \\ b_1 & b_2 & \cdots & b_n \end{pmatrix}.$$

7.3.8.2 Scaling, Reflection, Rotation, Shearing of 2D Figures

- Let T be a scaling/reflection/rotation/shearing on \mathbb{R}^2
 - Then T is a **linear transformation** with **standard matrix** \mathbf{A} .
- Let $\mathbf{M} = \begin{pmatrix} \mathbf{v}_1 & \mathbf{v}_2 & \cdots & \mathbf{v}_n \end{pmatrix}$ be a 2D figure.
- Then the resulting figure by T is
 - $\begin{pmatrix} T(\mathbf{v}_1) & \cdots & T(\mathbf{v}_n) \end{pmatrix} = \begin{pmatrix} \mathbf{A}\mathbf{v}_1 & \cdots & \mathbf{A}\mathbf{v}_n \end{pmatrix} = \mathbf{A} \begin{pmatrix} \mathbf{v}_1 & \cdots & \mathbf{v}_n \end{pmatrix} = \mathbf{A}\mathbf{M}.$

7.3.8.3 Translating 2D Figure with Homogeneous Coordinate System

- **Homogeneous coordinate system** is formed by identifying \mathbb{R}^2 with plane $z = 1$ in \mathbb{R}^3 :

$$\begin{pmatrix} a \\ b \end{pmatrix} \leftrightarrow \begin{pmatrix} a \\ b \\ 1 \end{pmatrix}$$

- A figure $(a_1, b_1), \dots, (a_n, b_n)$ is identified by $(a_1, b_1, 1), \dots, (a_n, b_n, 1)$.

- The associated matrix $\mathbf{M} = \begin{pmatrix} a_1 & \cdots & a_n \\ b_1 & \cdots & b_n \\ 1 & \cdots & 1 \end{pmatrix}$

- Suppose we want to do a translation by $(a, b)^T$.

- Define shear $T \left(\begin{pmatrix} x \\ y \\ z \end{pmatrix} \right) = \begin{pmatrix} x + az \\ y + bz \\ z \end{pmatrix}$ with **standard matrix** $\mathbf{A} = \begin{pmatrix} 1 & 0 & a \\ 0 & 1 & b \\ 0 & 0 & 1 \end{pmatrix}$

- $\mathbf{A}\mathbf{M} = \begin{pmatrix} a_1 + a & \cdots & a_n + a \\ b_1 + b & \cdots & b_n + b \\ 1 & \cdots & 1 \end{pmatrix}$ represent the translation by $(a, b)^T$.