

Referred By

Bangalore

Ahmedabad

Registration Id: 1670660003

: Cisco

Patient Name: Mr. Fredrick Antony 855519

Collection Date / Time

Registration Date/Time :05/06/2017 / 08:13AM :05/06/2017 / 12:00AM

Reporting Date / Time

:05/06/2017 / 12:20PM

Age /Sex

: 47 Years

/ Male

Thyroid Function Test

Sample Type: SERUM

Ultrasensitive TSH

OBSERVED VALUE TEST

UNITS

Reference Range

Total T3 (Tri-iodothyronine)

75.5

ng/dl

58 - 159

Method : CMIA)

5.76

ug/dl

4.51 - 11.66

Total T4 (Thyroxine) Method : CMIA)

1.3947

uIU/ml

Euthyroid: 0.5 to 5.0

- In cases of primary hypothyroidism, T3 and T4 levels are low and TSH is significantly elevated. In the case of pituatary dysfunction, either due to intrinsic hypothalamic or pituatary disease i.e central hypothyroidism, normal or marginally elevated basal TSH levels are often seen despite significant reduction in T4 and T3 levels.

- Primary hyperthyroidism (eg: Grave~s disease, nodular goiter) is associated with high levels of thyroid hormones and depressed or undetectable levels of TSH.

END OF REPORT

Dr. Sangeeta Sharma M.D. **Pathologist**

MAHESH B

Verified By

Page 5 of 5

Corporate Office:

☐ Mumbai - Jn of 14th and 33rd Rd, Khar (W), Mumbai - 52. Tel : 91-22-4342 8888 Fax : 91-22-4342 8880

□ Pune - Fortaleza, Central Avenue, Kalyani Nagar, Pune - 06 Tel : 91-20-2660 0505 Fax : 91-20-2660 0506

□ Bangalore - Corporate Block-A, Airport Road, Bangalore - 08. Tel : 91-80-4046 6464 Fax : 91-80-4046 6465

info@nmmedical.com | www.nmmedical.com