

Exchangeable Array and Random Graph

Dongwoo Kim
ANU

April 13, 2016

1 Exchangeable array for graph

Definition 1 (Jointly exchangeable array). A random 2-array $(X_{ij})_{i,j \in \mathbb{N}}$ is jointly exchangeable if

$$(X_{ij}) \stackrel{d}{=} (X_{\pi(i)\pi(j)}) \quad \text{for } i, j \in \mathbb{N}^2 \quad (1)$$

for any permutation π of \mathbb{N} .

Theorem 1 (Aldous, Hoover Theorem [Aldous, 1981, Hoover, 1979]). *A random 2-array $(X_{ij})_{i,j \in \mathbb{N}}$ is jointly exchangeable if and only if there exists a random measurable function $f : [0, 1]^3 \rightarrow \mathcal{X}$ such that*

$$(X_{ij}) \stackrel{d}{=} (f(U_i, U_j, U_{ij})), \quad (2)$$

where $(U_i)_{i \in \mathbb{N}}$ and $(U_{ij})_{ij \in \mathbb{N}^2}$ with $U_{ij} = U_{ji}$ are a sequence and matrix of i.i.d. Uniform $[0, 1]$ random variables.

If the function f is a symmetric in its first two argument, then the matrix X is symmetric. In general, however, it does not need to be a symmetric (e.g. directed graph).

For the undirected graph case $\mathcal{X} = \{0, 1\}$, the theorem can be simplified further through a random function called *graphon* $W : [0, 1]^2 \rightarrow [0, 1]$, symmetric in its arguments, where

$$f(U_i, U_j, U_{ij}) = \begin{cases} 1 & U_{ij} < W(U_i, U_j) \\ 0 & \text{otherwise} \end{cases}$$

Example 1.1 (Random function prior on function f [Lloyd et al., 2013]). Lloyd et al. use a Gaussian process prior to define function f for an undirected graph. They define $W(U_i, U_j) = \phi(\Theta(U_i, U_j))$ where ϕ is a logistic function, and $\Theta(\cdot, \cdot)$ is a continuous function with a Gaussian process prior. Thus, the probability of edge between node i and j is equal to Bernoulli($\phi(\Theta(U_i, U_j))$).

Example 1.2 (Stochastic block model). The stochastic block model is another generative model for random graphs. The model assumes that each node belongs to a community and there is a edge probability between communities. Then the probability of edge between two different nodes are determined by the edge probability between communities where the two nodes belong. In stochastic block model with Aldous, Hoover Theorem, U_i and U_j determine the communities of node i and j , respectively, and U_{ij} determines the existence of edge between node i and j as done in graphon. The stick breaking process is one way to partition unit length interval into an infinite number of communities [Sethuraman, 1994].

Definition 2 (Sparse Graph). Let the number of nodes in a graph be n . The graph is sparse if the number of edges are $o(n^2)$ or dense if the number of edges are $\Theta(n^2)$.

Remark 1 (Graphon is trivially dense). Every graph represented by graphon W are either empty or dense. The asymptotic proportion of edges is $p = \frac{1}{2} \int W(x, y) dx dy$ and the graph is hence either empty ($p = 0$) or dense (since $O(pn^2) = O(n^2)$).

In [Lloyd et al., 2013], the authors place a Gaussian process prior over graphon $W : [0, 1]^2 \rightarrow \mathbb{R}$ and transform the output through the logistic function to model the edge probability between nodes.

For the undirected graph case where $X_{ij} = X_{ji}$, one can sample upper triangle of the adjacency matrix, and use the same result for the lower triangle. For the directed graph case, however, the adjacency matrix is no longer symmetric, also, by the theorem, both X_{ij} and X_{ji} rely on single parameter U_{ij} . Therefore, one should jointly sample (X_{ij}, X_{ji}) together from three parameters U_i, U_j, U_{ij} , which means X_{ij} and X_{ji} are not conditionally independent. In addition, asymmetric graphon W could be employed for directed random graph. However, [Cai et al., 2015] show that asymmetric graphon is inappropriate to impose certain structures on a graph such as the partial ordering and propose a class of priors for directed graphs.

2 Exchangeable array for matrix factorisation

Jointly exchangeable array assume that both rows and columns index the same entities. However, the assumption does not hold for the general matrix where rows and columns represent different entities (e.g. users and items in recommendation system).

Definition 3 (Separately exchangeable array). A random 2-array $(X_{ij})_{i,j \in \mathbb{N}}$ is separately exchangeable if

$$(X_{ij}) \stackrel{d}{=} (X_{\pi(i)\sigma(j)}) \quad \text{for } i, j \in \mathbb{N}^2 \quad (3)$$

for any permutation π and σ of \mathbb{N} .

Corollary 1.1 (Separately exchangeable array). A random 2-array (X_{ij}) is separately exchangeable if and only if there exists a random measurable function $f : [0, 1]^3 \rightarrow \mathcal{X}$ such that

$$(X_{ij}) \stackrel{d}{=} (f(U_i, U_j, U_{ij})), \quad (4)$$

where $(U_i)_{i \in \mathbb{N}}$, $(U_j)_{j \in \mathbb{N}}$, and $(U_{ij})_{i,j \in \mathbb{N}^2}$ are i.i.d. Uniform $[0, 1]$ random variables.

In the separately exchangeable array case, function f is not a symmetric in its first two arguments.

Example 3.1 (Probabilistic matrix factorisation (PMF) [Salakhutdinov and Mnih, 2008]). PMF is one instantiation of Corollary 1.1. Let the generative process of PMF be

$$U_i \sim \mathcal{MN}_d(0, \Sigma_U) \quad (5)$$

$$V_j \sim \mathcal{MN}_d(0, \Sigma_V) \quad (6)$$

$$X_{ij} \sim \mathcal{N}(U_i^\top V_j, \sigma_x^2) \quad (7)$$

where \mathcal{MN}_d is a d -dimensional zero-mean multivariate normal distribution with covariance Σ . This corresponds to random measurable function $f(U_i, U_j, U_{ij}) = \Phi_1(U_{ij}; \Phi_d(U_i; 0, \Sigma_U)^\top \Phi_d(U_j; 0, \Sigma_U), \sigma_x^2)$, where $\Phi_d(\cdot; \mu, \sigma^2)$ is an inverse-CDF of d -dimensional multivariate function with mean μ and variance σ^2 . Note that σ here differs from the permutation notation.

3 Exchangeable array for knowledge graph

A knowledge graph can be represented as a 3-array tensor X_{ijk} , where the first two dimensions represent entities, and the third dimension represent relations. Therefore, X_{ijk} is jointly exchangeable in its first two dimensions, and the third dimension is separately exchangeable to the first two dimensions.

Definition 4 (π -exchangeable array for knowledge graph [Orbanz and Roy, 2015]). A random 3-array $(X_{ijk})_{i,j,k \in \mathbb{N}}$ is jointly exchangeable in its first two dimensions and separately exchangeable with the third dimension if

$$(X_{ijk}) \stackrel{d}{=} (X_{\pi(i)\pi(j)\sigma(k)}) \quad \text{for } i, j, k \in \mathbb{N}^3 \quad (8)$$

for any permutation π and σ of \mathbb{N} .

Corollary 1.2 (π -exchangeable array for knowledge graph). *A random 3-array X_{ijk} is jointly exchangeable in its first two dimension and separately exchangeable for the third dimension if and only if there exists a random measurable function $f : [0, 1]^7 \rightarrow \mathcal{X}$ such that*

$$(X_{ijk}) \stackrel{d}{=} (f(U_i, U_j, U_k, U_{ij}, U_{ik}, U_{jk}, U_{ijk})), \quad (9)$$

where $(U_i)_{i \in \mathbb{N}}$, $(U_k)_{k \in \mathbb{N}}$, $(U_{ij})_{ij \in \mathbb{N}^2}$, $(U_{ik})_{ik \in \mathbb{N}^2}$, and $(U_{ijk})_{ijk \in \mathbb{N}^3}$ are i.i.d. Uniform $[0, 1]$ random variables.

Remark 2 (Compositional model [Gu et al., 2015]). Compositional vector space models cannot be modelled through the exchangeable array theorem because the dependency between triples breaks the exchangeability.

Example 4.1 (Bayesian RESCAL (BRESICAL)). The generative process of BRESICAL with normally distributed output variable X_{ijk} is as follows:

$$e_i, e_j \sim \mathcal{MN}_d(0, \sigma_e^2 I_d) \quad (10)$$

$$R_k \sim \mathcal{MN}_{d^2}(0, \sigma_r^2 I_{d^2}) \quad (11)$$

$$X_{ijk} \sim \mathcal{N}(R_k^\top (e_i \otimes e_j), \sigma_x^2). \quad (12)$$

Again, we can transform uniform random variables U_i, U_j, U_k to e_i, e_j, R_k via the inverse-CDF of zero-mean multivariate normal distribution. Then X_{ijk} is determined by U_{ijk} via the inverse-CDF of normal distribution with mean $R_k^\top (e_i \otimes e_j)$ and variance σ_x^2 , i.e. $X_{ijk} = \Phi_1(U_{ijk}; R_k^\top (e_i \otimes e_j), \sigma_x^2)$. The model is invariant w.r.t the changes in the other uniform variables U_{ij}, U_{ik} , and U_{jk} .

4 Sparse exchangeable array

Definition 5 (Jointly exchangeable random measure on \mathbb{R}^2). A random measure ξ on \mathbb{R}^2 is jointly exchangeable if for every measure preserving transformation f on \mathbb{R}_+ , we have

$$\xi \circ (f \otimes f)^{-1} \stackrel{d}{=} \xi \quad (13)$$

Theorem 2 (Kallenberg’s exchangeable theorem [Kallenberg, 1990]). *A random measure ξ on \mathbb{R}^2 is jointly exchangeable if and only if almost surely*

$$\xi = \sum_{i,j} f(\alpha, \vartheta_i, \vartheta_j, \zeta_{i,j}) \delta_{\theta_i, \theta_j} \quad (14)$$

$$+ \sum_{j,k} (g(\alpha, \vartheta_j, \chi_{jk}) \delta_{\theta_j, \sigma_{jk}} + g'(\alpha, \vartheta_j, \chi_{jk}) \delta_{\sigma_{jk}, \theta_j}) \quad (15)$$

$$+ \sum_k (l(\alpha, \eta_k) \delta_{\rho_k, \rho'_k} + l'(\alpha, \eta_k) \delta_{\rho'_k, \rho_k}) \quad (16)$$

$$+ \sum_j (h(\alpha, \vartheta_j) (\delta_{\theta_j} \otimes \Lambda) + h'(\alpha, \vartheta_j) (\Lambda \otimes \delta_{\theta_j})) + \beta \Lambda_D + \gamma \Lambda^2 \quad (17)$$

for some measurable function $f : \mathbb{R}_+^4 \rightarrow \mathbb{R}_+$, $g, g' : \mathbb{R}_+^3 \rightarrow \mathbb{R}_+$, and $l, l', h, h' : \mathbb{R}_+^2 \rightarrow \mathbb{R}_+$, some collection of independent uniformly distributed random variables $(\zeta_{i,j})$ on $[0, 1]$, some independent unit rate Poisson processes $\{(\theta_j, \vartheta_j)\}$ and $\{(\sigma_{ij}, \chi_{ij})\}$ on \mathbb{R}_+^2 and $\{(\rho_j, \rho'_j, \eta_j)\}$ on \mathbb{R}^3 , and some independent set of random variables $\alpha, \beta, \gamma \geq 0$. Λ is Lebesgue measure on \mathbb{R}_+ , and Λ_D is Lebesgue measure on the diagonal of \mathbb{R}_+^2 .

Matrix factorisation for the sparse bipartite graph has been partially discussed in [Caron, 2012].

References

- [Aldous, 1981] Aldous, D. J. (1981). Representations for partially exchangeable arrays of random variables. *Journal of Multivariate Analysis*, 11(4):581–598.
- [Cai et al., 2015] Cai, D., Ackerman, N., and Freer, C. (2015). Priors on exchangeable directed graphs. pages 1–17.
- [Caron, 2012] Caron, F. (2012). Bayesian nonparametric models for bipartite graphs. *Advances in Neural Information Processing Systems 25 (Proceedings of NIPS)*, pages 1–9.
- [Gu et al., 2015] Gu, K., Miller, J., and Liang, P. (2015). Traversing knowledge graphs in vector space. In *EMNLP*.
- [Hoover, 1979] Hoover, D. N. (1979). Relations on probability spaces and arrays of random variables. *Preprint, Institute for Advanced Study, Princeton, NJ*, 2.
- [Kallenberg, 1990] Kallenberg, O. (1990). Exchangeable random measures in the plane. *Journal of Theoretical Probability*.
- [Lloyd et al., 2013] Lloyd, J., Orbanz, P., Ghahramani, Z., and Roy, D. (2013). Random function priors for exchangeable arrays with applications to graphs and relational data. In *Advances in Neural Information Processing Systems 26*, pages 1–9.
- [Orbanz and Roy, 2015] Orbanz, P. and Roy, D. M. (2015). Bayesian Models of Graphs, Arrays and Other Exchangeable Random Structures. *TPAMI*, pages 1–25.
- [Salakhutdinov and Mnih, 2008] Salakhutdinov, R. and Mnih, A. (2008). Bayesian probabilistic matrix factorization using Markov chain Monte Carlo. In *Proceedings of the 25th International Conference on Machine Learning (2008)*, volume 25, pages 880–887.

[Sethuraman, 1994] Sethuraman, J. (1994). A constructive definition of dirichlet priors. *Statistica sinica*, pages 639–650.