

North York General

MSS Laboratory, 4001 Leslie Street 3rd Floor Southeast Toronto ON M2K 1E1 Fax: (416) 756-6108

**Prenatal Screening Requisition – North York General** 

for Down Syndrome, Trisomy 18 and ONTD

<u>Health Care Provider points to consider:</u> Prenatal screening requires patient education and should proceed only with informed choice of the patient.

<u>Instructions for patients:</u> Nuchal Translucency (NT) ultrasounds need to be ordered by your health care provider. The blood sample can be drawn at any community lab **after** the NT ultrasound, ideally on the same day. **The MSS Laboratory does not make arrangements for the NT ultrasound.** 

\*\*Accurate information is necessary for a valid interpretation\*\*

* Name:(SURNAME)	(GIVEN)
* Date of Birth:/	(DD)
* Health Card #:	
* Address:	
* Postal Code:Phor	ne:

Obtain this requisition online at: https://prenatalscreeningonta	ario.ca/en/pso/requisitions-and	-provider-tools/mms-requisitions.aspx
Test Requested (choose one only)	Clinical Information- pleas	e complete all sections
Only select the eFTS or Maternal Serum Screening below if:  • NIPT has not been ordered in this pregnancy  • NIPT has been ordered, but has been uninformative	Racial origin:	Weight kg or lbs
Enhanced First Trimester Screen (eFTS)  (eFTS: NT, PAPPA, FBHCG, PIGF, AFP)  [CRL 45-84 mm]; corresponding to approximately 11 weeks and 2 days to 13 weeks and 3 days gestation.  Requires nuchal translucency (NT) ultrasound and blood sample	☐ Black ☐ Asian ☐ South East Asian ☐ Indigenous ☐ Other:	Last Menstrual Period (LMP):  (Ultrasound dating is required for eFTS)
Maternal Serum Screen [14w – 20w6d] (AFP, hCG, UE3, inhibin A)		sulin prior to pregnancy? Yes
Ultrasound dating preferred to LMP dating  Maternal Serum AFP only [15w – 20w6d]  SOGC recommends AFP testing only when ultrasound examination has failed to provide a sufficiently clear image of the neural tube to make a decision regarding the likelihood of Open Neural Tube Defect  Poor visibility on anatomy scan	Complete the following	/ER during this pregnancy? Yes  ng if this is an IVF pregnancy  patient is donor):(YYYY/MM/DD)  (YYYY/MM/DD)
Ultrasound (U/S) Information Sonographer or ordering prov	vider to complete. Identify U/S	operator code only if doing NT Scan.
Singleton/Twin A:  U/S Date:  (YYYY/MM/DD)  Twin B:  dichorionic monochorionic uncertain IUFD  CRL: Crown-Rump Length Crown-Rump Length Crown-Rump Length	h Bi-Parietal Diar cm mm BPD:	neter Nuchal Translucency  CRL 45.0-84.0 mm  cm  mm NT: mm
Sonographer's information:		
Operator Code: Site:	Site phone #: Signature:	
Ordering Provider:Address:		:
Phone: Fax:	Phone:	Fax:
Signature : Billing # For Blood Collection Centre Use Only	Provider Billing #	
Send 2 mL of serum to the laboratory indicated above (serum separator tube preferred). <b>Do not anticoagulate or freeze blood. Centrifuge.</b>		
Send primary tube to laboratory if there is a gel barrier, otherwise aliquot.		
Collection Centre:  Specimen Date:(YYYY/MM/DD) Phone :	#:(	