

A Not So Simple Full-history Recurrence Relation

$$T(n) = n - 1 + \frac{2}{n} \sum_{i=1}^{n-1} T(i), \text{ (for } n \geq 2); T(1) = 0.$$

A Not So Simple Full-history Recurrence Relation

$$T(n) = n - 1 + \frac{2}{n} \sum_{i=1}^{n-1} T(i), \text{ (for } n \geq 2); T(1) = 0.$$

- **Idea:** Use the shifting and canceling terms technique, s.t., most of the $T(i)$ terms gets canceled out.

A Not So Simple Full-history Recurrence Relation (Cont.)

Multiplying both sides by n , we get:

$$nT(n) = n(n-1) + 2 \sum_{i=1}^{n-1} T(i),$$

$$(n+1)T(n+1) = n(n+1) + 2 \sum_{i=1}^n T(i).$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Multiplying both sides by n , we get:

$$nT(n) = n(n-1) + 2 \sum_{i=1}^{n-1} T(i),$$

$$(n+1)T(n+1) = n(n+1) + 2 \sum_{i=1}^n T(i).$$

Therefore,

$$(n+1)T(n+1) - nT(n) = n(n+1) - n(n-1) + 2T(n)$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Multiplying both sides by n , we get:

$$nT(n) = n(n-1) + 2 \sum_{i=1}^{n-1} T(i),$$

$$(n+1)T(n+1) = n(n+1) + 2 \sum_{i=1}^n T(i).$$

Therefore,

$$\begin{aligned}(n+1)T(n+1) - nT(n) &= n(n+1) - n(n-1) + 2T(n) \\ &= 2n + 2T(n)\end{aligned}$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Multiplying both sides by n , we get:

$$\begin{aligned}nT(n) &= n(n-1) + 2 \sum_{i=1}^{n-1} T(i), \\(n+1)T(n+1) &= n(n+1) + 2 \sum_{i=1}^n T(i).\end{aligned}$$

Therefore,

$$\begin{aligned}(n+1)T(n+1) - nT(n) &= n(n+1) - n(n-1) + 2T(n) \\&= 2n + 2T(n) \\ \Rightarrow T(n+1) &= \frac{n+2}{n+1}T(n) + \frac{2n}{n+1}\end{aligned}$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Multiplying both sides by n , we get:

$$\begin{aligned}nT(n) &= n(n-1) + 2 \sum_{i=1}^{n-1} T(i), \\(n+1)T(n+1) &= n(n+1) + 2 \sum_{i=1}^n T(i).\end{aligned}$$

Therefore,

$$\begin{aligned}(n+1)T(n+1) - nT(n) &= n(n+1) - n(n-1) + 2T(n) \\&= 2n + 2T(n) \\ \Rightarrow T(n+1) &= \frac{n+2}{n+1}T(n) + \frac{2n}{n+1} \\&\leq \frac{n+2}{n+1}T(n) + 2 \quad (\text{A close approx.})\end{aligned}$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Expanding, we get

$$T(n) \leq 2 + \frac{n+1}{n} \left[2 + \frac{n}{n-1} \left[2 + \frac{n-1}{n-2} \left[\dots \frac{4}{3} \right] \right] \right]$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Expanding, we get

$$\begin{aligned}T(n) &\leq 2 + \frac{n+1}{n} \left[2 + \frac{n}{n-1} \left[2 + \frac{n-1}{n-2} \left[\dots \frac{4}{3} \right] \right] \right] \\&= 2 \left[1 + \frac{n+1}{n} + \frac{n+1}{n} \cdot \frac{n}{n-1} + \frac{n+1}{n} \cdot \frac{n}{n-1} \cdot \frac{n-1}{n-2} + \right. \\&\quad \left. \dots + \frac{n+1}{n} \cdot \frac{n}{n-1} \cdot \frac{n-1}{n-2} \dots \frac{4}{3} \right]\end{aligned}$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Expanding, we get

$$\begin{aligned}T(n) &\leq 2 + \frac{n+1}{n} \left[2 + \frac{n}{n-1} \left[2 + \frac{n-1}{n-2} \left[\dots \frac{4}{3} \right] \right] \right] \\&= 2 \left[1 + \frac{n+1}{n} + \frac{n+1}{n} \cdot \frac{n}{n-1} + \frac{n+1}{n} \cdot \frac{n}{n-1} \cdot \frac{n-1}{n-2} + \right. \\&\quad \left. \dots + \frac{n+1}{n} \cdot \frac{n}{n-1} \cdot \frac{n-1}{n-2} \dots \frac{4}{3} \right] \\&= 2 \left[1 + \frac{n+1}{n} + \frac{n+1}{n-1} + \frac{n+1}{n-2} + \dots + \frac{n+1}{3} \right]\end{aligned}$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Expanding, we get

$$\begin{aligned}T(n) &\leq 2 + \frac{n+1}{n} \left[2 + \frac{n}{n-1} \left[2 + \frac{n-1}{n-2} \left[\dots \frac{4}{3} \right] \right] \right] \\&= 2 \left[1 + \frac{n+1}{n} + \frac{n+1}{n} \cdot \frac{n}{n-1} + \frac{n+1}{n} \cdot \frac{n}{n-1} \cdot \frac{n-1}{n-2} + \right. \\&\quad \left. \dots + \frac{n+1}{n} \cdot \frac{n}{n-1} \cdot \frac{n-1}{n-2} \dots \frac{4}{3} \right] \\&= 2 \left[1 + \frac{n+1}{n} + \frac{n+1}{n-1} + \frac{n+1}{n-2} + \dots + \frac{n+1}{3} \right] \\&= 2(n+1) \left[\frac{1}{n+1} + \frac{1}{n} + \frac{1}{n-1} + \frac{1}{n-2} + \dots + \frac{1}{3} \right]\end{aligned}$$


A Not So Simple Full-history Recurrence Relation (Cont.)

Expanding, we get

$$\begin{aligned}T(n) &\leq 2 + \frac{n+1}{n} \left[2 + \frac{n}{n-1} \left[2 + \frac{n-1}{n-2} \left[\cdots \frac{4}{3} \right] \right] \right] \\&= 2 \left[1 + \frac{n+1}{n} + \frac{n+1}{n} \cdot \frac{n}{n-1} + \frac{n+1}{n} \cdot \frac{n}{n-1} \cdot \frac{n-1}{n-2} + \right. \\&\quad \left. \cdots + \frac{n+1}{n} \cdot \frac{n}{n-1} \cdot \frac{n-1}{n-2} \cdots \frac{4}{3} \right] \\&= 2 \left[1 + \frac{n+1}{n} + \frac{n+1}{n-1} + \frac{n+1}{n-2} + \cdots + \frac{n+1}{3} \right] \\&= 2(n+1) \left[\frac{1}{n+1} + \frac{1}{n} + \frac{1}{n-1} + \frac{1}{n-2} + \cdots + \frac{1}{3} \right] \\&= 2(n+1)(H(n+1) - 1.5);\end{aligned}$$


where $H(n) = 1 + \frac{1}{2} + \frac{1}{3} + \frac{1}{4} + \cdots + \frac{1}{n}$ is the Harmonic series.

Harmonic Series Approximation



$$H(n) = 1 + \frac{1}{2} + \cdots + \frac{1}{n} > \int_1^{n+1} \frac{dx}{x} = \ln(n+1) \quad \text{and}$$


Harmonic Series Approximation



$$H(n) = 1 + \frac{1}{2} + \cdots + \frac{1}{n} > \int_1^{n+1} \frac{dx}{x} = \ln(n+1) \quad \text{and}$$

$$H(n) = 1 + \left(\frac{1}{2} + \cdots + \frac{1}{n} \right) < 1 + \int_1^n \frac{dx}{x} = 1 + \ln(n).$$

Harmonic Series Approximation



$$H(n) = 1 + \frac{1}{2} + \cdots + \frac{1}{n} > \int_1^{n+1} \frac{dx}{x} = \ln(n+1) \quad \text{and}$$
$$H(n) = 1 + \left(\frac{1}{2} + \cdots + \frac{1}{n} \right) < 1 + \int_1^n \frac{dx}{x} = 1 + \ln(n).$$

Combining: $\ln(n+1) < H(n) < 1 + \ln(n).$

Harmonic Series Approximation (Cont.)

Let $\delta_n = H(n) - \ln n$.

Harmonic Series Approximation (Cont.)

Let $\delta_n = H(n) - \ln n$.

Then,

$$\ln(1 + 1/n) < H(n) - \ln n < 1, \text{ and}$$

Harmonic Series Approximation (Cont.)

Let $\delta_n = H(n) - \ln n$.

Then,

$$\ln(1 + 1/n) < H(n) - \ln n < 1, \text{ and}$$

$$\delta_n - \delta_{n+1} = (H(n) - \ln n) - (H(n+1) - \ln(n+1))$$

Harmonic Series Approximation (Cont.)

Let $\delta_n = H(n) - \ln n$.

Then,

$$\ln(1 + 1/n) < H(n) - \ln n < 1, \text{ and}$$

$$\begin{aligned}\delta_n - \delta_{n+1} &= (H(n) - \ln n) - (H(n+1) - \ln(n+1)) \\ &= \ln(n+1) - \ln n - \frac{1}{n+1}\end{aligned}$$

Harmonic Series Approximation (Cont.)

Let $\delta_n = H(n) - \ln n$.

Then,

$$\ln(1 + 1/n) < H(n) - \ln n < 1, \text{ and}$$

$$\begin{aligned}\delta_n - \delta_{n+1} &= (H(n) - \ln n) - (H(n+1) - \ln(n+1)) \\ &= \ln(n+1) - \ln n - \frac{1}{n+1} \\ &= \int_n^{n+1} \frac{dx}{x} - \frac{1}{n+1} > 0 \quad (\text{See the picture!}),\end{aligned}$$

i.e., δ_n is monotone decreasing.

Harmonic Series Approximation (Cont.)

Let $\delta_n = H(n) - \ln n$.

Then,

$$\ln(1 + 1/n) < H(n) - \ln n < 1, \text{ and}$$

$$\begin{aligned}\delta_n - \delta_{n+1} &= (H(n) - \ln n) - (H(n+1) - \ln(n+1)) \\ &= \ln(n+1) - \ln n - \frac{1}{n+1} \\ &= \int_n^{n+1} \frac{dx}{x} - \frac{1}{n+1} > 0 \quad (\text{See the picture!}),\end{aligned}$$

i.e., δ_n is monotone decreasing. Therefore δ_n converges and let

$$\gamma = \lim_{n \rightarrow \infty} \delta_n = \lim_{n \rightarrow \infty} (H(n) - \ln n).$$

Harmonic Series Approximation (Cont.)

Let $\delta_n = H(n) - \ln n$.

Then,

$$\ln(1 + 1/n) < H(n) - \ln n < 1, \text{ and}$$

$$\begin{aligned}\delta_n - \delta_{n+1} &= (H(n) - \ln n) - (H(n+1) - \ln(n+1)) \\ &= \ln(n+1) - \ln n - \frac{1}{n+1} \\ &= \int_n^{n+1} \frac{dx}{x} - \frac{1}{n+1} > 0 \quad (\text{See the picture!}),\end{aligned}$$

i.e., δ_n is monotone decreasing. Therefore δ_n converges and let

$$\gamma = \lim_{n \rightarrow \infty} \delta_n = \lim_{n \rightarrow \infty} (H(n) - \ln n).$$

$\gamma \approx 0.5772$ is called the **Euler constant** (Euler, 1735).

Harmonic Series Approximation (Cont.)

Let $\delta_n = H(n) - \ln n$.

Then,

$$\ln(1 + 1/n) < H(n) - \ln n < 1, \text{ and}$$

$$\begin{aligned}\delta_n - \delta_{n+1} &= (H(n) - \ln n) - (H(n+1) - \ln(n+1)) \\ &= \ln(n+1) - \ln n - \frac{1}{n+1} \\ &= \int_n^{n+1} \frac{dx}{x} - \frac{1}{n+1} > 0 \quad (\text{See the picture!}),\end{aligned}$$

i.e., δ_n is monotone decreasing. Therefore δ_n converges and let

$$\gamma = \lim_{n \rightarrow \infty} \delta_n = \lim_{n \rightarrow \infty} (H(n) - \ln n).$$

$\gamma \approx 0.5772$ is called the **Euler constant** (Euler, 1735).

$$\therefore H(n) \approx \ln n + \gamma.$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Using the Harmonic series approximation, we get

$$T(n) \leq 2(n+1)(H(n+1) - 1.5)$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Using the Harmonic series approximation, we get

$$\begin{aligned} T(n) &\leq 2(n+1)(H(n+1) - 1.5) \\ &\approx 2(n+1)(\ln n + \gamma - 1.5) + \mathcal{O}(1) \end{aligned}$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Using the Harmonic series approximation, we get

$$\begin{aligned}T(n) &\leq 2(n+1)(H(n+1) - 1.5) \\&\approx 2(n+1)(\ln n + \gamma - 1.5) + \mathcal{O}(1) \\&= \mathcal{O}(n \log n).\end{aligned}$$

A Not So Simple Full-history Recurrence Relation (Cont.)

Using the Harmonic series approximation, we get

$$\begin{aligned}T(n) &\leq 2(n+1)(H(n+1) - 1.5) \\&\approx 2(n+1)(\ln n + \gamma - 1.5) + \mathcal{O}(1) \\&= \mathcal{O}(n \log n).\end{aligned}$$

\therefore Quicksort is indeed quick on the average!!