

# ON FREENESS OF COMPACTLY INDUCED MOD $p$ REPRESENTATIONS OF $\mathrm{SL}_2(\mathbb{F})$

ARPAN DAS

**ABSTRACT.** Let  $p$  be a prime, and  $\mathbb{F}$  a non-archimedean local field with residue characteristic  $p$  and ring of integers  $\mathcal{O}_{\mathbb{F}}$ . Set  $G_{\mathbb{S}} := \mathrm{SL}_2(\mathbb{F})$  and  $K_0 := \mathrm{SL}_2(\mathcal{O}_{\mathbb{F}})$ . For a smooth irreducible  $\bar{\mathbb{F}}_p$ -representation  $\sigma$  of  $K_0$ , we study the structure of the compact induction  $\mathrm{ind}_{K_0}^{G_{\mathbb{S}}}(\sigma)$  as a left module over the standard spherical Hecke algebra  $\mathrm{End}_{G_{\mathbb{S}}}(\mathrm{ind}_{K_0}^{G_{\mathbb{S}}}(\sigma))$ . We prove that it is free and of infinite rank.

## 1. INTRODUCTION

The study of smooth mod- $p$  representations of  $p$ -adic groups has been a central theme in the mod- $p$  and  $p$ -adic Langlands programs. Following the pioneering work of Barthel–Livné [5] on  $\mathrm{GL}_2$ , much effort has gone into understanding the structure and classification of such representations, particularly the supersingular ones. Among the four types—characters, principal series, Steinberg, and supersingular—much is known about the first three, but supersingular representations remain more mysterious in general. For  $\mathrm{GL}_2(\mathbb{Q}_p)$ , Breuil [6] achieved a complete classification by constructing explicit models, but for general local fields and for groups of higher ranks, the picture remains incomplete and technically subtle. However, the analogue of Barthel–Livné’s classification for the group  $\mathrm{SL}_2(\mathbb{F})$  in the mod- $p$  setting was subsequently established in [1, 7]. Also, Herzig extended this classification for  $p$ -adic  $\mathrm{GL}_n$  in his pioneering work [9, 10]. More recently, this classification has been further generalized to connected reductive  $p$ -adic groups through the far-reaching work of Abe–Henniart–Herzig–Vignéras (see [2, 3]).

In all of the above works the structure of the compactly induced representations and their pro- $p$ -Iwahori invariants plays a crucial role. For example, if  $G$  is split reductive  $p$ -adic group and  $\sigma$  is a weight of some hyperspecial maximal compact subgroup  $K$  then  $\mathrm{ind}_K^G(\sigma)$  is torsion free as a module over  $\mathcal{H}(G, K, \sigma) := \mathrm{End}_G(\mathrm{ind}_K^G(\sigma))$  (see [9, Corollary 6.5]). When  $G$  has semisimple rank 1 and is split or quasi-split, say  $\mathrm{GL}_2, \mathrm{SL}_2$  or the unramified  $\mathrm{U}(2,1)$ , finer structural results are known. For instance, in the above three examples the explicit right action of the pro- $p$ -Iwahori Hecke algebra on the pro- $p$ -Iwahori invariants of the compactly induced representations can be computed (see [5] for  $\mathrm{GL}_2$ , [8] for  $\mathrm{SL}_2$ , and [13] for  $\mathrm{U}(2,1)$ ).

In the present paper we consider the  $p$ -adic group  $\mathrm{SL}_2$  and we show that the compactly induced representations as left modules over their spherical Hecke algebras are free and of infinite rank, thus refining [9, Corollary 6.5] in this special case. Our main theorem is as follows.

**Theorem 1.1** (Theorem 3.2). *The compactly induced representation  $\mathrm{ind}_{K_0}^{G_{\mathbb{S}}}(\sigma_{\bar{\tau}})$  is a free module of infinite rank over the spherical Hecke algebra  $\mathcal{H}(G_{\mathbb{S}}, K_0, \sigma_{\bar{\tau}})$ .*

Our result complements the analogous freeness results previously established for  $\mathrm{GL}_2$  (see [5, Section 5]) and for the unramified unitary group  $\mathrm{U}(2,1)$  (see [12]). There are, however, two notable distinctions. First, we isolate and formalize the key combinatorial–linear algebraic argument that underlies both proofs in [5, 12]; this is stated abstractly as Proposition 2.4. This conceptual clarification separates the structural core of the argument from the group-specific computations, making the mechanism of freeness more transparent and potentially applicable to other  $p$ -adic groups. Second, we give an explicit proof of the non-vanishing of the standard Hecke operator on the subspace of functions supported on the central vertex of the Bruhat–Tits tree of  $\mathrm{SL}_2$ . Although this fact is expected on general grounds, its verification requires a careful analysis of the Hecke action and, to our knowledge, has not appeared explicitly in the literature. It was used implicitly, for example, in the proof of the main theorem of [12].

Finally, we note that Lemma 3.5, together with [8, Propositions 4.9 and 4.10], provides a detailed description of the explicit Hecke actions on the pro- $p$ -Iwahori invariants of compactly induced representations of  $\mathrm{SL}_2$  in the mod  $p$  setting. The significance of this analysis is underscored both by the main theorem of the present paper and by the existence of Hecke eigenvalues for  $\mathrm{SL}_2$  (see [8, Proposition 4.14]), analogous to the  $\mathrm{GL}_2$  case [5, Proposition 32]; see also [4, Question 8].

**Outline of the proof of the main theorem.** The proof proceeds in two stages.

- (1) In subsection 2.6 we prove a general combinatorial–linear algebra assertion : if a vector space  $V$  over any field admits a graded decomposition  $V = \bigoplus_{k \geq 0} C_k$  and there is a linear operator  $T$  satisfying certain *injectivity*, *triangularity*, and *filtration compatibility* conditions, then we can choose subsets  $A_k \subset C_k$  inductively so that the family  $\{T^i(A_j) \mid 0 \leq i, j \leq n, i + j \leq n\}$  of sets are mutually disjoint and their union forms a basis of the truncated sum  $B_n := \bigoplus_{k \leq n} C_k$ .

This yields an explicit combinatorial construction of a free  $T$ -module basis  $\sqcup_{k \geq 0} A_k$  of  $V$ .

- (2) In the subsequent section we apply this framework to  $V = \mathrm{ind}_{K_0}^{\mathrm{GS}}(\sigma_{\vec{r}})$  and  $T = \tau$ . The relevant subspaces  $C_k := C_{k, \sigma_{\vec{r}}}$  are obtained from the Cartan-Iwahori decomposition

$$G_S = \bigsqcup_{n \in \mathbb{Z}} K_0 \alpha_0^n I_S(1).$$

See the next section for precise meaning of the notations. Then we use the explicit formula for the action of  $\tau$  on a standard function (eq. 2.4.1) to find the possible support points of the image of an element of  $C_k$  in the tree of  $\mathrm{SL}_2$ . This yields the inclusions

$$\tau(C_0) \subset C_1 \quad \text{and} \quad \tau(C_k) \subset C_{k-1} \oplus C_k \oplus C_{k+1} \quad \text{for } k \geq 1.$$

The injectivity of  $\tau : C_0 \rightarrow C_1$  is proved in the technical Lemma 3.4. Finally, we compute the explicit action of  $\tau$  on the pro- $p$ -Iwahori invariants and use this computation to establish the *filtration compatibility* condition of the abstract setup. The hypothesis of the Proposition 2.4 is satisfied, and the freeness follows.

## 2. PRELIMINARIES

**2.1. General notions.** We take  $p$  to be a prime throughout, and  $\bar{\mathbf{F}}_p$  a fixed algebraic closure of the finite field  $\mathbf{F}_p$  with  $p$  elements. All representations, unless otherwise mentioned, are considered over  $\bar{\mathbf{F}}_p$ . We recall some generalities on the abstract representation theory of locally profinite groups. We let  $G$  be any locally profinite group, and  $H$  some closed subgroup. A representation  $\pi$  of  $G$  is called *smooth* if every vector  $v \in \pi$  is fixed by some compact open subgroup of  $G$ . Let  $\sigma$  be a smooth representation of  $H$ . We consider the following space of functions :

$$\mathrm{IND}_H^G(\sigma) := \{f : G \rightarrow \sigma \mid f(hg) = \sigma(h)(f(g)), \forall g \in G, h \in H\}.$$

Then,  $G$  acts on  $\mathrm{IND}_H^G(\sigma)$  via  $(g \cdot f)(g') := f(g'g)$ . The *smooth part* of  $\mathrm{IND}_H^G(\sigma)$ , that is, vectors that have open stabilizers, is denoted by  $\mathrm{Ind}_H^G(\sigma)$ , and this subrepresentation is called the *smooth induction* of  $\sigma$ . The subrepresentation of  $\mathrm{Ind}_H^G(\sigma)$  consisting of functions  $f$  such that the image of its support  $\mathrm{Supp}(f)$  inside  $H \backslash G$  is compact (equivalently, finite, whenever  $H$  is also open) is denoted by  $\mathrm{c-Ind}_H^G(\sigma)$  or  $\mathrm{ind}_H^G(\sigma)$ , and is called the *compact induction* of  $\sigma$ .

In practice, whenever we use compact induction the subgroup  $H$  is typically considered to be open as well. So, for the remaining part of this subsection we take  $H$  to be an open subgroup of  $G$ . Then, by virtue of the  $H$ -linearity, the support of any  $f \in \mathrm{ind}_H^G(\sigma)$  can be written as a finite disjoint union of right  $H$ -cosets. We define some standard functions in  $\mathrm{ind}_H^G(\sigma)$ . For  $g \in G$  and  $v \in \sigma$  we define :

$$[g, v](x) := \begin{cases} \sigma(xg)(v) & \text{if } x \in Hg^{-1} \\ 0 & \text{otherwise} \end{cases}.$$

It can be checked that  $g \cdot [g', v] = [gg', v]$  and  $[gh, v] = [g, \sigma(h)(v)]$  for every  $g, g' \in \mathbf{G}$  and  $h \in \mathbf{H}$ . Also, any  $f \in \mathrm{ind}_{\mathbf{H}}^{\mathbf{G}}(\sigma)$  can be written as

$$f = \sum_{Hg \in \mathrm{Supp}(f)} [g^{-1}, f(g)].$$

**2.2. Some standard notations related to  $\mathrm{SL}_2(\mathbf{F})$ .** Let  $\mathbf{F}$  be a non-archimedean local field of residue characteristic  $p$ . We denote its valuation ring by  $\mathcal{O}_{\mathbf{F}}$  and its valuation ideal by  $\mathfrak{p}_{\mathbf{F}}$  and we fix a *uniformizer*  $\varpi_{\mathbf{F}}$  that generates  $\mathfrak{p}_{\mathbf{F}}$ . The residue field is  $k_{\mathbf{F}} := \mathcal{O}_{\mathbf{F}}/\mathfrak{p}_{\mathbf{F}}$  is a finite field of cardinality  $q$  which is a power of  $p$ . We set  $\mathbf{G}_{\mathbf{S}} := \mathrm{SL}_2(\mathbf{F})$  throughout this article. It is known that  $\mathbf{G}_{\mathbf{S}}$  has two maximal compact subgroups  $\mathbf{K}_0 := \mathrm{SL}_2(\mathcal{O}_{\mathbf{F}})$  and  $\mathbf{K}_1 = \alpha \mathbf{K}_0 \alpha^{-1}$  where  $\alpha := \mathrm{diag}(1, \varpi_{\mathbf{F}}) \in \mathrm{GL}_2(\mathbf{F})$ . Let  $\mathbf{I}_{\mathbf{S}}(1)$  denote the *pro- $p$ -Iwahori subgroup* of  $\mathbf{G}_{\mathbf{S}}$ . We have

$$\mathbf{I}_{\mathbf{S}}(1) = \begin{pmatrix} 1 + \mathfrak{p}_{\mathbf{F}} & \mathcal{O}_{\mathbf{F}} \\ \mathfrak{p}_{\mathbf{F}} & 1 + \mathfrak{p}_{\mathbf{F}} \end{pmatrix} \cap \mathbf{K}_0.$$

We finally let  $\mathbf{U}_{\mathbf{S}}(\mathfrak{p}_{\mathbf{F}}^n)$  (resp.  $\bar{\mathbf{U}}(\mathfrak{p}_{\mathbf{F}}^n)$ ) denote the upper triangular (resp. lower triangular) matrices in  $\mathbf{G}_{\mathbf{S}}$  with top right (resp. bottom left) entry in the fractional ideal  $\mathfrak{p}_{\mathbf{F}}^n$  for  $n \in \mathbf{Z}$ .

**2.3. Generators of the weights of  $\mathrm{SL}_2(\mathcal{O}_{\mathbf{F}})$ .** Let  $k_{\mathbf{F}} = \mathbf{F}_q$  and  $q = p^n$ . Then it is well known that the irreducible  $\bar{\mathbf{F}}_p$ -representations of  $\mathrm{SL}_2(\mathbf{F}_q)$  (or equivalently, smooth irreducible  $\bar{\mathbf{F}}_p$ -representations of  $\mathrm{SL}_2(\mathcal{O}_{\mathbf{F}})$ ) are precisely of the form

$$\mathrm{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2 := \mathrm{Sym}^{r_0} \bar{\mathbf{F}}_p^2 \otimes \cdots \otimes \mathrm{Sym}^{r_{n-1}} \bar{\mathbf{F}}_p^2,$$

where  $\vec{r} = (r_0, \dots, r_{n-1}) \in \{0, \dots, p-1\}^n$  and  $\mathrm{SL}_2(\mathbf{F}_q)$  acts on  $\mathrm{Sym}^{r_i} \bar{\mathbf{F}}_p^2 := \bigoplus_{l=0}^{r_i} \bar{\mathbf{F}}_p X^{r_i-l} Y^l$  via :

$$\begin{pmatrix} a & b \\ c & d \end{pmatrix} \cdot (X^{r_i-l} Y^l) := (a^{p^i} X + c^{p^i} Y)^{r_i-l} (b^{p^i} X + d^{p^i} Y)^l.$$

Consider the element  $X^{\vec{r}} := X^{r_0} \otimes X^{r_1} \otimes \cdots \otimes X^{r_{n-1}} \in \mathrm{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$ . The following result is well known. We will however give a proof to keep things self-contained, and also because some arguments in the proof will be used later.

**Proposition 2.1.** *The representation  $\mathrm{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$  is generated by  $X^{\vec{r}}$  as an  $\bar{\mathbf{F}}_p[\bar{\mathbf{U}}]$ -module, and the line  $\bar{\mathbf{F}}_p \cdot X^{\vec{r}}$  is the unique  $\bar{\mathbf{U}}$ -invariant line in  $\mathrm{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$ . Here,  $\bar{\mathbf{U}}$  (resp.  $\bar{\mathbf{U}}$ ) denotes the upper (resp. lower) unipotent subgroup of  $\mathrm{SL}_2(\mathbf{F}_q)$ .*

*Consequently, the set  $\{\bar{u} \cdot X^{\vec{r}} \mid \bar{u} \in \bar{\mathbf{U}}_{\mathbf{S}}(\mathcal{O}_{\mathbf{F}})\}$  spans  $\mathrm{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$ , and the line  $\bar{\mathbf{F}}_p \cdot X^{\vec{r}}$  is the unique  $\mathbf{I}_{\mathbf{S}}(1)$ -invariant line in  $\sigma_{\vec{r}}$ .*

*Proof.* Write  $r = r_0 + r_1 p + \cdots + r_{n-1} p^{n-1}$ ; then  $0 \leq r < q = p^n$ . Consider  $\mathrm{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$  as an  $\mathrm{SL}_2(\mathbf{F}_q)$ -subrepresentation of  $\mathrm{Sym}^r \bar{\mathbf{F}}_p^2$  via the map  $v_0 \otimes v_1 \otimes \cdots \otimes v_{n-1} \mapsto v_0 v_1^{p^1} \cdots v_{n-1}^{p^{n-1}}$ . Then, a basis of  $\mathrm{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$  is given by the family of monomials  $X^{\sum_{j=0}^{n-1} i_j p^j} Y^{r - (\sum_{j=0}^{n-1} i_j p^j)}$ , with  $0 \leq i_j \leq r_j$  for all  $j \in \{0, \dots, n-1\}$ .

Now, we recall that for  $i = i_0 + i_1 p + \cdots + i_{n-1} p^{n-1} \leq r = r_0 + r_1 p + \cdots + r_{n-1} p^{n-1}$ , the binomial coefficient  $\binom{r}{i}$  in  $\bar{\mathbf{F}}_p$  is given by

$$\binom{r}{i} = \begin{cases} \binom{r_0}{i_0} \binom{r_1}{i_1} \cdots \binom{r_{n-1}}{i_{n-1}} & \text{; if } i_l \leq r_l \text{ for all } l \\ 0 & \text{; otherwise} \end{cases}.$$

This simply follows by writing  $(X+1)^r \in \bar{\mathbf{F}}_p[X]$  as  $(X+1)^{r_0} (X^p+1)^{r_1} \cdots (X^{p^{n-1}}+1)^{r_{n-1}}$  and then expanding and comparing the binomial coefficients.

Consequently, the vector  $X^i Y^{r-i}$  (for  $0 \leq i \leq r$ ) is in the above basis if and only if  $\binom{r}{i} \neq 0$ . Therefore, for any  $\lambda \in \mathbf{F}_q$  we take  $\bar{u}(\lambda) \in \bar{\mathbf{U}}$ , and then we have

$$\bar{u}(\lambda) \cdot X^r = \sum_{i=0}^r \binom{r}{i} \lambda^{r-i} X^i Y^{r-i} \in \bar{\mathbf{F}}_p[\bar{\mathbf{U}}] \cdot X^r,$$

and the non-zero terms of the above sum correspond to those  $i$  for which  $\binom{r}{i} \neq 0$ . Since  $\lambda$  has  $q$  many possible values and  $r < q$ , an elementary argument in linear algebra shows that for each  $i \leq r$  with  $\binom{r}{i} \neq 0$ , we have  $X^i Y^{r-i} \in \bar{\mathbf{F}}_p[\bar{U}] \cdot X^r$ , and hence  $\text{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2 = \bar{\mathbf{F}}_p[\bar{U}] \cdot X^{\vec{r}}$ .

We now turn to the second statement to show that  $\bar{\mathbf{F}}_p \cdot X^{\vec{r}}$  is the unique  $\bar{U}$ -invariant line in  $\text{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$ . As before, we think of  $\text{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$  inside  $\text{Sym}^r \bar{\mathbf{F}}_p^2$ , and consider a vector  $v \in \text{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2 \setminus \{\bar{\mathbf{F}}_p \cdot X^{\vec{r}}\}$ . We can write  $v = \sum_{\vec{i} \preceq \vec{r}} \mu_{\vec{i}} X^{\sum_{j=0}^{n-1} i_j p^j} Y^{r - (\sum_{j=0}^{n-1} i_j p^j)}$ , where for the tuples  $\vec{i} = (i_0, \dots, i_{n-1})$  and  $\vec{r} = (r_0, \dots, r_{n-1})$  the notation  $\vec{i} \preceq \vec{r}$  means  $i_j \leq r_j$  for every  $j \in \{0, \dots, n-1\}$ , and  $\mu_{\vec{i}}$  are scalars such that  $\mu_{\vec{i}} \neq 0$  for some  $\vec{i}$ . Then, for any  $\lambda \in \mathbf{F}_q$ , we have for  $u(\lambda) \in \bar{U}$  the following

$$\begin{aligned} u(\lambda) \cdot v &= \sum_{\vec{i} \preceq \vec{r}} \mu_{\vec{i}} X^{\sum_{j=0}^{n-1} i_j p^j} (\lambda X + Y)^{r - (\sum_{j=0}^{n-1} i_j p^j)} \\ &= \sum_{\vec{i} \preceq \vec{r}} \sum_{\vec{j} + \vec{j} \preceq \vec{r}} \mu_{\vec{i}} \lambda^j \binom{r-i}{j} X^{i+j} Y^{r-(i+j)} \\ &= \sum_{\vec{i} \preceq \vec{r}} \lambda^i \underbrace{\sum_{\vec{j} + \vec{j} \preceq \vec{r}} \mu_{\vec{j}} \binom{r-j}{i} X^{i+j} Y^{r-(i+j)}}_{:= v_{\vec{i}}}, \end{aligned}$$

where in the second equality the integers  $i$  and  $j$  are the  $p$ -ary sums corresponding to the tuples  $\vec{i}$  and  $\vec{j}$  respectively, and in the third equality we have interchanged the tuples  $\vec{i}$  and  $\vec{j}$ . Now, since  $v \notin \bar{\mathbf{F}}_p \cdot X^{\vec{r}}$ , there is some tuple  $\vec{j}_0 \not\preceq \vec{r}$  such that  $\mu_{\vec{j}_0} \neq 0$ . We take such a  $\vec{j}_0$  with the corresponding  $p$ -ary sum  $j_0$  being least. Consider two distinct tuples  $\vec{i}_1, \vec{i}_2 \preceq \vec{r} - \vec{j}_0$  so that the polynomials  $v_{\vec{i}_1}$  and  $v_{\vec{i}_2}$  are non-zero and hence linearly independent (by comparing the monomials of least degree in  $X$ ). As a result, the subspace  $\bar{\mathbf{F}}_p[\bar{U}] \cdot v$  has dimension at least two.  $\square$

*Notation 2.2.* We write  $\sigma_{\vec{r}}$  for the representation  $\text{Sym}^{\vec{r}} \bar{\mathbf{F}}_p^2$ , and  $v_{\sigma_{\vec{r}}}$  for the generating vector  $X^{\vec{r}}$  throughout this article. When  $0 \leq r < p$  we will write  $r$  instead of  $\vec{r}$ .

*Remark 2.3.* In a similar manner one can show that  $Y^{\vec{r}}$  also generates  $\sigma_{\vec{r}}$  as an  $\bar{\mathbf{F}}_p[\bar{U}]$ -module, and the line  $\bar{\mathbf{F}}_p \cdot Y^{\vec{r}}$  is the unique  $\bar{U}$ -invariant (equivalently  $\text{I}_S(1)^{\bar{U}}$ -invariant) line in  $\sigma_{\vec{r}}$ . Hence, the set  $\{u \cdot (w_0 \cdot v_{\sigma_{\vec{r}}}) \mid u \in \text{U}_S(\mathcal{O}_F)\}$  spans  $\sigma_{\vec{r}}$ .

**2.4. Spherical Hecke algebra.** We recall that for a weight  $\sigma_{\vec{r}}$  of  $K_0 := \text{SL}_2(\mathcal{O}_F)$ , the *spherical Hecke algebra*  $\mathcal{H}(\text{G}_S, K_0, \sigma_{\vec{r}}) := \text{End}_{\text{G}_S}(\text{ind}_{K_0}^{\text{G}_S}(\sigma_{\vec{r}}))$  is generated by a single operator  $\tau$  as a polynomial algebra in one variable i.e.  $\text{End}_{\text{G}_S}(\text{ind}_{K_0}^{\text{G}_S}(\sigma_{\vec{r}})) = \bar{\mathbf{F}}_p[\tau]$ . In fact, we can compute the explicit action of  $\tau$  on a standard function  $[g, v] \in \text{ind}_{K_0}^{\text{G}_S}(\sigma_{\vec{r}})$ , as follows (see [1, Corollaire 3.12]) :

(2.4.1)

$$\tau([g, v]) = \sum_{\lambda \in k_F^2} \left[ g \left( \begin{pmatrix} 1 & A(\lambda) \\ 0 & 1 \end{pmatrix} \right) \alpha_0^{-1}, w_0 U_{\vec{r}} \sigma_{\vec{r}} \left( \begin{pmatrix} 0 & 1 \\ -1 & A(\lambda) \end{pmatrix} \right) v \right] + \sum_{\mu \in k_F} \left[ g \left( \begin{pmatrix} 1 & 0 \\ \varpi_F A(\mu) & 1 \end{pmatrix} \right) \alpha_0, U_{\vec{r}} v \right].$$

Here, for  $\lambda = (\lambda_0, \lambda_1, \dots, \lambda_{m-1}) \in k_F^m$  the notation  $A(\lambda) := \sum_{j=0}^{m-1} [\lambda_j] \varpi_F^j$ , and  $\alpha_0 := \text{diag}(\varpi_F^{-1}, \varpi_F)$ . The operator  $U_{\vec{r}} := U_{r_0} \otimes \dots \otimes U_{r_{n-1}} \in \text{End}_{\bar{\mathbf{F}}_p}(\sigma_{\vec{r}})$  is such that  $U_{r_j} \in \text{End}_{\bar{\mathbf{F}}_p}(\sigma_{r_j})$  and is defined as follows :

$$(2.4.2) \quad U_{r_j}(X^l Y^{r_j-l}) = \begin{cases} Y^{r_j} & ; l = 0 \\ 0 & ; l \neq 0 \end{cases}.$$

Now the compact induction  $\text{ind}_{K_0}^{\text{G}_S}(\sigma_{\vec{r}})$  is a left  $\mathcal{H}(\text{G}_S, K_0, \sigma_{\vec{r}})$ -module. The mod  $p$  irreducibles of  $K_1$  are denoted by  $\sigma_{\vec{r}}^\alpha$ . These have  $\sigma_{\vec{r}}$  as the underlying representation space on which an element  $k_1 = \alpha k_0 \alpha^{-1} \in K_1$  (where  $k_0 \in K_0$ ) acts by  $k_0$ . Since canonically we have  $\mathcal{H}(\text{G}_S, K_0, \sigma_{\vec{r}}) \simeq \mathcal{H}(\text{G}_S, K_1, \sigma_{\vec{r}}^\alpha)$  and  $\text{ind}_{K_1}^{\text{G}_S}(\sigma_{\vec{r}}^\alpha) \simeq \text{ind}_{K_0}^{\text{G}_S}(\sigma_{\vec{r}})^\alpha$  (see

[1, Propositions 3.6 and 3.23]) it suffices to consider only the representation  $\mathrm{ind}_{K_0}^{\mathrm{G}_S}(\sigma_{\bar{\tau}})$  as a left  $\mathcal{H}(\mathrm{G}_S, K_0, \sigma_{\bar{\tau}})$ -module. The  $\mathcal{H}(\mathrm{G}_S, K_1, \sigma_{\bar{\tau}}^\alpha)$ -module structure of  $\mathrm{ind}_{K_1}^{\mathrm{G}_S}(\sigma_{\bar{\tau}}^\alpha)$  is essentially same.

**2.5. Pro- $p$ -Iwahori invariants of  $\mathrm{ind}_{K_0}^{\mathrm{G}_S}(\sigma_{\bar{\tau}})$ .** We now recall some facts about the invariants  $\mathrm{ind}_{K_0}^{\mathrm{G}_S}(\sigma_{\bar{\tau}})^{\mathrm{I}_S(1)}$ . It is known that it has a basis consisting of functions  $\{f_n \mid n \in \mathbb{Z}\}$  such that the function  $f_n$  is supported on  $K_0\alpha_0^{-n}\mathrm{I}_S(1)$  and satisfies

$$f_n(\alpha_0^{-n}) = \begin{cases} w_0 \cdot v_{\sigma_{\bar{\tau}}} & ; n > 0 \\ v_{\sigma_{\bar{\tau}}} & ; n \leq 0 \end{cases}.$$

Now, as  $\mathrm{I}_S(1) = \mathrm{U}_S(\mathcal{O}_F) \times \mathrm{T}_S(1 + \mathfrak{p}_F) \times \bar{\mathrm{U}}_S(\mathfrak{p}_F)$ , and  $\alpha_0$  normalizes  $\mathrm{T}_S$ , we have  $K_0\alpha_0^{-n}\mathrm{I}_S(1) = K_0\alpha_0^{-n}\bar{\mathrm{U}}_S(\mathfrak{p}_F)$  and  $K_0\alpha_0^n\mathrm{I}_S(1) = K_0\alpha_0^n\mathrm{U}_S(\mathcal{O}_F)$  for  $n > 0$ . Therefore, we have the decompositions

$$K_0\alpha_0^{-n}\mathrm{I}_S(1) = \bigcup_{\bar{u} \in \bar{\mathrm{U}}_S(\mathfrak{p}_F)/\bar{\mathrm{U}}_S(\mathfrak{p}_F^{2^n})} K_0\alpha_0^{-n}\bar{u}$$

and

$$K_0\alpha_0^n\mathrm{I}_S(1) = \bigcup_{u \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^{2^n})} K_0\alpha_0^n u$$

for  $n > 0$ , and hence for  $n \in \mathbb{Z}$  we can write these  $f_n$  as follows :

$$(2.5.1) \quad f_n = \begin{cases} \sum_{\bar{u} \in \bar{\mathrm{U}}_S(\mathfrak{p}_F)/\bar{\mathrm{U}}_S(\mathfrak{p}_F^{2^n})} [\bar{u}\alpha_0^n, w_0 \cdot v_{\sigma_{\bar{\tau}}}] & ; n > 0 \\ \sum_{u \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^{2^n})} [u\alpha_0^n, v_{\sigma_{\bar{\tau}}}] & ; n \leq 0 \end{cases}.$$

**2.6. A technical result from linear algebra.** We will now prove a technical fact which is purely linear algebraic in nature. The conditions in the statement of the following proposition form the basis of the results we prove in the next section. The conclusion also plays a crucial role in the proof of the main result.

**Proposition 2.4.** *Let  $V$  be a vector space and  $T$  an operator on  $V$ . Suppose  $C_0, C_1, \dots$  are non-zero subspaces with  $\dim C_{k+1} > \dim C_k$  for all  $k \geq 0$  such that  $V = \bigoplus_{k \geq 0} C_k$  and satisfying the following properties :*

- (1)  $T(C_0) \subset C_1$  and  $T : C_0 \rightarrow C_1$  is injective.
- (2)  $T(C_k) \subset C_{k-1} \oplus C_k \oplus C_{k+1}$  for all  $k \geq 1$ .
- (3) Let  $B_n := \bigoplus_{k \leq n} C_k$  for  $n \geq 0$ . If  $f \in B_{n+1}$  and  $T(f) \in B_{n+1}$  then  $f \in B_n$ .

Then for a fixed  $n \geq 0$  we can choose non-empty subsets  $A_k \subset C_k$  for each  $k \leq n$  such that the sets  $\{T^i(A_j) \mid 0 \leq i, j \leq n, i+j \leq n\}$  are mutually disjoint and  $\bigsqcup_{i+j \leq n} T^i(A_j)$  is a basis of  $B_n$ .

*Proof.* Let  $\pi_m : V \rightarrow C_m$  be the projection maps. For a non-zero  $f \in V$  we define

$$\mathrm{top}(f) := \max\{m \mid \pi_m(f) \neq 0\}.$$

If  $\mathrm{top}(f) = 0$  then  $\mathrm{top}(T(f)) = 1$  by condition (1). Suppose  $n = \mathrm{top}(f) \geq 1$ . By condition (2) we have  $\mathrm{top}(T(f)) \leq n+1$ . But if  $\mathrm{top}(T(f)) \leq n$  then  $f \in B_n$  and  $T(f) \in B_n$  and by condition (3) we have  $f \in B_{n-1}$ . This is a contradiction to  $\mathrm{top}(f) = n$ . Thus  $\mathrm{top}(T(f)) = n+1$ . Thus for any  $i \geq 1$  and any  $f$  with  $\mathrm{top}(f) \geq 0$  we have  $\mathrm{top}(T^i(f)) = \mathrm{top}(f) + i$ . In particular, if  $S_m : C_m \rightarrow C_{m+1}$  is defined by  $S_m(c) := \pi_{m+1}(T(c))$ , then for  $c \neq 0$  we know  $\mathrm{top}(T(c)) = m+1$  and thus  $S_m$  is injective. Also since  $T$  strictly increases the top value, it is an injective linear map.

We take  $A_0 \subset C_0$  to be a basis. Since  $S_0 = T|_{C_0} : C_0 \rightarrow C_1$  is injective we can choose  $A_1 \subset C_1$  so that  $A_1 \sqcup T(A_0)$  is a basis of  $C_1$ . Thus  $|A_1| > 0$ . Now suppose  $n \geq 1$  and we have chosen  $A_j$  for all  $j \leq n$  so that  $\bigsqcup_{j+i \leq k} T^i(A_j)$  is a basis of  $B_k$  for all  $k \leq n$ . Now, note that the family  $\{T^{n+1-j}(A_j) \mid j \leq n\}$  is mutually disjoint. To see this take  $j < j_1 \leq n$  and suppose  $T^{n+1-j}(a) = T^{n+1-j_1}(b)$  for some  $a \in A_j$  and  $b \in A_{j_1}$ . This means  $T(T^{n-j}(a)) = T(T^{n-j_1}(b))$  and so we have  $T^{n-j}(a) = T^{n-j_1}(b)$  which is a contradiction by the induction

hypothesis. At first, we show that the following set

$$\bigsqcup_{i+j \leq n+1, j \leq n} T^i(A_j) = \bigsqcup_{i+j \leq n} T^i(A_j) \sqcup \bigsqcup_{j \leq n} T^{n+1-j}(A_j)$$

is linearly independent. We proceed by way of contradiction. Suppose we have a non-trivial linear combination of elements of the above set written as  $f_1 + f_2 = 0$  where  $f_1$  is a linear combination of elements from the first set and  $f_2$  is a linear combination of elements from the second set. We can write  $f_2$  as  $T(f'_2)$  where  $f'_2$  is a linear combination of elements from  $\bigsqcup_{j+i=n} T^i(A_j)$  with same coefficients that appear in the combination representing  $f_2$ . Thus  $f'_2 \in B_n$  and  $T(f'_2) = -f_1 \in B_n$  and by condition (3) we have  $f'_2 \in B_{n-1}$  since we have assumed  $n \geq 1$ . But  $\bigsqcup_{j+i \leq n} T^i(A_j) = \bigsqcup_{j+i \leq n-1} T^i(A_j) \sqcup \bigsqcup_{j+i=n} T^i(A_j)$  is a basis of  $B_n$  and  $\bigsqcup_{j+i \leq n-1} T^i(A_j)$  is a basis of  $B_{n-1}$  by the induction hypothesis. Therefore, all coefficients in the linear combination representing  $f'_2$  (and hence  $f_2$ ) are zero. Consequently, all coefficients representing  $f_1$  are also zero. Thus we have established the linear independence.

We will now construct  $A_{n+1}$  so that  $\bigsqcup_{j+i \leq n+1} T^i(A_j)$  is a basis of  $B_{n+1}$ . To this end we first observe that

$$\bigsqcup_{i+j=n+1, j \leq n} T^i(A_j) = T \left( \bigsqcup_{i+j=n, j \leq n} T^i(A_j) \right)$$

and

$$\pi_{n+1} \left( T \left( \bigsqcup_{i+j=n, j \leq n} T^i(A_j) \right) \right) \subset S_n(C_n) \subsetneq C_{n+1}.$$

We choose  $A_{n+1} \subset C_{n+1}$  so that it extends a maximal linearly independent subset of  $\pi_{n+1} \left( T \left( \bigsqcup_{i+j=n, j \leq n} T^i(A_j) \right) \right)$  to a basis of  $C_{n+1}$ . Thus  $|A_{n+1}| > 0$  and the set

$$\bigsqcup_{i+j \leq n} T^i(A_j) \sqcup \bigsqcup_{i+j=n+1, j \leq n} T^i(A_j) \sqcup A_{n+1}$$

forms a basis of  $B_{n+1} = B_n \oplus C_{n+1}$ . This completes the induction process.  $\square$

### 3. FREENESS OF $\text{ind}_{K_0}^{\text{Gs}}(\sigma_{\vec{r}})$ AS A HECKE MODULE

At first we introduce some new objects and notations. We follow the notations of [11, 12]. First recall that we have the well known Cartan-Iwahori decomposition :  $G_S = \bigsqcup_{n \in \mathbb{Z}} K_0 \alpha_0^n I_S(1)$ . For  $n \geq 0$ , we denote by  $R_n^+(\sigma_{\vec{r}})$  (resp.  $R_n^-(\sigma_{\vec{r}})$ ) the subspace of functions in  $\text{ind}_{K_0}^{\text{Gs}}(\sigma_{\vec{r}})$  which are supported on  $K_0 \alpha_0^n I_S(1) = K_0 \alpha_0^n U_S(\mathcal{O}_F)$  (resp.  $K_0 \alpha_0^{-(n+1)} I_S(1) = K_0 \alpha_0^{-(n+1)} \bar{U}_S(\mathfrak{p}_F)$ ). We write  $R_n^+(\sigma_{\vec{r}}) = [U_S(\mathcal{O}_F) \alpha_0^{-n}, \sigma_{\vec{r}}]$  for  $n \geq 0$ , and  $R_{n-1}^-(\sigma_{\vec{r}}) = [\bar{U}_S(\mathfrak{p}_F) \alpha_0^n, \sigma_{\vec{r}}]$  for  $n \geq 1$ . By convention  $R_{-1}^-(\sigma_{\vec{r}}) := R_0^+(\sigma_{\vec{r}})$ . We also set  $C_{0, \sigma_{\vec{r}}} := R_0^+(\sigma_{\vec{r}})$ ,  $C_{n, \sigma_{\vec{r}}} := R_n^+(\sigma_{\vec{r}}) \oplus R_{n-1}^-(\sigma_{\vec{r}})$  for  $n \geq 1$  and  $B_{n, \sigma_{\vec{r}}} := \bigoplus_{k \leq n} C_{k, \sigma_{\vec{r}}}$  for  $n \geq 0$ .

*Remark 3.1.* It is often useful to interpret the subspaces  $C_{n, \sigma_{\vec{r}}}$  and  $B_{n, \sigma_{\vec{r}}}$  in terms of the Bruhat-Tits tree of  $\text{SL}_2(F)$ . Note that  $C_{0, \sigma_{\vec{r}}}$  is the subspace of all functions supported on the central vertex corresponding to the subgroup  $K_0$ . For  $n \geq 1$  the subspace  $C_{n, \sigma_{\vec{r}}}$  consists of all functions supported on vertices which are at a distance of  $2n$  from the central vertex. Next, note that for  $n \geq 0$  both the spaces  $R_n^+(\sigma_{\vec{r}})$  and  $R_{n-1}^-(\sigma_{\vec{r}})$  are  $I_S(1)$ -stable. For  $n \geq 0$  we have  $f_{-n} \in R_n^+(\sigma_{\vec{r}})^{I_S(1)}$ , and for  $n \geq 1$  we have  $f_n \in R_{n-1}^-(\sigma_{\vec{r}})^{I_S(1)}$ . Also, every function in  $R_n^+(\sigma_{\vec{r}})^{I_S(1)}$  (resp.  $R_{n-1}^-(\sigma_{\vec{r}})^{I_S(1)}$ ) is determined by its value on  $\alpha_0^n$  (resp.  $\alpha_0^{-n}$ ). Hence, the spaces  $R_n^+(\sigma_{\vec{r}})^{I_S(1)}$  and  $R_{n-1}^-(\sigma_{\vec{r}})^{I_S(1)}$  are one dimensional and generated by  $f_{-n}$  and  $f_n$  respectively.

We now state the main theorem of this article.

**Theorem 3.2.** *The compactly induced representation  $\text{ind}_{K_0}^{\text{Gs}}(\sigma_{\vec{r}})$  is a free module of infinite rank over the spherical Hecke algebra  $\mathcal{H}(G_S, K_0, \sigma_{\vec{r}})$ .*

To prove this we will show that the Hecke operator  $\tau$  satisfies the conditions (1)-(3) of Proposition 2.4.

**3.1. Action of  $\tau$  on the subspaces  $C_{n,\sigma_{\bar{r}}}$ .** We will now show that the Hecke operator  $\tau$  satisfies the condition (1) and (2) of Proposition 2.4.

**Lemma 3.3.** *We have the following :*

- (i)  $\tau(R_0^+(\sigma_{\bar{r}})) \subseteq R_1^+(\sigma_{\bar{r}}) \oplus R_0^-(\sigma_{\bar{r}})$ .
- (ii)  $\tau(R_n^+(\sigma_{\bar{r}})) \subseteq R_{n-1}^+(\sigma_{\bar{r}}) \oplus R_n^+(\sigma_{\bar{r}}) \oplus R_{n+1}^+(\sigma_{\bar{r}})$ , for  $n \geq 1$ .
- (iii)  $\tau(R_n^-(\sigma_{\bar{r}})) \subseteq R_{n-1}^-(\sigma_{\bar{r}}) \oplus R_n^-(\sigma_{\bar{r}}) \oplus R_{n+1}^-(\sigma_{\bar{r}})$ , for  $n \geq 0$ .

Thus  $\tau(C_{0,\sigma_{\bar{r}}}) \subseteq C_{1,\sigma_{\bar{r}}}$  and  $\tau(C_{n,\sigma_{\bar{r}}}) \subseteq C_{n-1,\sigma_{\bar{r}}} \oplus C_{n,\sigma_{\bar{r}}} \oplus C_{n+1,\sigma_{\bar{r}}}$  for  $n \geq 1$ ,

*Proof.* (i) It suffices to compute the action of  $\tau$  on a standard function  $[u, v] \in R_0^+(\sigma_{\bar{r}})$ , where  $u \in U_S(\mathcal{O}_{\mathbb{F}})$ . Using the formula 2.4.1 the required containment follows from the simple observation that if  $\bar{u} \in \bar{U}_S(\mathfrak{p}_{\mathbb{F}})$ , then  $u\bar{u} \in I_S(1)$ , so that we can write  $u\bar{u} = \bar{u}_1 t u_2$ , where  $\bar{u}_1 \in \bar{U}_S(\mathfrak{p}_{\mathbb{F}})$ ,  $t \in T_S(1 + \mathfrak{p}_{\mathbb{F}})$ , and  $u_2 \in U_S(\mathcal{O}_{\mathbb{F}})$ . But then again we can write  $u_2 \alpha_0 = \alpha_0 u'_2$  with  $u'_2 \in U_S(\mathfrak{p}_{\mathbb{F}}^2) \subset K_0$ . Hence, the sum  $\sum_{\mu} [u\bar{u}(\varpi_{\mathbb{F}} A(\mu)) \alpha_0, U_{\bar{r}} v] \in R_0^-(\sigma_{\bar{r}})$ .

(ii) Again, take a standard function  $[u(x) \alpha_0^{-n}, v] \in R_n^+(\sigma_{\bar{r}})$  where  $u(x)$  denotes an upper unipotent matrix with the top right entry  $x \in \mathcal{O}_{\mathbb{F}}$ . Here, we look at the formula 2.4.1 and consider the elements of the form  $u(x) \alpha_0^{-n} u(A(\lambda)) \alpha_0^{-1}$  and  $u(x) \alpha_0^{-n} \bar{u}(\varpi_{\mathbb{F}} A(\mu)) \alpha_0$  to determine which  $U_S(\mathcal{O}_{\mathbb{F}})$ - $K_0$  double coset they belong to. At first, note that  $u(x) \alpha_0^{-n} u(A(\lambda)) \alpha_0^{-1} = u(x + \varpi_{\mathbb{F}}^{2n} A(\lambda)) \alpha_0^{-(n+1)} \in U_S(\mathcal{O}_{\mathbb{F}}) \alpha_0^{-(n+1)}$ . Next, for  $\mu \in k_{\mathbb{F}}$  we can easily see by using elementary row and column reduction that :

$$\alpha_0^{-n} \bar{u}(\varpi_{\mathbb{F}} A(\mu)) \alpha_0 = \begin{pmatrix} \varpi_{\mathbb{F}}^{n-1} & 0 \\ \varpi_{\mathbb{F}}^{-n} A(\mu) & \varpi_{\mathbb{F}}^{-n+1} \end{pmatrix} \in \begin{cases} U_S(\mathcal{O}_{\mathbb{F}}) \alpha_0^{-(n-1)} K_0 & ; \mu = 0 \\ U_S(\mathcal{O}_{\mathbb{F}}) \alpha_0^{-n} K_0 & ; \mu \neq 0 \end{cases}.$$

As a result we have  $\tau(R_n^+(\sigma_{\bar{r}})) \subseteq R_{n-1}^+(\sigma_{\bar{r}}) \oplus R_n^+(\sigma_{\bar{r}}) \oplus R_{n+1}^+(\sigma_{\bar{r}})$ , for  $n \geq 1$ .

(iii) As before we take a standard function  $[\bar{u}(y) \alpha_0^{n+1}, v] \in R_n^-(\sigma_{\bar{r}})$  where  $\bar{u}(y)$  denotes a lower unipotent matrix with the bottom left entry  $y \in \mathfrak{p}_{\mathbb{F}}$ . In this case also we look at the formula 2.4.1 and consider the elements  $\bar{u}(y) \alpha_0^{n+1} u(A(\lambda)) \alpha_0^{-1}$  and  $\bar{u}(y) \alpha_0^{n+1} \bar{u}(\varpi_{\mathbb{F}} A(\mu)) \alpha_0$ . Then we determine which  $\bar{U}_S(\mathfrak{p}_{\mathbb{F}})$ - $K_0$  double coset they belong to. First, note that  $\bar{u}(y) \alpha_0^{n+1} \bar{u}(\varpi_{\mathbb{F}} A(\mu)) \alpha_0 = \bar{u}(y + \varpi_{\mathbb{F}}^{2n+3} A(\mu)) \alpha_0^{n+2} \in \bar{U}_S(\mathfrak{p}_{\mathbb{F}}) \alpha_0^{n+2}$ . Next, for  $\lambda \in k_{\mathbb{F}}^2$  we have by row and column reduction the following :

$$\alpha_0^{n+1} u(A(\lambda)) \alpha_0^{-1} = \begin{pmatrix} \varpi_{\mathbb{F}}^{-n} & \varpi_{\mathbb{F}}^{-n-2} A(\lambda) \\ 0 & \varpi_{\mathbb{F}}^n \end{pmatrix} \in \begin{cases} \bar{U}_S(\mathfrak{p}_{\mathbb{F}}) \alpha_0^{n+2} K_0 & ; A(\lambda) \in \mathcal{O}_{\mathbb{F}}^{\times} \\ \bar{U}_S(\mathfrak{p}_{\mathbb{F}}) \alpha_0^{n+1} K_0 & ; A(\lambda) \in \mathfrak{p}_{\mathbb{F}} \setminus \mathfrak{p}_{\mathbb{F}}^2 \\ \bar{U}_S(\mathfrak{p}_{\mathbb{F}}) \alpha_0^n K_0 & ; A(\lambda) \in \mathfrak{p}_{\mathbb{F}}^2 \end{cases}.$$

Hence, we have  $\tau(R_n^-(\sigma_{\bar{r}})) \subseteq R_{n-1}^-(\sigma_{\bar{r}}) \oplus R_n^-(\sigma_{\bar{r}}) \oplus R_{n+1}^-(\sigma_{\bar{r}})$ , for  $n \geq 0$ . □

**Lemma 3.4.** *The map  $\tau : C_{0,\sigma_{\bar{r}}} \rightarrow C_{1,\sigma_{\bar{r}}}$  is injective.*

*Proof.* Note that  $\bar{U}_S(\mathcal{O}_{\mathbb{F}})$ -translates of  $[I, X^{\bar{r}}]$  generates the space  $C_{0,\sigma_{\bar{r}}}$ , by Proposition 2.1. Since the right hand side of the formula 2.4.1 has standard functions supported on distinct cosets, it suffices to show that for any non-zero  $v \in \sigma_{\bar{r}}$  there exists some  $\lambda \in k_{\mathbb{F}}$  such that

$$U_{\bar{r}} \left( \sigma_{\bar{r}} \left( \begin{pmatrix} 0 & 1 \\ -1 & A(\lambda) \end{pmatrix} \right) v \right) \neq 0.$$

As in the proof of Proposition 2.1 we consider  $\mathrm{Sym}^{\bar{r}} \bar{\mathbf{F}}_p^2$  as an  $\mathrm{SL}_2(k_{\mathbb{F}})$ -subrepresentation of  $\mathrm{Sym}^{\bar{r}} \bar{\mathbf{F}}_p^2$  via the map  $v_0 \otimes v_1 \otimes \cdots \otimes v_{n-1} \mapsto v_0 v_1^p \cdots v_{n-1}^{p^{n-1}}$  and work with the monomial basis

$$\left\{ X^{r-i} Y^i : i = \sum_{j=0}^{n-1} i_j p^j \leq r = \sum_{j=0}^{n-1} r_j p^j, i_j, r_j \in \{0, \dots, p-1\}, i_j \leq r_j \text{ for every } j \right\}.$$

Now we write  $v = \sum_i c_i X^{r-i} Y^i$  as a linear combination with the admissible monomials and coefficients  $c_i \in \bar{\mathbf{F}}_p$  not all zero. For any  $a \in k_{\mathbb{F}}$  the coefficient of  $X^r$  in  $u(a) \cdot v$  is  $P_v(a) = \sum_i c_i a^i$  which is a polynomial (in  $a$ ) of degree  $r < q$  with coefficients  $c_i \in \bar{\mathbf{F}}_p$  not all zero. Thus there exists some  $a \in k_{\mathbb{F}}$  so that  $P_v(a) \neq 0$ . We let



$A(\lambda) := [a]$  where  $[\cdot]$  denotes the Teichmüller lift as usual. Therefore, the coefficient of  $X^r$  in  $u(A(\lambda)) \cdot v$  is  $P_v(a)$  which is non-zero. Hence, the  $Y^r$  coefficient of

$$\begin{pmatrix} 0 & 1 \\ -1 & A(\lambda) \end{pmatrix} \cdot v = w_0 u(A(\lambda)) \cdot v$$

is non-zero. So  $U_{\vec{r}} \left( \sigma_{\vec{r}} \left( \begin{pmatrix} 0 & 1 \\ -1 & A(\lambda) \end{pmatrix} \right) v \right) \neq 0$  by eq. 2.4.2 as required.  $\square$

**3.2. Action of  $\tau$  strictly increases the top value.** Next, we will show that  $\tau$  satisfies the condition (3) of Proposition 2.4. But we first need to explicitly compute the action of  $\tau$  on  $\text{ind}_{K_0}^{\text{Gs}}(\sigma_{\vec{r}})^{\text{Is}(1)}$  using the explicit formulae 2.4.1 and 2.5.1.

**Lemma 3.5.** *We have*

- (i)  $\tau(f_0) = f_{-1} + \lambda_{\sigma_{\vec{r}}} f_1$ , where  $\lambda_{\sigma_{\vec{r}}} = \begin{cases} 0 & ; \vec{r} \neq \vec{0} \\ 1 & ; \vec{r} = \vec{0} \end{cases}$ .
- (ii) For  $n \neq 0$  we have  $\tau(f_n) = c_n f_n + f_{n+\delta(n)}$ , where  $c_n$  is a scalar and  $\delta(n) = \begin{cases} 1 & ; n > 0 \\ -1 & ; n < 0 \end{cases}$ .

*Proof.* (i) At first, we consider the case when  $\sigma_{\vec{r}} \neq 1$ . Then  $f_0 = [I_2, X^{\vec{r}}]$ , and we have :

$$\begin{aligned} \tau(f_0) &= \tau([I_2, X^{\vec{r}}]) \\ &= \sum_{\lambda \in k_{\mathbb{F}}^2} \left[ \begin{pmatrix} 1 & A(\lambda) \\ 0 & 1 \end{pmatrix} \alpha_0^{-1}, w_0 U_{\vec{r}} \sigma_{\vec{r}} \left( \begin{pmatrix} 0 & 1 \\ -1 & A(\lambda) \end{pmatrix} \right) (X^{\vec{r}}) \right] + \sum_{\mu \in k_{\mathbb{F}}} \left[ \begin{pmatrix} 1 & 0 \\ \varpi_{\mathbb{F}} A(\mu) & 1 \end{pmatrix} \alpha_0, U_{\vec{r}} X^{\vec{r}} \right] \\ &= \sum_{\lambda \in k_{\mathbb{F}}^2} \left[ \begin{pmatrix} 1 & A(\lambda) \\ 0 & 1 \end{pmatrix} \alpha_0^{-1}, w_0 U_{\vec{r}} ((-1)^r Y^{\vec{r}}) \right] \\ &= \sum_{\lambda \in k_{\mathbb{F}}^2} \left[ \begin{pmatrix} 1 & A(\lambda) \\ 0 & 1 \end{pmatrix} \alpha_0^{-1}, (-1)^r w_0 \cdot Y^{\vec{r}} \right] \\ &= \sum_{\lambda \in k_{\mathbb{F}}^2} \left[ \begin{pmatrix} 1 & A(\lambda) \\ 0 & 1 \end{pmatrix} \alpha_0^{-1}, (-1)^{2r} X^{\vec{r}} \right] \\ &= f_{-1}. \end{aligned}$$

Next, let  $\sigma_{\vec{r}} = 1$ . Then, we have :

$$\begin{aligned} \tau(f_0) &= \tau([I_2, 1]) \\ &= \sum_{\lambda \in k_{\mathbb{F}}^2} \begin{pmatrix} 1 & A(\lambda) \\ 0 & 1 \end{pmatrix} \alpha_0^{-1} \cdot [I_2, 1] + \sum_{\mu \in k_{\mathbb{F}}} \begin{pmatrix} 1 & 0 \\ \varpi_{\mathbb{F}} A(\mu) & 1 \end{pmatrix} \alpha_0 \cdot [I_2, 1] \\ &= \sum_{u \in U_{\mathbb{S}}(\mathcal{O}_{\mathbb{F}})/U_{\mathbb{S}}(\mathfrak{p}_{\mathbb{F}}^2)} [u \alpha_0^{-1}, 1] + \sum_{\bar{u} \in \bar{U}_{\mathbb{S}}(\mathfrak{p}_{\mathbb{F}})/\bar{U}_{\mathbb{S}}(\mathfrak{p}_{\mathbb{F}}^2)} [\bar{u} \alpha_0, 1] \\ &= f_1 + f_{-1}. \end{aligned}$$

(ii) At first, we take  $n = -m$  to be a negative integer. We will show that

$$\tau(f_n) = c_n f_n + f_{n-1}$$

for some  $c_n \in \bar{\mathbb{F}}_p$ . Now,  $f_{-m}$  is  $\text{Is}(1)$ -invariant and as  $\tau$  is a  $\text{Gs}$ -intertwiner,  $\tau(f_{-m})$  is also  $\text{Is}(1)$ -invariant. Since each  $R_j^+(\sigma_{\vec{r}})$  for  $j \geq 0$  is  $\text{Is}(1)$ -stable we have

$$\tau(f_{-m}) \in (R_{m-1}^+(\sigma_{\vec{r}}) \oplus R_m^+(\sigma_{\vec{r}}) \oplus R_{m+1}^+(\sigma_{\vec{r}}))^{\text{Is}(1)} = R_{m-1}^+(\sigma_{\vec{r}})^{\text{Is}(1)} \oplus R_m^+(\sigma_{\vec{r}})^{\text{Is}(1)} \oplus R_{m+1}^+(\sigma_{\vec{r}})^{\text{Is}(1)},$$



and so we can write

$$\tau(f_{-m}) = c_{m-1}f_{-m+1} + c_m f_{-m} + c_{m+1}f_{-m-1}$$

for some  $c_{m-1}, c_m, c_{m+1} \in \bar{\mathbb{F}}_p$ . Now, by the formula 2.5.1 and part (1), we have

$$\tau(f_{-m}) = \tau\left(\sum_{u \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^{2m})} u\alpha_0^{-m} \cdot f_0\right) = \sum_{u \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^{2m})} u\alpha_0^{-m}(f_{-1} + \lambda_{\bar{r}}f_1).$$

So, to find the constants  $c_{m-1}$  and  $c_{m+1}$  we evaluate  $\tau(f_{-m})$  above at  $\alpha_0^{m-1}$  and  $\alpha_0^{m+1}$ . For this we at first need to find which  $K_0\text{-I}_S(1)$  double coset the elements  $\alpha_0^{m-1}u\alpha_0^{-m}$  and  $\alpha_0^{m+1}u\alpha_0^{-m}$  lie in, for  $u \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^{2m})$ . Let  $u$  be an upper unipotent with the top right entry  $x \in \mathcal{O}_F/\mathfrak{p}_F^{2m}$ , and we have

$$\alpha_0^{m-1} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \alpha_0^{-m} = \begin{pmatrix} \varpi_F & x\varpi_F^{1-2m} \\ 0 & \varpi_F^{-1} \end{pmatrix} \in \begin{cases} K_0\alpha_0^{-1}\mathrm{I}_S(1) & ; x \in \mathfrak{p}_F^{2m-2} \\ K_0\alpha_0^{l(x)}\mathrm{I}_S(1) & ; x \in \mathcal{O}_F \setminus \mathfrak{p}_F^{2m-2} \end{cases},$$

where  $l(x) < -1$ ; the above containments can be seen by elementary row and column reduction. For instance if  $x = a\varpi_F^{2m-2}$  with  $a \in \mathcal{O}_F$  then :

$$\begin{pmatrix} 1 & -a \\ 0 & 1 \end{pmatrix} \begin{pmatrix} \varpi_F & a\varpi_F^{-1} \\ 0 & \varpi_F^{-1} \end{pmatrix} = \alpha_0^{-1},$$

and if  $x = a\varpi_F^{2m-d} \in \mathcal{O}_F$  with  $d \geq 3$  and  $a \in \mathcal{O}_F^\times$  then :

$$\begin{pmatrix} a & 0 \\ 0 & a^{-1} \end{pmatrix} \begin{pmatrix} 0 & -1 \\ 1 & 0 \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -a^{-1}\varpi_F^{d-2} & 1 \end{pmatrix} \begin{pmatrix} \varpi_F & a\varpi_F^{1-d} \\ 0 & \varpi_F^{-1} \end{pmatrix} \begin{pmatrix} 1 & 0 \\ -a^{-1}\varpi_F^d & 1 \end{pmatrix} = \alpha_0^{1-d}.$$

Then, we have

$$\begin{aligned} \tau(f_{-m})(\alpha_0^{m-1}) &= \sum_{u \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^{2m})} f_{-1}(\alpha_0^{m-1}u\alpha_0^{-m}) + \lambda_{\bar{r}} \sum_{u \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^{2m})} f_1(\alpha_0^{m-1}u\alpha_0^{-m}) \\ &= \lambda_{\bar{r}} \sum_{u \in \mathrm{U}_S(\mathfrak{p}_F^{2m-2})/\mathrm{U}_S(\mathfrak{p}_F^{2m})} f_1 \left( \underbrace{\alpha_0^{m-1}u\alpha_0^{-m}}_{\text{write as } u_1\alpha_0^{-1} \text{ with } u_1 \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^2)} \right) \\ &= \lambda_{\bar{r}} \sum_{u_1 \in \mathrm{U}_S(\mathcal{O}_F)/\mathrm{U}_S(\mathfrak{p}_F^2)} f_1(u_1\alpha_0^{-1}) \\ &= \lambda_{\bar{r}} \sum_{\mu \in k_F^2} \begin{pmatrix} 1 & [\mu_0] + [\mu_1]\varpi_F \\ 0 & 1 \end{pmatrix} \cdot f_1(\alpha_0^{-1}) \\ &= \lambda_{\bar{r}} \sum_{\mu \in k_F^2} \begin{pmatrix} 1 & [\mu_0] + [\mu_1]\varpi_F \\ 0 & 1 \end{pmatrix} \cdot (w_0 \cdot v_{\sigma_{\bar{r}}}) \\ &= \lambda_{\bar{r}} \sum_{(\mu_0, \mu_1) \in k_F \times k_F} \begin{pmatrix} 1 & [\mu_0] \\ 0 & 1 \end{pmatrix} \cdot (w_0 \cdot v_{\sigma_{\bar{r}}}) \\ &= 0. \end{aligned}$$

Hence,  $c_{m-1} = 0$ . Next, we take  $u$  to be an upper unipotent with the top right entry  $x \in \mathcal{O}_F/\mathfrak{p}_F^{2m}$ , and we have

$$\alpha_0^{m+1} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \alpha_0^{-m} = \begin{pmatrix} \varpi_F^{-1} & x\varpi_F^{-2m-1} \\ 0 & \varpi_F \end{pmatrix} \in \begin{cases} K_0\alpha_0\mathrm{I}_S(1) & ; x \in \mathfrak{p}_F^{2m} \\ K_0\alpha_0^{l'(x)}\mathrm{I}_S(1) & ; x \in \mathcal{O}_F \setminus \mathfrak{p}_F^{2m} \end{cases},$$

where  $l'(x) < -1$ . The above containments can be proved as before. Therefore, we have :

$$\begin{aligned}\tau(f_{-m})(\alpha_0^{m+1}) &= \sum_{u \in U_S(\mathcal{O}_F)/U_S(\mathfrak{p}_F^{2m})} f_{-1}(\alpha_0^{m+1} u \alpha_0^{-m}) + \lambda_{\bar{\tau}} \sum_{u \in U_S(\mathcal{O}_F)/U_S(\mathfrak{p}_F^{2m})} f_1(\alpha_0^{m+1} u \alpha_0^{-m}) \\ &= f_{-1}(\alpha_0) = v_{\sigma_{\bar{\tau}}}.\end{aligned}$$

Hence,  $c_{m+1} = 1$ .

Next, we consider the case when  $n = m + 1$  is positive i.e.  $m \geq 0$ . Then,  $f_n = f_{m+1} \in R_m^-(\sigma_{\bar{\tau}})$ , hence

$$\tau(f_{m+1}) = c_{m-1}f_m + c_{m+1}f_{m+1} + c_{m+2}f_{m+2},$$

and we want to show that  $c_{m-1} = 0$  and  $c_{m+2} = 1$ . Now, by the formula 2.5.1 and part (1), we get

$$\begin{aligned}\tau(f_{m+1}) &= \tau\left(\sum_{\bar{u} \in \bar{U}_S(\mathfrak{p}_F)/\bar{U}_S(\mathfrak{p}_F^{2m+2})} [\bar{u}\alpha_0^{m+1}, w_0 \cdot v_{\sigma_{\bar{\tau}}}] \right) = \sum_{\bar{u} \in \bar{U}_S(\mathfrak{p}_F)/\bar{U}_S(\mathfrak{p}_F^{2m+2})} \bar{u}\alpha_0^{m+1} w_0 \tau(f_0) \\ &= \sum_{\bar{u} \in \bar{U}_S(\mathfrak{p}_F)/\bar{U}_S(\mathfrak{p}_F^{2m+2})} \bar{u}\alpha_0^{m+1} w_0 (f_{-1} + \lambda_{\sigma_{\bar{\tau}}} f_1)\end{aligned}$$

So, to find the constants  $c_{m-1}$  and  $c_{m+2}$  we evaluate  $\tau(f_{m+1})$  at  $\alpha_0^{-m}$  and  $\alpha_0^{-m-2}$  respectively. As before, we at first find which  $K_0$ - $I_S(1)$  double coset the elements  $\alpha_0^m u \alpha_0^{-m-1}$  and  $\alpha_0^{m+2} u \alpha_0^{-m-1}$  lie in, for  $u \in U_S(\mathfrak{p}_F)/U_S(\mathfrak{p}_F^{2m+2})$ . Let  $u$  be an upper unipotent with the top right entry  $x \in \mathfrak{p}_F/\mathfrak{p}_F^{2m+2}$ . Then, we have

$$\alpha_0^m \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \alpha_0^{-m-1} = \begin{pmatrix} \varpi_F & x\varpi_F^{-2m-1} \\ 0 & \varpi_F^{-1} \end{pmatrix} \in \begin{cases} K_0 \alpha_0^{-1} I_S(1) & ; x \in \mathfrak{p}_F^{2m} \\ K_0 \alpha_0^{l(x)} I_S(1) & ; x \in \mathfrak{p}_F \setminus \mathfrak{p}_F^{2m} \end{cases},$$

where  $l(x) < -1$ . As a result we have

$$\begin{aligned}\tau(f_{m+1})(\alpha_0^{-m}) &= \sum_{\bar{u} \in \bar{U}_S(\mathfrak{p}_F)/\bar{U}_S(\mathfrak{p}_F^{2m+2})} f_{-1}(\alpha_0^{-m} \bar{u} \alpha_0^{m+1} w_0) + \lambda_{\bar{\tau}} \sum_{\bar{u} \in \bar{U}_S(\mathfrak{p}_F)/\bar{U}_S(\mathfrak{p}_F^{2m+2})} f_1(\alpha_0^{-m} \bar{u} \alpha_0^{m+1} w_0) \\ &= \lambda_{\bar{\tau}} \sum_{u \in U_S(\mathfrak{p}_F^{2m})/U_S(\mathfrak{p}_F^{2m+2})} w_0 \cdot f_1 \left( \underbrace{\alpha_0^m u \alpha_0^{-m-1}}_{\text{write as } u_1 \alpha_0^{-1} \text{ with } u_1 \in U_S(\mathcal{O}_F)/U_S(\mathfrak{p}_F^2)} \right) \\ &= \lambda_{\bar{\tau}} \sum_{u_1 \in U_S(\mathcal{O}_F)/U_S(\mathfrak{p}_F^2)} (w_0 u_1 w_0) \cdot v_{\sigma_{\bar{\tau}}} \\ &= \lambda_{\bar{\tau}} \sum_{(\mu_0, \mu_1) \in k_F^2} \left( w_0 \begin{pmatrix} 1 & [\mu_0] \\ 0 & 1 \end{pmatrix} w_0 \right) \cdot v_{\sigma_{\bar{\tau}}} = 0,\end{aligned}$$

and hence  $c_{m-1} = 0$ . Next, for  $u$  an upper unipotent with the top right entry  $x \in \mathfrak{p}_F/\mathfrak{p}_F^{2m+2}$ , we have

$$\alpha_0^{m+2} \begin{pmatrix} 1 & x \\ 0 & 1 \end{pmatrix} \alpha_0^{-m-1} = \begin{pmatrix} \varpi_F^{-1} & x\varpi_F^{-2m-3} \\ 0 & \varpi_F \end{pmatrix} \in \begin{cases} K_0 \alpha_0 I_S(1) & ; x \in \mathfrak{p}_F^{2m+2} \\ K_0 \alpha_0^{l'(x)} I_S(1) & ; x \in \mathfrak{p}_F \setminus \mathfrak{p}_F^{2m+2} \end{cases},$$

where  $l'(x) < -1$ . Hence, we get

$$\tau(f_{m+1})(\alpha_0^{-m-2}) = f_{-1}(\alpha_0^{-m-2} \alpha_0^{m+1} w_0) = w_0 \cdot f_{-1}(\alpha_0) = w_0 \cdot v_{\sigma_{\bar{\tau}}},$$

that is to say,  $c_{m+2} = 1$  as required. This completes the proof.  $\square$

Finally we are ready to show that  $\tau$  satisfies the condition (3) of Proposition 2.4. Given the results we have established, the proof is a formal argument essentially same as [12, Lemma 4.4].

**Lemma 3.6.** *For  $n \geq 0$ , let  $f \in B_{n+1, \sigma_{\bar{\tau}}}$ . If  $\tau(f) \in B_{n+1, \sigma_{\bar{\tau}}}$  then  $f \in B_{n, \sigma_{\bar{\tau}}}$ .*

*Proof.* Let  $M_{n+1, \sigma_{\bar{\tau}}}$  be the subspace of  $B_{n+1, \sigma_{\bar{\tau}}}$  consisting of functions  $f$  such that  $\tau(f) \in B_{n+1, \sigma_{\bar{\tau}}}$ . So we have to show that  $M_{n+1, \sigma_{\bar{\tau}}} \subset B_{n, \sigma_{\bar{\tau}}}$ .

Suppose for contradiction that there exists some  $f \in M_{n+1, \sigma_{\bar{r}}} \setminus B_{n, \sigma_{\bar{r}}}$ . Then  $f \in B_{n+1, \sigma_{\bar{r}}} = B_{n, \sigma_{\bar{r}}} \oplus C_{n+1, \sigma_{\bar{r}}}$  and so we write  $f = f' + f''$  for some  $f' \in B_{n, \sigma_{\bar{r}}}$ ,  $f'' \in C_{n+1, \sigma_{\bar{r}}}$ . Now note that  $B_{n, \sigma_{\bar{r}}} \subset M_{n+1, \sigma_{\bar{r}}}$  since  $B_{n, \sigma_{\bar{r}}} \subset B_{n+1, \sigma_{\bar{r}}}$  and  $\tau(B_{n, \sigma_{\bar{r}}}) \subset B_{n+1, \sigma_{\bar{r}}}$  by Lemma 3.3. Hence  $f'' = f - f' \in M_{n+1, \sigma_{\bar{r}}}$ . But since  $f \notin B_{n, \sigma_{\bar{r}}}$  we have  $f'' \neq 0$ . Thus  $C_{n+1, \sigma_{\bar{r}}} \cap M_{n+1, \sigma_{\bar{r}}} \neq \emptyset$ .

By Remark 3.1 we know that  $C_{n+1, \sigma_{\bar{r}}}$  is  $\mathrm{Is}(1)$ -stable for  $n \geq 0$  and so  $B_{n+1, \sigma_{\bar{r}}}$  is  $\mathrm{Is}(1)$ -stable for  $n \geq 0$ . Also since  $\tau$  is  $G_S$ -invariant  $M_{n+1, \sigma_{\bar{r}}}$  is  $\mathrm{Is}(1)$ -stable. But since  $\mathrm{Is}(1)$  is a pro- $p$  group there exists some non-zero vector  $f^* \in C_{n+1, \sigma_{\bar{r}}} \cap M_{n+1, \sigma_{\bar{r}}}$  which is fixed by  $\mathrm{Is}(1)$ . Now  $C_{n+1, \sigma_{\bar{r}}} := R_{n+1}^+(\sigma_{\bar{r}}) \oplus R_n^-(\sigma_{\bar{r}})$  and hence  $C_{n+1, \sigma_{\bar{r}}}^{\mathrm{Is}(1)} = R_{n+1}^+(\sigma_{\bar{r}})^{\mathrm{Is}(1)} \oplus R_n^-(\sigma_{\bar{r}})^{\mathrm{Is}(1)} = \bar{\mathbf{F}}_p \cdot f_{-(n+1)} \oplus \bar{\mathbf{F}}_p \cdot f_{n+1}$  by Remark 3.1. We write  $f^* = \lambda f_{-(n+1)} + \mu f_{n+1}$  and using Lemma 3.5(ii) we obtain that  $\tau(f^*)$  is of the form

$$c_{-(n+1)} f_{-(n+1)} + d_{n+1} f_{n+1} + f_{-(n+2)} + f_{n+2}, \quad \text{where } c_{-(n+1)}, d_{n+1} \text{ are scalars.}$$

Hence  $\tau(f^*)$  does not lie in  $B_{n+1, \sigma_{\bar{r}}}$  because  $f_{-(n+2)}, f_{n+2} \in C_{n+2, \sigma_{\bar{r}}}$ . Thus we have a contradiction.  $\square$

We now complete the proof of our main theorem.

*Proof of Theorem 3.2.* To apply Proposition 2.4 we take  $V := \mathrm{ind}_{K_0}^{G_S}(\sigma_{\bar{r}})$  and  $C_n := C_{n, \sigma_{\bar{r}}}$  for each  $n \geq 0$ . The operator  $T := \tau$ . By Lemmas 3.3, 3.4, and 3.6 we have ensured that the conditions (1)-(3) of Proposition 2.4 are satisfied. Thus we can choose non-empty subsets  $A_k \subset C_k$  for each  $k \geq 0$  such that  $\bigcup_{n \geq 0} \bigsqcup_{i+j \leq n} \tau^i(A_j)$  is a basis of  $\mathrm{ind}_{K_0}^{G_S}(\sigma_{\bar{r}})$ . Consequently, the set  $\bigcup_{n \geq 0} A_n$  forms a basis for the module action of  $\mathcal{H}(G_S, K_0, \sigma_{\bar{r}}) = \bar{\mathbf{F}}_p[\tau]$  on  $\mathrm{ind}_{K_0}^{G_S}(\sigma_{\bar{r}})$ .  $\square$

*Remark 3.7.* The freeness result provides a transparent and conceptual understanding of the compactly induced representations of  $\mathrm{SL}_2$  in the mod- $p$  setting. Beyond its intrinsic interest, the formal framework of the inductive combinatorial argument presented in Proposition 2.4 may be adapted to split reductive groups of higher ranks when the corresponding Hecke action exhibits a graded or triangular behavior. Such constructions may play a role in the study of supersingular representations and their mod- $p$  local-global compatibility properties.

## REFERENCES

- [1] R. Abdellatif, *Classification des représentations modulo  $p$  de  $\mathrm{SL}(2, \mathbb{F})$* , Bull. Soc. Math. France **142** (2014), no. 3, 537–589; <https://doi.org/10.24033/bsmf.2673>
- [2] N. Abe, *On a classification of irreducible admissible modulo  $p$  representations of a  $p$ -adic split reductive group*, Compositio Math. **149** (2013), 2139–2168; <https://doi.org/10.1112/S0010437X13007379>
- [3] N. Abe, G. Henniart, F. Herzig, M.-F. Vignéras, *A classification of irreducible admissible mod  $p$  representations of  $p$ -adic reductive groups*, J. Amer. Math. Soc. **30** (2017), no. 2, 495–559; <https://doi.org/10.1090/jams/862>
- [4] N. Abe, G. Henniart, F. Herzig, M.-F. Vignéras, *Questions on mod  $p$  representations of reductive  $p$ -adic groups*. 2017. <https://doi.org/10.48550/arXiv.1703.02063>
- [5] L. Barthel and R. A. Livné, *Irreducible modular representations of  $\mathrm{GL}_2$  of a local field*, Duke Math. J. **75** (1994), no. 2, 261–292; <https://doi.org/10.1215/S0012-7094-94-07508-X>
- [6] C. Breuil, *Sur quelques représentations modulaires et  $p$ -adiques de  $\mathrm{GL}_2(\mathbb{Q}_p)$* . I, Compositio Math. **138** (2003), no. 2, 165–188; <https://doi.org/10.1023/A:1026191928449>
- [7] C. Cheng, *Mod  $p$  representations of  $\mathrm{SL}_2(\mathbb{Q}_p)$* , J. Number Theory **133** (2013), no. 4, 1312–1330; <https://doi.org/10.1016/j.jnt.2012.09.010>
- [8] A. Das, *A restriction problem for mod- $p$  representations of  $\mathrm{SL}_2(\mathbb{F})$* . 2024. <https://doi.org/10.48550/arXiv.2412.11751>
- [9] F. Herzig, *The classification of irreducible admissible mod  $p$  representations of a  $p$ -adic  $\mathrm{GL}_n$* , Invent. math. **186** (2011), 373–434; <https://doi.org/10.1007/s00222-011-0321-z>
- [10] F. Herzig, *A Satake isomorphism in characteristic  $p$* , Compositio Math. **147** (2011), no. 1, 263–283; <https://doi.org/10.1112/S0010437X10004951>
- [11] Y. Hu, *Diagrammes canoniques et représentations modulo  $p$  de  $\mathrm{GL}_2(\mathbb{F})$* , J. Inst. Math. Jussieu **11** (2012), no. 1, 67–118; <https://doi.org/10.1017/S1474748010000265>
- [12] P. Xu, *Freeness of spherical Hecke modules of unramified  $\mathrm{U}(2,1)$  in characteristic  $p$* , J. Number Theory **195** (2019), 293–311; <https://doi.org/10.1016/j.jnt.2018.06.006>
- [13] P. Xu, *Hecke eigenvalues in  $p$ -modular representations of unramified  $\mathrm{U}(2,1)$* , Proc. Amer. Math. Soc. **153** (2025), no. 1, 437–450; <https://doi.org/10.1090/proc/17042>