




GRAPH THEORY

~By Sparsh Jain

111601026



Under the guidance of
Dr. Deepak Rajendraprasad






Objective

To characterize the **outerplanar graphs**
which are also **B_0 -VPG graphs**



B₀-VPG Graph




Planar and Outerplanar Graphs





Weak Dual Graph





Asteroidal Triple



Permutation Graph



Counter Examples!



Prior Works and Important Theorems

- Bipartite permutation graphs, i.e. permutation graphs which are also bipartite graphs, are B_0 VPG graphs [1]
- A graph G is a permutation graph if and only if both G and its complement G^c are comparability graphs [3]



Prior Works and Important Theorems

- Permutation graphs have no induced cycles of lengths 5 or greater [2]
- The graph C_4 has a unique B0 VPG representation; it consists of two horizontal parallel paths intersecting with two vertical parallel paths [4]

Prior Works and Important Theorems

- A planar graph is outerplanar if and only if its weak dual is a forest [5]. A simple corollary is that the weak dual of a 2-connected outerplanar graph is a tree.
- If a graph G is the compliment of a comparability graph, then G contains no asteroidal triple (i.e. G is AT-free) [6]. Thus, permutation graphs are also AT-free.





Define: Chain






2-Connected Outerplanar Permutation Graphs are Chains

- **Claim:** Weak dual of a 2-Connected Outerplanar Permutation Graphs G is a path.
- **Proof:** Weak dual of a 2-Connected Outerplanar Graph is known to be a tree [5]. Thus it is sufficient to show that the maximum degree of weak dual of G is 2. Note that a bounded face of the graph G can either be a C_3 or a C_4 since Permutation Graphs cannot have induced cycles of length 5 or more [2].


Case 1




Case 2



Example of a B_0 -VPG



Example of a B_0 -VPG



Conclusion and Future Work

- Devise a concrete algorithm to construct B_0 -VPG Diagram for 2-Connected Outerplanar Permutation Graphs
- Since we are using only AT-Freeness and restriction of induced cycles to only C_3 and C_4 , we may be able to target a bigger subclass of 2-Connected Outerplanar Graphs
- It will be interesting to further expand the idea and categorize Outerplanar Graphs which are also B_0 -VPG graphs

References

1. E. Cohen, M. C. Golumbic, W. T. Trotter, and R. Wang, “Posets and vpg graphs,” *Order*, vol. 33, pp. 39–49, 2016.
2. M. C. Golumbic, “Algorithmic graph theory and perfect graphs,” 1980.
3. A. Pnueli, A. Lempel, and S. Even, “Transitive orientation of graphs and identification of permutation graphs,” *Canadian Journal of Mathematics*, vol. 23, no. 1, p. 160–175, 1971.
4. A. Asinowski, E. Cohen, M. C. Golumbic, V. Limouzy, M. Lipshteyn, and M. Stern, “Vertex intersection graphs of paths on a grid,” *J. Graph Algorithms Appl.*, vol. 16, pp. 129–150, 2012.
5. M. Syslo, “Characterizations of outerplanar graphs,” *Discrete Mathematics*, vol. 26, p. 47–53, 12 1979.
6. M. C. Golumbic, C. L. Monma, and W. T. Trotter, “Tolerance graphs,” *Discrete Applied Mathematics*, vol. 9, no. 2, pp. 157 – 170, 1984
7. M. C. Golumbic, D. Rotem, and J. Urrutia, “Comparability graphs and intersection graphs,” *Discrete Mathematics*, vol. 43, pp. 37–46, 1983.

Thank You!