





# Towards a Semantics of Unsatisfiability Proofs with Inprocessing

Tobias Philipp (TU Dresden) Adrián Rebola-Pardo (TU Wien)

LPAR-21 Maun, Botswana May 8th, 2017 Supported by FWF W1255-N23, WWTF VRG11-005, Microsoft Research and Graduate Academy TU Dresden

# **Inprocessing for SAT Solving**

Symmetry breaking clause  $\overline{x}y$  can be added without affecting satisfiability

# **Inprocessing for SAT Solving**

Symmetry breaking clause  $\overline{x}y$  can be added without affecting satisfiability

In-/preprocessing techniques replace the formula by an equisatisfiable one

symmetry clause parity breaking elimination reasoning

bounded variable cardinality addition resolution

How to certify the result of a SAT solver with inprocessing?

# **Satisfiability-preserving proof systems**

How to certify the result of a SAT solver with inprocessing? RUP cannot be used!

#### **Theorem**

If C is a RAT in F, then whenever F is satisfiable, so is  $F \cup \{C\}$ .

# Satisfiability-preserving proof systems

How to certify the result of a SAT solver with inprocessing? RUP cannot be used!

#### **Theorem**

If C is a RAT in F, then whenever F is satisfiable, so is  $F \cup \{C\}$ .

$$F \vDash_{\mathsf{sat}} F \cup \{C\}$$

### Satisfiability-preserving proof systems

How to certify the result of a SAT solver with inprocessing? RUP cannot be used!

#### **Theorem**


If C is a RAT in F, then whenever F is satisfiable, so is  $F \cup \{C\}$ .

$$F \vDash_{\mathsf{sat}} F \cup \{C\}$$


Idea build a proof system with ⊨<sub>sat</sub> as an invariant

- If C can be inferred, then  $F \vDash_{sat} C$ .
- Rules must be simple and efficient to check.
- Inprocessing rules should be easy to express.


The RAT property satisfies these conditions!




#### Clause C is a RAT in F



# Clause C is a RAT in F





# Clause C is a RAT in F non-RAT, non-consequence non-RAT, non-consequence non-RAT, non-consequence

# Clause C is a RAT in F non-RAT, non-consequence non-RAT, non-consequence consequence!


Clause C is a RAT in F

non-RAT, non-consequence

non-RAT, non-consequence

consequence!

consequence



Some invariant seems to be preserved...

Question what is the invariant preserved along a DRAT proof? if derived clauses are not consequences, then what are they?

# DRAT proofs

#### **RUP Introduction**

Most derived clauses are simply consequences of previously introduced clauses.

#### **Definition**

A clause C is a reverse unit propagation (RUP) clause in a CNF formula F if boolean constraint propagation over  $F \cup \overline{C}$  leads to contradiction.

#### **RUP Introduction**

Most derived clauses are simply consequences of previously introduced clauses.

#### **Definition**

A clause C is a reverse unit propagation (RUP) clause in a CNF formula F if boolean constraint propagation over  $F \cup \overline{C}$  leads to contradiction.

#### Characterization

C is a RUP in F if and only if  $\square \in F$ , or the least fixed point of  $ala_F$  contains complementary clauses, where:

$$\operatorname{ala}_F(C) = C \cup \{\overline{l} \mid \text{for some } D \in F, \ l \in D \text{ and } D \setminus \{l\} \subseteq C\}$$

#### **RUP Introduction**

Most derived clauses are simply consequences of previously introduced clauses.

#### **Definition**

A clause C is a reverse unit propagation (RUP) clause in a CNF formula F if boolean constraint propagation over  $F \cup \overline{C}$  leads to contradiction.

#### Characterization

C is a RUP in F if and only if  $\square \in F$ , or the least fixed point of  $ala_F$  contains complementary clauses, where:

$$\operatorname{ala}_F(C) = C \cup \{\bar{l} \mid \text{for some } D \in F, \ l \in D \text{ and } D \setminus \{l\} \subseteq C\}$$

#### Characterization

*C* is a RUP in *F* if and only if it can be proved from *F* through an SSSR chain:

$$\frac{C_0}{\operatorname{ssr}} \frac{C_0}{D_0} \qquad C_1 \\
\vdots \\
\operatorname{ssr} \frac{D_{n-1}}{D_n} \qquad C_n \\
= C_0$$

Introducing a RAT clause preserves satisfiability


#### **Definition**

A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $I \in C$  if every resolvent  $C \otimes_I D$  for  $D \in F$  is a RUP in F.

Introducing a RAT clause preserves satisfiability

#### **Definition**


A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.



Introducing a RAT clause preserves satisfiability

#### **Definition**


A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.



Introducing a RAT clause preserves satisfiability

#### **Definition**


A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.



Introducing a RAT clause preserves satisfiability

#### **Definition**


A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $I \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.



Introducing a RAT clause preserves satisfiability

#### **Definition**


A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.



Introducing a RAT clause preserves satisfiability

#### **Definition**

A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.



Introducing a RAT clause preserves satisfiability

#### **Definition**

A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.



Introducing a RAT clause preserves satisfiability

#### **Definition**

A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.



Clause deletion preserves satisfiability for arbitrary clauses

- Deletion information is needed for efficiency in BCP
- Arbitrary clause deletion can be expressed

#### Delete Resolution Asymmetric Tautology (DRAT) proof system

$$F \Rightarrow_{\mathsf{DRAT}} G \qquad \left\{ egin{array}{l} G = F \cup \{C\} \ & C \text{ is a RUP in } F \ & C \text{ is a RAT in } F \ & G = F \setminus \{C\} \ & G = F \setminus \{C\} \ & C \text{ is a RAT in } F \ & C \text{ is a$$

A DRAT proof of G from F is a sequence of DRAT inferences:

$$F = F_0 \Rightarrow_{\mathsf{DRAT}} F_1 \Rightarrow_{\mathsf{DRAT}} \dots \Rightarrow_{\mathsf{DRAT}} F_{n-1} \Rightarrow_{\mathsf{DRAT}} F_n = G$$



Clauses containing variable x can be regarded as a definition:

$$x \vee C_1$$

$$x \vee C_1$$
 ...  $x \vee C_n$   $\overline{x} \vee D_1$  ...  $\overline{x} \vee D_m$ 

$$\overline{x} \vee D_1$$

$$\overline{x}\vee D_m$$

Clauses containing variable x can be regarded as a definition:

$$x \leftarrow \overline{C_1}$$

$$x \leftarrow \overline{C_n}$$

$$\overline{x} \leftarrow \overline{D_1}$$

$$x \leftarrow \overline{C_1}$$
 ...  $x \leftarrow \overline{C_n}$   $\overline{x} \leftarrow \overline{D_1}$  ...  $\overline{x} \leftarrow \overline{D_m}$ 

Clauses containing variable x can be regarded as a definition:

$$x \leftarrow \overline{C_1} \qquad \dots \qquad x \leftarrow \overline{C_n} \qquad \overline{x} \leftarrow \overline{D_1} \qquad \dots \qquad \overline{x} \leftarrow \overline{D_m}$$
 
$$x = \begin{cases} 1 & \text{if } \overline{C_1} \\ \vdots \\ 1 & \text{if } \overline{C_n} \\ 0 & \text{if } \overline{D_1} \\ \vdots \\ 0 & \text{if } \overline{D_m} \\ \text{don't care} & \text{otherwise} \end{cases}$$

Clauses containing variable  $\boldsymbol{x}$  can be regarded as a definition:

$$x \leftarrow \overline{C_1} \qquad \dots \qquad x \leftarrow \overline{C_n} \qquad \overline{x} \leftarrow \overline{D_1} \qquad \dots \qquad \overline{x} \leftarrow \overline{D_m}$$
 
$$x = \begin{cases} & 1 & \text{if } \overline{C_1} \\ & \vdots \\ & 1 & \text{if } \overline{C_n} \\ & 0 & \text{if } \overline{D_1} \\ & \vdots \\ & 0 & \text{if } \overline{D_m} \\ & \text{don't care} \qquad \text{otherwise} \end{cases}$$

**Definition refinement** to add rule  $x \leftarrow C_{n+1}$ , we require:

 $\overline{C_{n+1}} \wedge \overline{D_i}$  is unsatisfiable for  $1 \leq i \leq m$ 

Clauses containing variable x can be regarded as a definition:

$$x \leftarrow \overline{C_1} \qquad \dots \qquad x \leftarrow \overline{C_n} \qquad \overline{x} \leftarrow \overline{D_1} \qquad \dots \qquad \overline{x} \leftarrow \overline{D_m}$$
 
$$x = \begin{cases} & 1 & \text{if } \overline{C_1} \\ & \vdots \\ & 1 & \text{if } \overline{C_n} \\ & 0 & \text{if } \overline{D_1} \\ & \vdots \\ & 0 & \text{if } \overline{D_m} \\ & \text{don't care} \qquad \text{otherwise} \end{cases}$$

**Definition refinement** to add rule  $x \leftarrow C_{n+1}$ , we require:

$$F \vDash \neg (\overline{C_{n+1}} \wedge \overline{D_i}) \text{ for } 1 \leq i \leq m$$

Clauses containing variable x can be regarded as a definition:

$$x \leftarrow \overline{C_1} \qquad \dots \qquad x \leftarrow \overline{C_n} \qquad \overline{x} \leftarrow \overline{D_1} \qquad \dots \qquad \overline{x} \leftarrow \overline{D_m}$$
 
$$x = \begin{cases} & 1 & \text{if } \overline{C_1} \\ & \vdots \\ & 1 & \text{if } \overline{C_n} \\ & 0 & \text{if } \overline{D_1} \\ & \vdots \\ & 0 & \text{if } \overline{D_m} \\ & \text{don't care} \qquad \text{otherwise} \end{cases}$$

**Definition refinement** to add rule  $x \leftarrow C_{n+1}$ , we require:

$$F \vDash \overline{C_{n+1}} \vee \overline{D_i} \text{ for } 1 \leq i \leq m$$

Clauses containing variable x can be regarded as a definition:

$$x \leftarrow \overline{C_1} \qquad \dots \qquad x \leftarrow \overline{C_n} \qquad \overline{x} \leftarrow \overline{D_1} \qquad \dots \qquad \overline{x} \leftarrow \overline{D_m}$$
 
$$x = \begin{cases} & 1 & \text{if } \overline{C_1} \\ & \vdots \\ & 1 & \text{if } \overline{C_n} \\ & 0 & \text{if } \overline{D_1} \\ & \vdots \\ & 0 & \text{if } \overline{D_m} \\ & \text{don't care} \qquad \text{otherwise} \end{cases}$$

**Definition refinement** to add rule  $x \leftarrow C_{n+1}$ , we require:

$$F \vDash (x \vee \overline{C_{n+1}}) \otimes_x (\overline{x} \vee \overline{D_i}) \text{ for } 1 \leq i \leq m$$

# **Defining a Definition**

Clauses containing variable x can be regarded as a definition:

$$x \leftarrow \overline{C_1} \qquad \dots \qquad x \leftarrow \overline{C_n} \qquad \overline{x} \leftarrow \overline{D_1} \qquad \dots \qquad \overline{x} \leftarrow \overline{D_m}$$
 
$$x = \begin{cases} & 1 & \text{if } \overline{C_1} \\ & \vdots \\ & 1 & \text{if } \overline{C_n} \\ & 0 & \text{if } \overline{D_1} \\ & \vdots \\ & 0 & \text{if } \overline{D_m} \\ & \text{don't care} \qquad \text{otherwise} \end{cases}$$

**Definition refinement** to add rule  $x \leftarrow C_{n+1}$ , we require:

$$F \vDash (x \lor \overline{C_{n+1}}) \otimes_x D$$
 for every clause  $D \in F$ 

Clauses containing variable x can be regarded as a definition:

$$x \leftarrow \overline{C_1} \qquad \dots \qquad x \leftarrow \overline{C_n} \qquad \overline{x} \leftarrow \overline{D_1} \qquad \dots \qquad \overline{x} \leftarrow \overline{D_m}$$
 
$$x = \begin{cases} & 1 & \text{if } \overline{C_1} \\ & \vdots \\ & 1 & \text{if } \overline{C_n} \\ & 0 & \text{if } \overline{D_1} \\ & \vdots \\ & 0 & \text{if } \overline{D_m} \\ & \text{don't care} \qquad \text{otherwise} \end{cases}$$

**Definition refinement** to add rule  $x \leftarrow C_{n+1}$ , we require:

$$F \vDash (x \lor \overline{C_{n+1}}) \otimes_x D$$
 for every clause  $D \in F$ 

### **Definition**

A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.

# **Defining a Definition**

Clauses containing variable  $\boldsymbol{x}$  can be regarded as a definition:

$$x \leftarrow \overline{C_1} \qquad \dots \qquad x \leftarrow \overline{C_n} \qquad \overline{x} \leftarrow \overline{D_1} \qquad \dots \qquad \overline{x} \leftarrow \overline{D_m}$$
 
$$x = \begin{cases} & 1 & \text{if } \overline{C_1} \\ & \vdots \\ & 1 & \text{if } \overline{C_n} \\ & 0 & \text{if } \overline{D_1} \\ & \vdots \\ & 0 & \text{if } \overline{D_m} \\ & \text{don't care} \qquad \text{otherwise} \end{cases}$$

**Definition refinement** to add rule  $x \leftarrow C_{n+1}$ , we require:

$$F \vDash (x \lor \overline{C_{n+1}}) \otimes_x D$$
 for every clause  $D \in F$ 

### **Definition**

### resolution consequence (RC)

A clause C is a resolution asymmetric tautology (RAT) in a CNF formula F upon a literal  $l \in C$  if every resolvent  $C \otimes_l D$  for  $D \in F$  is a RUP in F.

consequence of F

### **The Semantics of Definitions**

Question when is a definition correct from the semantic perspective?

- A definition changes the interpretation to force a truth value under some conditions.
- After the change, the interpretation still satisfies the original constraints.

### **The Semantics of Definitions**

Question when is a definition correct from the semantic perspective?

- A definition changes the interpretation to force a truth value under some conditions.
- After the change, the interpretation still satisfies the original constraints.

Mutation rule l := Q

Interpretation mutation

$$I \triangleleft (l :- Q) = \begin{cases} I + l & \text{if } I \vDash Q \\ I & \text{if } I \not \succeq Q \end{cases}$$

#### **Definition**

l := Q is a definition refinement of F if, whenever  $I \models F$  holds, then  $I \triangleleft (l := Q) \models F$  holds as well.

### **Definition**

A clause C is a resolution consequence in F upon I if every resolvent  $C \otimes_I D$  for  $D \in F$  is a consequence of F.

### **Definition**

l := Q is a definition refinement of F if, whenever  $I \models F$  holds, then  $I \triangleleft (l := Q) \models F$  holds as well.

### **Definition**

A clause C is a resolution consequence in F upon I if every resolvent  $C \otimes_I D$  for  $D \in F$  is a consequence of F.

#### **Definition**

l := Q is a definition refinement of F if, whenever  $I \models F$  holds, then  $I \triangleleft (l := Q) \models F$  holds as well.

### **Theorem**

 $C \vee l$  is a RC in F upon l if and only if  $l := \overline{C}$  is a definition refinement.

### **Definition**

A clause C is a resolution consequence in F upon I if every resolvent  $C \otimes_I D$  for  $D \in F$  is a consequence of F.

### **Definition**

l := Q is a definition refinement of F if, whenever  $I \models F$  holds, then  $I \triangleleft (l := Q) \models F$  holds as well.

### **Theorem**

 $C \vee l$  is a RC in F upon l if and only if  $l := \overline{C}$  is a definition refinement.

Example z is defined as true when x holds, and false when y holds correct only if  $\overline{x} \wedge \overline{y}$  is disallowed

$$\overline{x}\overline{y} \xrightarrow{z := \overline{x}} zx \xrightarrow{\overline{z} := \overline{y}} \overline{z}$$

#### Definition

A clause C is a resolution consequence in F upon I if every resolvent  $C \otimes_I D$  for  $D \in F$  is a consequence of F.


#### **Definition**

l := Q is a definition refinement of F if, whenever  $I \models F$  holds, then  $I \triangleleft (l := Q) \models F$  holds as well.


#### **Theorem**

 $C \vee l$  is a RC in F upon l if and only if  $l := \overline{C}$  is a definition refinement.

Example z is defined as true when x holds, and false when y holds correct only if  $\overline{x} \land \overline{y}$  is disallowed




## **Resolution Consequences as Proof Invariants**



Theorem If C is a RC in F upon l, and  $F \land C \models D$ , then either:

- $\blacksquare F \vDash D$
- D is an RC in F upon l


## **Resolution Consequences as Proof Invariants**



Theorem If C is a RC in F upon l, and  $F \land C \models D$ , then either:

- $\blacksquare F \vDash D$
- lacksquare D is an RC in F upon l

## **Resolution Consequences as Proof Invariants**



**Theorem** If C is a RC in F upon l, and  $F \land C \models D$ , then either:

- $\blacksquare F \vDash D$
- lacksquare D is an RC in F upon l

**Question** which definitions are required for each derived clause?

### Formula trees

$$\begin{array}{ccc} xyz & \overline{x}yz \\ x\overline{y}z & \overline{x}\overline{y}z & & & w := \overline{y} \\ \end{array} \longrightarrow \begin{array}{c} wy \\ wz & & & \overline{w} := \overline{z} \\ \end{array} \longrightarrow \overline{w}z$$

### Formula trees

$$I \models \frac{xyz}{x\overline{y}z} \xrightarrow{\overline{x}yz} \xrightarrow{w := \overline{y}} wy \xrightarrow{\overline{w} := \overline{z}} \overline{w}z$$

### Formula trees

$$I \models \begin{bmatrix} xyz & \overline{x}yz \\ x\overline{y}z & \overline{x}yz \end{bmatrix} \xrightarrow{w := \overline{y}} \xrightarrow{wy} \xrightarrow{\overline{w} := \overline{z}} \overline{w}z$$

### Formula trees

$$I \triangleleft (w :- \overline{y}) \models \begin{bmatrix} xyz & \overline{x}yz & w :- \overline{y} \\ x\overline{y}z & \overline{x}\overline{y}z \end{bmatrix} \xrightarrow{w :- \overline{y}} wy \xrightarrow{\overline{w} :- \overline{z}} \overline{w}z$$

### Formula trees

$$I \triangleleft (w := \overline{y}) \triangleleft (\overline{w} := \overline{z}) \models \underbrace{xyz \quad \overline{x}yz}_{x\overline{y}z} \xrightarrow{w := \overline{y}} \underbrace{wy}_{wz} \xrightarrow{\overline{w} := \overline{z}} \overline{w}z$$

#### Formula trees

Stratification derived clauses can be split based on which definitions they depend upon

Example 
$$z = wz \otimes_w \overline{w}z$$

#### Formula trees

$$\begin{array}{ccc} xyz & \overline{x}yz \\ x\overline{y}z & \overline{x}\overline{y}z \end{array} \xrightarrow{w := \overline{y}} \begin{array}{c} wy & \overline{w} := \overline{z} \\ wz & \end{array} \xrightarrow{\overline{w}}$$

Stratification derived clauses can be split based on which definitions they depend upon

Example 
$$z = wz \otimes_w \overline{w}z$$

#### Formula trees

$$\begin{array}{ccc} xyz & \overline{x}yz \\ x\overline{y}z & \overline{x}\overline{y}z & \end{array} \xrightarrow{w := \overline{y}} \begin{array}{c} wy \\ wz & \end{array} \xrightarrow{\overline{w} := \overline{z}} \xrightarrow{\overline{w}z}$$

Stratification derived clauses can be split based on which definitions they depend upon

Example  $z = wz \otimes_w \overline{w}z$  actually requires no definition

#### Formula trees

$$\begin{array}{ccc}
xyz & \overline{x}yz \\
x\overline{y}z & \overline{x}\overline{y}z
\end{array} \xrightarrow{w := \overline{y}} \begin{array}{c}
w := \overline{y} \\
wz & \overline{w} := \overline{z}
\end{array} \longrightarrow \overline{w}z$$

Stratification derived clauses can be split based on which definitions they depend upon

Example  $z = wz \otimes_w \overline{w}z$  actually requires no definition

#### Formula trees

$$\underbrace{\frac{xyz}{x\overline{y}z}\frac{\overline{x}yz}{x\overline{y}z}}_{x\overline{y}z}\underbrace{-\frac{w \div \overline{y}}{z}}_{wz}\xrightarrow{wy}\underbrace{-\overline{w} \div \overline{z}}_{wz}$$

Stratification derived clauses can be split based on which definitions they depend upon

Example  $z = wz \otimes_w \overline{w}z$  actually requires no definition

### Formula trees

$$\begin{array}{ccc}
xyz & \overline{x}yz \\
x\overline{y}z & \overline{x}yz
\end{array} \xrightarrow{w := \overline{y}} wy & \overline{w} := \overline{z} \\
wz & \overline{w}z & \overline{w}z
\end{array}$$

Stratification derived clauses can be split based on which definitions they depend upon

Example  $z = wz \otimes_w \overline{w}z$  actually requires no definition

### **Equivalence- and definitionality-preserving inferences**

- Consequence introduction simulates RUP introduction
- Definition extension simulates RAT introduction
- Clause upgrade applied on resolvents upon pivot literals

### **Definitions and deletion**

Clause deletion can make correct definitions incorrect:

$$\begin{array}{ccc} xyz & \overline{x}yz & & w : \overline{y} & wy & \overline{w} : \overline{z} & \overline{w}z \\ x\overline{y}z & \overline{x}\overline{y}z & & wz & & z \end{array}$$

### **Definitions and deletion**

Clause deletion can make correct definitions incorrect:

$$\begin{array}{ccc}
xyz & \overline{x}yz & \underline{w} & -\overline{y} \\
x\overline{y}z & & & wz & -\overline{z} \\
\end{array} \xrightarrow{w} \begin{array}{c}
w & -\overline{z} \\
w & & z
\end{array}$$

## **Definitions and deletion**

Clause deletion can make correct definitions incorrect:

Clause deletion can make correct definitions incorrect:

Question is there any invariant throughout DRAT proofs stronger than  $\vDash_{sat}$ ?

**Theorem** G is DRAT-derivable from F if and only if  $F \vDash_{\mathsf{sat}} G$  no stronger invariant within propositional logic and unrestricted deletion

Clause deletion can make correct definitions incorrect:

Question is there any invariant throughout DRAT proofs stronger than  $\vDash_{sat}$ ?

**Theorem** G is DRAT-derivable from F if and only if  $F \vDash_{sat} G$  no stronger invariant within propositional logic and unrestricted deletion

#### Possible workarounds

- Restrict deletion to realistic situations
- Construe a DRAT proof as a derivation on DQBF clauses.
- Change mutation rules upon deletion.



DRAT proofs are widely used for validation of SAT solvers' results, but its semantics are not well understood.

DRAT proofs are widely used for validation of SAT solvers' results, but its semantics are not well understood.

In the absence of deletion, proofs correspond to derivations on definitional formulas. In particular, resolution consequences correspond exactly to a form of permissive definition.

DRAT proofs are widely used for validation of SAT solvers' results, but its semantics are not well understood.

In the absence of deletion, proofs correspond to derivations on definitional formulas. In particular, resolution consequences correspond exactly to a form of permissive definition.

#### **Future work**

can we generate resolution/RUP proofs from a DRAT proof? yes!

DRAT proofs are widely used for validation of SAT solvers' results, but its semantics are not well understood.

In the absence of deletion, proofs correspond to derivations on definitional formulas. In particular, resolution consequences correspond exactly to a form of permissive definition.

#### **Future work**

can we generate resolution/RUP proofs from a DRAT proof? yes!

#### Actual future work

can we find a semantics stronger than satisfiability preservation for DRAT proofs?