

Proofs (that contain programs that contain proofs)*

Adrián Rebola-Pardo

TU Wien, JKU Linz

Timisoara, Romania

26 September 2025

Supported by FWF 10.55776/COE12

Proofs in SAT solving: the good ol' times

Resolution [Davis, Putnam '60]

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D}$$

$$\{C \vee x, D \vee \bar{x}\} \models C \vee D$$

Proofs in SAT solving: the good ol' times

Resolution [Davis, Putnam '60]

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D} \quad \{C \vee x, D \vee \bar{x}\} \models C \vee D$$

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a resolvent of clauses in $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Proofs in SAT solving: the good ol' times

Resolution [Davis, Putnam '60]

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D} \quad \{C \vee x, D \vee \bar{x}\} \models C \vee D$$

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a resolvent of clauses in $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Reverse unit propagation (RUP) [Goldberg, Novikov '03]

C is a RUP over F if C can be obtained by iterated resolution over clauses in F

Proofs in SAT solving: the good ol' times

Resolution [Davis, Putnam '60]

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D} \quad \{C \vee x, D \vee \bar{x}\} \models C \vee D$$

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a resolvent of clauses in $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Reverse unit propagation (RUP) [Goldberg, Novikov '03]

C is a RUP over F if C can be obtained by iterated resolution over clauses in F

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a RUP over $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Proofs in SAT solving: the good ol' times

Resolution [Davis, Putnam '60]

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D} \quad \{C \vee x, D \vee \bar{x}\} \models C \vee D$$

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a resolvent of clauses in $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Reverse unit propagation (RUP) [Goldberg, Novikov '03]

C is a RUP over F if C can be obtained by iterated resolution over clauses in F

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a RUP over $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Deletions [Heule, Hunt, Wetzler '14]

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

Proofs in SAT solving: the good ol' times

Resolution [Davis, Putnam '60]

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D} \quad \{C \vee x, D \vee \bar{x}\} \models C \vee D$$

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a resolvent of clauses in $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Reverse unit propagation (RUP) [Goldberg, Novikov '03]

C is a RUP over F if C can be obtained by iterated resolution over clauses in F

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a RUP over $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Deletions [Heule, Hunt, Wetzler '14]

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

- If the i -th instruction is i: C , then C must be RUP in F_{i-1} , and $F_i = F_{i-1} \cup \{C\}$

Proofs in SAT solving: the good ol' times

Resolution [Davis, Putnam '60]

$$\frac{C \vee x \quad D \vee \bar{x}}{C \vee D} \quad \{C \vee x, D \vee \bar{x}\} \models C \vee D$$

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a resolvent of clauses in $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Reverse unit propagation (RUP) [Goldberg, Novikov '03]

C is a RUP over F if C can be obtained by iterated resolution over clauses in F

$\pi = C_1, \dots, C_n$ is a proof of $F \models G$ if

- C_i is a RUP over $F \cup \{C_1, \dots, C_{i-1}\}$
- $G \subseteq F \cup \{C_1, \dots, C_n\}$

Deletions [Heule, Hunt, Wetzler '14]

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

- If the i -th instruction is i: C , then C must be RUP in F_{i-1} , and $F_i = F_{i-1} \cup \{C\}$
- If the i -th instruction is d: C , then $F_i = F_{i-1} \setminus \{C\}$

Proofs in SAT solving: the redundance era

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

- If the i -th instruction is i: C , then C must be RUP in F_{i-1} , and $F_i = F_{i-1} \cup \{C\}$
- If the i -th instruction is d: C , then $F_i = F_{i-1} \setminus \{C\}$

Proofs in SAT solving: the redundance era

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

- If the i -th instruction is i: C , then C must be RUP in F_{i-1} , and $F_i = F_{i-1} \cup \{C\}$
- If the i -th instruction is d: C , then $F_i = F_{i-1} \setminus \{C\}$

$$F = F_0 \models F_1 \models F_2 \models \dots \models F_n \ni \perp$$

Proofs in SAT solving: the redundance era

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

- If the i -th instruction is i: C , then C must be RUP in F_{i-1} , and $F_i = F_{i-1} \cup \{C\}$
- If the i -th instruction is d: C , then $F_i = F_{i-1} \setminus \{C\}$

$$F = F_0 \models F_1 \models F_2 \models \dots \models F_n \ni \perp$$

... **but why \models ?** [Järvisalo, Heule, Biere '12]

C is redundant in F if F satisfiable implies $F \cup \{C\}$ satisfiable

If the proof ends in \perp , then we can replace RUP by any redundance criterion!

Proofs in SAT solving: the redundance era

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

- If the i -th instruction is i: C , then C must be RUP in F_{i-1} , and $F_i = F_{i-1} \cup \{C\}$
- If the i -th instruction is d: C , then $F_i = F_{i-1} \setminus \{C\}$

$$F = F_0 \models F_1 \models F_2 \models \dots \models F_n \ni \perp$$

... **but why \models ?** [Järvisalo, Heule, Biere '12]

C is redundant in F if F satisfiable implies $F \cup \{C\}$ satisfiable

If the proof ends in \perp , then we can replace RUP by any redundance criterion!

$$F = F_0 \models_{\text{sat}} F_1 \models_{\text{sat}} F_2 \models_{\text{sat}} \dots \models_{\text{sat}} F_n \ni \perp$$

Proofs in SAT solving: the redundance era

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

- If the i -th instruction is i: C , then C must be RUP in F_{i-1} , and $F_i = F_{i-1} \cup \{C\}$
- If the i -th instruction is d: C , then $F_i = F_{i-1} \setminus \{C\}$

$$F = F_0 \models F_1 \models F_2 \models \dots \models F_n \ni \perp$$

... **but why \models ?** [Järvisalo, Heule, Biere '12]

C is redundant in F if F satisfiable implies $F \cup \{C\}$ satisfiable

If the proof ends in \perp , then we can replace RUP by any redundance criterion!

$$F = F_0 \models_{\text{sat}} F_1 \models_{\text{sat}} F_2 \models_{\text{sat}} \dots \models_{\text{sat}} F_n \ni \perp$$

... **why though?** proof complexity considerations [Tseitin '83] [Haken '85]

Proofs in SAT solving: the redundance era

Proofs are sequences of either introductions i: C or deletions d: C .

At every point in the proof we have an **accumulated formula** F_i .

- If the i -th instruction is i: C , then C must be RUP in F_{i-1} , and $F_i = F_{i-1} \cup \{C\}$
- If the i -th instruction is d: C , then $F_i = F_{i-1} \setminus \{C\}$

$$F = F_0 \models F_1 \models F_2 \models \dots \models F_n \ni \perp$$

... **but why \models ?** [Järvisalo, Heule, Biere '12]

C is redundant in F if F satisfiable implies $F \cup \{C\}$ satisfiable

If the proof ends in \perp , then we can replace RUP by any redundance criterion!

$$F = F_0 \models_{\text{sat}} F_1 \models_{\text{sat}} F_2 \models_{\text{sat}} \dots \models_{\text{sat}} F_n \ni \perp$$

... **why though?** proof complexity considerations [Tseitin '83] [Haken '85]

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]
but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, T or \perp)

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]

but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, \top or \perp)

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$ (simplified)

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]

but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, \top or \perp)

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$ (simplified)

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]
but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, \top or \perp)

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$ (simplified)

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

Pros

Succinctly captures a lot (but not all!) of beyond-CDCL reasoning

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]
but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, \top or \perp)

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$ (simplified)

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

Pros

Succinctly captures a lot (but not all!) of beyond-CDCL reasoning

Cons [RP, Suda '17] [RP '23] [RP '25]

Fixpoints in trimming *huh, that's weird*

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]
but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, \top or \perp)

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$ (simplified)

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

Pros

Succinctly captures a lot (but not all!) of beyond-CDCL reasoning

Cons [RP, Suda '17] [RP '23] [RP '25]

Fixpoints in trimming *huh, that's weird*

Subproofs cannot be merged *that's ok i guess?*

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]
but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, \top or \perp)

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$ (simplified)

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

Pros

Succinctly captures a lot (but not all!) of beyond-CDCL reasoning

Cons [RP, Suda '17] [RP '23] [RP '25]

Fixpoints in trimming *huh, that's weird*

Subproofs cannot be merged *that's ok i guess?*

Deriving lemmas prevents inferences **slams desk in anger**

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]
but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, \top or \perp)

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$ (simplified)

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

Pros

Succintly captures a lot (but not all!) of beyond-CDCL reasoning

Cons [RP, Suda '17] [RP '23] [RP '25]

Fixpoints in trimming *huh, that's weird*

Subproofs cannot be merged *that's ok i guess?*

Deriving lemmas prevents inferences **slams desk in anger**

Deletions are now semantically relevant **yells at clouds**

Proofs in SAT solving: substitution redundancy and interference

Substitution redundancy (SR) [Buss, Thapen '19] [Gocht, Nordström, '21]
but also DRAT [Wetzler, Heule, Hunt '14], DPR [Heule, Kiesl, Biere '17]

F is a CNF formula

C is a clause

σ is an atomic substitution (variables map to literals, \top or \perp)

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$ (simplified)

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

Pros

Succinctly captures a lot (but not all!) of beyond-CDCL reasoning

Cons [RP, Suda '17] [RP '23] [RP '25]

Fixpoints in trimming *huh, that's weird*

Subproofs cannot be merged *that's ok i guess?*

Deriving lemmas prevents inferences **slams desk in anger**

Deletions are now semantically relevant **yells at clouds**

The real issue non-monotonicity and global dependence a.k.a. **interference**

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_\sigma$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

This is a monotonic property! **bangs head against the wall**

Solution [RP, Suda '17] [RP '23]

Operate over constraints $\nabla(\sigma_1 :- T_1). \dots \nabla(\sigma_n :- T_n). C$

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

This is a monotonic property! **bangs head against the wall**

Solution [RP, Suda '17] [RP '23]

Operate over constraints $\nabla(\sigma_1 :- T_1) \dots \nabla(\sigma_n :- T_n).C$

$I \models \nabla(T :- \sigma).\varphi$ iff $I' \models \varphi$, where $I' = I \circ \sigma$ if $I \models T$, and $I' = I$ otherwise

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_\sigma$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

This is a monotonic property! **bangs head against the wall**

Solution [RP, Suda '17] [RP '23]

Operate over constraints $\nabla(\sigma_1 :- T_1) \dots \nabla(\sigma_n :- T_n).C$

$I \models \nabla(T :- \sigma).\varphi$ iff $I' \models \varphi$, where $I' = I \circ \sigma$ if $I \models T$, and $I' = I$ otherwise

If C is SR over F upon σ then $F \models \nabla(\bar{C} :- \sigma).F \cup \{C\}$

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_\sigma$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

This is a monotonic property! **bangs head against the wall**

Solution [RP, Suda '17] [RP '23]

Operate over constraints $\nabla(\sigma_1 :- T_1) \dots \nabla(\sigma_n :- T_n).C$

$I \models \nabla(T :- \sigma).\varphi$ iff $I' \models \varphi$, where $I' = I \circ \sigma$ if $I \models T$, and $I' = I$ otherwise

If C is SR over F upon σ then $F \models \nabla(\bar{C} :- \sigma).F \cup \{C\}$

Fixpoints in trimming are now gone!

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

This is a monotonic property! **bangs head against the wall**

Solution [RP, Suda '17] [RP '23]

Operate over constraints $\nabla(\sigma_1 :- T_1) \dots \nabla(\sigma_n :- T_n).C$

$I \models \nabla(T :- \sigma).\varphi$ iff $I' \models \varphi$, where $I' = I \circ \sigma$ if $I \models T$, and $I' = I$ otherwise

If C is SR over F upon σ then $F \models \nabla(\bar{C} :- \sigma).F \cup \{C\}$

Fixpoints in trimming are now gone! **Lemmas** can now safely derived!

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_\sigma$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

This is a monotonic property! **bangs head against the wall**

Solution [RP, Suda '17] [RP '23]

Operate over constraints $\nabla(\sigma_1 :- T_1) \dots \nabla(\sigma_n :- T_n).C$

$I \models \nabla(T :- \sigma).\varphi$ iff $I' \models \varphi$, where $I' = I \circ \sigma$ if $I \models T$, and $I' = I$ otherwise

If C is SR over F upon σ then $F \models \nabla(\bar{C} :- \sigma).F \cup \{C\}$

Fixpoints in trimming are now gone! **Lemmas** can now safely derived!

Deletions are now semantically irrelevant!

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

This is a monotonic property! **bangs head against the wall**

Solution [RP, Suda '17] [RP '23]

Operate over constraints $\nabla(\sigma_1 :- T_1) \dots \nabla(\sigma_n :- T_n).C$

$I \models \nabla(T :- \sigma).\varphi$ iff $I' \models \varphi$, where $I' = I \circ \sigma$ if $I \models T$, and $I' = I$ otherwise

If C is SR over F upon σ then $F \models \nabla(\bar{C} :- \sigma).F \cup \{C\}$

Fixpoints in trimming are now gone! **Lemmas** can now safely derived!

Deletions are now semantically irrelevant! All inferences have clearly defined **dependencies**!

Accumulated formulas are not your friend

C is SR over F upon σ if $F \cup \{\bar{C}\} \models (F \cup \{C\})|_{\sigma}$

If I had an assignment I satisfying F , then check if it satisfies C .

If it does, then I satisfies $F \cup \{C\}$; otherwise, $I \circ \sigma$ satisfies $F \cup \{C\}$.

If π is a proof of $F \vdash G$ then, given any assignment I satisfying F , I know how to construct $\text{mut}(I)$ satisfying G

This is a monotonic property! **bangs head against the wall**

Solution [RP, Suda '17] [RP '23]

Operate over constraints $\nabla(\sigma_1 :- T_1) \dots \nabla(\sigma_n :- T_n).C$

$I \models \nabla(T :- \sigma).\varphi$ iff $I' \models \varphi$, where $I' = I \circ \sigma$ if $I \models T$, and $I' = I$ otherwise

If C is SR over F upon σ then $F \models \nabla(\bar{C} :- \sigma).F \cup \{C\}$

Fixpoints in trimming are now gone! **Lemmas** can now safely derived!

Deletions are now semantically irrelevant! All inferences have clearly defined **dependencies**!

... but subproofs still can't be merged *I must be missing something...*

Dominance: interference with a vengeance

Dominance needs **three** accumulated formulas *please stop...*

$K(x)$, $R(x)$, $O(x, x')$ are CNF formulas

Dominance: interference with a vengeance

Dominance needs **three** accumulated formulas *please stop...*

$K(x), R(x), O(x, x')$ are CNF formulas

$O(x, x')$ encodes a preorder: $O(x, x') \equiv \sum_{i=0}^n 2^i x_i - \sum_{i=0}^n 2^i x'_i \leq 0$

$$F \xrightarrow[\kappa]{\sim} K \xrightarrow[\delta]{\sim} K \cup R \qquad K(x) \models O(x, \delta(x))$$

RUP and SR add clauses in $R(x)$

SR now requires proving $K(x) \models O(x, \sigma(x))$, and σ is added to δ

Dominance: interference with a vengeance

Dominance needs **three** accumulated formulas *please stop...*

$K(x), R(x), O(x, x')$ are CNF formulas

$O(x, x')$ encodes a preorder: $O(x, x') \equiv \sum_{i=0}^n 2^i x_i - \sum_{i=0}^n 2^i x'_i \leq 0$

$$F \xrightarrow[\kappa]{\sim} K \xrightarrow[\delta]{\sim} K \cup R \qquad K(x) \models O(x, \delta(x))$$

RUP and SR add clauses in $R(x)$

SR now requires proving $K(x) \models O(x, \sigma(x))$, and σ is added to δ

Dominance [Boggaerts, Gocht, McCreesh, Nordström '23]

C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

Dominance: interference with a vengeance

Dominance needs **three** accumulated formulas *please stop...*

$K(x), R(x), O(x, x')$ are CNF formulas

$O(x, x')$ encodes a preorder: $O(x, x') \equiv \sum_{i=0}^n 2^i x_i - \sum_{i=0}^n 2^i x'_i \leq 0$

$$F \xrightarrow[\kappa]{\sim} K \xrightarrow[\delta]{\sim} K \cup R \qquad K(x) \models O(x, \delta(x))$$

RUP and SR add clauses in $R(x)$

SR now requires proving $K(x) \models O(x, \sigma(x))$, and σ is added to δ

Dominance [Boggaerts, Gocht, McCreesh, Nordström '23]

C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\bar{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\bar{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If C is dominance-redundant over K, R, O upon σ , then C is redundant on $K \cup R$

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder

$$F \xrightarrow[\kappa]{\sim} K \xrightarrow[\delta]{\sim} K \cup R \qquad K(x) \models O(x, \delta(x))$$

Dominance-based strengthening (*very simplified*) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder

$$F \xrightarrow[\kappa]{\sim} K \xrightarrow[\delta]{\sim} K \cup R \qquad K(x) \models O(x, \delta(x))$$

Dominance-based strengthening (*very simplified*) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder

$$F \xrightarrow[\kappa]{\sim} K \xrightarrow[\delta]{\sim} K \cup R \qquad K(x) \models O(x, \delta(x))$$

Dominance-based strengthening (*very simplified*) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas O encodes a preorder

$$F \overset{\kappa}{\rightsquigarrow} K \overset{\delta}{\rightsquigarrow} K \cup R \qquad K(x) \models O(x, \delta(x))$$

Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder



Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder



Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\bar{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\bar{C}(x)\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder



Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

If J_1 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder



Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

If J_1 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_2 blah blah blah.

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder



Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\bar{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\bar{C}(x)\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

If J_1 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_2 blah blah blah.

This process cannot continue forever because
 O is non-increasing on δ and strictly decreasing on σ .

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder



Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\bar{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\bar{C}(x)\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

If J_1 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_2 blah blah blah.

This process cannot continue forever because
 O is non-increasing on δ and strictly decreasing on σ .

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder



$$K(x) \cup R(x) \cup \overline{C}(x) \models x > \sigma(x)$$

$$K(x) \models x \geq \delta(x)$$

Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

If J_1 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_2 blah blah blah.

This process cannot continue forever because
 O is non-increasing on δ and strictly decreasing on σ .

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder



$$K(x) \cup R(x) \cup \overline{C}(x) \models x > \sigma(x)$$

$$K(x) \models x \geq \delta(x)$$

Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C}(x)\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

If J_1 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_2 blah blah blah.

This process cannot continue forever because
 O is non-increasing on δ and strictly decreasing on σ .

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder

termination order



$$K(x) \cup R(x) \cup \overline{C}(x) \models x > \sigma(x)$$

$$K(x) \models x \geq \delta(x)$$

Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C(x)}\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

If J_1 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_2 blah blah blah.

This process cannot continue forever because
 O is non-increasing on δ and strictly decreasing on σ .

Look who's hiding under dominance

$K(x), R(x), O(x, x')$ are CNF formulas

O encodes a preorder

invariant

$F \xrightarrow{\delta} K \cup R$
MODEL CHECKING BY ANY OTHER NAME

termination order

$$K(x) \cup R(x) \cup \overline{C}(x) \models x > \sigma(x)$$

$$K(x) \models x \geq \delta(x)$$

Dominance-based strengthening (very simplified) C is dominance-redundant over K, R, O upon σ if:

- $K \cup R \cup \{\overline{C}\} \models K|_{\sigma}$
- $K(x) \cup R(x) \cup \{\overline{C}(x)\} \models O(x, \sigma(x)) \cup \overline{O(\sigma(x), x)}$

If I had an assignment I_0 satisfying K ,
then by δ I have an assignment J_0 satisfying $K \cup R$.

If J_0 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_1 satisfying K ,
and by δ I have an assignment J_1 satisfying $K \cup R$.

If J_1 satisfies C , then $K \cup R \cup \{C\}$ is satisfiable;
otherwise by σ I have an assignment I_2 blah blah blah.

This process cannot continue forever because
 O is non-increasing on δ and strictly decreasing on σ .

The swiss army knife of the non-monotonicist

Why do we keep having global conditions on classical reasoning when *we know* classical logic is monotonic? [\[Martin Suda over lunch in 2017\]](#)

$\nabla(T : - \sigma)(I)$ is $I \circ \sigma$ if $I \models T$, or I otherwise.

The swiss army knife of the non-monotonicist

Why do we keep having global conditions on classical reasoning when *we know* classical logic is monotonic? [Martin Suda over lunch in 2017]

$\nabla(T : - \sigma)(I)$ is $I \circ \sigma$ if $I \models T$, or I otherwise.

I maps variables to bits \rightsquigarrow memory states

The swiss army knife of the non-monotonicist

Why do we keep having global conditions on classical reasoning when *we know* classical logic is monotonic? [Martin Suda over lunch in 2017]

$\nabla(T : - \sigma)(I)$ is $I \circ \sigma$ if $I \models T$, or I otherwise.

I maps variables to bits \rightsquigarrow **memory states**

$\nabla(T : - \sigma)$ transforms a memory state into a memory state \rightsquigarrow **programs**

The swiss army knife of the non-monotonicist

Why do we keep having global conditions on classical reasoning when *we know* classical logic is monotonic? [Martin Suda over lunch in 2017]

$\nabla(T : - \sigma)(I)$ is $I \circ \sigma$ if $I \models T$, or I otherwise.

I maps variables to bits \rightsquigarrow **memory states**

$\nabla(T : - \sigma)$ transforms a memory state into a memory state \rightsquigarrow **programs**

if we want to make this work for dominance, we must be even more general:

- programs may be **partial maps** (to allow while loops)
- programs may be **non-deterministic** (to encode preorders)

The swiss army knife of the non-monotonicist

Why do we keep having global conditions on classical reasoning when *we know* classical logic is monotonic? [Martin Suda over lunch in 2017]

$\nabla(T : - \sigma)(I)$ is $I \circ \sigma$ if $I \models T$, or I otherwise.

I maps variables to bits \rightsquigarrow **memory states**

$\nabla(T : - \sigma)$ transforms a memory state into a memory state \rightsquigarrow **programs**

if we want to make this work for dominance, we must be even more general:

- programs may be **partial maps** (to allow while loops)
- programs may be **non-deterministic** (to encode preorders)

(Static) constraints semantics given by a set of (satisfying) assignments

$$J \models C$$

The swiss army knife of the non-monotonicist

Why do we keep having global conditions on classical reasoning when *we know* classical logic is monotonic? [Martin Suda over lunch in 2017]

$\nabla(T : - \sigma)(I)$ is $I \circ \sigma$ if $I \models T$, or I otherwise.

I maps variables to bits \rightsquigarrow **memory states**

$\nabla(T : - \sigma)$ transforms a memory state into a memory state \rightsquigarrow **programs**

if we want to make this work for dominance, we must be even more general:

- programs may be **partial maps** (to allow while loops)
- programs may be **non-deterministic** (to encode preorders)

(Static) constraints semantics given by a set of (satisfying) assignments

Programs semantics given by a binary relation of (transitioning) assignments

$$J \models C$$

$$I \otimes J \models \epsilon$$

The swiss army knife of the non-monotonicist

Why do we keep having global conditions on classical reasoning when *we know* classical logic is monotonic? [Martin Suda over lunch in 2017]

$\nabla(T : - \sigma)(I)$ is $I \circ \sigma$ if $I \models T$, or I otherwise.

I maps variables to bits \rightsquigarrow **memory states**

$\nabla(T : - \sigma)$ transforms a memory state into a memory state \rightsquigarrow **programs**

if we want to make this work for dominance, we must be even more general:

- programs may be **partial maps** (to allow while loops)
- programs may be **non-deterministic** (to encode preorders)

(Static) constraints semantics given by a set of (satisfying) assignments

Programs semantics given by a binary relation of (transitioning) assignments

Dynamic constraints a static constraint must hold after executing a program

$$J \models C$$

$$I \otimes J \models \epsilon$$

The swiss army knife of the non-monotonicist

Why do we keep having global conditions on classical reasoning when *we know* classical logic is monotonic? [Martin Suda over lunch in 2017]

$\nabla(T : - \sigma)(I)$ is $I \circ \sigma$ if $I \models T$, or I otherwise.

I maps variables to bits \rightsquigarrow **memory states**

$\nabla(T : - \sigma)$ transforms a memory state into a memory state \rightsquigarrow **programs**

if we want to make this work for dominance, we must be even more general:

- programs may be **partial maps** (to allow while loops)
- programs may be **non-deterministic** (to encode preorders)

(Static) constraints semantics given by a set of (satisfying) assignments

Programs semantics given by a binary relation of (transitioning) assignments

Dynamic constraints a static constraint must hold after executing a program

$$I \models \varepsilon.C \quad \text{iff} \quad J \models C \quad \text{for all } J \text{ such that } I \otimes J \models \varepsilon$$

What's in the box? Programs!

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

Branch $I \otimes J \models \nabla(T : \varepsilon_1 \parallel \varepsilon_0)$ iff $I \otimes J \models (T? \varepsilon_1) \sqcup (\bar{T}? \varepsilon_0)$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

Branch $I \otimes J \models \nabla(T : \varepsilon_1 \parallel \varepsilon_0)$ iff $I \otimes J \models (T? \varepsilon_1) \sqcup (\bar{T}? \varepsilon_0)$

Repeat $I \otimes J \models \varepsilon^*$ iff $\exists I_i, I_0 = I, I_n = J, I_{i-1} \otimes I_i \models \varepsilon$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

Branch $I \otimes J \models \nabla(T : \varepsilon_1 \parallel \varepsilon_0)$ iff $I \otimes J \models (T? \varepsilon_1) \sqcup (\bar{T}? \varepsilon_0)$

Repeat $I \otimes J \models \varepsilon^*$ iff $\exists I_i, I_0 = I, I_n = J, I_{i-1} \otimes I_i \models \varepsilon$

Loop $I \otimes J \models \square(T : \varepsilon)$ iff $I \otimes J \models (\bar{T}? \varepsilon)^* T?$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

Branch $I \otimes J \models \nabla(T : \varepsilon_1 \parallel \varepsilon_0)$ iff $I \otimes J \models (T? \varepsilon_1) \sqcup (\bar{T}? \varepsilon_0)$

Repeat $I \otimes J \models \varepsilon^*$ iff $\exists I_i, I_0 = I, I_n = J, I_{i-1} \otimes I_i \models \varepsilon$

Loop $I \otimes J \models \square(T : \varepsilon)$ iff $I \otimes J \models (\bar{T}? \varepsilon)^* T?$

Rendezvous $I \otimes J \models \diamond(V : \varepsilon_1 \parallel \varepsilon_0)$ iff $\exists J_1, J_0, I \otimes J_i \models \varepsilon_i$ and $J = J_1 +_V J_0$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

Branch $I \otimes J \models \nabla(T : \varepsilon_1 \parallel \varepsilon_0)$ iff $I \otimes J \models (T? \varepsilon_1) \sqcup (\bar{T}? \varepsilon_0)$

Repeat $I \otimes J \models \varepsilon^*$ iff $\exists I_i, I_0 = I, I_n = J, I_{i-1} \otimes I_i \models \varepsilon$

Loop $I \otimes J \models \square(T : \varepsilon)$ iff $I \otimes J \models (\bar{T}? \varepsilon)^* T?$

Rendezvous $I \otimes J \models \diamond(V : \varepsilon_1 \parallel \varepsilon_0)$ iff $\exists J_1, J_0, I \otimes J_i \models \varepsilon_i$ and $J = J_1 +_V J_0$

Solve $I \otimes J \models [R]$ iff $I \otimes J \models R$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

Branch $I \otimes J \models \nabla(T : \varepsilon_1 \parallel \varepsilon_0)$ iff $I \otimes J \models (T? \varepsilon_1) \sqcup (\bar{T}? \varepsilon_0)$

Repeat $I \otimes J \models \varepsilon^*$ iff $\exists I_i, I_0 = I, I_n = J, I_{i-1} \otimes I_i \models \varepsilon$

Loop $I \otimes J \models \square(T : \varepsilon)$ iff $I \otimes J \models (\bar{T}? \varepsilon)^* T?$

Rendezvous $I \otimes J \models \diamond(V : \varepsilon_1 \parallel \varepsilon_0)$ iff $\exists J_1, J_0, I \otimes J_i \models \varepsilon_i$ and $J = J_1 +_V J_0$

Solve $I \otimes J \models [R]$ iff $I \otimes J \models R$

Havoc $I \otimes J \models \forall V$ iff $I \otimes J \models \diamond(V : [T] \parallel 1)$

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

Branch $I \otimes J \models \nabla(T : \varepsilon_1 \parallel \varepsilon_0)$ iff $I \otimes J \models (T? \varepsilon_1) \sqcup (\bar{T}? \varepsilon_0)$

Repeat $I \otimes J \models \varepsilon^*$ iff $\exists I_i, I_0 = I, I_n = J, I_{i-1} \otimes I_i \models \varepsilon$

Loop $I \otimes J \models \square(T : \varepsilon)$ iff $I \otimes J \models (\bar{T}? \varepsilon)^* T?$

Rendezvous $I \otimes J \models \diamond(V : \varepsilon_1 \parallel \varepsilon_0)$ iff $\exists J_1, J_0, I \otimes J_i \models \varepsilon_i$ and $J = J_1 +_V J_0$

Solve $I \otimes J \models [R]$ iff $I \otimes J \models R$

Havoc $I \otimes J \models \forall V$ iff $I \otimes J \models \diamond(V : [T] \parallel 1)$

Not even a new thing! [Fischer, Ladner '79] [Balbiani, Herzig, Troquard '13]

Propositional dynamic logic (PDL) defines modalities for each program

What's in the box? Programs!

Noop $I \otimes J \models \mathbf{1}$ iff $I = J$

Composition $I \otimes J \models \varepsilon_1 \varepsilon_2$ iff $\exists K, I \otimes K \models \varepsilon_1$ and $K \otimes J \models \varepsilon_2$

Assignment $I \otimes J \models \langle \sigma \rangle$ iff $J = I \circ \sigma$

Choice $I \otimes J \models \varepsilon_1 \sqcup \varepsilon_2$ iff $I \otimes J \models \varepsilon_1$ or $I \otimes J \models \varepsilon_2$

Test $I \otimes J \models T?$ iff $I = J$ and $I \models T$

Branch $I \otimes J \models \nabla(T : \varepsilon_1 \parallel \varepsilon_0)$ iff $I \otimes J \models (T? \varepsilon_1) \sqcup (\bar{T}? \varepsilon_0)$

Repeat $I \otimes J \models \varepsilon^*$ iff $\exists I_i, I_0 = I, I_n = J, I_{i-1} \otimes I_i \models \varepsilon$

Loop $I \otimes J \models \square(T : \varepsilon)$ iff $I \otimes J \models (\bar{T}? \varepsilon)^* T?$

Rendezvous $I \otimes J \models \diamond(V : \varepsilon_1 \parallel \varepsilon_0)$ iff $\exists J_1, J_0, I \otimes J_i \models \varepsilon_i$ and $J = J_1 +_V J_0$

Solve $I \otimes J \models [R]$ iff $I \otimes J \models R$

Havoc $I \otimes J \models \forall V$ iff $I \otimes J \models \diamond(V : [T] \parallel 1)$

Not even a new thing! [Fischer, Ladner '79] [Balbiani, Herzig, Troquard '13]

Propositional dynamic logic (PDL) defines modalities for each program

Necessitation law (a.k.a. I can apply programs to a proof)

If $F \models G$ holds, then $\varepsilon.F \models \varepsilon.G$ holds too

What can I do with this?

Proving unsatisfiability F is unsatisfiable if $F \vdash \varepsilon. \perp$ and $\varepsilon. \perp \vdash \perp$

What can I do with this?

Proving unsatisfiability F is unsatisfiable if $F \vdash \varepsilon. \perp$ and $\varepsilon. \perp \vdash \perp$

Proving satisfiability F is satisfiable if $\top \vdash \varepsilon. F$ and $\varepsilon. \perp \vdash \perp$

What can I do with this?

Proving unsatisfiability F is unsatisfiable if $F \vdash \varepsilon. \perp$ and $\varepsilon. \perp \vdash \perp$

Proving satisfiability F is satisfiable if $\top \vdash \varepsilon.F$ and $\varepsilon. \perp \vdash \perp$

Parametric proofs If I have proven $F \vdash G$, then I can prove $F|_\sigma \vdash G|_\sigma$ as $\langle \sigma \rangle.F \vdash \langle \sigma \rangle.G$

What can I do with this?

Proving unsatisfiability F is unsatisfiable if $F \vdash \varepsilon. \perp$ and $\varepsilon. \perp \vdash \perp$

Proving satisfiability F is satisfiable if $\top \vdash \varepsilon.F$ and $\varepsilon. \perp \vdash \perp$

Parametric proofs If I have proven $F \vdash G$, then I can prove $F|_{\sigma} \vdash G|_{\sigma}$ as $\langle \sigma \rangle.F \vdash \langle \sigma \rangle.G$

Proving a safety property P always holds in ε assuming A if $A \vdash \varepsilon^*.P$

What can I do with this?

Proving unsatisfiability F is unsatisfiable if $F \vdash \varepsilon. \perp$ and $\varepsilon. \perp \vdash \perp$

Proving satisfiability F is satisfiable if $\top \vdash \varepsilon.F$ and $\varepsilon. \perp \vdash \perp$

Parametric proofs If I have proven $F \vdash G$, then I can prove $F|_{\sigma} \vdash G|_{\sigma}$ as $\langle \sigma \rangle.F \vdash \langle \sigma \rangle.G$

Proving a safety property P always holds in ε assuming A if $A \vdash \varepsilon^*.P$

Proving a liveness property P eventually holds in ε if $\varepsilon^*.\overline{P} \vdash \perp$

What can I do with this?

Proving unsatisfiability F is unsatisfiable if $F \vdash \varepsilon. \perp$ and $\varepsilon. \perp \vdash \perp$

Proving satisfiability F is satisfiable if $\top \vdash \varepsilon.F$ and $\varepsilon. \perp \vdash \perp$

Parametric proofs If I have proven $F \vdash G$, then I can prove $F|_{\sigma} \vdash G|_{\sigma}$ as $\langle \sigma \rangle.F \vdash \langle \sigma \rangle.G$

Proving a safety property P always holds in ε assuming A if $A \vdash \varepsilon^*.P$

Proving a liveness property P eventually holds in ε if $\varepsilon^*.\overline{P} \vdash \perp$

Refinements ε refines δ if $\varepsilon \vdash [R_{\varepsilon}]$ and $[R_{\delta}] \models \delta$ and $R_{\varepsilon} \vdash R_{\delta}$

What can I do with this?

Proving unsatisfiability F is unsatisfiable if $F \vdash \varepsilon. \perp$ and $\varepsilon. \perp \vdash \perp$

Proving satisfiability F is satisfiable if $\top \vdash \varepsilon.F$ and $\varepsilon. \perp \vdash \perp$

Parametric proofs If I have proven $F \vdash G$, then I can prove $F|_{\sigma} \vdash G|_{\sigma}$ as $\langle \sigma \rangle.F \vdash \langle \sigma \rangle.G$

Proving a safety property P always holds in ε assuming A if $A \vdash \varepsilon^*.P$

Proving a liveness property P eventually holds in ε if $\varepsilon^*.\overline{P} \vdash \perp$

Refinements ε refines δ if $\varepsilon \vdash [R_{\varepsilon}]$ and $[R_{\delta}] \models \delta$ and $R_{\varepsilon} \vdash R_{\delta}$

I think this could be a good foundation for a general, versatile, unified certificate system for propositional reasoning beyond SAT

What can I do with this?

Proving unsatisfiability F is unsatisfiable if $F \vdash \varepsilon. \perp$ and $\varepsilon. \perp \vdash \perp$

Proving satisfiability F is satisfiable if $\top \vdash \varepsilon.F$ and $\varepsilon. \perp \vdash \perp$

Parametric proofs If I have proven $F \vdash G$, then I can prove $F|_{\sigma} \vdash G|_{\sigma}$ as $\langle \sigma \rangle.F \vdash \langle \sigma \rangle.G$

Proving a safety property P always holds in ε assuming A if $A \vdash \varepsilon^*.P$

Proving a liveness property P eventually holds in ε if $\varepsilon^*.\overline{P} \vdash \perp$

Refinements ε refines δ if $\varepsilon \vdash [R_{\varepsilon}]$ and $[R_{\delta}] \models \delta$ and $R_{\varepsilon} \vdash R_{\delta}$

I think this could be a good foundation for a general, versatile, unified certificate system for propositional reasoning beyond SAT

An appetizer: [Rebola-Pardo '25, SYNASC]