

Interpolants and interference

Adrián Rebola-Pardo

TU Wien, JKU Linz (Austria)

4th Workshop on Interpolation: From Proofs to Applications

Haifa (Israel)

August 11th, 2022

Supported by FWF W1255-N23, WWTF VRG11-005, WWTF ICT15-103, and Microsoft Research PhD Programme

SAT
solver



theorem provers
many industrial applications

[Davis, Putnam '60]

SAT
solver

theorem provers
many industrial applications

[Davis, Putnam '60]

model
checking

formal guarantees on
the correctness of
hardware systems

[McMillan '03]

SAT
solver

theorem provers
many industrial applications

[Davis, Putnam '60]

[Craig '57]

abstraction of
information relevant
for correctness

interpolant

model
checking

formal guarantees on
the correctness of
hardware systems


[McMillan '03]

SAT
solver

interpolant

model
checking


Proofs for SAT solving



interpolant

model
checking


Proofs for SAT solving



interpolant

model
checking


Proofs for SAT solving



interpolant

model
checking


Proofs for SAT solving



interpolant

model
checking


Proofs for SAT solving



interpolant

model
checking


Proofs for SAT solving




interpolant

model
checking


Proofs for SAT solving




Proofs for SAT solving




Proofs for SAT solving



Proofs for SAT solving



Proofs for SAT solving



Inference-based proofs

$$\otimes_y \frac{\frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}}}{\otimes_z \frac{\overline{xzt} \quad \overline{xzt}}{\overline{xt}}}$$

Inference-based proofs

$$\otimes_y \frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}} \quad \otimes_y \frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}} \\ \otimes_z \frac{\overline{xzt} \quad \overline{xzt}}{\overline{xt}}$$

any model satisfying the premises...

Inference-based proofs

$$\begin{array}{c} \otimes_y \frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}} \quad \otimes_y \frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}} \\ \otimes_z \frac{\quad}{\overline{xt}} \end{array}$$

any model satisfying the premises...

... also satisfies the conclusion

Inference-based proofs

$$\otimes_y \frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}} \quad \otimes_y \frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}} \\ \otimes_z \frac{\overline{xzt} \quad \overline{xzt}}{\overline{xt}}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}} \quad \otimes_y \frac{\overline{xyz}t \quad \overline{xyz}t}{\overline{xzt}} \\ \otimes_z \frac{\overline{xzt} \quad \overline{xzt}}{\overline{xt}}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\bar{x}yzt}{\bar{x}zt}}{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t} \quad \frac{}{\bar{x}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

$$\begin{array}{c} \otimes_b \frac{\bar{u}ab}{\bar{u}a} \quad \frac{\bar{b}}{a} \quad \frac{\bar{b}}{\bar{a}} \quad \otimes_{\bar{b}} \\ \otimes_a \frac{}{\bar{u}} \\ \otimes_{\bar{u}} \frac{\bar{x}yu}{\bar{x}y} \quad \frac{\bar{y}v}{\bar{y}v} \quad \otimes_{\bar{y}} \\ \otimes_{\bar{x}} \frac{yv}{yv} \quad \frac{\bar{y}v}{\bar{y}v} \quad \frac{\bar{b}}{\bar{v}} \quad \frac{\bar{v}\bar{b}}{\bar{v}} \quad \otimes_{\bar{b}} \\ \otimes_y \frac{}{v} \\ \otimes_v \frac{}{\perp} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\bar{x}yzt}{\bar{x}zt}}{\otimes_z \frac{\bar{x}zt}{\bar{x}t}} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \otimes_b \frac{\bar{u}ab}{\bar{u}a} \quad \frac{\bar{b}}{a} \quad \frac{\bar{b}}{\bar{a}} \quad \otimes_{\bar{b}} \\ \otimes_a \frac{\bar{u}a}{\bar{u}} \quad \frac{\bar{b}}{a} \\ \otimes_{\bar{u}} \frac{\bar{u}}{\bar{x}y} \quad \frac{\bar{x}yu}{\bar{y}v} \quad \frac{\bar{y}v}{xy} \quad \otimes_{\bar{y}} \\ \otimes_{\bar{x}} \frac{\bar{x}y}{y} \quad \frac{y}{\bar{y}v} \\ \otimes_y \frac{y}{v} \quad \frac{\bar{y}v}{\bar{v}} \quad \frac{\bar{b}}{\bar{v}} \quad \frac{\bar{v}\bar{b}}{\bar{b}} \quad \otimes_{\bar{b}} \\ \otimes_v \frac{v}{\perp} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\overline{xyz}t}{\overline{xzt}}}{\otimes_z \frac{\overline{xyz}t}{\overline{xzt}}} \quad \otimes_y \frac{\overline{xyz}t}{\overline{xzt}} \quad \overline{xt}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{c} \text{T} \quad \text{T} \quad \text{T} \quad \text{T} \\ \overline{uab} \quad \overline{b} \quad \overline{b} \quad \overline{ab} \end{array} \\ \otimes_b \frac{\overline{uab} \quad \overline{b} \quad \overline{b} \quad \overline{ab}}{\overline{ua} \quad \overline{a}} \otimes_{\overline{b}} \\ \otimes_a \frac{\overline{ua} \quad \overline{a}}{\overline{u}} \\ \otimes_{\overline{u}} \frac{\overline{u} \quad \overline{xy} \quad \overline{y}v \quad xy}{\overline{xy} \quad \overline{y}v} \otimes_{\overline{y}} \\ \otimes_{\overline{x}} \frac{\overline{xy} \quad \overline{y}v}{y \quad v} \\ \otimes_y \frac{y \quad v}{v} \quad \overline{y}v \quad \overline{b} \quad \overline{v} \quad \overline{b} \\ \otimes_v \frac{v}{\perp} \otimes_{\overline{b}} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}z\bar{t}}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{\bar{u}a \quad \text{T} \quad \bar{a} \quad \text{T}}{\bar{u}} \quad \otimes_{\bar{b}} \\ \otimes_a \frac{\bar{u}}{\bar{x}y} \quad \bar{x}yu \quad \bar{y}v \quad xy \quad \otimes_{\bar{y}} \\ \otimes_{\bar{x}} \frac{yv}{v} \quad \bar{y}v \quad \bar{b} \quad \bar{v} \quad \bar{b} \quad \otimes_{\bar{b}} \\ \otimes_y \frac{v}{\perp} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}z\bar{t}}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{\bar{u}a \quad \text{T} \quad \bar{a} \quad \text{T}}{\bar{u} \quad \text{T}} \quad \otimes_{\bar{b}} \\ \otimes_a \frac{\bar{u} \quad \text{T}}{\bar{x}y} \quad \bar{x}yu \quad \bar{y}v \quad xy \quad \otimes_{\bar{y}} \\ \otimes_{\bar{x}} \frac{yv}{v} \quad \bar{y}v \quad \bar{b} \quad \bar{v} \quad \bar{b} \quad \otimes_{\bar{b}} \\ \otimes_y \frac{v}{\perp} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}z\bar{t}}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{\bar{u}a \quad \text{T} \quad \bar{a} \quad \text{T}}{\bar{u} \quad \text{T}} \quad \otimes_{\bar{b}} \frac{\perp}{\bar{x}yu} \\ \otimes_a \frac{\bar{u} \quad \text{T}}{\bar{x}y} \quad \frac{\bar{y}v \quad xy}{\bar{y}v} \quad \otimes_{\bar{y}} \\ \otimes_{\bar{x}} \frac{yv}{v} \quad \frac{\bar{y}v \quad \bar{b} \quad \bar{v}\bar{b}}{\bar{v}} \quad \otimes_{\bar{b}} \\ \otimes_y \frac{v}{\perp} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\bar{x}yzt}{\bar{x}zt}}{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t} \quad \frac{\quad}{\bar{x}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{\quad}{\bar{u}a} \quad \text{T} \quad \frac{\quad}{\bar{a}} \quad \text{T} \quad \otimes_{\bar{b}} \\ \otimes_a \frac{\quad}{\bar{u}} \quad \text{T} \quad \frac{\quad}{\bar{x}y} \quad u \quad \frac{\bar{x}yu}{\bar{y}v} \quad \otimes_{\bar{y}} \\ \otimes_{\bar{u}} \quad \frac{\quad}{\bar{x}} \quad \frac{yv}{\bar{y}v} \quad \otimes_y \\ \otimes_v \frac{\quad}{\bar{v}} \quad \frac{\bar{b} \quad \bar{v}\bar{b}}{\bar{v}} \quad \otimes_{\bar{b}} \\ \hline \perp \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \otimes_b \frac{\bar{u}ab}{\bar{u}a} & \frac{\bar{b}}{\text{T}} & \frac{\bar{b}}{\bar{a}} & \frac{\bar{a}b}{\text{T}} \\ \hline \otimes_a \frac{\bar{u}a}{\bar{u}} & \text{T} & \bar{a} & \text{T} \\ \hline \otimes_{\bar{u}} \frac{\bar{u}}{\bar{x}y} & \text{T} & & \\ \hline \otimes_{\bar{x}} \frac{\bar{x}y}{y} & u & & \\ \hline \otimes_y \frac{y}{v} & & \bar{y}v & \\ \hline \otimes_v \frac{v}{\perp} & & \bar{v} & \frac{\bar{b}}{\bar{v}} \end{array} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{}{\bar{u}a \quad \text{T} \quad \bar{a} \quad \text{T}} \\ \otimes_a \frac{}{\bar{u} \quad \text{T}} \\ \otimes_{\bar{u}} \frac{}{\bar{x}y \quad u} \\ \otimes_x \frac{}{yv} \\ \otimes_y \frac{}{v} \\ \otimes_v \frac{}{\perp} \end{array} \quad \begin{array}{ccc} \perp & \perp & \perp \\ \hline \bar{x}yu & \bar{y}v & xy \\ \hline \bar{y}v & \bar{b} & \bar{v}\bar{b} \\ \hline \bar{v} \end{array} \quad \begin{array}{c} \otimes_{\bar{y}} \\ \otimes_{\bar{b}} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{c} \text{T} \quad \text{T} \quad \text{T} \quad \text{T} \\ \hline \bar{u}ab \quad \bar{b} \quad \bar{b} \quad \bar{a}b \\ \hline \otimes_b \frac{\bar{u}a \quad \text{T}}{\bar{u}} \quad \frac{\bar{a} \quad \text{T}}{\bar{a}} \quad \otimes_{\bar{b}} \end{array} \\ \begin{array}{c} \perp \quad \perp \quad \perp \\ \hline \bar{x}yu \quad \bar{y}v \quad xy \\ \hline \otimes_{\bar{u}} \frac{\bar{x}y \quad u}{\bar{x}} \quad \frac{\bar{y}v \quad \perp}{\bar{y}} \quad \otimes_{\bar{y}} \end{array} \\ \begin{array}{c} \perp \\ \hline yv \quad u \quad \bar{y}v \quad \bar{b} \quad \bar{v}\bar{b} \\ \hline \otimes_y \frac{yv \quad u}{y} \quad \frac{\bar{y}v}{\bar{y}} \quad \frac{\bar{b} \quad \bar{v}\bar{b}}{\bar{v}} \quad \otimes_{\bar{b}} \end{array} \\ \begin{array}{c} \perp \\ \hline v \\ \hline \otimes_v \end{array} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\overline{xyz}t}{\overline{xzt}}}{\otimes_z \frac{\overline{xyz}t}{\overline{xzt}}} \quad \otimes_y \frac{\overline{xyz}t}{\overline{xzt}} \quad \overline{xt}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \overline{uab} & \overline{b} & \overline{b} & \overline{ab} \end{array} \\ \otimes_b \frac{\overline{uab} \quad \overline{b} \quad \overline{b} \quad \overline{ab}}{\overline{ua} \quad \text{T} \quad \overline{a} \quad \text{T}} \quad \otimes_{\overline{b}} \\ \otimes_a \frac{\overline{ua} \quad \text{T} \quad \overline{a} \quad \text{T}}{\overline{u} \quad \text{T}} \quad \perp \quad \perp \quad \perp \\ \otimes_{\overline{u}} \frac{\overline{u} \quad \text{T}}{\overline{xy} \quad u} \quad \overline{xy}u \quad \overline{y}v \quad xy \\ \otimes_x \frac{\overline{xy} \quad u}{y \quad v} \quad \overline{y}v \quad \perp \quad \otimes_{\overline{y}} \\ \otimes_y \frac{y \quad v}{v} \quad \overline{y}v \quad \perp \quad \overline{b} \quad \overline{v} \quad \overline{b} \\ \otimes_v \frac{v}{\perp} \quad \overline{v} \quad \otimes_{\overline{b}} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{\bar{u}a \quad \text{T} \quad \bar{a} \quad \text{T}}{\bar{u} \quad \text{T}} \quad \otimes_{\bar{b}} \frac{\perp}{\bar{x}yu} \quad \frac{\perp}{\bar{y}v} \quad \frac{\perp}{xy} \\ \otimes_a \frac{\bar{u} \quad \text{T}}{\bar{x}y \quad u} \quad \otimes_{\bar{y}} \frac{\perp}{\bar{y}v} \quad \frac{\perp}{v \quad u} \\ \otimes_u \frac{\bar{x}y \quad u}{y \quad v} \quad \frac{\perp}{\bar{y}v} \quad \frac{\bar{b} \quad \bar{v} \quad \bar{b}}{\bar{v}} \quad \otimes_{\bar{v}} \frac{\perp}{\perp} \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\bar{x}yzt}{\bar{x}zt}}{\otimes_z \frac{\bar{x}zt}{\bar{x}t}}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{}{\bar{u}a \text{ T} \quad \bar{a} \text{ T}} \\ \otimes_a \frac{}{\bar{u} \text{ T}} \\ \otimes_{\bar{u}} \frac{}{\bar{x}y \text{ u}} \\ \otimes_x \frac{}{y \text{ v} \text{ u}} \\ \otimes_y \frac{}{v \text{ u}} \\ \otimes_v \frac{}{\perp} \end{array} \quad \begin{array}{ccc} \perp & \perp & \perp \\ \hline \bar{x}yu & \bar{y}v & xy \\ \hline \bar{y}v & \perp & \perp \\ \hline \bar{y}v & \bar{b} & \bar{v} \bar{b} \\ \hline \perp \end{array} \quad \begin{array}{ccc} \perp & \text{T} & \text{T} \\ \hline \bar{y}v & \bar{b} & \bar{v} \bar{b} \\ \hline \perp \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{}{} \quad \otimes_a \frac{}{} \quad \otimes_{\bar{u}} \frac{}{} \quad \otimes_{\bar{a}} \frac{}{} \quad \otimes_{\bar{b}} \frac{}{} \\ \hline \bar{u}a \quad \text{T} \quad \bar{a} \quad \text{T} \\ \hline \bar{u} \quad \text{T} \\ \hline \bar{x}y \quad u \\ \hline yv \quad u \\ \hline v \quad u \\ \hline \perp \end{array}$$

Inference-based proofs and recursive interpolation

Inference-based proofs

$$\otimes_y \frac{\frac{\otimes_z \frac{\bar{x}yzt}{\bar{x}zt}}{\bar{x}t}}{\bar{x}t} \quad \otimes_y \frac{\bar{x}y\bar{z}t}{\bar{x}\bar{z}t}$$

any model satisfying the premises...

... also satisfies the conclusion

Resolution proofs [Davis, Putnam '60]

RUP proofs [Goldberg, Novikov '03]

Generating interpolants from inference-based proofs [Huang '95]

Partial interpolant

$$A \models P(C) \vee C|_{\text{Var}(A)}$$

$$B \models \neg P(C) \vee C|_{\text{Var}(B)}$$

$$\text{Var}(P(C)) \subseteq \text{Var}(A) \cap \text{Var}(B)$$

$$\begin{array}{c} \begin{array}{cccc} \text{T} & \text{T} & \text{T} & \text{T} \\ \hline \bar{u}ab & \bar{b} & \bar{b} & \bar{a}b \end{array} \\ \otimes_b \frac{}{} \quad \otimes_a \frac{}{} \quad \otimes_{\bar{u}} \frac{}{} \quad \otimes_{\bar{a}} \frac{}{} \quad \otimes_{\bar{b}} \frac{}{} \\ \hline \bar{u}a \quad \text{T} \quad \bar{a} \quad \text{T} \\ \hline \bar{u} \quad \text{T} \\ \hline \bar{x}y \quad u \\ \hline yv \quad u \\ \hline v \quad u \\ \hline \perp \quad \boxed{u \vee v} \end{array}$$

Interference-based proofs

$i:\bar{x}y$

$i:\bar{z}t$

$d:\bar{x}yzt$

$d:xy\bar{z}t$

$i:\bar{x}t$

Interference and non-monotonicity

Interference-based proofs

$i:\bar{x}y$

$i:\bar{z}t$

$d:\bar{x}yzt$

$d:xy\bar{z}t$

$i:\bar{x}t$

if *some* model satisfies
the original formula...

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$

d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$

d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

[Reckhow '75; Kullman '99]

ER

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$

d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

[Wetzler, Heule, Hunt '14]



Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$

d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

[Heule, Kiesl, Biere '17]



Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

[Buss, Thapen '19]



Interference and non-monotonicity

Interference-based proofs

$i:\bar{x}y$

$i:\bar{z}t$

$d:\bar{x}yzt$


$d:xy\bar{z}t$

$i:\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

[Rebola-Pardo '22]



Interference and non-monotonicity

Interference-based proofs

[Kiesl, Rebola-Pardo, Heule '18; Heule, Biere '18]

$i:\bar{x}y$

$i:\bar{z}t$

$d:\bar{x}yzt$


$d:xy\bar{z}t$

$i:\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Interference and non-monotonicity

Interference-based proofs

$i:\bar{x}y$

$i:\bar{z}t$

$d:\bar{x}yzt$


$d:xy\bar{z}t$

$i:\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

A clause C can be introduced in a formula F if a witness substitution σ is given such that

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

A clause C can be introduced in a formula F if a witness substitution σ is given such that for each clause $D \in F$, the clause $C \vee \sigma(D)$ can be derived from F through chained resolution.

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

A clause C can be introduced in a formula F if a witness substitution σ is given such that for each clause $D \in F$, the clause $C \vee \sigma(D)$ can be derived from F through chained resolution.

Theorem If C is SR over F , then F is satisfiability-equivalent to $F \wedge C$

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

A clause C can be introduced in a formula F if a witness substitution σ is given such that for each clause $D \in F$, the clause $C \vee \sigma(D)$ can be derived from F through chained resolution.

Theorem If C is SR over F , then F is satisfiability-equivalent to $F \wedge C$

Interference DSR depends on the presence *and absence* of clauses in F

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

A clause C can be introduced in a formula F if a witness substitution σ is given such that **for each clause $D \in F$** , the clause $C \vee \sigma(D)$ can be derived from F through chained resolution.

Theorem If C is SR over F , then F is satisfiability-equivalent to $F \wedge C$

Interference DSR depends on the presence *and absence* of clauses in F

Non-monotonicity adding clauses to F may make an introduction incorrect

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

A clause C can be introduced in a formula F if a witness substitution σ is given such that for each clause $D \in F$, the clause $C \vee \sigma(D)$ can be derived from F through chained resolution.

Theorem If C is SR over F , then F is satisfiability-equivalent to $F \wedge C$

Interference DSR depends on the presence *and absence* of clauses in F

Non-monotonicity adding clauses to F may make an introduction incorrect

no tree-shaped proofs for interference

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

A clause C can be introduced in a formula F if a witness substitution σ is given such that for each clause $D \in F$, the clause $C \vee \sigma(D)$ can be derived from F through chained resolution.

Theorem If C is SR over F , then F is satisfiability-equivalent to $F \wedge C$

Interference DSR depends on the presence *and absence* of clauses in F

Non-monotonicity adding clauses to F may make an introduction incorrect

no tree-shaped proofs for interference

no recursive interpolation for interference

Interference and non-monotonicity

Interference-based proofs

i: $\bar{x}y$

i: $\bar{z}t$

d: $\bar{x}yzt$


d: $xy\bar{z}t$

i: $\bar{x}t$

if *some* model satisfies
the original formula...

... then *some* model satisfies
the accumulated formula

all these have the same proof complexity



Subsumption redundancy (SR) [Buss, Thapen '19]

A clause C can be introduced in a formula F if a witness substitution σ is given such that for each clause $D \in F$, the clause $C \vee \sigma(D)$ can be derived from F through chained resolution.

Theorem If C is SR over F , then F is satisfiability-equivalent to $F \wedge C$

Interference DSR depends on the presence *and absence* of clauses in F

Non-monotonicity adding clauses to F may make an introduction incorrect

no tree-shaped proofs for interference

no recursive interpolation for interference

unless...

atomic substitution σ

maps variables to literals, T or \perp

atomic substitution σ

cube Q

maps variables to literals, \top or \perp

conjunction of literals

Mutation logic

atomic substitution σ

cube Q

mutation rule $\sigma :- Q$

maps variables to literals, \top or \perp

conjunction of literals

“if Q is satisfied, then apply σ ”

Mutation logic

atomic substitution	σ	maps variables to literals, \top or \perp
cube	Q	conjunction of literals
mutation rule	$\sigma :- Q$	“if Q is satisfied, then apply σ ”
model mutation	$I \circ (\sigma :- Q)$	if $I \models Q$, then $I \circ \sigma$; otherwise, I

Mutation logic

atomic substitution	σ	maps variables to literals, \top or \perp
cube	Q	conjunction of literals
mutation rule	$\sigma :- Q$	“if Q is satisfied, then apply σ ”
model mutation	$I \circ (\sigma :- Q)$	if $I \models Q$, then $I \circ \sigma$; otherwise, I
mutated formula	$\nabla(\sigma :- Q). \varphi$	$I \models \nabla(\sigma :- Q). \varphi$ iff $I \circ (\sigma :- Q) \models \varphi$

atomic substitution	σ	maps variables to literals, \top or \perp
cube	Q	conjunction of literals
mutation rule	$\sigma :- Q$	“if Q is satisfied, then apply σ ”
model mutation	$I \circ (\sigma :- Q)$	if $I \models Q$, then $I \circ \sigma$; otherwise, I
mutated formula	$\nabla(\sigma :- Q). \varphi$	$I \models \nabla(\sigma :- Q). \varphi$ iff $I \circ (\sigma :- Q) \models \varphi$

Theorem [Rebola-Pardo, Suda '18] if C is a substitution-redundant clause over F witnessed by σ , then $F \models \nabla(\sigma :- \overline{C}). F \wedge C$

atomic substitution	σ	maps variables to literals, \top or \perp
cube	Q	conjunction of literals
mutation rule	$\sigma :- Q$	“if Q is satisfied, then apply σ ”
model mutation	$I \circ (\sigma :- Q)$	if $I \models Q$, then $I \circ \sigma$; otherwise, I
mutated formula	$\nabla(\sigma :- Q). \varphi$	$I \models \nabla(\sigma :- Q). \varphi$ iff $I \circ (\sigma :- Q) \models \varphi$

Theorem [Rebola-Pardo, Suda '18] if C is a substitution-redundant clause over F witnessed by σ , then $F \models \nabla(\sigma :- \overline{C}). F \wedge C$

atomic substitution	σ	maps variables to literals, \top or \perp
cube	Q	conjunction of literals
mutation rule	$\sigma :- Q$	“if Q is satisfied, then apply σ ”
model mutation	$I \circ (\sigma :- Q)$	if $I \models Q$, then $I \circ \sigma$; otherwise, I
mutated formula	$\nabla(\sigma :- Q). \varphi$	$I \models \nabla(\sigma :- Q). \varphi$ iff $I \circ (\sigma :- Q) \models \varphi$

Theorem [Rebola-Pardo, Suda '18] if C is a substitution-redundant clause over F witnessed by σ , then $F \models \nabla(\sigma :- \overline{C}). F \wedge C$

Intuition interference encodes reasoning without loss of generality
 C holds; otherwise we apply σ and get a similar problem where it does

atomic substitution	σ	maps variables to literals, \top or \perp
cube	Q	conjunction of literals
mutation rule	$\sigma :- Q$	“if Q is satisfied, then apply σ ”
model mutation	$I \circ (\sigma :- Q)$	if $I \models Q$, then $I \circ \sigma$; otherwise, I
mutated formula	$\nabla(\sigma :- Q). \varphi$	$I \models \nabla(\sigma :- Q). \varphi$ iff $I \circ (\sigma :- Q) \models \varphi$

Theorem [Rebola-Pardo, Suda '18] if C is a substitution-redundant clause over F witnessed by σ , then $F \models \nabla(\sigma :- \overline{C}). F \wedge C$

Intuition interference encodes reasoning without loss of generality
 C holds; otherwise we apply σ and get a similar problem where it does

Mutation resolution an inference-based proof system over mutated clauses that fully encodes all interference-based proof systems

atomic substitution	σ	maps variables to literals, \top or \perp
cube	Q	conjunction of literals
mutation rule	$\sigma :- Q$	“if Q is satisfied, then apply σ ”
model mutation	$I \circ (\sigma :- Q)$	if $I \models Q$, then $I \circ \sigma$; otherwise, I
mutated formula	$\nabla(\sigma :- Q). \varphi$	$I \models \nabla(\sigma :- Q). \varphi$ iff $I \circ (\sigma :- Q) \models \varphi$

Theorem [Rebola-Pardo, Suda '18] if C is a substitution-redundant clause over F witnessed by σ , then $F \models \nabla(\sigma :- \overline{C}). F \wedge C$

Intuition interference encodes reasoning without loss of generality
 C holds; otherwise we apply σ and get a similar problem where it does

Mutation resolution an inference-based proof system over mutated clauses that fully encodes all interference-based proof systems

tree-shaped proofs for interference (within ML)

atomic substitution	σ	maps variables to literals, \top or \perp
cube	Q	conjunction of literals
mutation rule	$\sigma :- Q$	“if Q is satisfied, then apply σ ”
model mutation	$I \circ (\sigma :- Q)$	if $I \models Q$, then $I \circ \sigma$; otherwise, I
mutated formula	$\nabla(\sigma :- Q). \varphi$	$I \models \nabla(\sigma :- Q). \varphi$ iff $I \circ (\sigma :- Q) \models \varphi$

Theorem [Rebola-Pardo, Suda '18] if C is a substitution-redundant clause over F witnessed by σ , then $F \models \nabla(\sigma :- \overline{C}). F \wedge C$


Intuition interference encodes reasoning without loss of generality
 C holds; otherwise we apply σ and get a similar problem where it does

Mutation resolution an inference-based proof system over mutated clauses that fully encodes all interference-based proof systems


tree-shaped proofs for interference (within ML)

no recursive interpolation for interference?


Interpolants in mutation logic?



Interpolants in mutation logic?



Interpolants in mutation logic?




Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment

linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Interpolants in mutation logic?




Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment

linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Interpolants in mutation logic?




Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment

linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Interpolants in mutation logic?




Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment

linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Interpolants in mutation logic?




Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment

linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Interpolants in mutation logic?



Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment


linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Propositional interpolants from MRes proofs

- ER p-simulates DRAT [Kiesl, Rebola-Pardo, Heule '18]
- EF p-simulates ER [Reckhow '75]
- RSA is insecure \Rightarrow EF lacks feasible interpolation [Krajíček, Pudlák '98]

Interpolants in mutation logic?



Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment


linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Propositional interpolants from MRes proofs

- ER p-simulates DRAT [Kiesl, Rebola-Pardo, Heule '18]
- EF p-simulates ER [Reckhow '75]
- RSA is insecure \Rightarrow EF lacks feasible interpolation [Krajíček, Pudlák '98]

Interpolants in mutation logic?



Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment


linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Propositional interpolants from MRes proofs

- ER p-simulates DRAT [Kiesl, Rebola-Pardo, Heule '18]
- EF p-simulates ER [Reckhow '75]
- RSA is insecure \Rightarrow EF lacks feasible interpolation [Krajíček, Pudlák '98]

Interpolants in mutation logic?



Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment

linearly as expressive as full mutation propositional logic


satisfiability problem reduces linearly to SAT

Propositional interpolants from MRes proofs

- ER p-simulates DRAT [Kiesl, Rebola-Pardo, Heule '18]
- EF p-simulates ER [Reckhow '75]
- RSA is insecure \Rightarrow EF lacks feasible interpolation [Krajíček, Pudlák '98]

Mutation-propositional interpolants from MRes proofs

Interpolants in mutation logic?



Universally mutated CNF $\nabla(\sigma_1 :- Q_1) \dots \nabla(\sigma_n :- Q_n).F$

DRAT operates naturally in this fragment

linearly as expressive as full mutation propositional logic

satisfiability problem reduces linearly to SAT

Propositional interpolants from MRes proofs

- ER p-simulates DRAT [Kiesl, Rebola-Pardo, Heule '18]
- EF p-simulates ER [Reckhow '75]
- RSA is insecure \Rightarrow EF lacks feasible interpolation [Krajíček, Pudlák '98]

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Is y a variable in $\nabla(\{y \mapsto \top\} :- x). \nabla(\{y \mapsto \perp\} :- \bar{x}). y$?

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Is y a variable in $\nabla(\{y \mapsto \top\} :- x). \nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x$?

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Is y a variable in $\nabla(\{y \mapsto \top\} :- x). \nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x$?

Is y a variable in $\nabla(\{y \mapsto \perp\} :- \bar{x}). y$?

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Is y a variable in $\nabla(\{y \mapsto \top\} :- x). \nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x$?

Is y a variable in $\nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x \wedge y$?

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Is y a variable in $\nabla(\{y \mapsto \top\} :- x). \nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x$?

Is y a variable in $\nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x \wedge y$?

Semantic variables

A variable x is a **semantic variable** of a formula φ whenever we can find two models I^+ and I^- differing only on their truth value on x such that $I^+ \models \varphi$ and $I^- \not\models \varphi$. \Rightarrow semantic interpolants

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Is y a variable in $\nabla(\{y \mapsto \top\} :- x). \nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x$?

Is y a variable in $\nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x \wedge y$?

Semantic variables

A variable x is a **semantic variable** of a formula φ whenever we can find two models I^+ and I^- differing only on their truth value on x such that $I^+ \models \varphi$ and $I^- \not\models \varphi$. \Rightarrow semantic interpolants

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Is y a variable in $\nabla(\{y \mapsto \top\} :- x). \nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x$?

Is y a variable in $\nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x \wedge y$?

Semantic variables

A variable x is a **semantic variable** of a formula φ whenever we can find two models I^+ and I^- differing only on their truth value on x such that $I^+ \models \varphi$ and $I^- \not\models \varphi$. \Rightarrow **semantic interpolants**

What gives?

- Semantic interpolants are stable under logical equivalence
- Finding if x is a variable of φ is NP-complete
but they are bounded by syntactic variables
- Enables extending **some** recursive interpolation rules to mutation logic

Redefining interpolants

Mutation-propositional interpolants from MRes proofs

What are the variables of a mutation formula?

Is y a variable in $\nabla(\{y \mapsto \top\} :- x). \nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x$?

Is y a variable in $\nabla(\{y \mapsto \perp\} :- \bar{x}). y \equiv x \wedge y$?

Semantic variables

A variable x is a **semantic variable** of a formula φ whenever we can find two models I^+ and I^- differing only on their truth value on x such that $I^+ \models \varphi$ and $I^- \not\models \varphi$. \Rightarrow **semantic interpolants**

What gives?

- Semantic interpolants are stable under logical equivalence
- Finding if x is a variable of φ is NP-complete
but they are bounded by syntactic variables
- Enables extending **some** recursive interpolation rules to mutation logic

Questions

- Does this affect the unfeasible interpolation [Krajíček, Pudlák '98]?
- Do other similar variations on the notion of interpolant exist?
- Does this shed any light on the connection between feasible interpolation and cryptography?