

# MATH 131 SECTION, IV: COMPLETION OF METRIC SPACES

ARPON RAKSIT

## 1. PRELIMINARIES

Let's first remember what convergent and Cauchy sequences and completeness are in metric spaces.

**1.1. Notation.** Throughout we let  $(X, d_X)$  be a metric space.

**1.2. Definition.** A sequence  $(x_k) \in X^{\mathbb{N}}$  is called *convergent* if there exists  $x \in X$  such that for each  $\epsilon > 0$  there exists some  $n \in \mathbb{N}$  such that  $d_X(x, x_k) < \epsilon$  for all  $k \geq n$ . In this case, we say  $(x_k)$  *converges to*  $x$  and write  $\lim_{k \rightarrow \infty} x_k = x$ , or equivalently  $x_k \rightarrow x$  as  $k \rightarrow \infty$ .

**1.3. Definition.** A sequence  $(x_k) \in X^{\mathbb{N}}$  is called *Cauchy* if for each  $\epsilon > 0$  there exists some  $n \in \mathbb{N}$  such that  $d_X(x_j, x_k) < \epsilon$  for all  $j, k \geq n$ .

**1.4.** You might have seen Cauchy sequences defined in the past for the case  $X = \mathbb{R}$ . And you might have learned that a sequence being Cauchy is *equivalent* to the sequence being convergent. But this equivalence is something special to  $\mathbb{R}$ —it says that  $\mathbb{R}$  is a complete metric space, which we will define in a second. But first check that we always have one implication.

**1.5. Exercise.** *All convergent sequences are Cauchy.*

**1.6. Definition.** We say  $X$  is *complete* if all Cauchy sequences in  $X$  are convergent in  $X$ .

**1.7. Examples.** As stated above  $\mathbb{R}$  is complete. This is sort of a key fact when we're learning real analysis. So maybe it's counterintuitive that there are spaces which are *not* complete. But we don't have to stray too far for some examples!

- $(0, 1)$  is not complete: it's easy to check that the sequence given by  $x_k := 1/k$  for  $k \in \mathbb{N}$  is Cauchy, but for any  $x \in (0, 1)$  there exists  $n \in \mathbb{N}$  such that  $1/k < x$  for all  $k \geq n$ , whence  $(x_k)$  can't converge to  $x$ .
- $\mathbb{Q}$  is not complete: I won't say this is as rigourously as the previous example, but rational approximations to irrational numbers (e.g., 1.4, 1.41, 1.414, ... tending to  $\sqrt{2}$ ) are Cauchy but cannot be convergent in  $\mathbb{Q}$ .

For another example of a complete metric space, look at the fourth problem set: we can take  $X$  the space of bounded sequences in  $\mathbb{R}$ , equipped with the sup metric.

**1.8.** Perhaps you've noticed something a little subtle about the difference between convergent and Cauchy sequences. Namely, the notion of convergence depends on the space  $X$  in a way that notion of Cauchyness does not. What I mean is: we said that  $(1/k)_{k \in \mathbb{N}}$  doesn't converge in  $(0, 1)$  even though it's Cauchy—but certainly it

converges in  $[0, 1]$  or  $\mathbb{R}$ ! On the other hand, passing to a larger ambient space doesn't change anything about what it means to be a Cauchy sequence.

This is essentially because our definition of convergence requires us to actually produce a limit  $x \in X$  (so of course it depends on  $X$  in the way described above), whereas our definition of Cauchy is completely intrinsic, in that it only refers to the elements of our given sequence.

Finally let's recall a couple of properties that maps between metric spaces can have.

**1.9. Definition.** Let  $(Y, d_Y)$  another metric space. A map  $\phi: X \rightarrow Y$  is:

- (1) *uniformly continuous* if for each  $\epsilon > 0$  there exists a  $\delta > 0$  such that  $d_X(x, y) < \delta \implies d_Y(\phi(x), \phi(y)) < \epsilon$  for all  $x, y \in X$ ;
- (2) an *isometry* (or “distance-preserving”) if  $d_X(x, y) = d_Y(\phi(x), \phi(y))$  for all  $x, y \in X$ ;
- (3) an *isomorphism (of metric spaces)* if  $\phi$  is a bijective isometry. (Note the inverse of an isomorphism is clearly automatically an isometry.)

**1.10. Exercise.** Let  $\phi: X \rightarrow Y$  a map of metric spaces.

- (1) If  $\phi$  is uniformly continuous and  $(x_k) \in X^{\mathbb{N}}$  is a Cauchy sequence, then  $(\phi(x_k)) \in Y^{\mathbb{N}}$  is also Cauchy.
- (2) If  $\phi$  is an isometry then  $\phi$  is uniformly continuous and an embedding of topological spaces.

## 2. METRIC COMPLETION

The goal for the rest of these notes is to prove the following.

**2.1. Proposition.** There exist a metric space  $X_c$  and an isometry  $\phi: X \rightarrow X_c$  such that:

- (1)  $X_c$  is complete;
- (2)  $\phi(X)$  is dense in  $X_c$ ;
- (3) for any complete metric space  $Y$  and uniformly continuous map  $\psi: X \rightarrow Y$ , there exists a unique uniformly continuous map  $\psi_c: X_c \rightarrow Y$  such that  $\psi_c \circ \phi = \psi$ ;
- (4) if  $Y_c$  is a complete metric space and  $\psi: X \rightarrow Y_c$  an isometry such that  $\psi(X)$  is dense in  $Y_c$ , then there exists a unique isomorphism  $\eta: X_c \rightarrow Y_c$  such that  $\eta \circ \phi = \psi$ .

**2.2. Definition.** Since  $X_c$  is unique up to unique isomorphism, we are justified in calling  $X_c$  the *completion* of  $X$ .

There are a number of steps in proving the proposition, so I won't even try to put them all inside one proof environment! (Actually, as—well, if—you read along, you should try to prove the claims before reading on. This is a decent exercise in knowing how to prove fundamental things about metric spaces.) Here we go.

**2.3 (Construction).** First define  $C := \{(x_k) \in X^{\mathbb{N}} \mid (x_k) \text{ is Cauchy}\}$  the set of all Cauchy sequences in  $X$ . We then define an equivalence relation on  $C$  by saying

$$(x_k) \sim (y_k) \iff \lim_{k \rightarrow \infty} d_X(x_k, y_k) = 0.$$

Reflexivity and symmetry of this relation are obvious. For transitivity we just use nonnegativity of the metric and the triangle inequality: if  $(x_k) \sim (y_k)$  and  $(y_k) \sim (z_k)$

then

$$\begin{aligned} 0 \leq \lim_{k \rightarrow \infty} d_X(x_k, z_k) &\leq \lim_{k \rightarrow \infty} d_X(x_k, y_k) + d_X(y_k, z_k) \\ &\leq \lim_{k \rightarrow \infty} d_X(x_k, y_k) + \lim_{k \rightarrow \infty} d_X(y_k, z_k) = 0, \end{aligned}$$

which implies  $(x_k) \sim (z_k)$ .

We then define  $X_c := C/\sim$  to be the set of equivalence classes of  $C$  under this relation, and define a metric  $d_{X_c}$  on  $X_c$  as follows. For  $\sigma, \tau \in X_c$ , with equivalence class representatives  $(x_k), (y_k) \in C$ , respectively, we set

$$d_{X_c}(\sigma, \tau) := \lim_{k \rightarrow \infty} d_X(x_k, y_k).$$

Before we define  $\phi$  or prove properties (1) and (2) from the proposition, we should  $d_{X_c}$  is well-defined and satisfies the axioms of a metric.

**2.4** (The metric is well-defined). Let  $\sigma, \tau \in X_c$  with representatives  $(x_n), (y_n) \in C$ , respectively. Since  $(x_n), (y_n)$  are Cauchy, there exists  $n \in \mathbb{N}$  such that

$$d_X(x_j, x_k) < \epsilon \quad \text{and} \quad d_E(y_j, y_k) < \epsilon \quad \text{if } j, k \geq n.$$

By the triangle inequality we have

$$\begin{aligned} d_X(x_j, y_j) &\leq d_X(x_j, x_k) + d_X(x_k, y_k) + d_X(y_j, y_k) \\ \implies d_X(x_j, y_j) - d_X(x_k, y_k) &< 2\epsilon, \end{aligned}$$

and similarly  $d_X(x_k, y_k) - d_X(x_j, y_j) < 2\epsilon$ , for  $j, k \geq n$ . It follows that the sequence  $(d_E(x_n, y_n))$  in  $\mathbb{R}$  is Cauchy. Since  $\mathbb{R}$  is complete,

$$d_{X_c}(\sigma, \tau) = \lim_{n \rightarrow \infty} d_E(x_n, y_n)$$

exists.

Now suppose  $(x_k) \sim (x'_k)$  and  $(y_k) \sim (y'_k)$ . Then by the triangle inequality

$$\begin{aligned} \lim_{k \rightarrow \infty} d_X(x'_k, y'_k) &\leq \lim_{k \rightarrow \infty} d_X(x'_k, x_k) + \lim_{k \rightarrow \infty} d_X(x_k, y_k) + \lim_{k \rightarrow \infty} d_X(y_k, y'_k) \\ &= \lim_{k \rightarrow \infty} d_X(x_k, y_k). \end{aligned}$$

By symmetry the reverse inequality holds as well, so  $d_{X_c}$  is well-defined.

**2.5** (The metric is a metric). Next we show  $d_{X_c}$  is in fact a metric on  $X_c$ . Let  $\sigma, \tau \in X_c$  with representatives  $(x_n), (y_n) \in C$ , respectively. Nonnegativity symmetry, and the triangle inequality for  $d_{X_c}$  follow immediately from these holding for  $d_X$ . And that  $d_{X_c}(\sigma, \tau) = 0$  if and only if  $\sigma = \tau$  follows by definition of  $d_{X_c}$  and the relation  $\sim$ .

**2.6** (The isometric embedding). Define  $\phi: X \rightarrow X_c$  by letting  $\phi(x)$  be the equivalence class of the constant sequence given by  $x_k := x$  for  $k \in \mathbb{N}$ . Then for  $x, y \in X$  we have

$$d_X(\phi(x), \phi(y)) = \lim_{k \rightarrow \infty} d_X(x, y) = d_X(x, y),$$

so  $\phi$  is an isometry.

Next we show that  $\phi(X)$  is dense in  $X_c$ . Let  $\sigma \in X_c$  with representative  $(x_n) \in C$ . Let  $\epsilon > 0$ . Since  $(x_n)$  is Cauchy there exists  $n \in \mathbb{N}$  such that  $d_X(x_k, x_n) < \epsilon$  for all  $k \geq N$ . It follows that

$$d_{X_c}(\sigma, \phi(x_n)) = \lim_{k \rightarrow \infty} d_E(x_k, x_n) < \epsilon.$$

I.e.,  $B_{d_{X_c}}(\sigma, \epsilon)$  intersects  $\phi(X)$  for any  $\sigma \in X_c$  and  $\epsilon > 0$ . So indeed  $\phi(X)$  is dense in  $X_c$ .

**2.7** (The completion is complete). Finally we show that  $X_c$  is complete. Let  $(\sigma_k) \in X_c^{\mathbb{N}}$  a Cauchy sequence in  $X_c$ . Then for each  $r \in \mathbb{N}$  there exists  $n_r \in \mathbb{N}$  such that  $d_{X_c}(\sigma_k, \sigma_{n_r}) < 1/r$  for  $k \geq n_r$ . Since  $\phi(X)$  is dense in  $X_c$ , there exists  $x_r \in E$  such that  $d_{X_c}(\sigma_{n_r}, \phi(x_r)) < 1/r$  for each  $r \in \mathbb{N}$ .<sup>1</sup> We claim  $(x_r) \in X^{\mathbb{N}}$  is Cauchy. Let  $\epsilon > 0$ , and choose  $n \in \mathbb{N}$  such that  $3/n < \epsilon$ . Then for  $k \geq j \geq n$  we have by the triangle inequality and the fact that  $\phi$  is an isometry that

$$\begin{aligned} d_X(x_j, x_k) &= d_{X_c}(\phi(x_j), \phi(x_k)) \\ &\leq d_{X_c}(\phi(x_j), \sigma_{n_j}) + d_{X_c}(\sigma_{n_j}, \sigma_{n_k}) + d_{X_c}(\sigma_{n_k}, \phi(x_k)) \\ &\leq 1/j + 1/j + 1/k \leq 3/n < \epsilon, \end{aligned}$$

so indeed  $(x_r)$  is Cauchy.

Then let  $\sigma \in X_c$  be the equivalence class of  $(x_r)$ . We will show  $\lim_{k \rightarrow \infty} \sigma_k = \sigma$ , and since  $(\sigma_k)$  was arbitrary this will show that  $X_c$  is indeed complete. So let  $\epsilon > 0$ . Choose  $r \in \mathbb{N}$  such that  $4/r < \epsilon$ . We have by the triangle inequality that

$$d_{X_c}(\sigma_{n_r}, \sigma) \leq d_{X_c}(\sigma_{n_r}, \phi(x_r)) + d_{X_c}(\phi(x_r), \sigma).$$

We know  $d_{X_c}(\sigma_{n_r}, \phi(x_r)) < 1/r$ , and we showed above that  $d_X(x_k, x_r) < 3/r$  for  $k \geq r$ , which implies

$$d_{X_c}(\phi(x_r), \sigma) = \lim_{k \rightarrow \infty} d_X(x_r, x_k) < 3/r.$$

It follows that  $d_{X_c}(\sigma_{n_r}, \sigma) < 4/r < \epsilon$ . Then since  $(\sigma_{n_r})$  is Cauchy, it is easy to see we must then have  $\lim_{k \rightarrow \infty} \sigma_k = \sigma$ . So we have proven (2.1.1, 2.1.2).

We now prove (2.1.3, 2.1.4). These will follow from the following more general facts.<sup>2</sup>

**2.8. Lemma.** *Let  $A, B$  be topological spaces, and  $A_0 \subseteq A$  a dense subspace. Let  $f: A_0 \rightarrow B$  a continuous map.*

- (1) *If  $B$  is Hausdorff, there exists at most one extension of  $f$  to  $A$ , that is, there exists at most one continuous map  $g: A \rightarrow B$  such that  $g|_{A_0} = f$ .*
- (2) *If  $A$  and  $B$  are metric spaces,  $B$  is complete, and  $f$  is uniformly continuous, then there exists a unique uniformly continuous extension  $g: A \rightarrow B$  of  $f$ . If  $f$  is an isometry, then so is  $g$ .*

*Proof.* (1) Let  $g_1, g_2: A \rightarrow B$  two extensions of  $f$ . Suppose  $g_1(a) \neq g_2(a)$  for some  $a \in A$ . Then we can choose  $V_1$  and  $V_2$  disjoint neighbourhoods of  $g_1(a)$  and  $g_2(a)$ , respectively, by the Hausdorff hypothesis. Let  $U_i := g_i^{-1}(V_i)$ , open by continuity of  $g_i$ , for  $i \in \{1, 2\}$ . Then  $a \in U_1 \cap U_2$ , so  $U_1 \cap U_2$  is a nonempty open set in  $X$ , and hence there exists  $a_0 \in U_1 \cap U_2 \cap A_0$  by density of  $A_0$ . Now observe that since  $g_1, g_2$  extend  $f$  we must have

$$V_1 \ni g_1(a_0) = f(a_0) = g_2(a_0) \in V_2,$$

contradicting the disjointness of  $V_1$  and  $V_2$ .

(2) Uniqueness is immediate from (1), so it suffices to show existence. Define  $g: A \rightarrow B$  as follows. Let  $a \in A$ . By density of  $A_0$  there is a sequence  $(a_k) \in A_0^{\mathbb{N}}$  such that  $a_k \rightarrow a$  as  $k \rightarrow \infty$ . By (1.5),  $(a_k)$  is Cauchy, whence  $(f(a_k)) \in B^{\mathbb{N}}$  is Cauchy by (1.10). Then  $f(a_k) \rightarrow b$  as  $k \rightarrow \infty$  for some  $b \in B$  by completeness. We define

<sup>1</sup>Observe (for intuition) that we start with a sequence of sequences, which can visualise as a grid, and are producing a new sequence by taking some type of rapidly converging diagonal from this grid.

<sup>2</sup>The presentation here is essentially taken from Pete Clark's notes, [math.uga.edu/~pete/8410Chapter2v2.pdf](http://math.uga.edu/~pete/8410Chapter2v2.pdf).

$g(a) := b$ . We claim this is independent of the choice of sequence  $(a_k)$ . Indeed let  $(a'_k) \in A_0^{\mathbb{N}}$  another sequence converging to  $a$ . Then for any  $\delta > 0$  there exists  $n \in \mathbb{N}$  such that, by the triangle inequality,

$$d(a, a_k) < \delta/2 \text{ and } d(a, a'_k) < \delta/2 \implies d(a_k, a'_k) < \delta$$

for  $k \geq n$ . Therefore by uniform continuity there exists  $n \in \mathbb{N}$  such that

$$d(f(a_k), f(a'_k)) < \epsilon \text{ for all } k \geq n.$$

It follows that  $\lim_{k \rightarrow \infty} f(a_k) = \lim_{k \rightarrow \infty} f(a'_k)$ , proving our claim. In particular, if  $a \in A_0$  then we can choose the constant sequence  $a_k := a_0$  for  $k \in \mathbb{N}$ , from which we see  $g(a) = \lim_{k \rightarrow \infty} f(a_0) = f(a_0)$ , so  $g$  in fact extends  $f$ .

We now show  $g$  is uniformly continuous. Let  $\epsilon > 0$ . Since  $f$  is uniformly continuous there exists  $\delta > 0$  such that  $d_B(f(a), f(a')) < \epsilon/2$  whenever  $a_0, a'_0 \in A_0$  are such that  $d_A(a_0, a'_0) < \delta$ . Suppose  $a, a' \in A$  are such that  $d(a, a') < \delta/3$ . Choose sequences  $(a_k)$  and  $(a'_k)$  in  $A_0$  converging to  $a$  and  $a'$ , respectively. Let  $n \in \mathbb{N}$  such that, by the triangle inequality,

$$\begin{aligned} d_X(a_k, a) < \delta/3 \text{ and } d_X(a'_k, a') < \delta/3 &\implies d_X(a_k, a'_k) < \delta \\ &\implies d_Y(f(a_k), f(a'_k)) < \epsilon/2 \end{aligned}$$

for all  $k \geq n$ . Then by continuity<sup>3</sup> of  $d_Y$  and definition of  $g$  we have

$$(2.9) \quad d_Y(g(a), g(a')) = \lim_{k \rightarrow \infty} d_Y(f(a_k), f(a'_k)) \leq \epsilon/2 < \epsilon.$$

Thus  $g$  is uniformly continuous. If  $f$  is moreover an isometry, the first equality in (2.9) also clearly implies that  $g$  is an isometry.  $\square$

**2.10** (The universal property). We now apply the lemma to finish the proof of the proposition. Suppose  $Y$  is a complete metric space and  $\psi: X \rightarrow Y$  uniformly continuous. Since  $\phi: X \rightarrow X_c$  is an isometry with dense image, the lemma gives that there is a unique uniformly continuous map  $\psi_c: X_c \rightarrow Y$  such that  $\psi_c \circ \phi = \psi$ , proving (2.1.3).

Finally, (2.1.4) is proven by the general argument giving uniqueness of objects satisfying a universal property like (2.1.3). Suppose  $Y_c$  is a complete metric space and  $\psi: X \rightarrow Y_c$  an isometry with dense image. Then by (2.1.3) there is a unique uniformly continuous map  $\psi_c: X_c \rightarrow Y_c$  such that  $\psi_c \circ \phi = \psi$ , and the lemma tells us that  $\psi_c$  is in fact an isometry. But our proof of (2.1.3) applies to  $Y_c$  equipped with  $\psi$  as well, so we symmetrically have a unique isometry  $\phi_c: Y_c \rightarrow X_c$  such that  $\phi_c \circ \psi = \phi$ . I.e., we have the diagram

$$\begin{array}{ccc} X & \xrightarrow{\phi} & X_c \\ \downarrow \psi & \nearrow \phi_c & \uparrow \psi_c \\ Y_c & & \end{array}$$

Observe then that

$$\phi_c \circ \psi_c \circ \phi = \phi = \text{id}_{X_c} \circ \phi \quad \text{and} \quad \psi_c \circ \phi_c \circ \psi = \psi = \text{id}_{Y_c} \circ \psi.$$

---

<sup>3</sup>One could also argue more directly, but see, e.g., my solutions to problem set 4 for a proof of the continuity of the metric.

Then the *uniqueness* statement in (2.1.3) implies that we must have  $\phi_c \circ \psi_c = \text{id}_{X_c}$  and  $\psi_c \circ \phi_c = \text{id}_{Y_c}$ .<sup>4</sup> I.e.,  $\phi_c, \psi_c$  are inverse isomorphisms of metric spaces. And thus we are done!

---

<sup>4</sup>This argument takes a couple of reads to absorb and understand, I think (it certainly did for me!). But it's extremely general and very useful so try to do so!