

Structural Calculus: From Discrete Structural Derivatives to the Structural Spectrum of Integers

Abel Liu

September 9, 2026

Abstract

This paper establishes a discrete calculus framework acting on the internal structure of formal mathematical expressions. The fundamental premise is that mathematical expressions possess not only numerical semantics but also an internal syntax determined by their construction. Consequently, two expressions that are numerically equivalent are not necessarily identical as differential objects. To capture this distinction, we construct a free term algebra and define a structural derivative $\mathcal{D}$ upon it. Unlike classical calculus, $\mathcal{D}$ does not rely on limits; rather, it is recursively determined by local structural rules on expression trees.

We establish four fundamental axioms of structural calculus. The cornerstone of this framework is the **power node rule**,

$$\boxed{\mathcal{D}(U^V) = U^{V-1} + (U-1)^V,}$$

which highlights exponentiation as the true bedrock of the theory by decomposing a power structure into two parallel discrete evolution directions. Under a strict structural normalization principle, trivial multiplications by the identity element are forbidden, while atomic integers are fundamentally embedded as degenerate power structures (e.g., $8 = 8^1$). This leads to a striking departure from classical calculus: the structural derivative of an integer is not zero, but possesses its own non-trivial value (e.g., $\mathcal{D}(8) = 8$).

By introducing two commuting discrete shift operators, we express the structural derivative as their sum and derive a binomial expansion for its higher-order iterations. As an application, we obtain an explicit formula for the n -th order structural derivative of x^x , yielding a novel combinatorial integer sequence. Furthermore, we extend the scope from individual expressions to the complete structural decompositions of integers. We define the *structural spectrum* $\Sigma(n)$ and the *total structural response* $R(n)$ for an integer n . This reveals a profound structural dichotomy between primes and composites. Finally, we propose two natural arithmetic conjectures concerning the integrality and critical self-consistency of $R(n)$, opening a new pathway from local structural rules to the global arithmetic properties of integers.

Keywords: Structural Calculus; Formal Expressions; Free Term Algebra; Discrete Derivative; Expression Trees; Power Node; Leibniz Rule; Structural Spectrum; Integer Decompositions.

Contents

1	Introduction	3
2	The Algebra of Formal Expressions and Fundamental Rules	3
2.1	Expression Trees and Structural Equivalence	3
2.2	The Four Fundamental Axioms	3
3	The Structural Derivative and the Primacy of Exponentiation	4
3.1	Why the Derivative of an Integer is Not Zero	4

4	Higher-Order Structural Derivatives and Combinatorial Paths	4
4.1	Discrete Shift Operators	4
4.2	The n -th Order Structural Derivative	4
4.3	Explicit Examples and Combinatorial Interpretation	5
5	Structural States and the Structural Spectrum of Integers	5
5.1	Structural States and Response Rates	5
5.2	The Structural Spectrum	5
6	Total Structural Response and Arithmetic Conjectures	6
6.1	Total Structural Response	6
6.2	Two Fundamental Conjectures	6
7	Conclusion	6

1 Introduction

In traditional mathematics, an expression is primarily identified by its numerical value. For instance, 6 , 2×3 , and $1 + 5$ all denote the same numerical entity. Furthermore, in classical calculus, the derivative of a constant is identically zero. However, this perspective entirely discards the syntactic and structural information inherent in how a mathematical object is constructed.

This paper introduces **Structural Calculus**, a novel discrete mathematical framework that treats the internal construction of expressions as the primary object of study. In this framework, expressions are modeled as finite rooted trees, and differentiation is redefined as a local structural evolution on these trees. We demonstrate that exponentiation serves as the fundamental bedrock of this calculus, leading to the counterintuitive but mathematically consistent result that the structural derivative of an integer is not zero. By analyzing the complete set of structural decompositions of an integer, we define its *structural spectrum* and propose new arithmetic conjectures that bridge discrete operator calculus and number theory.

2 The Algebra of Formal Expressions and Fundamental Rules

2.1 Expression Trees and Structural Equivalence

Let $\mathcal{A}$ be an alphabet of symbols containing variables, constants, and formal parameters. We define the set of terms $\mathcal{T}$ recursively:

- (i) Every basic symbol in $\mathcal{A}$ is a term in $\mathcal{T}$.
- (ii) If $F, G \in \mathcal{T}$, then $(F + G)$, $(F - G)$, (FG) , and (F^G) are in $\mathcal{T}$.

Every term can be uniquely represented as a finite rooted tree, where leaves correspond to basic symbols and internal nodes correspond to the operators $+$, $-$, $\times$, and $^{\wedge}$.

Definition 2.1 (Structural Equivalence). *Two terms $F, G \in \mathcal{T}$ are structurally equivalent, denoted $F \cong_{\text{str}} G$, if and only if they are isomorphic as labeled rooted trees.*

It is evident that numerical equality does not imply structural equivalence. If we define the standard numerical evaluation map $\text{ev} : \mathcal{T} \rightarrow \mathbb{R}$, we have $\text{ev}(6) = \text{ev}(2 \times 3) = 6$, but $6 \not\cong_{\text{str}} 2 \times 3$. This separation of numerical semantics and structural syntax is the cornerstone of our theory.

2.2 The Four Fundamental Axioms

The structural derivative $\mathcal{D}$ is a linear operator on the free term algebra $\mathcal{E} = \mathbb{K}(\mathcal{T})$. Its action is entirely governed by the following four fundamental axioms:

Axiom 2.2 (Linearity for Addition and Subtraction).

$$\mathcal{D}(F \pm G) = \mathcal{D}(F) \pm \mathcal{D}(G). \quad (1)$$

Axiom 2.3 (Leibniz Rule for Multiplication).

$$\mathcal{D}(F \cdot G) = \mathcal{D}(F) \cdot G + F \cdot \mathcal{D}(G). \quad (2)$$

Axiom 2.4 (The Power Node Rule).

$$\boxed{\mathcal{D}(U^V) = U^{V-1} + (U - 1)^V.} \quad (3)$$

This is the most critical and disruptive rule. It decomposes a power node into two parallel structural evolution directions: the “exponent descent” (U^{V-1}) and the “base descent” ($(U - 1)^V$).

Axiom 2.5 (Structural Normalization). *The expression trees must be in their most reduced form. The system strictly eliminates all redundant parenthesization and trivial multiplications by the identity element (1).*

3 The Structural Derivative and the Primacy of Exponentiation

3.1 Why the Derivative of an Integer is Not Zero

In classical calculus, the derivative of a constant is zero. In structural calculus, we assert that **the structural derivative of an integer is not zero; it inherently possesses its own non-trivial value.**

This profound departure stems from the fact that **exponentiation is the true bedrock of this calculus.** Under Axiom 4 (Structural Normalization), trivial multiplicative structures involving the identity element, such as 8×1 or $8 \times 1 \times 1$, are strictly excluded from the set of valid structural states. An integer cannot be artificially inflated by multiplying it by 1.

However, representing an integer as a degenerate power structure with an exponent of 1 is fundamentally permitted and required as the base structural form. That is, any atomic integer N is intrinsically embedded as:

$$N = N^1$$

Applying Axiom 3 (The Power Node Rule) to $8 = 8^1$, we obtain:

$$\mathcal{D}(8) = \mathcal{D}(8^1) = 8^{1-1} + (8 - 1)^1 = 8^0 + 7^1 = 1 + 7 = 8. \quad (4)$$

Thus, $\mathcal{D}(8) = 8$. We do not need to artificially postulate that “the derivative of a constant is itself.” This result emerges naturally and inevitably as the degenerate case of the Power Node Rule when the exponent is 1. This demonstrates the profound self-consistency of structural calculus: a single core rule governs the entire universe of expressions.

4 Higher-Order Structural Derivatives and Combinatorial Paths

4.1 Discrete Shift Operators

Consider the formal subspace generated by power terms u^v . We define two discrete shift operators:

$$E_u^{-1}(u^v) = (u - 1)^v, \quad E_v^{-1}(u^v) = u^{v-1}. \quad (5)$$

The Power Node Rule can then be elegantly rewritten as:

$$\mathcal{D} = E_v^{-1} + E_u^{-1}. \quad (6)$$

Crucially, these two operators commute: $E_u^{-1}E_v^{-1} = E_v^{-1}E_u^{-1}$. This commutativity implies that the order of structural evolution (whether we descend the base first or the exponent first) does not affect the final structural state.

4.2 The n -th Order Structural Derivative

Because the operators commute, we can directly apply the binomial theorem to compute higher-order derivatives.

Theorem 4.1. *For any $n \in \mathbb{N}$, the n -th order structural derivative is given by:*

$$\boxed{\mathcal{D}^n = \sum_{k=0}^n \binom{n}{k} (E_u^{-1})^k (E_v^{-1})^{n-k}.} \quad (7)$$

Applying this to the self-referential expression x^x , we obtain a completely explicit formula:

$$\boxed{\mathcal{D}^n[x^x] = \sum_{k=0}^n \binom{n}{k} (x-k)^{x-n+k}.} \quad (8)$$

4.3 Explicit Examples and Combinatorial Interpretation

The binomial coefficients $\binom{n}{k}$ are not arbitrary; they represent the exact number of discrete structural evolution paths. In n steps, choosing k times to descend the base and $n-k$ times to descend the exponent yields $\binom{n}{k}$ identical final structures.

Let us examine two explicit examples:

Example 4.2 (First-Order Derivative, $n = 1$).

$$\mathcal{D}[x^x] = \binom{1}{0} (x-0)^{x-1} + \binom{1}{1} (x-1)^x = x^{x-1} + (x-1)^x. \quad (9)$$

This perfectly mirrors the core Power Node Rule.

Example 4.3 (Second-Order Derivative, $n = 2$).

$$\begin{aligned} \mathcal{D}^2[x^x] &= \binom{2}{0} (x-0)^{x-2} + \binom{2}{1} (x-1)^{x-1} + \binom{2}{2} (x-2)^x \\ &= x^{x-2} + 2(x-1)^{x-1} + (x-2)^x. \end{aligned} \quad (10)$$

Evaluating this at $x = n$ generates a novel combinatorial integer sequence $A_n = \mathcal{D}^n[n^n] = \sum_{k=0}^n \binom{n}{k} (n-k)^k$, with initial terms $A_1 = 1, A_2 = 3, A_3 = 10, A_4 = 41, A_5 = 196, \dots$

5 Structural States and the Structural Spectrum of Integers

5.1 Structural States and Response Rates

We now extend our focus from individual expressions to the integer itself. For an integer $n > 1$, let $\text{States}(n)$ be the set of all valid, normalized structural representations of n (e.g., for $n = 4$, the states are $4, 2 \times 2$, and 2^2). Note that 4×1 is excluded by Axiom 4.

To measure the “structural activity” of a state $F \in \text{States}(n)$, we define its *structural response rate*:

$$\boxed{\rho(F) = \frac{\text{ev}(\mathcal{D}(F))}{\text{ev}(F)} = \frac{\text{ev}(\mathcal{D}(F))}{n}.} \quad (11)$$

For multiplicative structures $F = G \cdot H$, the Leibniz rule implies a strict additivity of response rates:

$$\rho(G \cdot H) = \rho(G) + \rho(H). \quad (12)$$

Conversely, power structures exhibit non-linear responses. For $F = a^m$ ($a, m \geq 2$):

$$\rho(a^m) = \frac{a^{m-1} + (a-1)^m}{a^m} = \frac{1}{a} + \left(1 - \frac{1}{a}\right)^m. \quad (13)$$

5.2 The Structural Spectrum

Definition 5.1 (Structural Spectrum). *The structural spectrum of an integer n is defined as the set of response rates of all its valid structural states:*

$$\boxed{\Sigma(n) = \{\rho(F) : F \in \text{States}(n)\}.} \quad (14)$$

This definition reveals a profound structural dichotomy in number theory:

- **Primes** (e.g., 2, 3, 5) possess no non-trivial multiplicative decompositions. Their structural spectrum is always the singleton $\Sigma(p) = \{1\}$, representing an absolute “ground state.”
- **Composites** (e.g., 4, 8, 27) possess rich non-trivial decompositions. For example, $\Sigma(4) = \{1, 2, \frac{3}{4}\}$ and $\Sigma(8) = \{1, 2, 3, \frac{5}{8}, \frac{7}{4}\}$. They exhibit multiple “excited states.”

6 Total Structural Response and Arithmetic Conjectures

6.1 Total Structural Response

We define the *total structural response* $R(n)$ as the sum of the response rates over all valid structural states of n :

$$R(n) = \sum_{F \in \text{States}(n)} \rho(F). \quad (15)$$

Let us compute $R(n)$ for small integers:

- For $n = 4$: $R(4) = 1 + 2 + \frac{3}{4} = \frac{15}{4} = 3.75$.
- For $n = 8$: $R(8) = 1 + 2 + 3 + \frac{5}{8} + \frac{7}{4} = \frac{67}{8} = 8.375$.

Notice that $R(4) < 4$, while $R(8) > 8$. The function $R(n) - n$ changes sign, which naturally motivates the following arithmetic conjectures.

6.2 Two Fundamental Conjectures

Conjecture 6.1 (Integrality of the Structural Spectrum (Conjecture A)). *Does there exist an integer $n > 1$ such that its total structural response $R(n)$ is an exact integer? That is,*

$$\exists n > 1, \quad R(n) \in \mathbb{Z}. \quad (16)$$

Conjecture 6.2 (Structural Criticality (Conjecture B)). *Does there exist a special integer $n > 1$ such that its total structural response exactly equals its numerical value? That is,*

$$\exists n > 1, \quad R(n) = n. \quad (17)$$

If such an integer exists, we term it a **Structural Critical Integer**. Conjecture B implies Conjecture A. The existence of a structural critical integer would mean that the numerical scale of the integer perfectly balances the total “structural energy” generated by all its possible internal decompositions. While finite computations show $R(n) - n$ crossing zero between $n = 4$ and $n = 8$, the discrete nature of n prevents us from concluding the existence of a root without further rigorous analysis.

7 Conclusion

This paper has established Structural Calculus, a discrete mathematical framework that differentiates expressions based on their internal syntactic trees rather than their numerical values. We formalized the theory through four fundamental axioms, establishing exponentiation as the foundational bedrock. By enforcing structural normalization, we demonstrated that the structural derivative of an integer is inherently non-zero (e.g., $\mathcal{D}(8) = 8$), emerging naturally from the degenerate power rule.

We derived explicit formulas for higher-order structural derivatives, revealing a beautiful combinatorial path-counting structure. Furthermore, by extending the derivative to the space

of all valid structural decompositions of an integer, we introduced the *structural spectrum* and the *total structural response* $R(n)$. This led to the proposal of two novel arithmetic conjectures regarding the integrality and critical self-consistency of $R(n)$.

The research pathway established herein transitions from local expression tree rules to global arithmetic properties of integers:

Formal Expressions $\longrightarrow$ Structural Derivative $\longrightarrow$ Structural States $\longrightarrow$ Structural Spectrum $\longrightarrow R(n) \longrightarrow A$

Future work will focus on rigorously enumerating States(n), analyzing the asymptotic growth of $R(n)$, and computationally searching for structural critical integers.

References

- [1] G.-C. Rota, *Finite Operator Calculus*, Academic Press, 1975.
- [2] S. Roman, *The Umbral Calculus*, Academic Press, 1984.
- [3] J. Riordan, *Combinatorial Identities*, John Wiley & Sons, 1968.
- [4] R. P. Stanley, *Enumerative Combinatorics, Vol. 1*, Cambridge University Press, 2012.
- [5] P. Flajolet and R. Sedgewick, *Analytic Combinatorics*, Cambridge University Press, 2009.
- [6] R. L. Graham, D. E. Knuth and O. Patashnik, *Concrete Mathematics*, Addison-Wesley, 1994.
- [7] H. S. Wilf, *Generatingfunctionology*, A K Peters, 2006.
- [8] T. M. Apostol, *Introduction to Analytic Number Theory*, Springer, 1976.
- [9] OEIS Foundation Inc., *The On-Line Encyclopedia of Integer Sequences*, <https://oeis.org>, 2026.