

Quantum algorithms for multivariate Monte Carlo estimation

A.J. Cornelissen¹

Joint work with S. Jerbi² & Y. Hamoudi³

¹QuSoft, University of Amsterdam

²Institute for Theoretical Physics, University of Innsbruck

³Department of Electrical Engineering and Computer Sciences, University of California, Berkeley

September 15th, 2021



- 1 Prior work on Monte Carlo estimation:

	Classically	Quantumly
Univariate	Textbook	Textbook
Multivariate	Textbook	??

- 1 Prior work on Monte Carlo estimation:

	Classically	Quantumly
Univariate	Textbook	Textbook
Multivariate	Textbook	??

- 2 Core technique: Quantum analog computation

- 1 Prior work on Monte Carlo estimation:

	Classically	Quantumly
Univariate	Textbook	Textbook
Multivariate	Textbook	??

- 2 Core technique: Quantum analog computation
- 3 Generalization of the univariate quantum algorithm to the multivariate setting

- 1 Prior work on Monte Carlo estimation:

	Classically	Quantumly
Univariate	Textbook	Textbook
Multivariate	Textbook	??

- 2 Core technique: Quantum analog computation
- 3 Generalization of the univariate quantum algorithm to the multivariate setting
- 4 Improvements & further directions

Univariate Monte Carlo estimation

Univariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]$ a random variable.

Univariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]$ a random variable.

Example: Suppose we have two dice:

① $\Omega = \{1, 2, 3, 4, 5, 6\}^2$.

② $X((i, j)) = (i + j)/12$.

Univariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]$ a random variable.

Example: Suppose we have two dice:

① $\Omega = \{1, 2, 3, 4, 5, 6\}^2$.

② $X((i, j)) = (i + j)/12$.

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Univariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]$ a random variable.

Example: Suppose we have two dice:

① $\Omega = \{1, 2, 3, 4, 5, 6\}^2$.

② $X((i, j)) = (i + j)/12$.

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Classical algorithm:

① Take samples $\omega_1, \dots, \omega_N$ from Ω .

② Compute $\frac{1}{N} \sum_{j=1}^N X(\omega_j)$.

Univariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]$ a random variable.

Example: Suppose we have two dice:

① $\Omega = \{1, 2, 3, 4, 5, 6\}^2$.

② $X((i, j)) = (i + j)/12$.

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Classical algorithm:

① Take samples $\omega_1, \dots, \omega_N$ from Ω .

② Compute $\frac{1}{N} \sum_{j=1}^N X(\omega_j)$.

This estimates $\mathbb{E}[X]$ up to $\sim 1/\sqrt{N}$.

Univariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]$ a random variable.

Example: Suppose we have two dice:

① $\Omega = \{1, 2, 3, 4, 5, 6\}^2$.

② $X((i, j)) = (i + j)/12$.

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Classical algorithm:

① Take samples $\omega_1, \dots, \omega_N$ from Ω .

② Compute $\frac{1}{N} \sum_{j=1}^N X(\omega_j)$.

This estimates $\mathbb{E}[X]$ up to $\sim 1/\sqrt{N}$.

To obtain precision ε , we need $N = \mathcal{O}(1/\varepsilon^2)$.

Quantum univariate Monte Carlo estimation

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Quantum univariate Monte Carlo estimation

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Quantum algorithm: Let $\{|\omega\rangle : \omega \in \Omega\}$ be an orthonormal set of states.

$$U : |0\rangle |0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle |0\rangle$$

Quantum univariate Monte Carlo estimation

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Quantum algorithm: Let $\{|\omega\rangle : \omega \in \Omega\}$ be an orthonormal set of states.

$$U : |0\rangle |0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle |0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{X(\omega)} |1\rangle + \sqrt{1 - X(\omega)} |0\rangle \right)$$

Quantum univariate Monte Carlo estimation

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Quantum algorithm: Let $\{|\omega\rangle : \omega \in \Omega\}$ be an orthonormal set of states.

$$\begin{aligned} U : |0\rangle |0\rangle &\mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle |0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{X(\omega)} |1\rangle + \sqrt{1 - X(\omega)} |0\rangle \right) \\ &= \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)} |\omega\rangle |1\rangle + \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) (1 - X(\omega))} |\omega\rangle |0\rangle \end{aligned}$$

Quantum univariate Monte Carlo estimation

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Quantum algorithm: Let $\{|\omega\rangle : \omega \in \Omega\}$ be an orthonormal set of states.

$$\begin{aligned} U : |0\rangle |0\rangle &\mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle |0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{X(\omega)} |1\rangle + \sqrt{1 - X(\omega)} |0\rangle \right) \\ &= \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)} |\omega\rangle |1\rangle + \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) (1 - X(\omega))} |\omega\rangle |0\rangle \\ &= \sqrt{\sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega)} \underbrace{\left(\frac{\sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)}}{\sqrt{\sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega)}} |\omega\rangle \right)}_{|\psi_1\rangle} |1\rangle + (\dots) |0\rangle \end{aligned}$$

Quantum univariate Monte Carlo estimation

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Quantum algorithm: Let $\{|\omega\rangle : \omega \in \Omega\}$ be an orthonormal set of states.

$$\begin{aligned} U : |0\rangle |0\rangle &\mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle |0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{X(\omega)} |1\rangle + \sqrt{1 - X(\omega)} |0\rangle \right) \\ &= \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)} |\omega\rangle |1\rangle + \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) (1 - X(\omega))} |\omega\rangle |0\rangle \\ &= \sqrt{\sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega)} \underbrace{\left(\frac{\sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)}}{\sqrt{\sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega)}} |\omega\rangle \right)}_{|\psi_1\rangle} |1\rangle + (\dots) |0\rangle \\ &= \sqrt{\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \mathbb{E}[X]} |\psi_0\rangle |0\rangle. \end{aligned}$$

Quantum univariate Monte Carlo estimation

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Quantum algorithm: Let $\{|\omega\rangle : \omega \in \Omega\}$ be an orthonormal set of states.

$$\begin{aligned} U : |0\rangle |0\rangle &\mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle |0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{X(\omega)} |1\rangle + \sqrt{1 - X(\omega)} |0\rangle \right) \\ &= \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)} |\omega\rangle |1\rangle + \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) (1 - X(\omega))} |\omega\rangle |0\rangle \\ &= \sqrt{\sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega)} \underbrace{\left(\frac{\sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)}}{\sqrt{\sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega)}} |\omega\rangle \right)}_{|\psi_1\rangle} |1\rangle + (\dots) |0\rangle \\ &= \sqrt{\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \mathbb{E}[X]} |\psi_0\rangle |0\rangle. \end{aligned}$$

Amplitude estimation computes $\mathbb{E}[X] \pm \varepsilon$ with $\mathcal{O}(1/\varepsilon)$ calls to U .

Quantum univariate Monte Carlo estimation

Question: How to compute $\mathbb{E}[X]$ up to precision ε ?

Quantum algorithm: Let $\{|\omega\rangle : \omega \in \Omega\}$ be an orthonormal set of states.

$$\begin{aligned} U : |0\rangle |0\rangle &\mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle |0\rangle \mapsto \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{X(\omega)} |1\rangle + \sqrt{1 - X(\omega)} |0\rangle \right) \\ &= \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)} |\omega\rangle |1\rangle + \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) (1 - X(\omega))} |\omega\rangle |0\rangle \\ &= \sqrt{\sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega)} \underbrace{\left(\frac{\sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega) X(\omega)}}{\sqrt{\sum_{\omega \in \Omega} \mathbb{P}(\omega) X(\omega)}} |\omega\rangle \right)}_{|\psi_1\rangle} |1\rangle + (\dots) |0\rangle \\ &= \sqrt{\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \mathbb{E}[X]} |\psi_0\rangle |0\rangle. \end{aligned}$$

Amplitude estimation computes $\mathbb{E}[X] \pm \varepsilon$ with $\mathcal{O}(1/\varepsilon)$ calls to U .

Quadratic speed-up!

Multivariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]^d$ a random variable.

Question: How to compute $\mathbb{E}[X]$ up to ℓ_∞ -error ε ?

Multivariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]^d$ a random variable.

Question: How to compute $\mathbb{E}[X]$ up to ℓ_∞ -error ε ?

Classical algorithm:

- ① Run the classical algorithm with N samples.
 - ① Take samples $\omega_1, \dots, \omega_N$ from Ω .
 - ② Compute $\frac{1}{N} \sum_{j=1}^N X(\omega_j) \in [0, 1]^d$.

Multivariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]^d$ a random variable.

Question: How to compute $\mathbb{E}[X]$ up to ℓ_∞ -error ε ?

Classical algorithm:

- ① Run the classical algorithm with N samples.
 - ① Take samples $\omega_1, \dots, \omega_N$ from Ω .
 - ② Compute $\frac{1}{N} \sum_{j=1}^N X(\omega_j) \in [0, 1]^d$.
- ② *Median trick:* Repeat $\mathcal{O}(\log(d))$ times and take the coordinate-wise median.

Multivariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]^d$ a random variable.

Question: How to compute $\mathbb{E}[X]$ up to ℓ_∞ -error ε ?

Classical algorithm:

- ① Run the classical algorithm with N samples.
 - ① Take samples $\omega_1, \dots, \omega_N$ from Ω .
 - ② Compute $\frac{1}{N} \sum_{j=1}^N X(\omega_j) \in [0, 1]^d$.
- ② *Median trick:* Repeat $\mathcal{O}(\log(d))$ times and take the coordinate-wise median.

This computes $\mathbb{E}[X]$, with ℓ_∞ -error $1/\sqrt{N} \Rightarrow N = \mathcal{O}(1/\varepsilon^2)$.

Multivariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]^d$ a random variable.

Question: How to compute $\mathbb{E}[X]$ up to ℓ_∞ -error ε ?

Classical algorithm:

- ① Run the classical algorithm with N samples.
 - ① Take samples $\omega_1, \dots, \omega_N$ from Ω .
 - ② Compute $\frac{1}{N} \sum_{j=1}^N X(\omega_j) \in [0, 1]^d$.
- ② *Median trick:* Repeat $\mathcal{O}(\log(d))$ times and take the coordinate-wise median.

This computes $\mathbb{E}[X]$, with ℓ_∞ -error $1/\sqrt{N} \Rightarrow N = \mathcal{O}(1/\varepsilon^2)$.

Total number of samples: $\mathcal{O}(\log(d)/\varepsilon^2)$.

Multivariate Monte Carlo estimation

Let Ω a finite set, $(\Omega, 2^\Omega, \mathbb{P})$ a probability space, and $X : \Omega \rightarrow [0, 1]^d$ a random variable.

Question: How to compute $\mathbb{E}[X]$ up to ℓ_∞ -error ε ?

Classical algorithm:

- ① Run the classical algorithm with N samples.
 - ① Take samples $\omega_1, \dots, \omega_N$ from Ω .
 - ② Compute $\frac{1}{N} \sum_{j=1}^N X(\omega_j) \in [0, 1]^d$.
- ② *Median trick:* Repeat $\mathcal{O}(\log(d))$ times and take the coordinate-wise median.

This computes $\mathbb{E}[X]$, with ℓ_∞ -error $1/\sqrt{N} \Rightarrow N = \mathcal{O}(1/\varepsilon^2)$.

Total number of samples: $\mathcal{O}(\log(d)/\varepsilon^2)$.

Can we get a similar quadratic quantum speed-up here?

Overview

Goal:

		Classically	Quantumly
Univariate	$X : \Omega \rightarrow [0, 1]$	$\mathcal{O}(\frac{1}{\varepsilon^2})$	$\mathcal{O}(\frac{1}{\varepsilon})$
Multivariate	$X : \Omega \rightarrow [0, 1]^d$	$\tilde{\mathcal{O}}(\frac{1}{\varepsilon^2})$	$\tilde{\mathcal{O}}(\frac{1}{\varepsilon})??$

Overview

Goal:

		Classically	Quantumly
Univariate	$X : \Omega \rightarrow [0, 1]$	$\mathcal{O}(\frac{1}{\varepsilon^2})$	$\mathcal{O}(\frac{1}{\varepsilon})$
Multivariate	$X : \Omega \rightarrow [0, 1]^d$	$\tilde{\mathcal{O}}(\frac{1}{\varepsilon^2})$	$\tilde{\mathcal{O}}(\frac{1}{\varepsilon})??$

Difficulty is that

$$U : |0\rangle |0\rangle \mapsto \sqrt{\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \mathbb{E}[X]} |\psi_0\rangle |0\rangle$$

has no clear multidimensional generalization.

Quantum analog computation

Let D a finite set, $f : D \rightarrow [0, 1]$, and $x \in D$.

Quantum analog computation

Let D a finite set, $f : D \rightarrow [0, 1]$, and $x \in D$.

① *Binary oracle*: $B_{f,\varepsilon} : |x\rangle |0\rangle \mapsto |x\rangle \left| \tilde{f}(x) \right\rangle$, where $|f(x) - \tilde{f}(x)| < \varepsilon$.

Quantum analog computation

Let D a finite set, $f : D \rightarrow [0, 1]$, and $x \in D$.

- ① *Binary oracle*: $B_{f,\varepsilon} : |x\rangle |0\rangle \mapsto |x\rangle \left| \tilde{f}(x) \right\rangle$, where $|f(x) - \tilde{f}(x)| < \varepsilon$.
- ② *Probability oracle*: $U_f : |x\rangle |0\rangle |0\rangle \mapsto |x\rangle \left(\sqrt{f(x)} |\psi_1\rangle |1\rangle + \sqrt{1 - f(x)} |\psi_0\rangle |0\rangle \right)$.

Quantum analog computation

Let D a finite set, $f : D \rightarrow [0, 1]$, and $x \in D$.


- ① *Binary oracle*: $B_{f,\varepsilon} : |x\rangle |0\rangle \mapsto |x\rangle \left| \tilde{f}(x) \right\rangle$, where $|f(x) - \tilde{f}(x)| < \varepsilon$.
- ② *Probability oracle*: $U_f : |x\rangle |0\rangle |0\rangle \mapsto |x\rangle \left(\sqrt{f(x)} |\psi_1\rangle |1\rangle + \sqrt{1-f(x)} |\psi_0\rangle |0\rangle \right)$.
- ③ *Phase oracle*: $O_f : |x\rangle \mapsto e^{if(x)} |x\rangle$.

Quantum analog computation

Let D a finite set, $f : D \rightarrow [0, 1]$, and $x \in D$.

- ① *Binary oracle*: $B_{f,\varepsilon} : |x\rangle |0\rangle \mapsto |x\rangle |\tilde{f}(x)\rangle$, where $|f(x) - \tilde{f}(x)| < \varepsilon$.
- ② *Probability oracle*: $U_f : |x\rangle |0\rangle |0\rangle \mapsto |x\rangle \left(\sqrt{f(x)} |\psi_1\rangle |1\rangle + \sqrt{1-f(x)} |\psi_0\rangle |0\rangle \right)$.
- ③ *Phase oracle*: $O_f : |x\rangle \mapsto e^{if(x)} |x\rangle$.


Oracle conversion graph:



Modification of the quantum univariate Monte Carlo estimation algorithm

Let $X : \Omega \rightarrow [0, 1]$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}$, and $f : G \rightarrow [0, 1]$, defined by $f(a) = a\mathbb{E}[X]$.




Modification of the quantum univariate Monte Carlo estimation algorithm

Let $X : \Omega \rightarrow [0, 1]$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}$, and $f : G \rightarrow [0, 1]$, defined by $f(a) = a\mathbb{E}[X]$.

$$\begin{aligned} U_f : |a\rangle |0\rangle |0\rangle &\mapsto |a\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{aX(\omega)} |1\rangle + \sqrt{1 - aX(\omega)} |0\rangle \right) \\ &= |a\rangle \left(\sqrt{a\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - a\mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$




Modification of the quantum univariate Monte Carlo estimation algorithm

Let $X : \Omega \rightarrow [0, 1]$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}$, and $f : G \rightarrow [0, 1]$, defined by $f(a) = a\mathbb{E}[X]$.

$$\begin{aligned} U_f : |a\rangle |0\rangle |0\rangle &\mapsto |a\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{aX(\omega)} |1\rangle + \sqrt{1 - aX(\omega)} |0\rangle \right) \\ &= |a\rangle \left(\sqrt{a\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - a\mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |a\rangle \mapsto e^{ia\mathbb{E}[X]} |a\rangle$ with $\tilde{O}(1)$ calls to U_f .



Modification of the quantum univariate Monte Carlo estimation algorithm


Let $X : \Omega \rightarrow [0, 1]$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}$, and $f : G \rightarrow [0, 1]$, defined by $f(a) = a\mathbb{E}[X]$.

$$\begin{aligned} U_f : |a\rangle |0\rangle |0\rangle &\mapsto |a\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{aX(\omega)} |1\rangle + \sqrt{1 - aX(\omega)} |0\rangle \right) \\ &= |a\rangle \left(\sqrt{a\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - a\mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |a\rangle \mapsto e^{ia\mathbb{E}[X]} |a\rangle$ with $\tilde{O}(1)$ calls to U_f .

$$\frac{1}{\sqrt{|G|}} \sum_{a \in G} |a\rangle \xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{a \in G} e^{ia\mathbb{E}[X]N} |a\rangle$$



Modification of the quantum univariate Monte Carlo estimation algorithm


Let $X : \Omega \rightarrow [0, 1]$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}$, and $f : G \rightarrow [0, 1]$, defined by $f(a) = a\mathbb{E}[X]$.

$$\begin{aligned} U_f : |a\rangle |0\rangle |0\rangle &\mapsto |a\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{aX(\omega)} |1\rangle + \sqrt{1 - aX(\omega)} |0\rangle \right) \\ &= |a\rangle \left(\sqrt{a\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - a\mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |a\rangle \mapsto e^{ia\mathbb{E}[X]} |a\rangle$ with $\tilde{O}(1)$ calls to U_f .

$$\frac{1}{\sqrt{|G|}} \sum_{a \in G} |a\rangle \xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{a \in G} e^{ia\mathbb{E}[X]N} |a\rangle$$



For $k \in \{0, \dots, 2^n - 1\}$,

$$\text{QFT}_G : |k\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{a \in G} e^{2\pi i a k} |a\rangle.$$

Modification of the quantum univariate Monte Carlo estimation algorithm


Let $X : \Omega \rightarrow [0, 1]$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}$, and $f : G \rightarrow [0, 1]$, defined by $f(a) = a\mathbb{E}[X]$.

$$\begin{aligned} U_f : |a\rangle |0\rangle |0\rangle &\mapsto |a\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{aX(\omega)} |1\rangle + \sqrt{1 - aX(\omega)} |0\rangle \right) \\ &= |a\rangle \left(\sqrt{a\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - a\mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |a\rangle \mapsto e^{ia\mathbb{E}[X]} |a\rangle$ with $\tilde{O}(1)$ calls to U_f .

$$\begin{aligned} \frac{1}{\sqrt{|G|}} \sum_{a \in G} |a\rangle &\xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{a \in G} e^{ia\mathbb{E}[X]N} |a\rangle \\ &\xrightarrow{\text{QFT}_G^\dagger} \left| \text{round} \left(\frac{\mathbb{E}[X]N}{2\pi} \right) \right\rangle. \end{aligned}$$



For $k \in \{0, \dots, 2^n - 1\}$,

$$\text{QFT}_G : |k\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{a \in G} e^{2\pi i a k} |a\rangle.$$

Modification of the quantum univariate Monte Carlo estimation algorithm

Let $X : \Omega \rightarrow [0, 1]$.


Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}$, and $f : G \rightarrow [0, 1]$, defined by $f(a) = a\mathbb{E}[X]$.

$$\begin{aligned} U_f : |a\rangle |0\rangle |0\rangle &\mapsto |a\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{aX(\omega)} |1\rangle + \sqrt{1 - aX(\omega)} |0\rangle \right) \\ &= |a\rangle \left(\sqrt{a\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - a\mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |a\rangle \mapsto e^{ia\mathbb{E}[X]} |a\rangle$ with $\tilde{O}(1)$ calls to U_f .

$$\begin{aligned} \frac{1}{\sqrt{|G|}} \sum_{a \in G} |a\rangle &\xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{a \in G} e^{ia\mathbb{E}[X]N} |a\rangle \\ &\xrightarrow{\text{QFT}_G^\dagger} \left| \text{round} \left(\frac{\mathbb{E}[X]N}{2\pi} \right) \right\rangle. \end{aligned}$$

We can choose $N = \mathcal{O}(1/\varepsilon) \Rightarrow \tilde{O}(1/\varepsilon)$ calls to U_f .



For $k \in \{0, \dots, 2^n - 1\}$,

$$\text{QFT}_G : |k\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{a \in G} e^{2\pi i a k} |a\rangle.$$

Modification of the quantum univariate Monte Carlo estimation algorithm

Let $X : \Omega \rightarrow [0, 1]$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}$, and $f : G \rightarrow [0, 1]$, defined by $f(a) = a\mathbb{E}[X]$.


$$\begin{aligned} U_f : |a\rangle |0\rangle |0\rangle &\mapsto |a\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{aX(\omega)} |1\rangle + \sqrt{1 - aX(\omega)} |0\rangle \right) \\ &= |a\rangle \left(\sqrt{a\mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - a\mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |a\rangle \mapsto e^{ia\mathbb{E}[X]} |a\rangle$ with $\tilde{O}(1)$ calls to U_f .

$$\begin{aligned} \frac{1}{\sqrt{|G|}} \sum_{a \in G} |a\rangle &\xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{a \in G} e^{ia\mathbb{E}[X]N} |a\rangle \\ &\xrightarrow{\text{QFT}_G^\dagger} \left| \text{round} \left(\frac{\mathbb{E}[X]N}{2\pi} \right) \right\rangle. \end{aligned}$$

We can choose $N = \mathcal{O}(1/\varepsilon) \Rightarrow \tilde{O}(1/\varepsilon)$ calls to U_f .

Tiny tweak of amplitude estimation.




For $k \in \{0, \dots, 2^n - 1\}$,

$$\text{QFT}_G : |k\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{a \in G} e^{2\pi i a k} |a\rangle.$$

Quantum multivariate Monte Carlo estimation

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}^d$, and $f : G \rightarrow [0, 1]$, defined by $f(\vec{a}) = \vec{a}^T \mathbb{E}[X]$.




For $\vec{k} \in \{0, \dots, 2^n - 1\}^d$,

$$\text{QFT}_G : |\vec{k}\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{2\pi i \vec{a}^T \vec{k}} |\vec{a}\rangle.$$

Quantum multivariate Monte Carlo estimation

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}^d$, and $f : G \rightarrow [0, 1]$, defined by $f(\vec{a}) = \vec{a}^T \mathbb{E}[X]$.

$$\begin{aligned} U_f : |\vec{a}\rangle |0\rangle |0\rangle &\mapsto |\vec{a}\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{\vec{a}^T X(\omega)} |1\rangle + \sqrt{1 - \vec{a}^T X(\omega)} |0\rangle \right) \\ &= |\vec{a}\rangle \left(\sqrt{\vec{a}^T \mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \vec{a}^T \mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$



For $\vec{k} \in \{0, \dots, 2^n - 1\}^d$,

$$\text{QFT}_G : |\vec{k}\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{2\pi i \vec{a}^T \vec{k}} |\vec{a}\rangle.$$


Quantum multivariate Monte Carlo estimation

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}^d$, and $f : G \rightarrow [0, 1]$, defined by $f(\vec{a}) = \vec{a}^T \mathbb{E}[X]$.

$$\begin{aligned} U_f : |\vec{a}\rangle |0\rangle |0\rangle &\mapsto |\vec{a}\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{\vec{a}^T X(\omega)} |1\rangle + \sqrt{1 - \vec{a}^T X(\omega)} |0\rangle \right) \\ &= |\vec{a}\rangle \left(\sqrt{\vec{a}^T \mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \vec{a}^T \mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |\vec{a}\rangle \mapsto e^{i\vec{a}^T \mathbb{E}[X]} |\vec{a}\rangle$ with $\tilde{O}(1)$ calls to U_f , and

$$\begin{aligned} \frac{1}{\sqrt{|G|}} \sum_{\vec{a} \in G} |\vec{a}\rangle &\xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{i\vec{a}^T \mathbb{E}[X]N} |\vec{a}\rangle \\ &\stackrel{\text{QFT}_G^\dagger}{\rightsquigarrow} \left| \text{round} \left(\frac{\mathbb{E}[X]N}{2\pi} \right) \right\rangle. \end{aligned}$$



For $\vec{k} \in \{0, \dots, 2^n - 1\}^d$,

$$\text{QFT}_G : |\vec{k}\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{2\pi i \vec{a}^T \vec{k}} |\vec{a}\rangle.$$

Quantum multivariate Monte Carlo estimation


Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}^d$, and $f : G \rightarrow [0, 1]$, defined by $f(\vec{a}) = \vec{a}^T \mathbb{E}[X]$.

$$\begin{aligned} U_f : |\vec{a}\rangle |0\rangle |0\rangle &\mapsto |\vec{a}\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{\vec{a}^T X(\omega)} |1\rangle + \sqrt{1 - \vec{a}^T X(\omega)} |0\rangle \right) \\ &= |\vec{a}\rangle \left(\sqrt{\vec{a}^T \mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \vec{a}^T \mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |\vec{a}\rangle \mapsto e^{i\vec{a}^T \mathbb{E}[X]} |\vec{a}\rangle$ with $\tilde{O}(1)$ calls to U_f , and

$$\begin{aligned} \frac{1}{\sqrt{|G|}} \sum_{\vec{a} \in G} |\vec{a}\rangle &\xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{i\vec{a}^T \mathbb{E}[X]N} |\vec{a}\rangle \\ &\stackrel{\text{QFT}_G^\dagger}{\rightsquigarrow} \left| \text{round} \left(\frac{\mathbb{E}[X]N}{2\pi} \right) \right\rangle. \end{aligned}$$

We can choose $N = \mathcal{O}(1/\varepsilon) \Rightarrow \tilde{O}(1/\varepsilon)$ calls to U_f .



For $\vec{k} \in \{0, \dots, 2^n - 1\}^d$,

$$\text{QFT}_G : |\vec{k}\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{2\pi i \vec{a}^T \vec{k}} |\vec{a}\rangle.$$

Quantum multivariate Monte Carlo estimation

Let $X : \Omega \rightarrow \{\vec{x} \in \mathbb{R}^d : \|\vec{x}\|_1 \leq 1 \wedge \vec{x} \geq 0\}$.


Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}^d$, and $f : G \rightarrow [0, 1]$, defined by $f(\vec{a}) = \vec{a}^T \mathbb{E}[X]$.

$$\begin{aligned} U_f : |\vec{a}\rangle |0\rangle |0\rangle &\mapsto |\vec{a}\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{\vec{a}^T X(\omega)} |1\rangle + \sqrt{1 - \vec{a}^T X(\omega)} |0\rangle \right) \\ &= |\vec{a}\rangle \left(\sqrt{\vec{a}^T \mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \vec{a}^T \mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |\vec{a}\rangle \mapsto e^{i\vec{a}^T \mathbb{E}[X]} |\vec{a}\rangle$ with $\tilde{O}(1)$ calls to U_f , and

$$\begin{aligned} \frac{1}{\sqrt{|G|}} \sum_{\vec{a} \in G} |\vec{a}\rangle &\xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{i\vec{a}^T \mathbb{E}[X]N} |\vec{a}\rangle \\ &\stackrel{\text{QFT}_G^\dagger}{\rightsquigarrow} \left| \text{round} \left(\frac{\mathbb{E}[X]N}{2\pi} \right) \right\rangle. \end{aligned}$$

We can choose $N = \mathcal{O}(1/\varepsilon) \Rightarrow \tilde{O}(1/\varepsilon)$ calls to U_f .



For $\vec{k} \in \{0, \dots, 2^n - 1\}^d$,

$$\text{QFT}_G : |\vec{k}\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{2\pi i \vec{a}^T \vec{k}} |\vec{a}\rangle.$$

Quantum multivariate Monte Carlo estimation

Let $X : \Omega \rightarrow \{\vec{x} \in \mathbb{R}^d : \|\vec{x}\|_1 \leq 1 \wedge \vec{x} \geq 0\}$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}^d$, and $f : G \rightarrow [0, 1]$, defined by $f(\vec{a}) = \vec{a}^T \mathbb{E}[X]$.


$$\begin{aligned} U_f : |\vec{a}\rangle |0\rangle |0\rangle &\mapsto |\vec{a}\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{\vec{a}^T X(\omega)} |1\rangle + \sqrt{1 - \vec{a}^T X(\omega)} |0\rangle \right) \\ &= |\vec{a}\rangle \left(\sqrt{\vec{a}^T \mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \vec{a}^T \mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$

This can be converted into $O_f : |\vec{a}\rangle \mapsto e^{i\vec{a}^T \mathbb{E}[X]} |\vec{a}\rangle$ with $\tilde{O}(1)$ calls to U_f , and

$$\begin{aligned} \frac{1}{\sqrt{|G|}} \sum_{\vec{a} \in G} |\vec{a}\rangle &\xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{i\vec{a}^T \mathbb{E}[X]N} |\vec{a}\rangle \\ &\stackrel{\text{QFT}_G^\dagger}{\rightsquigarrow} \left| \text{round} \left(\frac{\mathbb{E}[X]N}{2\pi} \right) \right\rangle. \end{aligned}$$

We can choose $N = \mathcal{O}(1/\varepsilon) \Rightarrow \tilde{O}(1/\varepsilon)$ calls to U_f .

+ Same quadratic speed-up in the multidimensional case!



For $\vec{k} \in \{0, \dots, 2^n - 1\}^d$,

$$\text{QFT}_G : |\vec{k}\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{2\pi i \vec{a}^T \vec{k}} |\vec{a}\rangle.$$

Quantum multivariate Monte Carlo estimation

Let $X : \Omega \rightarrow \{\vec{x} \in \mathbb{R}^d : \|\vec{x}\|_1 \leq 1 \wedge \vec{x} \geq 0\}$.

Let $G = \{\frac{j}{2^n} : j \in \{0, \dots, 2^n - 1\}\}^d$, and $f : G \rightarrow [0, 1]$, defined by $f(\vec{a}) = \vec{a}^T \mathbb{E}[X]$.

$$\begin{aligned} U_f : |\vec{a}\rangle |0\rangle |0\rangle &\mapsto |\vec{a}\rangle \sum_{\omega \in \Omega} \sqrt{\mathbb{P}(\omega)} |\omega\rangle \left(\sqrt{\vec{a}^T X(\omega)} |1\rangle + \sqrt{1 - \vec{a}^T X(\omega)} |0\rangle \right) \\ &= |\vec{a}\rangle \left(\sqrt{\vec{a}^T \mathbb{E}[X]} |\psi_1\rangle |1\rangle + \sqrt{1 - \vec{a}^T \mathbb{E}[X]} |\psi_0\rangle |0\rangle \right). \end{aligned}$$


This can be converted into $O_f : |\vec{a}\rangle \mapsto e^{i\vec{a}^T \mathbb{E}[X]} |\vec{a}\rangle$ with $\tilde{O}(1)$ calls to U_f , and

$$\begin{aligned} \frac{1}{\sqrt{|G|}} \sum_{\vec{a} \in G} |\vec{a}\rangle &\xrightarrow{O_f^N} \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{i\vec{a}^T \mathbb{E}[X]N} |\vec{a}\rangle \\ &\stackrel{\text{QFT}_G^\dagger}{\rightsquigarrow} \left| \text{round} \left(\frac{\mathbb{E}[X]N}{2\pi} \right) \right\rangle. \end{aligned}$$

We can choose $N = \mathcal{O}(1/\varepsilon) \Rightarrow \tilde{O}(1/\varepsilon)$ calls to U_f .

+ Same quadratic speed-up in the multidimensional case!

– Weird assumption on X required.



For $\vec{k} \in \{0, \dots, 2^n - 1\}^d$,

$$\text{QFT}_G : |\vec{k}\rangle \mapsto \frac{1}{\sqrt{G}} \sum_{\vec{a} \in G} e^{2\pi i \vec{a}^T \vec{k}} |\vec{a}\rangle.$$

Improvements

Improvements on the algorithm: [CJ21]

Improvements

Improvements on the algorithm: [CJ21]

- 1 By modifying the construction, we can relax that $X \geq 0$.

Improvements

Improvements on the algorithm: [CJ21]

- 1 By modifying the construction, we can relax that $X \geq 0$.
- 2 By moving the grid to be centered around the origin and observing that

$$\mathbb{E}_{\vec{a} \sim G} \left[\left| \vec{a}^T \vec{v} \right| \right] = \mathcal{O}(\|\vec{v}\|_2),$$

we can relax the ℓ_1 -bound on X to be an ℓ_2 -bound.

Improvements

Improvements on the algorithm: [CJ21]

- 1 By modifying the construction, we can relax that $X \geq 0$.
- 2 By moving the grid to be centered around the origin and observing that

$$\mathbb{E}_{\vec{a} \sim G} \left[\left| \vec{a}^T \vec{v} \right| \right] = \mathcal{O}(\|\vec{v}\|_2),$$

we can relax the ℓ_1 -bound on X to be an ℓ_2 -bound.

- 3 By rotating the grid, we can compute $O\mathbb{E}[X]$, for some $O^T O = O O^T = I$.

Improvements

Improvements on the algorithm: [CJ21]

- 1 By modifying the construction, we can relax that $X \geq 0$.
- 2 By moving the grid to be centered around the origin and observing that

$$\mathbb{E}_{\vec{a} \sim G} \left[\left| \vec{a}^T \vec{v} \right| \right] = \mathcal{O}(\|\vec{v}\|_2),$$

we can relax the ℓ_1 -bound on X to be an ℓ_2 -bound.

- 3 By rotating the grid, we can compute $O\mathbb{E}[X]$, for some $O^T O = O O^T = I$.

Future directions:

- 1 What if $X : \Omega \rightarrow \mathbb{R}^d$, and all X_j 's have bounded variance?
Very recent paper by Hamoudi solves the univariate case. [Ham21]

Improvements

Improvements on the algorithm: [CJ21]

- 1 By modifying the construction, we can relax that $X \geq 0$.
- 2 By moving the grid to be centered around the origin and observing that

$$\mathbb{E}_{\vec{a} \sim G} \left[\left| \vec{a}^T \vec{v} \right| \right] = \mathcal{O}(\|\vec{v}\|_2),$$

we can relax the ℓ_1 -bound on X to be an ℓ_2 -bound.

- 3 By rotating the grid, we can compute $O\mathbb{E}[X]$, for some $O^T O = O O^T = I$.

Future directions:

- 1 What if $X : \Omega \rightarrow \mathbb{R}^d$, and all X_j 's have bounded variance?
Very recent paper by Hamoudi solves the univariate case. [Ham21]
- 2 Other ℓ_p estimates & lower bounds?

Improvements

Improvements on the algorithm: [CJ21]

- 1 By modifying the construction, we can relax that $X \geq 0$.
- 2 By moving the grid to be centered around the origin and observing that

$$\mathbb{E}_{\vec{a} \sim G} \left[\left| \vec{a}^T \vec{v} \right| \right] = \mathcal{O}(\|\vec{v}\|_2),$$

we can relax the ℓ_1 -bound on X to be an ℓ_2 -bound.

- 3 By rotating the grid, we can compute $O\mathbb{E}[X]$, for some $O^T O = O O^T = I$.

Future directions:

- 1 What if $X : \Omega \rightarrow \mathbb{R}^d$, and all X_j 's have bounded variance?
Very recent paper by Hamoudi solves the univariate case. [Ham21]
- 2 Other ℓ_p estimates & lower bounds?
- 3 Unbiased estimators?

Thanks for your attention!
arjan@cw.nl