

1 Binary Logistic Regression

Binary Logistic regression is used to determine the impact of multiple independent variables presented simultaneously to predict membership of one or other of the two dependent variable categories.

1.1 The Hosmer-Lemeshow Test

Performing the Hosmer-Lemeshow Goodness-of-Fit Test

- The Hosmer-Lemeshow test of goodness of fit is not automatically a part of the SPSS logistic regression output. To get this output, we need to go into **options** and tick the box marked Hosmer-Lemeshow test of goodness of fit.

In our example, this gives us the following output:

Step	Chi-square	df	Sig.
1	142.032	6	.000

Therefore, our model is significant, suggesting it does not fit the data. However, as we have a sample size of over 13,000, even very small divergencies of the model from the data would be flagged up and cause significance. Therefore, with samples of this size it is hard to find models that are parsimonious (i.e. that use the minimum amount of independent variables to explain the dependent variable) and fit the data. Therefore, other fit indices might be more appropriate.

2 Logistic Regression

Logistic regression, also called a logit model, is used to model **dichotomous outcome** variables. In the logit model the **log odds** of the outcome is modeled as a linear combination of the predictor variables.

In logistic regression theory, the predicted dependent variable is a function of the probability that a particular subject will be in one of the categories (for example, the probability that a patient has the disease, given his or her set of scores on the predictor variables).

2.1 Model Summary Table

The likelihood function can be thought of as a measure of how well a candidate model fits the data (although that is a very simplistic definition). The AIC criterion is based on the Likelihood function. The likelihood function of a fitted model is commonly re-expressed as -2LL (i.e. The log of the likelihood times minus 2). The 2LL value from the Model Summary table below is 17.359.

2.2 Hosmer and Lemeshow Statistic

An alternative to model chi square is the Hosmer and Lemeshow test which divides subjects into 10 ordered groups of subjects and then compares the number actually in the each group

(observed) to the number predicted by the logistic regression model (predicted). The 10 ordered groups are created based on their estimated probability; those with estimated probability below .1 form one group, and so on, up to those with probability .9 to 1.0.

Each of these categories is further divided into two groups based on the actual observed outcome variable (success, failure). The expected frequencies for each of the cells are obtained from the model. A probability (p) value is computed from the chi-square distribution with 8 degrees of freedom to test the fit of the logistic model.

If the H-L goodness-of-fit test statistic is greater than .05, as we want for well-fitting models, we fail to reject the null hypothesis that there is no difference between observed and model-predicted values, implying that the models estimates fit the data at an acceptable level. That is, well-fitting models show non-significance on the H-L goodness-of-fit test. This desirable outcome of non-significance indicates that the model prediction does not significantly differ from the observed.

2.3 Computing the Hosmer and Lemeshow Test Statistic

- The Hosmer and Lemeshow test divides subjects into 10 ordered groups of subjects and then compares the number actually in the each group (observed) to the number predicted by the logistic regression model (predicted).
- The 10 ordered groups are created based on their estimated probability; those with estimated probability below .1 form one group, and so on, up to those with probability .9 to 1.0.
- Each of these categories is further divided into two groups based on the actual observed outcome variable (success, failure). The expected frequencies for each of the cells are obtained from the model. A probability (p) value is computed from the chi-square distribution with 8 degrees of freedom to test the fit of the logistic model.

Contingency Table for Hosmer and Lemeshow Test						
		take solar panel offer = decline offer		take solar panel offer = take offer		Total
		Observed	Expected	Observed	Expected	
Step 1	1	3	2.996	0	.004	3
	2	3	2.957	0	.043	3
	3	3	2.629	0	.371	3
	4	2	2.136	1	.864	3
	5	2	1.881	1	1.119	3
	6	0	.833	3	2.167	3
	7	0	.376	3	2.624	3
	8	1	.171	2	2.829	3
	9	0	.019	3	2.981	3
	10	0	.000	3	3.000	3

Figure 1: Hosmer and Lemeshow Table

2.4 Sample Adequacy

The H-L statistic assumes sampling adequacy, with a rule of thumb being enough cases so that 95% of cells (typically, 10 decile groups times 2 outcome categories = 20 cells) have an expected frequency > 5 . Our H-L statistic has a significance of .605 which means that it is not statistically significant and therefore our model is quite a good fit.

Hosmer and Lemeshow Test			
Step	Chi-square	df	Sig.
1	6.378	8	.605

Figure 2: Hosmer and Lemeshow Statistic

Contingency Table for Hosmer and Lemeshow Test					
		take solar panel offer = decline offer		take solar panel offer = take offer	
		Observed	Expected	Observed	Expected
Step 1	1	3	2.996	0	.004
	2	3	2.957	0	.043
	3	3	2.629	0	.371
	4	2	2.136	1	.864
	5	2	1.881	1	1.119
	6	0	.833	3	2.167
	7	0	.376	3	2.624
	8	1	.171	2	2.829
	9	0	.019	3	2.981
	10	0	.000	3	3.000
				Total	
				3	
				3	
				3	
				3	
				3	
				3	
				3	
				3	
				3	

Figure 3: Hosmer and Lemeshow Table

3 Model Diagnostics for Logistic Regression

3.1 Logistic Regression

$$\pi(x) = \frac{e^{(\beta_0 + \beta_1 x)}}{e^{(\beta_0 + \beta_1 x)} + 1} = \frac{1}{e^{-(\beta_0 + \beta_1 x)} + 1},$$

and

$$g(x) = \ln \frac{\pi(x)}{1 - \pi(x)} = \beta_0 + \beta_1 x,$$

and

$$\frac{\pi(x)}{1 - \pi(x)} = e^{(\beta_0 + \beta_1 x)}.$$

3.2 Logistic Regression: Decision Rule

Our decision rule will take the following form: If the probability of the event is greater than or equal to some threshold, we shall predict that the event will take place. By default, SPSS sets this threshold to .5. While that seems reasonable, in many cases we may want to set it higher or lower than .5.

4 Model Diagnostics for Logistic Regression

Remark upon “Steps”

- This table has 1 step. This is because we are entering both variables and at the same time providing only one model to compare with the constant model.
- In stepwise logistic regression there are a number of steps listed in the table as each variable is added or removed, creating different models.
- The step is a measure of the improvement in the predictive power of the model since the previous step.
- I will revert to this next class.

4.1 Model Summary Table

- The likelihood function can be thought of as a measure of how well a candidate model fits the data (although that is a very simplistic definition).
- Later on in the course, we will meet the AIC Function. The AIC criterion is based on the Likelihood function.
- The likelihood function of a fitted model is commonly re-expressed as $-2LL$ (i.e. The log of the likelihood times minus 2). The 2LL value from the Model Summary table below is 17.359.
- It is not immediately clear how to interpret $-2LL$ yet. However this measure is useful in comparing various candidate models, and is therefore used in **Variable Selection Procedure**, such as backward and forward selection.

4.2 Hosmer-Lemeshow Goodness-of-Fit Test

- The Hosmer-Lemeshow Goodness-of-Fit test tells us whether we have constructed a valid overall model or not, and is an alternative to model chi square procedure.
- If the model is a good fit to the data then the Hosmer-Lemeshow Goodness-of-Fit test should have an associated p-value greater than 0.05.

Block 1: Method = Enter

Omnibus Tests of Model Coefficients

		Chi-square	df	Sig.
Step 1	Step	24.096	2	.000
	Block	24.096	2	.000
	Model	24.096	2	.000

Model Summary

Step	-2 Log likelihood	Cox & Snell R Square	Nagelkerke R Square
1	17.359 ^a	.552	.737

a. Estimation terminated at iteration number 8 because parameter estimates changed by less than .001.

Hosmer and Lemeshow Test

Step	Chi-square	df	Sig.
1	6.378	8	.605