

Binary Logistic Regression: Tutorial Sheet

1. What is logistic regression? How does it differ from linear regression? Under what circumstances would you use it?
2. What is the Likelihood Ratio Test? Describe how it used in Logistic regression.
3. Suppose the odds of an outcome are 4. What is the probability of that outcome?
4. Suppose the probability of an outcome is 70%. What is the odds of that outcome occurring? Answer this question for probabilities of 40%, 60% and 90% also.
5. The usual assumptions placed on the error terms in ordinary least squares regression are:
 - * independently distributed
 - * identically distributed (equal variance)
 - * normally distributed

Which of these assumptions are violated when dealing with binary response data? Explain briefly how each is violated.

6. What is a dummy variable? Explain how it is used in Logistic Regression. Support your answer with an example.
7. Suppose that, out of a sample of 100 women and 100 men, 80 men drank alcohol in the last week, while 20 women drank alcohol in past week. Compute the odds ratio for Women to men.
8. What is a logit? how is it computed into a probability?

9. A predictive model was used to predict onset of Coronary Heart Disease, based on a set of medical measurements and characteristics. Comment on the significance of each of the predictor variables used in this model.

sbp systolic blood pressure

tobacco cumulative tobacco (kg)

ldl low density lipoprotein cholesterol

adiposity A BMI index

famhist family history of heart disease, a factor with levels (Absent/Present)

typea type-A behavior

obesity A BMI index

alcohol current alcohol consumption

age age at onset

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	-6.1507209	1.3082600	-4.701	2.58e-06	***
sbp	0.0065040	0.0057304	1.135	0.256374	
tobacco	0.0793764	0.0266028	2.984	0.002847	**
ldl	0.1739239	0.0596617	2.915	0.003555	**
adiposity	0.0185866	0.0292894	0.635	0.525700	
famhistPresent	0.9253704	0.2278940	4.061	4.90e-05	***
typea	0.0395950	0.0123202	3.214	0.001310	**
obesity	-0.0629099	0.0442477	-1.422	0.155095	
alcohol	0.0001217	0.0044832	0.027	0.978350	
age	0.0452253	0.0121298	3.728	0.000193	***

10. *This question is a continuation of the previous question.* Using the following binary logistic model, which is used to predict the onset of Coronary Heart Disease, compute the predicted outcomes for the cases below;

Coefficients:

	Estimate	Std. Error	z value	Pr(> z)	
(Intercept)	-6.334452	0.897809	-7.055	1.72e-12	***
age	0.055040	0.009948	5.533	3.15e-08	***
ldl	0.179891	0.055027	3.269	0.00108	**
typea	0.037914	0.011885	3.190	0.00142	**
tobacco	0.075031	0.025699	2.920	0.00350	**

Signif. codes: 0 *** 0.001 ** 0.01 * 0.05 . 0.1 1

- (i) 52 year old person, $ldl = 5.73$, $typea = 49$, $tobacco = 12$
 - (ii) 63 year old person, $ldl = 4.41$, $typea = 55$, $tobacco = 0.01$
11. There are three variants for the forward selection procedures used by SPSS. Name these three.