
Monte Carlo Simulation



Photo by Sam Garza

Stanislaw Ulam




We've Been Doing this Already

- •A method of estimating the value of an unknown quantity using principles of inferential statistics.
- Inferential statistics
 - Population: a set of examples
 - Sample: a proper subset of a population
 - Key fact: a random sample tends to exhibit the same properties as the population from which it is drawn.

Finding Pi

3.1415926535897932384626433832795028841971693

Image from Tom Murphy



$$\frac{circumference}{diameter} = \Pi \quad area = \Pi * radius^2$$

Rhind Papyrus




The Bible


"And he made a molten sea, ten cubits from the one brim to the other: it was round all about, and his height was five cubits: and a line of thirty cubits did compass it round about."

—1 Kings 7.23

Archimedes



Buffon-Laplace



$$A_s = 2*2 = 4$$

 $A_c = \pi r^2 = \pi$

$$\frac{needles\ in\ circle}{needles\ in\ square} = \frac{area\ of\ circle}{area\ of\ square}$$

$$\frac{area\ of\ square*needles\ in\ circle}{needles\ in\ square}$$

$$area\ of\ circle = \frac{4*needles\ in\ circle}{needles\ in\ square}$$

Arrows Are More Fun than Needles



Photo Dharma

Not a Practical Method

In the next segment, we take Ana's advice and build a simulation