
A
Wild
Wild−like
Lab−like
Lab
PCoA2 (21.2%)
PCoA1 (58.5%)
B
Wild−like vs Wild
Lab−like vs Wild
Lab vs Wild
Lab−like vs Wild−like
Lab vs Wild−like
Lab vs Lab−like
Pairwise PERMANOVA R2
C
32/234 (13.7%)
117/234 (50%)
129/234 (55.1%)
144/234 (61.5%)
153/234 (65.4%)
89/234 (38%)
FDR−significant predicted KEGG pathways
D
Higher toward lab−associated
Higher toward wild−associated
Wild – Wild−like
Wild – Lab−like
Wild – Lab
Wild−like – Lab−like
Wild−like – Lab
Lab−like – Lab
Metabolism
Biosynthesis of other secondary metabolites (n = 25)
Xenobiotics biodegradation and metabolism (n = 21)
Glycan biosynthesis and metabolism (n = 18)
Metabolism of terpenoids and polyketides (n = 17)
Lipid metabolism (n = 16)
Carbohydrate metabolism (n = 15)
Amino acid metabolism (n = 14)
Metabolism of cofactors and vitamins (n = 12)
Energy metabolism (n = 8)
Metabolism of other amino acids (n = 7)
Nucleotide metabolism (n = 2)
Genetic information processing
Replication and repair (n = 6)
Folding, sorting and degradation (n = 5)
Translation (n = 4)
Transcription (n = 3)
Information processing in viruses (n = 2)
Environmental information processing
Signal transduction (n = 17)
Membrane transport (n = 3)
Signaling molecules and interaction (n = 1)
Cellular processes
Cell growth and death (n = 8)
Transport and catabolism (n = 7)
Cell motility (n = 4)
Cellular community − prokaryotes (n = 4)
Cellular community − eukaryotes (n = 1)
Human diseases
Infectious disease: bacterial (n = 11)
Drug resistance: antimicrobial (n = 3)
FDR−significant pathways within KEGG Level 2 category (%)