

An index formula for families of end-periodic Dirac operators

... and anomalies in QFT

Alex Taylor

University of Illinois at Urbana-Champaign

GLaMP 2025 – June 6, 2025

- 1 Index theory for Dirac operators
- 2 Index theory and anomalies
- 3 Index theory for end-periodic Dirac operators
- 4 A new index formula
- 5 Remarks on the proof depending on how much time is left

Geometric Dirac operators

Geometric Dirac operators

(M^n, g) = closed, even-dimensional Riemannian manifold.

$E \rightarrow M$ vector bundle with connection ∇ .

Δ = Laplace-type operator on $C^\infty(M; E)$.

Geometric Dirac operators

(M^n, g) = closed, even-dimensional Riemannian manifold.

$E \rightarrow M$ vector bundle with connection ∇ .

Δ = Laplace-type operator on $C^\infty(M; E)$.

Paul Dirac in 1928: can we find an operator D such that $D^2 = \Delta$?

Geometric Dirac operators

(M^n, g) = closed, even-dimensional Riemannian manifold.

$E \rightarrow M$ vector bundle with connection ∇ .

Δ = Laplace-type operator on $C^\infty(M; E)$.

Paul Dirac in 1928: can we find an operator D such that $D^2 = \Delta$?

Answer: define $D = \sum_j \text{cl}(e^j) \nabla_{e_j}$, where

$$\text{cl} : T^*M \rightarrow \text{End } E$$

satisfying $\text{cl}(e^i) \text{cl}(e^j) + \text{cl}(e^j) \text{cl}(e^i) = -2g^{ij} \text{id}$

(i.e. $E \rightarrow M$ is a Clifford module with Clifford action cl).

Geometric Dirac operators

(M^n, g) = closed, even-dimensional Riemannian manifold.

$E \rightarrow M$ vector bundle with connection ∇ .

Δ = Laplace-type operator on $C^\infty(M; E)$.

Paul Dirac in 1928: can we find an operator D such that $D^2 = \Delta$?

Answer: define $D = \sum_j \text{cl}(e^j) \nabla_{e_j}$, where

$$\text{cl} : T^*M \rightarrow \text{End } E$$

satisfying $\text{cl}(e^i) \text{cl}(e^j) + \text{cl}(e^j) \text{cl}(e^i) = -2g^{ij} \text{id}$

(i.e. $E \rightarrow M$ is a Clifford module with Clifford action cl).

ex. $D^+ = d + d^*$ acting on $\Omega^{\text{ev}}(M) \subseteq \Omega^\bullet(M)$ (Gauss-Bonnet operator).

Geometric Dirac operators

(M^n, g) = closed, even-dimensional Riemannian manifold.

$E \rightarrow M$ vector bundle with connection ∇ .

Δ = Laplace-type operator on $C^\infty(M; E)$.

Paul Dirac in 1928: can we find an operator D such that $D^2 = \Delta$?

Answer: define $D = \sum_j \text{cl}(e^j) \nabla_{e_j}$, where

$$\text{cl} : T^*M \rightarrow \text{End } E$$

satisfying $\text{cl}(e^i) \text{cl}(e^j) + \text{cl}(e^j) \text{cl}(e^i) = -2g^{ij} \text{id}$
(i.e. $E \rightarrow M$ is a Clifford module with Clifford action cl).

ex. $D^+ = d + d^*$ acting on $\Omega^{\text{ev}}(M) \subseteq \Omega^*(M)$ (Gauss-Bonnet operator).

ex. $D^+ = d + d^*$ acting on self-dual forms $\Omega_+^{n/2}(M)$, i.e. $*\omega = \omega$
(signature operator).

The Fredholm index

The Fredholm index

D^+ is a Fredholm operator on $H^1(M; E^+)$, meaning the Fredholm index

$$\operatorname{ind} D^+ = \dim \ker D^+ - \dim \operatorname{coker} D^+$$

is a finite integer $\operatorname{ind} D^+ \in \mathbb{Z}$.

The Fredholm index

D^+ is a Fredholm operator on $H^1(M; E^+)$, meaning the Fredholm index

$$\operatorname{ind} D^+ = \dim \ker D^+ - \dim \operatorname{coker} D^+$$

is a finite integer $\operatorname{ind} D^+ \in \mathbb{Z}$.

It turns out that the index of a Dirac operator is an interesting *topological invariant*!

The Fredholm index

D^+ is a Fredholm operator on $H^1(M; E^+)$, meaning the Fredholm index

$$\operatorname{ind} D^+ = \dim \ker D^+ - \dim \operatorname{coker} D^+$$

is a finite integer $\operatorname{ind} D^+ \in \mathbb{Z}$.

It turns out that the index of a Dirac operator is an interesting *topological invariant*!

ex. $\operatorname{ind}(d + d^*) = \chi(M) := \sum_{k=0}^n (-1)^k \dim H_{dR}^k(M).$

The Fredholm index

D^+ is a Fredholm operator on $H^1(M; E^+)$, meaning the Fredholm index

$$\operatorname{ind} D^+ = \dim \ker D^+ - \dim \operatorname{coker} D^+$$

is a finite integer $\operatorname{ind} D^+ \in \mathbb{Z}$.

It turns out that the index of a Dirac operator is an interesting *topological invariant*!

ex. $\operatorname{ind}(d + d^*) = \chi(M) := \sum_{k=0}^n (-1)^k \dim H_{dR}^k(M).$

Consider simple situation where $\dim M = 2$. Gauss-Bonnet theorem tells us that

$$\operatorname{ind}(d + d^*) = \chi(M) = \frac{1}{2\pi} \int_M K \, dA.$$

The Atiyah-Singer local index theorem

The Atiyah-Singer local index theorem

Natural question: in general, for a Dirac operator D , is there a local geometric formula for $\text{ind } D^+$?

The Atiyah-Singer local index theorem

Natural question: in general, for a Dirac operator D , is there a local geometric formula for $\text{ind } D^+$?

Answer: yes! (Atiyah, Getzler, Patodi, Singer ..., circa 1971)

The Atiyah-Singer local index theorem

Natural question: in general, for a Dirac operator D , is there a local geometric formula for $\text{ind } D^+$?

Answer: yes! (Atiyah, Getzler, Patodi, Singer ..., circa 1971)

$$\text{ind } D^+ = \frac{1}{(2\pi i)^{n/2}} \int_M \hat{A}(TM) \wedge \text{ch}'(E)$$

The Atiyah-Singer local index theorem

Natural question: in general, for a Dirac operator D , is there a local geometric formula for $\text{ind } D^+$?

Answer: yes! (Atiyah, Getzler, Patodi, Singer ..., circa 1971)

$$\text{ind } D^+ = \frac{1}{(2\pi i)^{n/2}} \int_M \hat{A}(TM) \wedge \text{ch}'(E)$$

Analysis \leftrightarrow Geometry/Topology

The Atiyah-Singer local index theorem

Natural question: in general, for a Dirac operator D , is there a local geometric formula for $\text{ind } D^+$?

Answer: yes! (Atiyah, Getzler, Patodi, Singer ..., circa 1971)

$$\text{ind } D^+ = \frac{1}{(2\pi i)^{n/2}} \int_M \hat{A}(TM) \wedge \text{ch}'(E)$$

Analysis \leftrightarrow Geometry/Topology

Notation: $\text{ind } D^+ = \int_M AS(D(M)).$

- 1 Index theory for Dirac operators
- 2 Index theory and anomalies**
- 3 Index theory for end-periodic Dirac operators
- 4 A new index formula
- 5 Remarks on the proof depending on how much time is left

The Atiyah-Patodi-Singer index theorem

Atiyah, Patodi, Singer, 1975: under the appropriate *global* boundary conditions, the Dirac operator $D^+(Z)$ is Fredholm, and the index is given by

The Atiyah-Patodi-Singer index theorem

Atiyah, Patodi, Singer, 1975: under the appropriate *global* boundary conditions, the Dirac operator $D^+(Z)$ is Fredholm, and the index is given by

$$\operatorname{ind} D^+(Z) = \int_Z AS(D(Z)) - \frac{1}{2}\eta(D(\partial Z))$$

The Atiyah-Patodi-Singer index theorem

Atiyah, Patodi, Singer, 1975: under the appropriate *global* boundary conditions, the Dirac operator $D^+(Z)$ is Fredholm, and the index is given by

$$\text{ind } D^+(Z) = \int_Z AS(D(Z)) - \frac{1}{2}\eta(D(\partial Z))$$

where the *eta invariant* is defined as

$$\eta(D(\partial Z)) = \frac{1}{\sqrt{\pi}} \int_0^\infty t^{-1/2} \text{Tr}(D(\partial Z)e^{-tD(\partial Z)^2}) dt.$$

Anomalies in QFT

Anomalies in QFT

(M, g_0) smooth Riemannian spin manifold. $\pi : M \rightarrow M$ a diffeomorphism. Given a metric g , Witten considers the *effective action* for a Majorana-Weyl fermion:

$$I(g) = \frac{1}{2} \ln(\det(i\partial)).$$

Anomalies in QFT

(M, g_0) smooth Riemannian spin manifold. $\pi : M \rightarrow M$ a diffeomorphism. Given a metric g , Witten considers the *effective action* for a Majorana-Weyl fermion:

$$I(g) = \frac{1}{2} \ln(\det(i\partial)).$$

Consider the variation ΔI of $I(g_t)$, where g_t interpolates between g_0 and $g_1 = \pi^* g_0$. If $\Delta I = 0$ then the theory is anomaly-free. How to compute ΔI ?

Anomalies in QFT

(M, g_0) smooth Riemannian spin manifold. $\pi : M \rightarrow M$ a diffeomorphism. Given a metric g , Witten considers the *effective action* for a Majorana-Weyl fermion:

$$I(g) = \frac{1}{2} \ln(\det(i\partial)).$$

Consider the variation ΔI of $I(g_t)$, where g_t interpolates between g_0 and $g_1 = \pi^* g_0$. If $\Delta I = 0$ then the theory is anomaly-free. How to compute ΔI ?

Consider the mapping torus $(M \times S^1)_\pi = (M \times [0, 1]) / \sim$, where $(x, 1) \sim (\pi(x), 0)$. Suppose $(M \times S^1)_\pi = \partial B$. Then using APS, Witten computes ΔI in terms of the eta invariant $\eta(D(\partial B))$.


Anomalies in QFT

(M, g_0) smooth Riemannian spin manifold. $\pi : M \rightarrow M$ a diffeomorphism. Given a metric g , Witten considers the *effective action* for a Majorana-Weyl fermion:

$$I(g) = \frac{1}{2} \ln(\det(i\partial)).$$

Consider the variation ΔI of $I(g_t)$, where g_t interpolates between g_0 and $g_1 = \pi^* g_0$. If $\Delta I = 0$ then the theory is anomaly-free. How to compute ΔI ?

Consider the mapping torus $(M \times S^1)_\pi = (M \times [0, 1]) / \sim$, where $(x, 1) \sim (\pi(x), 0)$. Suppose $(M \times S^1)_\pi = \partial B$. Then using APS, Witten computes ΔI in terms of the eta invariant $\eta(D(\partial B))$.



Anomalies in QFT

Global gravitational anomalies, Witten, 1985

where $H = dB + \omega_3^L$. Thus, the non-covariant objects dB and ω combine, as they must, into the covariant field strength H . ($\Delta\tilde{S}$ and ΔI_{Reg} were invariant under coordinate transformations that vanish at $t=0$ and $t=1$, but not otherwise. As far as I know, it does not make sense to identify $t=1$ with $t=0$ by means of a non-trivial diffeomorphism π until these expressions are combined into a covariant form.) Thus we finally get an expression for the change in the action in terms of topological invariants:

$$\begin{aligned}\Delta I_{\text{TOT}} &= \Delta I_{\text{det}} + \Delta\tilde{S} + \Delta I_{\text{Reg}} \\ &= -2\pi i \left[\frac{1}{(2\pi)^6} \int_B \left(\frac{1}{1536} (\text{Tr } R^2)^3 + \frac{1}{384} \text{Tr } R^2 \text{Tr } R^4 \right) \right. \\ &\quad \left. - \int_{(M \times S^1)_\pi} H \cdot \left(\frac{1}{1536} (\text{Tr } R^2)^2 + \frac{1}{384} \text{Tr } R^4 \right) \right] \\ &= -2\pi i \cdot \frac{1}{192} [-3p_1^3(B) + 4p_1 p_2(B)],\end{aligned}\tag{57}$$

where p_1^3 and $p_1 p_2$ were defined in Sect. II.

Families of Dirac operators

Families of Dirac operators

$D = (D^z)_{z \in B}$ = family of Dirac operators on family of Clifford modules $E_z \rightarrow M_z$.
 D defines a K -theory class $\text{Ind } D = [\ker D] - [\text{coker } D] \in K^0(B)$.

Families of Dirac operators

$D = (D^z)_{z \in B}$ = family of Dirac operators on family of Clifford modules $E_z \rightarrow M_z$.
 D defines a K -theory class $\text{Ind } D = [\ker D] - [\text{coker } D] \in K^0(B)$.

Chern character form of a (super)connection:

$$\text{ch}(\nabla) = \text{Str}(e^{-\nabla^2}) \in \Omega^{\text{even}}(B).$$

Families of Dirac operators

$D = (D^z)_{z \in B}$ = family of Dirac operators on family of Clifford modules $E_z \rightarrow M_z$.
 D defines a K -theory class $\text{Ind } D = [\ker D] - [\text{coker } D] \in K^0(B)$.

Chern character form of a (super)connection:

$$\text{ch}(\nabla) = \text{Str}(e^{-\nabla^2}) \in \Omega^{\text{even}}(B).$$

ch defines a ring homomorphism:

$$\text{ch} : K^0(B) \rightarrow H_{dR}^{\text{even}}(B)$$

$$\text{ch}([E] - [F]) = [\text{ch}(\nabla^E)] - [\text{ch}(\nabla^F)].$$

Families of Dirac operators

$D = (D^z)_{z \in B}$ = family of Dirac operators on family of Clifford modules $E_z \rightarrow M_z$.
 D defines a K -theory class $\text{Ind } D = [\ker D] - [\text{coker } D] \in K^0(B)$.

Chern character form of a (super)connection:

$$\text{ch}(\nabla) = \text{Str}(e^{-\nabla^2}) \in \Omega^{\text{even}}(B).$$

ch defines a ring homomorphism:

$$\text{ch} : K^0(B) \rightarrow H_{dR}^{\text{even}}(B)$$

$$\text{ch}([E] - [F]) = [\text{ch}(\nabla^E)] - [\text{ch}(\nabla^F)].$$

Goal: find a ‘nice’ representative for $\text{ch}(\text{Ind } D)$ in de Rham cohomology.

Families of Dirac operators

$D = (D^z)_{z \in B}$ = family of Dirac operators on family of Clifford modules $E_z \rightarrow M_z$.
 D defines a K -theory class $\text{Ind } D = [\ker D] - [\text{coker } D] \in K^0(B)$.

Chern character form of a (super)connection:

$$\text{ch}(\nabla) = \text{Str}(e^{-\nabla^2}) \in \Omega^{\text{even}}(B).$$

ch defines a ring homomorphism:

$$\text{ch} : K^0(B) \rightarrow H_{dR}^{\text{even}}(B)$$

$$\text{ch}([E] - [F]) = [\text{ch}(\nabla^E)] - [\text{ch}(\nabla^F)].$$

Goal: find a ‘nice’ representative for $\text{ch}(\text{Ind } D)$ in de Rham cohomology.

Important construction: the **Bismut superconnection**

$$A \circ C^\infty(M; E \otimes \wedge T^*B)$$

$$A = D + A_{[1]} + A_{[2]}.$$

Generalization of APS for families

Generalization of APS for families

Bismut-Cheeger 1990, Melrose-Piazza 1997: under the appropriate global boundary conditions, the family $(D^+(Z)^z)_{z \in B}$ is Fredholm, and the Chern character of the index bundle $\text{ch}(\text{Ind } D)$ is represented in de Rham cohomology by the differential form

Generalization of APS for families

Bismut-Cheeger 1990, Melrose-Piazza 1997: under the appropriate global boundary conditions, the family $(D^+(Z)^z)_{z \in B}$ is Fredholm, and the Chern character of the index bundle $\text{ch}(\text{Ind } D)$ is represented in de Rham cohomology by the differential form

$$\int_{Z/B} AS(D(Z)) - \frac{1}{2} \hat{\eta}(D(\partial Z)) \in \Omega(B)$$

Generalization of APS for families

Bismut-Cheeger 1990, Melrose-Piazza 1997: under the appropriate global boundary conditions, the family $(D^+(Z)^z)_{z \in B}$ is Fredholm, and the Chern character of the index bundle $\text{ch}(\text{Ind } D)$ is represented in de Rham cohomology by the differential form

$$\int_{Z/B} AS(D(Z)) - \frac{1}{2} \hat{\eta}(D(\partial Z)) \in \Omega(B)$$

where the *eta form* is defined as

$$\hat{\eta}(D(\partial Z)) = \frac{1}{\sqrt{\pi}} \int_0^\infty \text{Str} \left(\text{cl} \left(\frac{dx}{x} \right) \dot{A}_t(\partial Z) e^{-A_t^2(\partial Z)} \right) dt.$$

Anomaly as a determinant line bundle

Anomaly as a determinant line bundle

Witten's “effective action” is a special case of the Feynman path integral or pfaffian of a Dirac operator – the canonical section $\det(D_{M/B})$ of the determinant line bundle.

Anomaly as a determinant line bundle

Witten's “effective action” is a special case of the Feynman path integral or pfaffian of a Dirac operator – the canonical section $\det(D_{M/B})$ of the determinant line bundle.

Let $M \rightarrow B$ Riemannian spin fiber bundle, and $E_z \rightarrow M_z$ family of Clifford modules. Consider the determinant line bundle

$$Det(D_{M/B}) \rightarrow B.$$

Anomaly as a determinant line bundle

Witten's “effective action” is a special case of the Feynman path integral or pfaffian of a Dirac operator – the canonical section $\det(D_{M/B})$ of the determinant line bundle.

Let $M \rightarrow B$ Riemannian spin fiber bundle, and $E_z \rightarrow M_z$ family of Clifford modules. Consider the determinant line bundle

$$Det(D_{M/B}) \rightarrow B.$$

Bismut-Freed: the degree two component of $\text{ch}(\text{Ind } D)$ is the curvature of the Quillen connection on the determinant line bundle.

Anomaly as a determinant line bundle

Witten's “effective action” is a special case of the Feynman path integral or pfaffian of a Dirac operator – the canonical section $\det(D_{M/B})$ of the determinant line bundle.

Let $M \rightarrow B$ Riemannian spin fiber bundle, and $E_z \rightarrow M_z$ family of Clifford modules. Consider the determinant line bundle


$$\text{Det}(D_{M/B}) \rightarrow B.$$

Bismut-Freed: the degree two component of $\text{ch}(\text{Ind } D)$ is the curvature of the Quillen connection on the determinant line bundle.

A path of metrics g_t is a loop in the parameter space B and Witten's global anomaly is recovered as the holonomy around this loop of the connection on the determinant line bundle.


- 1 Index theory for Dirac operators
- 2 Index theory and anomalies
- 3 Index theory for end-periodic Dirac operators**
- 4 A new index formula
- 5 Remarks on the proof depending on how much time is left

Cylindrical end = boundary




$$g = dt^2 + g_{\partial Z}$$


Cylindrical end = boundary



$$g = dt^2 + g_{\partial Z}$$



$$x = e^{-t}$$



$$g = \frac{dx^2}{x^2} + g_{\partial Z}$$

End-periodic manifolds

End-periodic manifolds

(Taubes 1987)



End-periodic manifolds

(Taubes 1987)



End-periodic manifolds

(Taubes 1987)



End-periodic manifolds

(Taubes 1987)



⋮



End-periodic manifolds

(Taubes 1987)



⋮



$$\begin{array}{ccc}
 \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\
 \downarrow p & & \downarrow \exp \\
 Y & \xrightarrow{\bar{f}} & S^1
 \end{array}$$

End-periodic manifolds

(Taubes 1987)




...



$$\begin{array}{ccccc}
 \tilde{E} & \xrightarrow{\quad} & \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\
 \downarrow & & \downarrow p & & \downarrow \exp \\
 E_Y & \xrightarrow{\quad} & Y & \xrightarrow{\bar{f}} & S^1
 \end{array}$$

End-periodic manifolds

(Taubes 1987)



...



$$\begin{array}{ccccc}
 E|_{\text{end}(M)} & \xrightarrow{\quad} & \text{end}(M) & & \\
 \downarrow & & \downarrow & & \\
 \tilde{E} & \xrightarrow{\quad} & \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\
 \downarrow & & \downarrow p & & \downarrow \exp \\
 E_Y & \xrightarrow{\quad} & Y & \xrightarrow{\bar{f}} & S^1
 \end{array}$$

End-periodic Dirac operators

$$\begin{array}{ccccc} E|_{\text{end}(M)} & \xrightarrow{\quad} & \text{end}(M) & & \\ \downarrow & & \downarrow & & \\ \tilde{E} & \xrightarrow{\quad} & \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\ \downarrow & & \downarrow p & & \downarrow \exp \\ E_Y & \xrightarrow{\quad} & Y & \xrightarrow{\bar{f}} & S^1 \end{array}$$


End-periodic Dirac operators

$$\begin{array}{ccccc} E|_{\text{end}(M)} & \xrightarrow{\quad} & \text{end}(M) & & \\ \downarrow & & \downarrow & & \\ \tilde{E} & \xrightarrow{\quad} & \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\ \downarrow & & \downarrow p & & \downarrow \exp \\ E_Y & \xrightarrow{\quad} & Y & \xrightarrow{\bar{f}} & S^1 \\ & & \swarrow g_Y & & \end{array}$$

End-periodic Dirac operators

$$\begin{array}{ccccc}
 & & & & (p^* g_Y) \Big|_{\text{end}(M)} \\
 & & & \nearrow \text{dashed} & \\
 E|_{\text{end}(M)} & \xrightarrow{\text{blue}} & \text{end}(M) & & \\
 \downarrow \text{hook} & & \downarrow \text{hook} & & \\
 \tilde{E} & \xrightarrow{\text{blue}} & \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\
 \downarrow \text{green} & & \downarrow p & & \downarrow \text{exp} \\
 E_Y & \xrightarrow{\text{blue}} & Y & \xrightarrow{\bar{f}} & S^1 \\
 & & \nwarrow \text{dashed} & & \\
 & & g_Y & &
 \end{array}$$

End-periodic Dirac operators



End-periodic Dirac operators

$$\begin{array}{ccccc}
 & & & & (p^* g_Y) \Big|_{\text{end}(M)} \\
 & & & \nearrow & \\
 E|_{\text{end}(M)} & \xrightarrow{\quad} & \text{end}(M) & & \\
 \downarrow & & \downarrow & & \\
 \tilde{E} & \xrightarrow{\quad} & \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\
 \downarrow & & \downarrow p & & \downarrow \exp \\
 \tilde{D} = p^* D(Y) & & E_Y & \xrightarrow{\quad} & Y & \xrightarrow{\bar{f}} & S^1 \\
 \nwarrow & & \nwarrow g_Y & & & & \\
 D(Y) & & & & & &
 \end{array}$$


End-periodic Dirac operators

$$\begin{array}{ccccc}
 & (p^* D(Y))|_{\text{end}(M)} & & (p^* g_Y)|_{\text{end}(M)} & \\
 & \swarrow \text{dashed} & & \swarrow \text{dashed} & \\
 E|_{\text{end}(M)} & \xrightarrow{\text{blue}} & \text{end}(M) & & \\
 \downarrow \text{hook} & & \downarrow \text{hook} & & \\
 \tilde{E} & \xrightarrow{\text{blue}} & \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\
 \downarrow \text{green} & & \downarrow \text{green } p & & \downarrow \text{green exp} \\
 \tilde{D} = p^* D(Y) & E_Y \xrightarrow{\text{blue}} Y & \xrightarrow{\bar{f}} & S^1 & \\
 \swarrow \text{dashed } D(Y) & \swarrow \text{dashed } g_Y & & &
 \end{array}$$

End-periodic Dirac operators

$$\begin{array}{ccccc}
 (p^*D(Y))|_{\text{end}(M)} & & (p^*g_Y)|_{\text{end}(M)} & & \\
 \swarrow & & \swarrow & & \\
 E|_{\text{end}M} & \xrightarrow{\quad} & \text{end}(M) & & \\
 \downarrow & & \downarrow & & \\
 \tilde{E} & \xrightarrow{\quad} & \tilde{Y} & \xrightarrow{f} & \mathbb{R} \\
 \downarrow & & \downarrow p & & \downarrow \exp \\
 \tilde{D} = p^*D(Y) & \xrightarrow{E_Y} & Y & \xrightarrow{\bar{f}} & S^1 \\
 \swarrow & & \swarrow & & \\
 D(Y) & & g_Y & &
 \end{array}$$

End-periodic Dirac operators




D is an end periodic
Dirac operator on M , i.e.,

$$D|_Z = D(Z)$$

$$D|_{\text{end}(M)} = (p^*D(Y))|_{\text{end}(M)}$$

End-periodic Dirac operators



D is an end periodic
Dirac operator on M , i.e.,

$$D|_Z = D(Z)$$

$$D|_{\text{end}(M)} = (p^*D(Y))|_{\text{end}(M)}$$

Family context :

$$M \rightarrow B, Y \rightarrow B, \tilde{Y} \rightarrow B$$

$$\mathbb{E} = E \otimes \pi^* \Lambda T^* B, \text{ etc.}$$

Replace :

$$E_Y \rightsquigarrow \mathbb{E}, \tilde{E} \rightsquigarrow \tilde{\mathbb{E}}$$

$$E \rightsquigarrow \mathbb{E}, D \rightsquigarrow A, \text{ etc.}$$

Examples

Examples

- $g = dr^2 + h(r)^2 g_{\partial Z}$ on the cylinder.

Examples

- $g = dr^2 + h(r)^2 g_{\partial Z}$ on the cylinder.
- $g = u^{4/(n-2)}(dr^2 + g_{S^{n-1}})$ on $\mathbb{R} \times S^{n-1}$ (Mazzeo, Pollack, Uhlenbeck, 1994).

Examples

- $g = dr^2 + h(r)^2 g_{\partial Z}$ on the cylinder.
- $g = u^{4/(n-2)}(dr^2 + g_{S^{n-1}})$ on $\mathbb{R} \times S^{n-1}$ (Mazzeo, Pollack, Uhlenbeck, 1994).
- “Manifolds with periodic ends that are not products even topologically ... include manifolds whose ends arise from the infinite cyclic covers of 2-knot exteriors in the 4-sphere.” (Mrowka, Ruberman, Saveliev, 2014).

Examples

- $g = dr^2 + h(r)^2 g_{\partial Z}$ on the cylinder.
- $g = u^{4/(n-2)}(dr^2 + g_{S^{n-1}})$ on $\mathbb{R} \times S^{n-1}$ (Mazzeo, Pollack, Uhlenbeck, 1994).
- “Manifolds with periodic ends that are not products even topologically ... include manifolds whose ends arise from the infinite cyclic covers of 2-knot exteriors in the 4-sphere.” (Mrowka, Ruberman, Saveliev, 2014).
- *Gauge theory on asymptotically periodic 4-manifolds*, Taubes, 1987.

Examples

- $g = dr^2 + h(r)^2 g_{\partial Z}$ on the cylinder.
- $g = u^{4/(n-2)}(dr^2 + g_{S^{n-1}})$ on $\mathbb{R} \times S^{n-1}$ (Mazzeo, Pollack, Uhlenbeck, 1994).
- “Manifolds with periodic ends that are not products even topologically ... include manifolds whose ends arise from the infinite cyclic covers of 2-knot exteriors in the 4-sphere.” (Mrowka, Ruberman, Saveliev, 2014).
- *Gauge theory on asymptotically periodic 4-manifolds*, Taubes, 1987.
- *Seiberg-Witten equations, end-periodic Dirac operators, and a lift of Rohlin’s invariant*, Mrowka, Ruberman, Saveliev, 2011.

Examples

- $g = dr^2 + h(r)^2 g_{\partial Z}$ on the cylinder.
- $g = u^{4/(n-2)}(dr^2 + g_{S^{n-1}})$ on $\mathbb{R} \times S^{n-1}$ (Mazzeo, Pollack, Uhlenbeck, 1994).
- “Manifolds with periodic ends that are not products even topologically ... include manifolds whose ends arise from the infinite cyclic covers of 2-knot exteriors in the 4-sphere.” (Mrowka, Ruberman, Saveliev, 2014).
- *Gauge theory on asymptotically periodic 4-manifolds*, Taubes, 1987.
- *Seiberg-Witten equations, end-periodic Dirac operators, and a lift of Rohlin’s invariant*, Mrowka, Ruberman, Saveliev, 2011.
- *Smooth quantum gravity: exotic smoothness and quantum gravity*, Asselmeyer-Maluga, 2016.

Examples

- $g = dr^2 + h(r)^2 g_{\partial Z}$ on the cylinder.
- $g = u^{4/(n-2)}(dr^2 + g_{S^{n-1}})$ on $\mathbb{R} \times S^{n-1}$ (Mazzeo, Pollack, Uhlenbeck, 1994).
- “Manifolds with periodic ends that are not products even topologically ... include manifolds whose ends arise from the infinite cyclic covers of 2-knot exteriors in the 4-sphere.” (Mrowka, Ruberman, Saveliev, 2014).
- *Gauge theory on asymptotically periodic 4-manifolds*, Taubes, 1987.
- *Seiberg-Witten equations, end-periodic Dirac operators, and a lift of Rohlin’s invariant*, Mrowka, Ruberman, Saveliev, 2011.
- *Smooth quantum gravity: exotic smoothness and quantum gravity*, Asselmeyer-Maluga, 2016.
- *Metrics on end-periodic manifolds as models for dark matter*, Duston, 2021.

Index formula for end-periodic Dirac operators

Index formula for end-periodic Dirac operators

Mrowka, Ruberman, Saveliev, 2014: under the appropriate Fredholm condition, the index of the end-periodic Dirac operator D is given by

$$\operatorname{ind} D^+ = \int_Z AS(D(Z)) - \int_Y f AS(D(Y)) - \frac{1}{2} \eta_{\text{ep}}(D(Y))$$

where the end-periodic eta invariant is defined by

$$\widehat{\eta}_{\text{ep}}(D(Y)) = \frac{1}{\pi i} \int_0^\infty \oint_{S^1} \operatorname{Tr} \left(\operatorname{cl}(df) D_\xi^+ e^{-t D_\xi^- D_\xi^+} \right) \frac{d\xi}{\xi} dt$$

and $D_\xi = D(Y) - \ln(\xi) \operatorname{cl}(df)$, for $\xi \in S^1$, is the *indicial family* of D .

- 1 Index theory for Dirac operators
- 2 Index theory and anomalies
- 3 Index theory for end-periodic Dirac operators
- 4 A new index formula**
- 5 Remarks on the proof depending on how much time is left

Index for families of end-periodic Dirac operators

Index for families of end-periodic Dirac operators

Theorem (T.) *The Chern character of the index bundle, $\text{ch}(\text{Ind } D)$, is represented in de Rham cohomology by the differential form*

$$\int_{Z/B} AS(D(Z)) - \int_{W_0/B} f AS(D(Y)) - \frac{1}{2} \widehat{\eta}_{\text{ep}}(D(Y)) \in \Omega(B)$$

Index for families of end-periodic Dirac operators

Theorem (T.) *The Chern character of the index bundle, $\text{ch}(\text{Ind } D)$, is represented in de Rham cohomology by the differential form*

$$\int_{Z/B} AS(D(Z)) - \int_{W_0/B} f AS(D(Y)) - \frac{1}{2} \widehat{\eta}_{\text{ep}}(D(Y)) \in \Omega(B)$$

where the end-periodic eta form is given by

$$\begin{aligned} \widehat{\eta}_{\text{ep}}(D(Y)) = & \frac{-1}{\pi i} \int_0^\infty \oint_{|\xi|=1} t^{1/2} \text{Str} \left(\text{cl}(d_{Y/B} f) \dot{A}_t(\xi) e^{-A_t^2(\xi)} \right) \frac{d\xi}{\xi} dt \\ & - \frac{1}{\pi i} \int_0^\infty \oint_{|\xi|=1} \text{Str} \left(\frac{d}{dt} [A_t(\xi), \delta_t \text{cl}(df)] \mathcal{H}_t(\xi) \right) \frac{d\xi}{\xi} dt. \end{aligned}$$

where $\mathcal{H}_t(\xi) = \int_0^1 e^{-uA_t^2(\xi)} du$.

Index for families of end-periodic Dirac operators

Theorem (T.) *The Chern character of the index bundle, $\text{ch}(\text{Ind } D)$, is represented in de Rham cohomology by the differential form*

$$\int_{Z/B} AS(D(Z)) - \int_{W_0/B} f AS(D(Y)) - \frac{1}{2} \hat{\eta}_{\text{ep}}(D(Y)) \in \Omega(B)$$

where the end-periodic eta form is given by

$$\begin{aligned} \hat{\eta}_{\text{ep}}(D(Y)) = & \frac{-1}{\pi i} \int_0^\infty \oint_{|\xi|=1} t^{1/2} \text{Str} \left(\text{cl}(d_{Y/B} f) \dot{A}_t(\xi) e^{-A_t^2(\xi)} \right) \frac{d\xi}{\xi} dt \\ & - \frac{1}{\pi i} \int_0^\infty \oint_{|\xi|=1} \text{Str} \left(\frac{d}{dt} [A_t(\xi), \delta_t \text{cl}(df)] \mathcal{H}_t(\xi) \right) \frac{d\xi}{\xi} dt. \end{aligned}$$

where $\mathcal{H}_t(\xi) = \int_0^1 e^{-uA_t^2(\xi)} du$. Moreover, the degree zero part of the end-periodic eta form $\hat{\eta}_{\text{ep}}$ is the fiberwise end-periodic eta invariant: $(\hat{\eta}_{\text{ep}})_{[0]} \in C^\infty(B)$ given by $z \mapsto \eta_{\text{ep}}(D^z(Y_z))$.

Thank you

Extra slide

Some other important operators:

$$m_0 : T^*M \rightarrow \text{End } \mathbb{E}$$

$$\delta_t m_0(df) = t^{1/2} \text{cl}(d_{Y/B}f) + \pi^* d_B f$$

$$A_t(Y)(\xi) = A_t(Y) - \log(\xi) \delta_t m_0(df)$$

Extra slide

The Bismut superconnection on Y is

$$A(Y) = D(Y) + A_{[1]}(Y) + A_{[2]}(Y)$$

where

$$A_{[1]}(Y) = \sum_{\alpha} e^{\alpha} \wedge \left(\nabla_{e_{\alpha}}^{E_Y} + \frac{1}{2} k_Y(e_{\alpha}) \right)$$

$$A_{[2]}(Y) = -\frac{1}{4} \sum_{\alpha < \beta} \sum_j e^{\alpha} \wedge e^{\beta} \operatorname{cl}(e^j)(\Omega_Y(e_{\alpha}, e_{\beta}), e_j)$$

- 1 Index theory for Dirac operators
- 2 Index theory and anomalies
- 3 Index theory for end-periodic Dirac operators
- 4 A new index formula
- 5 Remarks on the proof depending on how much time is left

Heat equation proof of the local index theorem

Heat equation proof of the local index theorem

e^{-tD^2} = heat operator on $E \rightarrow M$

Heat equation proof of the local index theorem

e^{-tD^2} = heat operator on $E \rightarrow M$

$$\begin{cases} \left(\frac{\partial}{\partial t} + D^2 \right) e^{-tD^2} u(x) = 0, \\ \lim_{t \rightarrow 0^+} e^{-tD^2} u(x) = u(x). \end{cases}$$

Heat equation proof of the local index theorem

e^{-tD^2} = heat operator on $E \rightarrow M$

$$\begin{cases} \left(\frac{\partial}{\partial t} + D^2 \right) e^{-tD^2} u(x) = 0, \\ \lim_{t \rightarrow 0^+} e^{-tD^2} u(x) = u(x). \end{cases}$$

Consider the operator trace of the (chiral) heat operator

$$\mathrm{Tr}(e^{-tD^- D^+}) = \sum_j \langle e^{-tD^- D^+} \psi_j^+, \psi_j^+ \rangle$$

Heat equation proof of the local index theorem

e^{-tD^2} = heat operator on $E \rightarrow M$

$$\begin{cases} \left(\frac{\partial}{\partial t} + D^2 \right) e^{-tD^2} u(x) = 0, \\ \lim_{t \rightarrow 0^+} e^{-tD^2} u(x) = u(x). \end{cases}$$

Consider the operator trace of the (chiral) heat operator

$$\begin{aligned} \mathrm{Tr}(e^{-tD^- D^+}) &= \sum_j \langle e^{-tD^- D^+} \psi_j^+, \psi_j^+ \rangle \\ &= \int_M \mathrm{tr} (K_{e^{-tD^- D^+}}(x, x)) \, dx. \end{aligned}$$

Heat equation proof of the local index theorem

e^{-tD^2} = heat operator on $E \rightarrow M$

$$\begin{cases} \left(\frac{\partial}{\partial t} + D^2 \right) e^{-tD^2} u(x) = 0, \\ \lim_{t \rightarrow 0^+} e^{-tD^2} u(x) = u(x). \end{cases}$$

Consider the operator trace of the (chiral) heat operator

$$\begin{aligned} \mathrm{Tr}(e^{-tD^- D^+}) &= \sum_j \langle e^{-tD^- D^+} \psi_j^+, \psi_j^+ \rangle \\ &= \int_M \mathrm{tr} (K_{e^{-tD^- D^+}}(x, x)) \, dx. \end{aligned}$$

Define the supertrace of the heat operator

$$\mathrm{Str}(e^{-tD^2}) := \mathrm{Tr}(e^{-tD^- D^+}) - \mathrm{Tr}(e^{-tD^+ D^-}).$$

Heat equation proof of the local index theorem

Heat equation proof of the local index theorem

Transgression formula:

$$\frac{d}{dt} \text{Str}(e^{-tD^2}) = -\frac{1}{2} \text{Str}[D, D e^{-tD^2}] = 0.$$

Heat equation proof of the local index theorem

Transgression formula:

$$\frac{d}{dt} \text{Str}(e^{-tD^2}) = -\frac{1}{2} \text{Str}[D, De^{-tD^2}] = 0.$$

Integrate the transgression formula:

$$\text{ind } D^+ = \lim_{t \rightarrow \infty} \text{Str}(e^{-tD^2})$$

Heat equation proof of the local index theorem

Transgression formula:

$$\frac{d}{dt} \text{Str}(e^{-tD^2}) = -\frac{1}{2} \text{Str}[D, De^{-tD^2}] = 0.$$

Integrate the transgression formula:

$$\begin{aligned} \text{ind } D^+ &= \lim_{t \rightarrow \infty} \text{Str}(e^{-tD^2}) \\ &= \lim_{t \rightarrow 0^+} \text{Str}(e^{-tD^2}) \end{aligned}$$

Heat equation proof of the local index theorem

Transgression formula:

$$\frac{d}{dt} \text{Str}(e^{-tD^2}) = -\frac{1}{2} \text{Str}[D, De^{-tD^2}] = 0.$$

Integrate the transgression formula:

$$\begin{aligned} \text{ind } D^+ &= \lim_{t \rightarrow \infty} \text{Str}(e^{-tD^2}) \\ &= \lim_{t \rightarrow 0^+} \text{Str}(e^{-tD^2}) \\ &= \int_M AS(M). \end{aligned}$$

Renormalized trace defect

Renormalized trace defect

Cylindrical (Melrose)

$${}^b\mathrm{Tr}[P, Q] = \frac{-1}{2\pi i} \oint_{S^1} \mathrm{Tr} \left(\frac{\partial P_\xi}{\partial \xi} Q_\xi \right) d\xi.$$

Renormalized trace defect

Cylindrical (Melrose)

$${}^b\mathrm{Tr}[P, Q] = \frac{-1}{2\pi i} \oint_{S^1} \mathrm{Tr} \left(\frac{\partial P_\xi}{\partial \xi} Q_\xi \right) d\xi.$$

Periodic (MRS 2014)

$$\begin{aligned} {}^R\mathrm{Tr}[P, Q] &= \frac{-1}{2\pi i} \oint_{S^1} \mathrm{Tr} \left(\frac{\partial P_\xi}{\partial \xi} Q_\xi \right) d\xi \\ &\quad + \frac{1}{2\pi i} \oint_{S^1} \int_W f(x) \mathrm{tr} (K_{P_\xi Q_\xi}(x, x) - K_{Q_\xi P_\xi}(x, x)) dx \frac{d\xi}{\xi}. \end{aligned}$$

Renormalized trace defect

Cylindrical (Melrose)

$${}^b\mathrm{Tr}[P, Q] = \frac{-1}{2\pi i} \oint_{S^1} \mathrm{Tr} \left(\frac{\partial P_\xi}{\partial \xi} Q_\xi \right) d\xi.$$

Periodic (MRS 2014)

$$\begin{aligned} {}^R\mathrm{Tr}[P, Q] &= \frac{-1}{2\pi i} \oint_{S^1} \mathrm{Tr} \left(\frac{\partial P_\xi}{\partial \xi} Q_\xi \right) d\xi \\ &\quad + \frac{1}{2\pi i} \oint_{S^1} \int_W f(x) \mathrm{tr} (K_{P_\xi Q_\xi}(x, x) - K_{Q_\xi P_\xi}(x, x)) dx \frac{d\xi}{\xi}. \end{aligned}$$

Periodic + family + \mathbb{Z}_2 -graded (T. 2025)

$$\begin{aligned} {}^R\mathrm{Str}[P, Q] &= \frac{-1}{2\pi i} \oint_{S^1} \mathrm{Str} \left(\frac{\partial P_\xi}{\partial \xi} Q_\xi \right) d\xi \\ &\quad + \frac{1}{2\pi i} \oint_{S^1} \int_{W/B} f(x) \mathrm{str} (K_{[P_\xi, Q_\xi]}(x, x)) dx \frac{d\xi}{\xi}. \end{aligned}$$

EP & cylindrical family transgression formulas

EP & cylindrical family transgression formulas

End-periodic

Cylindrical

EP & cylindrical family transgression formulas

End-periodic

$$\begin{aligned}\frac{d}{dt} {}^R\text{ch}(A_t) &= - {}^R\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &\quad + \alpha_1(t) + \alpha_2(t)\end{aligned}$$

Cylindrical

$$\frac{d}{dt} {}^b\text{ch}(A_t) = - {}^b\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right]$$

EP & cylindrical family transgression formulas

End-periodic

$$\begin{aligned}\frac{d}{dt} {}^R\text{ch}(A_t) &= - {}^R\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &\quad + \alpha_1(t) + \alpha_2(t)\end{aligned}$$

Cylindrical

$$\begin{aligned}\frac{d}{dt} {}^b\text{ch}(A_t) &= - {}^b\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &= - {}^b\text{Str} \left[A_t - A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &\quad - {}^b\text{Str} \left[A_{[1]}, \dot{A}_t e^{-A_t^2} \right]\end{aligned}$$

EP & cylindrical family transgression formulas

End-periodic

$$\begin{aligned}\frac{d}{dt} {}^R\text{ch}(A_t) &= - {}^R\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &\quad + \alpha_1(t) + \alpha_2(t)\end{aligned}$$

Cylindrical

$$\begin{aligned}\frac{d}{dt} {}^b\text{ch}(A_t) &= - {}^b\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &= - {}^b\text{Str} \left[A_t - A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &\quad - {}^b\text{Str} \left[A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &= -\frac{1}{2} \hat{\eta}(t) - (\text{exact})\end{aligned}$$

EP & cylindrical family transgression formulas

End-periodic

$$\begin{aligned}\frac{d}{dt} {}^R\text{ch}(A_t) &= - {}^R\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &\quad + \alpha_1(t) + \alpha_2(t) \\ &= \beta_1(t) + \beta_2(t) + \gamma_1(t) + \gamma_2(t) \\ &\quad + \alpha_1(t) + \alpha_2(t)\end{aligned}$$

Cylindrical

$$\begin{aligned}\frac{d}{dt} {}^b\text{ch}(A_t) &= - {}^b\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &= - {}^b\text{Str} \left[A_t - A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &\quad - {}^b\text{Str} \left[A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &= -\frac{1}{2} \widehat{\eta}(t) - (\text{exact})\end{aligned}$$

EP & cylindrical family transgression formulas

End-periodic

$$\begin{aligned}\frac{d}{dt} {}^R\text{ch}(A_t) &= - {}^R\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &\quad + \alpha_1(t) + \alpha_2(t) \\ &= \beta_1(t) + \beta_2(t) + \gamma_1(t) + \gamma_2(t) \\ &\quad + \alpha_1(t) + \alpha_2(t) \\ &= -\frac{1}{2} \widehat{\eta}_{\text{ep}}(t)\end{aligned}$$

$$-\frac{d}{dt} \oint_{S^1} \int_{W/B} f(x) \text{str}(K_{e^{-A_t^2(\xi)}}(x, x)) dx \frac{d\xi}{\xi} + (\text{exact})$$

Cylindrical

$$\begin{aligned}\frac{d}{dt} {}^b\text{ch}(A_t) &= - {}^b\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &= - {}^b\text{Str} \left[A_t - A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &\quad - {}^b\text{Str} \left[A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &= -\frac{1}{2} \widehat{\eta}(t) - (\text{exact})\end{aligned}$$

EP & cylindrical family transgression formulas

End-periodic

$$\begin{aligned} \frac{d}{dt} {}^R\text{ch}(A_t) &= - {}^R\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &\quad + \alpha_1(t) + \alpha_2(t) \\ &= \beta_1(t) + \beta_2(t) + \gamma_1(t) + \gamma_2(t) \\ &\quad + \alpha_1(t) + \alpha_2(t) \\ &= -\frac{1}{2} \widehat{\eta}_{\text{ep}}(t) \end{aligned}$$

$$-\frac{d}{dt} \oint_{S^1} \int_{W/B} f(x) \text{str}(K_{e^{-A_t^2(\xi)}}(x, x)) dx \frac{d\xi}{\xi} + (\text{exact})$$

$$(*) \quad -\frac{1}{2} \widehat{\eta}_{\text{ep}}(t) = \alpha_1(t) + \beta_1(t)$$

$$= \alpha_1(t) - \frac{1}{2\pi i} \oint_{|\xi|=1} t^{1/2} \text{Str} \left(\text{cl}(d_{Y/B} f) \dot{A}_t(Y)(\xi) e^{-A_t^2(Y)(\xi)} \right) \frac{d\xi}{\xi}$$

Cylindrical

$$\begin{aligned} \frac{d}{dt} {}^b\text{ch}(A_t) &= - {}^b\text{Str} \left[A_t, \dot{A}_t e^{-A_t^2} \right] \\ &= - {}^b\text{Str} \left[A_t - A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &\quad - {}^b\text{Str} \left[A_{[1]}, \dot{A}_t e^{-A_t^2} \right] \\ &= -\frac{1}{2} \widehat{\eta}(t) - (\text{exact}) \end{aligned}$$

EP & cylindrical family transgression formulas

EP & cylindrical family transgression formulas

Melrose-Piazza (1997) first step: $\frac{d}{dt} {}^b\text{ch}(A_t) = - {}^b\text{Str}[A_t, \dot{A}_t e^{-A_t^2}]$.

EP & cylindrical family transgression formulas

Melrose-Piazza (1997) first step: $\frac{d}{dt} {}^b\text{ch}(A_t) = - {}^b\text{Str}[A_t, \dot{A}_t e^{-A_t^2}]$.

Proof. By Duhamel's formula, the derivative of the heat kernel is

$$(13.10) \quad \frac{d}{dt} e^{-A_t^2} = - \int_0^1 e^{-sA_t^2} \cdot \frac{dA_t^2}{dt} \cdot e^{-(1-s)A_t^2} ds.$$

bStr defect formula

A (The indicial family of A_t^2 is even in λ , so applying (12.5) to commute the leading term, from left to right, no boundary terms arise. Thus

$$(13.11) \quad \frac{d}{dt} {}^b\text{Ch}(A_t) = - {}^b\text{STr} \left(\frac{dA_t^2}{dt} e^{-A_t^2} \right).$$

Since A_t is odd and commutes with $\exp(-A_t^2)$, this can be written as a supercommutator:

$$(13.12) \quad \frac{d}{dt} {}^b\text{Ch}(A_t) = - {}^b\text{STr}[A_t, \frac{dA_t}{dt} e^{-A_t^2}].$$

Now, dA_t/dt is a fibre operator so (12.5) can be applied to give

Complication in the end-periodic case

Complication in the end-periodic case

Take $P = e^{-sA_t^2}$ and $Q = \partial_t(A_t^2)e^{-(1-s)A_t^2}$ in the supertrace defect formula:

$$\int_0^1 {}^R\text{Str}[e^{-sA_t^2}, \partial_t(A_t^2)e^{-(1-s)A_t^2}] ds = \alpha_1(t) + \alpha_2(t).$$

Complication in the end-periodic case

Take $P = e^{-sA_t^2}$ and $Q = \partial_t(A_t^2)e^{-(1-s)A_t^2}$ in the supertrace defect formula:

$$\int_0^1 {}^R\text{Str}[e^{-sA_t^2}, \partial_t(A_t^2)e^{-(1-s)A_t^2}] ds = \alpha_1(t) + \alpha_2(t).$$

The term $\alpha_2(t)$ is OK. More difficult is that $\alpha_1(t) \neq 0$ in the end-periodic case, because the parity argument made by M.-P. breaks down.

Complication in the end-periodic case

Take $P = e^{-sA_t^2}$ and $Q = \partial_t(A_t^2)e^{-(1-s)A_t^2}$ in the supertrace defect formula:

$$\int_0^1 {}^R\text{Str}[e^{-sA_t^2}, \partial_t(A_t^2)e^{-(1-s)A_t^2}] ds = \alpha_1(t) + \alpha_2(t).$$

The term $\alpha_2(t)$ is OK. More difficult is that $\alpha_1(t) \neq 0$ in the end-periodic case, because the parity argument made by M.-P. breaks down.

Recall that we have, by the supertrace defect formula,

$$\alpha_1(t) = \frac{-1}{2\pi i} \int_0^1 \oint_{S^1} \text{Str} \left(\partial_\xi(e^{-sA_t^2(\xi)}) \cdot \partial_t(A_t^2(\xi)) \cdot e^{-(1-s)sA_t^2(\xi)} \right) d\xi ds$$

Complication in the end-periodic case

Take $P = e^{-sA_t^2}$ and $Q = \partial_t(A_t^2)e^{-(1-s)A_t^2}$ in the supertrace defect formula:

$$\int_0^1 {}^R\text{Str}[e^{-sA_t^2}, \partial_t(A_t^2)e^{-(1-s)A_t^2}] ds = \alpha_1(t) + \alpha_2(t).$$

The term $\alpha_2(t)$ is OK. More difficult is that $\alpha_1(t) \neq 0$ in the end-periodic case, because the parity argument made by M.-P. breaks down.

Recall that we have, by the supertrace defect formula,

$$\begin{aligned}\alpha_1(t) &= \frac{-1}{2\pi i} \int_0^1 \oint_{S^1} \text{Str} \left(\partial_\xi(e^{-sA_t^2(\xi)}) \cdot \partial_t(A_t^2(\xi)) \cdot e^{-(1-s)sA_t^2(\xi)} \right) d\xi ds \\ &\stackrel{?}{=} \frac{-1}{2\pi i} \int_0^1 \oint_{S^1} \text{Str}(\text{odd} \cdot \text{even} \cdot \text{even}) \frac{d\xi}{\xi} ds \\ &= 0\end{aligned}$$

Complication in the end-periodic case

End-periodic

$$A_t(\xi) = A_t(Y) - \ln(\xi)\delta_t m_0(df)$$

Cylindrical

$$A_t(\xi) = A_t(Y) - \ln(\xi)\delta_t m_0(dr)$$

Complication in the end-periodic case

End-periodic

$$A_t(\xi) = A_t(Y) - \ln(\xi)\delta_t m_0(df)$$

$$A_t^2(\xi) = A_t^2(Y) - \ln(\xi)\delta_t[A(Y), m_0(df)]$$

$$- t \ln(\xi)^2 |d_{Y/B} f|^2$$

Cylindrical

$$A_t(\xi) = A_t(Y) - \ln(\xi)\delta_t m_0(dr)$$

$$A_t^2(\xi) = A_t^2(Y) - \ln(\xi)\delta_t[A(Y), m_0(dr)]$$

$$- t \ln(\xi)^2 |d_{Y/B} r|^2$$

Complication in the end-periodic case

End-periodic

$$A_t(\xi) = A_t(Y) - \ln(\xi)\delta_t m_0(df)$$

$$A_t^2(\xi) = A_t^2(Y) - \ln(\xi)\delta_t[A(Y), m_0(df)]$$

$$- t \ln(\xi)^2 |d_{Y/B} f|^2$$

Cylindrical

$$A_t(\xi) = A_t(Y) - \ln(\xi)\delta_t m_0(dr)$$

$$A_t^2(\xi) = A_t^2(Y) - \ln(\xi)\delta_t[A(Y), m_0(dr)]$$

$$- t \ln(\xi)^2 |d_{Y/B} r|^2$$

Proposition (T., 2025)

$$\begin{aligned} [A(Y), m_0(df)] &= m_0 \left(\nabla^{T^*Y} df \right) - 2 \nabla_{(d_{Y/B} f)^\#}^{\mathbb{E}_{Y,0}} \\ &= 0 \quad \text{iff the periodic end is cylindrical} \end{aligned}$$

Complication in the end-periodic case

End-periodic

$$A_t(\xi) = A_t(Y) - \ln(\xi) \delta_t m_0(df)$$

$$A_t^2(\xi) = A_t^2(Y) - \ln(\xi) \delta_t [A(Y), m_0(df)]$$

$$- t \ln(\xi)^2 |d_{Y/B} f|^2$$

Cylindrical

$$A_t(\xi) = A_t(Y) - \ln(\xi) \delta_t m_0(dr)$$

$$A_t^2(\xi) = A_t^2(Y) - \ln(\xi) \delta_t [A(Y), m_0(dr)]$$

$$- t \ln(\xi)^2 |d_{Y/B} r|^2$$

$$= A_t^2(Y) - t \ln(\xi)^2$$

Proposition (T., 2025)

$$\begin{aligned} [A(Y), m_0(df)] &= m_0 \left(\nabla^{T^*Y} df \right) - 2 \nabla_{(d_{Y/B} f)^\#}^{\mathbb{E}_Y, 0} \\ &= 0 \quad \text{iff the periodic end is cylindrical} \end{aligned}$$

Complication in the end-periodic case

End-periodic

$$A_t(\xi) = A_t(Y) - \ln(\xi)\delta_t m_0(df)$$

$$A_t^2(\xi) = A_t^2(Y) - \ln(\xi)\delta_t[A(Y), m_0(df)]$$

$$- t \ln(\xi)^2 |d_{Y/B} f|^2$$

$$e^{-A_t^2(\xi)} = ???$$

Cylindrical

$$A_t(\xi) = A_t(Y) - \ln(\xi)\delta_t m_0(dr)$$

$$A_t^2(\xi) = A_t^2(Y) - \ln(\xi)\delta_t[A(Y), m_0(dr)]$$

$$- t \ln(\xi)^2 |d_{Y/B} r|^2$$

$$= A_t^2(Y) - t \ln(\xi)^2$$

$$e^{-A_t^2(\xi)} = e^{-A_t^2(Y)} e^{-t \ln(\xi)^2}$$

Proposition (T., 2025)

$$\begin{aligned} [A(Y), m_0(df)] &= m_0 \left(\nabla^{T^* Y} df \right) - 2 \nabla_{(d_{Y/B} f)^\#}^{\mathbb{E}_Y, 0} \\ &= 0 \quad \text{iff the periodic end is cylindrical} \end{aligned}$$