

The roles of TGF β in the tumour microenvironment

Michael Pickup, Sergey Novitskiy and Harold L. Moses

Abstract | The influence of the microenvironment on tumour progression is becoming clearer. In this Review we address the role of an essential signalling pathway, that of transforming growth factor- β , in the regulation of components of the tumour microenvironment and how this contributes to tumour progression.

Epithelial-to-mesenchymal transition

The process by which epithelial cells lose their epithelial characteristics, such as cellular polarity and cell–cell junctions, in favour of mesenchymal characteristics, which include high cellular motility.

The transforming growth factor- β (TGF β) pathway has been established as essential for cancer progression, because of its prominent role in the regulation of cell growth, differentiation and migration. Through the canonical and non-canonical arms of the signalling pathway, TGF β instigates cellular phenotypical changes that mediate its role as both a tumour suppressor and a tumour promoter. Indeed, the first described phenotypical effect of TGF β signalling was the induction of a cellular cytostatic programme¹ and this provided the first evidence of the TGF β pathway being tumour suppressive². However, there was also evidence to the contrary, such as increased tumour progression in carcinomas that overexpressed the TGF β 1 ligand^{3,4}. The initial *in vitro* evidence for the pro-tumorigenic effects of TGF β consisted of the induction of a mesenchymal phenotype in epithelial tumour cells (commonly known as an epithelial-to-mesenchymal transition (EMT)) after prolonged exposure to TGF β ^{5,6}.

These early studies into the functional outcome of active TGF β signalling underlie the difficulties in implementing clinically efficacious treatment regimens that target the TGF β pathway. The contextual cues that drive the tumour suppressor and tumour promoter roles of TGF β , as well as the switch between these two phenotypes, are not fully understood. As the understanding of tumour progression has increased, the importance of the tumour microenvironment has been clearly shown. It is interesting to note that TGF β signalling, both endogenously in human disease and in genetically engineered mouse models of cancer, is associated with the characteristic epithelial changes outlined above, as well as with substantial changes in the tumour stroma^{7–9}; for example, TGF β 1 expression in invasive breast cancer correlates with markers of tumour progression, such as metastasis, extracellular matrix (ECM) deposition and the infiltration of immune cells⁷. Such findings have laid the

groundwork for recent studies that have established TGF β signalling as an important mediator not only of changes to the epithelial phenotype but also of changes in the stromal environment that are essential for tumour progression. In this Review, we address the functional role of TGF β signalling in modulating the tumour microenvironment and how these changes affect tumour progression.

The basics of TGF β signalling

TGF β 1, TGF β 2 and TGF β 3 ligands function as the primary mediators of TGF β signalling^{1,10,11} and are secreted as inactive homodimeric polypeptides that can bind to latent TGF β -binding proteins, which promote extracellular sequestration¹². On activation, the ligands bind to the type 2 TGF β receptor (TGFB β R2), which causes recruitment and phosphorylation of TGFB β R1 (also known as ALK5), resulting in downstream signalling activation¹³ (FIG. 1). The strength of this signal depends on which ligand is bound, as the ligands vary in their binding affinity for TGFB β R1. This variation in ligand binding promotes differential ligand presentation to TGFB β R2 (REF. 14). TGFB β R3 can augment the initialization of the signalling cascade by promoting differential ligand binding¹⁵. The final heterotetrameric form of the active receptors initiates downstream signalling through either SMAD-mediated canonical signalling or SMAD-independent non-canonical signalling¹³. The canonical signalling pathway involves phosphorylation of the carboxy-terminal serine residue of the internal modulator SMAD proteins, SMAD2 or SMAD3, by the activated receptors^{16,17}. This phosphorylation induces oligomerization of SMAD2 or SMAD3 with SMAD4, which is necessary for nuclear translocation¹⁸. Through interactions with a variety of transcription cofactors, the nuclear-localized SMAD complex initiates transcriptional activation or transcriptional repression of several genes. The non-canonical branch of the signalling

Vanderbilt University Medical Center, Vanderbilt-Ingram Comprehensive Cancer Center, Medicine and Pathology, Cancer Biology, 2220 Pierce Avenue, 691 Preston Research Building, Nashville, Tennessee 37232, USA. Correspondence to H.L.M. e-mail: hal.moses@vanderbilt.edu
doi:10.1038/nrc3603
Published online
17 October 2013

Key points

- During the early stages of tumorigenesis, transforming growth factor- β (TGF β) functions as a tumour suppressor. However, as tumours progress, tumour cells may lose their growth-inhibitory response to TGF β and may instead respond by initiating epithelial-to-mesenchymal transition and by increasing cell migration.
- Experimental data support the idea that both loss of and gain of TGF β signalling is pro-tumorigenic, as mouse models show that the overexpression of TGF β and the abrogation of signalling results in increased tumour cell metastasis. It is beginning to be appreciated that the effects of TGF β signalling in the tumour epithelium extend beyond tumour cell-autonomous mechanisms into the tumour microenvironment.
- Nearly every cell in the tumour microenvironment responds to TGF β in a unique way, and these diverse biological responses have a variety of effects on tumour progression.
- Given the pleiotropic nature of TGF β signalling, the numerous downstream signalling events of TGF β offer new and intriguing targets for therapeutic intervention.
- Successful therapeutic targeting of TGF β itself remains a highly desirable goal and the considerable advances in understanding TGF β signalling, not only in the tumour epithelium but also in the tumour microenvironment, bring the realization of this goal closer.

pathway¹⁹ involves activation of the PI3K–AKT, RHOA and MAPK pathways, among others, by the activated heterotetrameric receptors²⁰.

The outcome of these signalling pathways in epithelial cells is either suppression of cell proliferation or induction of cellular migration and invasion. Studies investigating the cytostatic phenotype induced by TGF β have established numerous pathways through which this is achieved, including repression of *MYC* and cyclin-dependent kinase 4 (*CDK4*), as well as the induced expression of CDK inhibitors p21 (also known as CIP1; encoded by *CDKN1A*) and INK4B (also known as p15)^{21–23}. Furthermore, the SMAD-dependent activation of Krueppel-like factor 10 (also known as TIEG1), death-associated protein kinase 1 (DAPK1) and the proapoptotic protein BCL-2-interacting mediator of cell death (BIM; also known as BCL2L11), among others, triggers programmed cell death²⁴. Studies elucidating the tumour suppressor role of TGF β corroborated evidence that the loss of TGF β -signalling components was associated with carcinoma progression^{25,26}. The signalling pathway downstream of EMT induction by TGF β has also been partly mapped. Inhibitor of DNA binding 1 (ID1; a transcriptional regulator) is inhibited by TGF β . This results in decreased expression of E-cadherin (also known as cadherin 1) and zona occludens 1 (ZO1; also known as TJP1), which are two factors known to help maintain an epithelial phenotype. TGF β signalling also induces the expression of EMT-associated transcription factors, such as snail family zinc finger 1 (*SNAI1*), *SNAI2*, zinc finger E-box binding homeobox 1 (*ZEB1*), *ZEB2*, and lymphoid enhancer-binding factor 1 (*LEF1*), which help to promote the loss of cellular adhesions and cytoskeletal rearrangement²⁷. Non-canonical signalling pathways activated by TGF β , particularly the RHO–ROCK1 (RHO-associated protein kinase 1) and the AKT pathways, were also shown to be essential in the promotion of the migratory and invasive cellular phenotypes observed after treatment of epithelial cells with TGF β ²⁸. RHO–ROCK1 signalling that is induced by TGF β is particularly important in single-cell migration.

Cohesive migratory phenotype

Contrary to single-cell invasion, this is a form of invasion by which cells maintain their cell–cell contacts and by which cells migrate as a unit.

MicroRNA

(miRNA). 20–30 nucleotide long non-coding RNA that modulates gene expression by binding to complementary sequences in the 3' untranslated region of target genes, which induces repression of gene translation or mRNA degradation.

Interestingly, abrogation of TGF β signalling did not completely stop epithelial cell migration, but rather it switched the cells towards a cohesive migratory phenotype²⁹. Thus, the dichotomous effects of TGF β signalling are associated with tumour suppression early in tumour development, through initiation of growth arrest, and with tumour promotion in late-stage tumours, through the induction of EMT and of cellular migration and invasion.

However, the effects of epithelial TGF β signalling on cells that constitute the tumour microenvironment have more recently aided our understanding of the diverse effects that TGF β has on tumour development and progression. These effects have also started to put in perspective why many studies cite increased metastasis and poor patient prognosis in cancers that lack intact TGF β signalling.

Disruption of TGF β signalling

Even without the introduction of oncogenic changes, altered TGF β -signalling in normal epithelial cells results in phenotypical changes in the stroma; for example, expression of constitutively active TGFBR1 in the mammary epithelium results in mammary glands with greater collagen deposition around the ductal epithelium³⁰. Notably, mammary and pancreatic carcinoma cells that harbour activated TGFBR1 induce a significant increase in angiogenesis following implantation into mice^{31,32} (FIG. 2). Active TGF β signalling in mammary tumour cells also promotes tumour progression through the SMAD-independent induction of the expression of matrix metalloproteinases (MMPs), which results in increased angiogenesis and in tumour cell invasion³¹. One of the more interesting effects of disrupted TGF β signalling in the malignant epithelium on the tumour microenvironment is related to the alteration of microRNA (miRNA) regulation. miRNAs, which are potent modulators of mRNA, are aberrantly expressed in numerous cancer types and are linked with several pro-tumorigenic and antitumorigenic functions³³. TGF β signalling can inhibit the function of specific miRNAs; for example, TGF β expression in hepatocellular carcinoma (HCC) cells induces the expression of CC-chemokine ligand 22 (CCL22) through the inhibition of miR-34a expression³⁴. This promotes the recruitment of regulatory T (T_{Reg}) cells. TGF β can also have direct effects on cells that form part of the tumour stroma; for example, in human vascular smooth muscle cells (VSMCs), TGF β can promote the cleavage and the maturation of pri-mir-21 sequences by inducing the formation of a complex that involves SMAD3 and ATP-dependent RNA helicase DDX5, which is a component of the DROSHA microprocessor complex³⁵. This increases the expression of miR-21, which alters the contractile properties of the VSMCs. TGF β signalling also increases the expression of miR-29a in human and mouse endothelial cells, which thereby increases the induction of angiogenesis in a non-tumour model³⁶. miR-29a targets phosphatase and tensin homolog (*PTEN*) mRNA, and it thereby increases the activity of the AKT signalling pathway. TGF β can also increase the expression of miR-494 in




Figure 1 | Epithelial TGFβ signalling during tumour progression. **a** | Normal transforming growth factor-β (TGFβ) signalling in TGFβ-responsive cells (blue cells) feeds through the type 2 TGFβ receptor (TGFBR2) to activate downstream signalling targets. Canonical signalling is activated through phosphorylation of TGFBR1 to induce nuclear localization and transcriptional activity of SMADs. Non-canonical signalling can occur independently of SMAD proteins and includes activation of RHOA, AKT and MAPK pathways. Early in tumorigenesis, TGFβ functions as a tumour suppressor, partly through the SMAD-dependent induction of cell cycle arrest. **b** | One hypothesis is that selective pressure leads to the expansion of the population of tumour cells that harbour inactivating mutations in the TGFβ pathway, thus allowing them to overcome the growth-inhibitory effects of active TGFβ signalling. As outlined by Levy *et al.*⁴⁸, loss of TGFβ responsiveness (green cells) can occur through loss-of-function mutations, loss of expression or promoter methylation of genes (shown by the grey circles) that encode TGFβ receptors or SMADs. JNK, JUN N-terminal kinase; mTOR, mammalian target of rapamycin; PAR6, partitioning defective 6 homologue; ROCK1, RHO-associated protein kinase 1; RSMAD, receptor SMAD; SBE, SMAD-binding element.

Myeloid-derived suppressor cells

(MDSCs). A heterogeneous population of cells that are defined by their myeloid origin, immature state and ability to potently suppress T cell responses.

Chemotaxis

The function of a cell responding to a chemokine gradient to promote directional migration towards sites of tissue injury or into tumour microenvironments.

myeloid-derived suppressor cells (MDSCs). This increases the CXC-chemokine receptor (CXCR4)-mediated chemotaxis of MDSCs to the tumour and it reduces the expression of PTEN. The reduced expression of PTEN results in increased AKT activity and in increased expression of pro-tumorigenic liver arginase 1 (ARG1), MMP2, MMP13 and MMP14, which contribute to increased tumour cell invasion and metastasis³⁷.

Induction of TGFβ signalling through genetic alterations or through the treatment of carcinoma cells with TGFβ has identified numerous gene targets, many of which are conserved in various cancer types^{38–41}. These experiments have begun to address the functional importance of gene expression induced by TGFβ by correlating these changes with patient data that predict poor

patient outcome^{38,42}. Importantly, many of these gene expression targets have well-characterized functions in the modification of tumour microenvironments and they link with phenotypical changes seen on alteration of TGFβ signalling in mouse models of cancer (FIG. 2). TGFβ signalling in epithelial cells induces the expression of numerous ECM genes, including collagen type 1 α1 (COL1A1) and COL4A1, as well as matrix modifying enzymes MMP2, MMP9 and lysyl oxidase homologue 4 (LOXL4). These same studies also provide supporting evidence for the observed pro-angiogenic phenotypes that are induced by TGFβ activation in epithelial cells. Gene expression analysis showed a significant increase in the expression levels of vascular endothelial growth factor A (VEGFA) and thrombospondin 1 (REFS 38–41). Given the




Figure 2 | TGFβ signalling in tumour cells determines microenvironmental modification. Transforming growth factor-β (TGFβ) signalling in tumour cells induces the expression of numerous mediators of extracellular change. Tumours in which cells show increased TGFβ activity are characterized by increased extracellular matrix (ECM) deposition; they show increased secretion of matrix proteins and maturation of these proteins through ECM-modifying enzymes such as lysyl oxidase homologue 4 (LOXL4). In addition, TGFβ signalling in tumour cells drives the induction of endothelial cell recruitment and proliferation, which promote increased angiogenesis. Conversely, TGFβ suppresses the expression of numerous cytokines and chemokines such as the CXC-chemokine ligand 1 (CXCL1) and CXCL5. Loss of TGFβ responsiveness relieves this suppression and results in enhanced immune cell infiltration. These microenvironmental changes promote epithelial cell and stromal cell phenotypal responses, which substantially affect tumour progression. The figure shows the phenotypal changes that specifically result from the epithelial-derived factors listed. ADAMTS9, a disintegrin and metalloproteinase with thrombospondin motifs 9; CTGF, connective tissue growth factor; EMT, epithelial-to-mesenchymal transition; G-CSF, granulocyte colony-stimulating factor; HGF, hepatocyte growth factor; IL-8, interleukin-8; MMP, matrix metalloproteinase; PDGFβ, platelet-derived growth factor-β; THBS1, thrombospondin 1; VEGFA, vascular endothelial growth factor A.

EMT and migratory phenotypal changes induced following activation of TGFβ signalling, as well as the epithelial cell TGFβ gene signature associated with tumour recurrence, the identification of stromal changes confirms epithelial cell TGFβ signalling as a pro-tumorigenic signalling pathway. However, such investigations do not distinguish the epithelial and stromal effects of epithelial cell TGFβ signalling on the promotion of tumour cell metastasis, and thus they do not address the specific contribution of each effect to the correlation of TGFβ signalling with poor patient prognosis.

As tumours progress, the growth-inhibitory effects of TGFβ are overcome by the loss of TGFβ pathway elements or of downstream signalling targets^{10,43} (FIG. 1). Loss of TGFβ responsiveness in tumour cells has substantial effects on tumour progression, not only through altered epithelial cell characteristics but also through gene expression changes that affect the tumour microenvironment. Activated TGFβ signalling has primarily been associated with an increase in metastasis and with poor patient prognosis, because EMT is induced^{29,44}. However, abrogation of TGFβ signalling in carcinoma cells can also result in increased metastasis⁴⁵. Loss of TGFβ signalling components in both mouse models of cancer and in human cancer has been associated with poor prognosis as a result of increased progression and metastasis^{25,46–49}. Similarly to receptor activation in the normal mammary epithelium, the pancreatic epithelium expressing

a dominant-negative TGFBR2 results in increased desmoplasia and angiogenesis in adult mice²⁵. The abrogation or attenuation of TGFβ signalling, either at the receptor or at the SMAD level in mouse models of cancer, has various expected epithelial effects, such as loss of growth inhibition, but it also induces several stromal changes, including activation of stromal fibroblasts, deposition of collagenous ECM, infiltration of a variety of immune cells and increased angiogenesis⁵⁰. Recent work has shown that the infiltration of MDSCs is increased in tumours in which epithelial TGFβ signalling is abrogated⁵¹ (FIG. 2). The increased recruitment of MDSCs is primarily associated with the increased expression of the CXC-chemokine ligand 1 (CXCL1) and CXCL5, the expression of which is normally inhibited by TGFβ. Although this altered chemokine expression pattern is associated with increased myeloid cell infiltration into the tumour microenvironment, these chemokines can also drive the activation of stromal fibroblasts through the induction of connective tissue growth factor (CTGF) expression⁵². Perhaps the most interesting induced change in gene expression to result from abrogation of TGFβ signalling in epithelial cells is the change in the expression of the TGFβ ligand itself^{53,54}. Given that the epithelium expressing TGFβ can no longer respond to it, any effects derived from its expression would either be on a separate population of epithelial cells that retain their ability to respond to TGFβ or on cells found in the stromal microenvironment.

Dominant-negative

A defective protein product that effectively inhibits the function of its wild-type counterpart by retaining essential interaction capabilities without having a desired effector function.

Desmoplasia

The accumulation of extracellular matrix proteins, usually through the enhanced activation of stromal fibroblasts.

The epithelial loss of TGFBR2 results in increased expression of CXCL1, CXCL5 and bone marrow stromal antigen 2 (BST2), and in a reduction in the expression of genes such as CXCL12, platelet-derived growth factor- β (PDGFB) and CTGF⁵⁵ (FIG. 2). These gene expression changes significantly correlate with a worse outcome in patients with lymph node-positive, oestrogen receptor (ER)-positive luminal A-type breast cancer. These findings introduce an interesting concept that explains many of the counter-intuitive findings regarding impairment of epithelial cell TGF β signalling that ultimately promotes metastasis. If, as many studies have shown, the stromal microenvironment is a prognostic factor for metastasis, it could override the previously established cell-autonomous signalling events that are associated with tumour cell migration, invasion and metastasis⁵⁶. Thus, even though tumour cells that lack TGF β responsiveness no longer gain any epithelial-centric metastatic advantages from TGF β production, stromal alterations as a result of altered TGF β expression might continue to promote metastasis. Tumours are heterogeneous and therefore probably contain cells with and without the ability to respond to TGF β — a concept that is yet to be fully appreciated in terms of its effects on tumour progression.

The effects of TGF β on stromal cells

Although the effects of increased TGF β 1 expression on epithelial tumour progression have been well established, consideration of the tumour stroma has recently garnered a considerable amount of momentum. Although they give no indication of whether they are causal or merely an effect of progression, specific gene expression changes in the stroma are noted at defined stages of breast cancer progression⁵⁷. Thus, much like the Vogelgram identified the specific genetic changes that are associated with the stages of colon cancer progression⁵⁸, we are beginning to appreciate that gene expression changes that correspond with tumour progression also occur in the stroma of tumours. A recent review examined the various effects of stromal components on each hallmark of cancer and highlighted the importance of stromal cells in influencing tumour progression⁵⁹. However, the mechanisms behind these changes and the pathways that drive specific gene expression changes have yet to be fully elucidated. Given the pleiotropic nature of TGF β and its established role in regulating the development and the function of numerous stromal cells that have an effect on tumour progression, in this section we discuss components of the tumour microenvironment and examine the role of TGF β signalling in each.

MSCs. TGF β has numerous roles in the maintenance and the function of mesenchymal stem cells (MSCs). MSCs have origins in both the bone marrow and a variety of other tissues and have been ascribed numerous functions in normal physiology, including maintaining a homeostatic bone marrow environment and differentiating into various cells that are required for the physiological function of the bone marrow⁶⁰. In terms of normal physiological function, TGF β promotes MSC proliferation through induction of the nuclear accumulation

of β -catenin, which is mediated by phosphorylation of SMAD3 (REF. 61). In addition, an essential function of MSCs in the bone marrow is the preservation of haematopoietic stem cells in an undifferentiated and inactive state⁶⁰. This is achieved through the secretion of numerous factors, including TGF β . Another known function of MSCs is that in various conditions they have the capacity to differentiate into osteoblasts, adipocytes, chondrocytes and fibroblasts⁶². This differentiation maintains the bone marrow microenvironment, mediating bone turnover and cartilage stability, and facilitating adaptability of tissue homeostasis through the modulation of cell population dynamics. In both normal and malignant conditions, TGF β mediates the differentiation process, primarily the induction of chondrocyte and fibroblast differentiation and the inhibition of adipocyte differentiation^{63–65}. Although many functions in MSCs are driven by TGF β , the contextual cues that mediate these differential responses remain unknown. As seen in MSC proliferation, it is probable that the induction of different responses is due to cooperation of the TGF β pathways with other signalling pathways. TGF β also modulates MSC function by promoting chemotaxis to sites of tissue damage and to neoplastic lesions^{63,66}.

In tumours, MSCs have been shown to promote tumour progression both on their own and through their ability to differentiate into other cell types^{67,68}. MSCs secrete numerous cytokines that increase tumour cell proliferation, that drive angiogenesis and that affect tumour cell resistance to chemotherapeutic drugs. MSCs in particular can increase tumour metastasis through interactions with carcinoma cells to induce CCL5 expression^{62,69}. Interestingly, this does not seem to be dependent on the ability of CCL5 to function as a chemoattractant for immune cells but rather on its ability to promote cell colonization in the lungs. In addition, as we discuss below, MSCs can inhibit T cell proliferation and activation, which substantially increases their tumour-promoting effects. Another normal physiological function of MSCs is their ability to differentiate into other cells of mesenchymal origin⁷⁰. Reports indicate that approximately 20% of cancer-associated fibroblasts (CAFs) have origins in the bone marrow, arising from the differentiation of MSCs⁷⁰. Once differentiated, these fibroblasts promote tumour progression in a TGF β -dependent manner; indeed, inhibition of TGF β -mediated MSC differentiation into CAFs through the expression of BAMBI (BMP and activin membrane-bound inhibitor homologue) has been shown to abolish the ability of MSCs to promote tumour progression⁷¹.

Fibroblasts. One of the most common features of carcinomas that overexpress TGF β is a desmoplastic stromal environment^{7,72–75}. Increased ECM deposition corresponds well with the known functions of TGF β in fibroblast activation and cancer^{11,50}. TGF β has previously been shown to induce myofibroblast differentiation in fibroblasts, which leads to increased collagen deposition and ECM remodelling^{76,77}. Expression of a constitutively active TGFBR1 in dermal fibroblasts in mice resulted in increased fibrosis and increased expression of known fibroblast-specific TGF β target genes in chemically

Vogelgram

A model for colorectal cancer development in which specific genetic changes are acquired, which mark progression of the disease from benign to malignant.

Myofibroblast

A generalized term for an activated fibroblast. In tumour biology, these cells function to deposit and to remodel the extracellular matrix as well as to alter tumour progression through paracrine protein signalling.

induced tumours⁷⁸. Thus, an abundance of TGF β in a tumour would probably elicit a similar response. This has been shown to be true as tumours that are characterized as having increased levels of TGF β are associated with increased fibroblast activation and collagen deposition. In addition, fibroblasts derived from small-cell lung carcinomas show enrichment in TGF β signalling compared with normal lung fibroblasts, and the TGF β -enriched gene signature of the carcinoma-derived fibroblasts predicts poor patient outcome⁷⁹. In fact, it has been shown that the increased TGF β activation is caused by interactions between stromal fibroblasts and colon carcinoma cells inducing the expression of numerous MMPs and known TGF β target genes⁸⁰. As discussed above, numerous studies show that TGF β -mediated signalling mechanisms are involved in the reciprocal interactions of tumour cells and the stroma. However, only recently have the effects of these interactions on the metastatic potential of tumours been appreciated. Overexpression of TGF β in a TGF β -non-responsive colon carcinoma cell line drives increased metastasis of these tumour cells⁸¹. In this case, TGF β expression altered the expression of TGF β target genes in fibroblasts, including interleukin-11 (*IL11*), angiopoietin-like 4 (*ANGPTL4*) and *CTGF* (TABLE 1). Importantly, the TGF β -driven gene signature from these fibroblasts was used to predict recurrence in human patients with colon cancer⁸¹. As Calon *et al.*⁸¹ comment, such TGF β -induced gene expression changes have also been ascribed to epithelial carcinoma cells; thus, regardless of the ability of the tumour cells to respond to TGF β , the pro-tumorigenic effects of the TGF β -induced gene expression profile are seen in tumours that express high levels of TGF β .

Much like epithelial cell TGF β signalling, work investigating the role of fibroblast TGF β on tumour progression has showed a contradictory role for the pathway with respect to tumour progression. Although there is a preponderance of evidence supporting a tumour-promoting role for TGF β in stromal fibroblasts, there is also a considerable amount of work showing a tumour-suppressive role. The foundation of this work lies in the paper from Bhowmick *et al.*⁸², which shows that specific deletion of *Tgfb2* in mouse fibroblasts results in spontaneous carcinoma initiation. Further work has shown that fibroblasts that lack TGF β signalling increase the progression of breast, prostate and squamous cell cancers, as well as melanoma^{83–85}. The major gene expression changes that are associated with the loss of this signalling axis in fibroblasts are in genes that encode cytokines and chemokines. In particular, increased expression of *CXCL1*, *CXCL5*, *CXCL12* and *TGFB1* is observed in fibroblasts that have abrogated TGF β signalling⁸⁶ (TABLE 1). This indicates that modulation of the tumour microenvironment through increased infiltration of immune cells is a primary driver of the enhanced tumour progression seen when fibroblasts lose TGF β responsiveness^{9,87}. Consistent with this idea, recent work shows that a significant increase in inflammation is observed in spontaneous forestomach carcinomas that develop as a result of abrogation of TGF β signalling in fibroblasts⁸⁸. Interestingly, this inflammatory response seems to drive tumour formation through the silencing of *CDKN1A* via epigenetic promoter methylation (BOX 1). Administration of anti-inflammatory drugs significantly delayed tumour onset and increased overall survival — a result that supports immune cell infiltration as a facilitator of tumour development. This work is in agreement with *TGFB2* expression having an important role during the progression of breast cancer from normal to ductal carcinoma *in situ* (DCIS) and then to invasive ductal carcinoma (IDC). Stromal *TGFB2* expression decreases as these tumours progress towards invasiveness⁵⁷. Interestingly, immunohistochemical studies in patients with colon cancer have shown a similar trend, but have gone one step further to show that low expression of *TGFB2* in the stroma is an independent predictor of poor patient prognosis⁸⁷.

The innate immune system. The innate immune system is the first line of defence during an immune response. It comprises various phagocytic and antigen-presenting cellular components, including macrophages, neutrophils, natural killer (NK) cells and dendritic cells. TGF β has a substantial role in regulating the recruitment, activation and function of these cells⁸⁹. TGF β functions as an antagonist of major immune functions, decreasing tumour cell recognition and clearing⁹⁰. One of the most notable functions of TGF β in this regard is the induction of M2 macrophage polarization⁹¹. The expression of inducible nitric oxide synthase (iNOS), which is a driver of the nitric oxide production that suppresses monocyte-mediated cell death, is associated with this polarization⁹². TGF β has a similar effect on

Table 1 | TGF β signalling affects central components of fibroblast function

Factors affected by TGF β	Expression levels increased	Expression levels decreased
Chemokines	<ul style="list-style-type: none"> IL-6 IL-11 LIF 	<ul style="list-style-type: none"> CXCL1 IFNγ TNF COX2 iNOS
Growth factors	<ul style="list-style-type: none"> CTGF VEGF CXCL12 TGFβ 	<ul style="list-style-type: none"> HGF EGF
Matrix deposition	<ul style="list-style-type: none"> Tenascin C Fibronectin Elastin Collagen type 1 	<ul style="list-style-type: none"> Collagen type 4 Decorin Vitronectin Fibrillin
ECM remodelling	<ul style="list-style-type: none"> MMP2 MMP3 MMP9 TIMP3 TIMP4 LOX THBS1 	<ul style="list-style-type: none"> ADAMTS1 ADAMTS5 MMP13

ADAMTS, a disintegrin and metalloproteinase with thrombospondin motifs; COX2, cyclooxygenase 2; CTGF, connective tissue growth factor; CXCL, CXC-chemokine ligand; EGF, pro-epidermal growth factor; HGF, hepatocyte growth factor; IL, interleukin; IFN γ , interferon- γ ; LIF, leukaemia inhibitory factor; LOX, lysyl oxidase; MMP, matrix metalloproteinase; iNOS, inducible nitric oxide synthase; TGF β , transforming growth factor- β ; THBS1, thrombospondin 1; TIMP, tissue inhibitor of metalloproteinase; TNF, tumour necrosis factor; VEGF, vascular endothelial growth factor.

Box 1 | TGF β induces epigenetic changes to modulate cellular functions

Epigenetic modulation of gene expression primarily occurs through modification of histones to promote access to specific stretches of DNA and through DNA methylation of CpG islands in gene promoter regions to modulate transcription factor binding and promoter activity¹³⁰. The occurrence of altered chromatin structure and of promoter hypermethylation is important in numerous cancer types. In this regard, tumour progression is associated with an increase in histone deacetylase (HDAC) and DNA methyltransferase (DNMT) activity, which drives these changes^{131,132}. Most transforming growth factor- β (TGF β) silencing in tumours is a direct result of promoter methylation of TGF β signalling components^{133–135}.

TGF β and the TGF β signalling pathway have been linked with histone and promoter modifications¹³⁶. The hypermethylation of promoter regions of a gene mainly results in the silencing of that gene. This seems to be essential to overcome the tumour-suppressor actions of TGF β signalling in epithelial cells. Numerous cancer types show hypermethylation of runt-related transcription factor 3 (RUNX3), which results in attenuation of its growth-suppressive functions¹³⁷. Active TGF β signalling is required for the maintenance of this methylation, as tumours overexpressing SMAD7 (which inhibits TGF β signalling), lose methylation of promoters and reverse the effects of gene silencing¹³⁸. Conversely, it has recently been shown that TGF β downregulates DNMT expression in fibroblasts¹³⁹. Fibroblasts have an inverse requirement for promoter methylation with regard to induction of TGF β target gene expression. DNMT inhibitors enhance the expression of α -smooth muscle actin, as well as collagen type 1, following TGF β treatment^{139,140}. Thus, in the context of TGF β signalling, the use of DNMT inhibitors might provide efficacious results with respect to epithelial cells but might ultimately result in increased microenvironmental desmoplasia in these tumours.

The transcriptional activation of SMAD2 target genes requires histone acetylation by p300 before assembly of transcriptional machinery to drive gene expression¹⁴¹. Conversely, induction of a myofibroblast phenotype by TGF β requires histone deacetylation by HDAC4 (REF. 142). Interestingly, although histone deacetylation is required for TGF β -mediated fibroblast activation, histone methylation is needed to drive the expression of the TGF β target genes connective tissue growth factor (CTGF), collagen type 1 α 1 (COL1A1) and plasminogen activator inhibitor type 1 (PAI1)¹⁴³. Such data support the hypothesis that tumour-promoting or -suppressing functions of TGF β are dependent not only on the microenvironmental context of the tumour but also on the epigenetic histone modifications found in these cells. Although there is a big push for the use of HDAC and DNMT inhibitors in tumour therapy, it is interesting to note that treatment with an HDAC inhibitor might inhibit TGF β activation in epithelial cells; however, this might increase the response of stromal cells to TGF β and promote stromal activation.

neutrophils as it has on macrophages⁹³. A recent publication shows that exposure of neutrophils to TGF β results in a pro-tumorigenic N2 neutrophil expression profile and that inhibition of TGF β signalling switches this cellular phenotype to an antitumorigenic N1 phenotype⁹⁴. N2 neutrophils are associated with the expression of pro-tumorigenic factors, such as MMP9, CXCL1 and ARG1.

Rather than inducing a different activation state in NK cells, TGF β has been identified as a factor that inhibits the functional maturation of this cell population⁹⁵. Immature NK cells do not respond to foreign antigens, thus exposure of NK cells to TGF β will impair the recognition and the clearance of tumour cells. The inhibition of maturation also prevents the systemic effects of NK cells, for example, activation of antigen-presenting dendritic cells and inhibition of interferon- γ (IFN γ) secretion, which drives T helper 1 cell (T_H1 cell) maturation^{96,97}. However, TGF β also has antagonistic effects on dendritic cells that inhibit their antitumorigenic properties⁹⁸. Recent work has shown that impaired TGF β signalling in dendritic cells significantly increases the ability of these cells to present foreign antigens and to activate the adaptive immune system⁹⁹. Such actions would promote the killing of tumour cells but, in TGF β -rich environments where dendritic cells respond to TGF β , their maturation would be impaired, thus promoting tumour growth and progression. Infiltrating dendritic cells in human lung cancer samples are characterized as immature compared with peritumoural dendritic cells, and this immature state has been shown to promote the differentiation of T_{Reg} cells, which increases tumour progression^{100,101}.

The adaptive immune system. Early work investigating the function of TGF β established that it is essential in the regulation of the adaptive immune system; indeed, systemic loss of TGF β 1 expression results in the development of autoimmune-mediated inflammation in numerous organs^{102,103}. In addition, abrogation of TGF β signalling specifically in T cells results in a similar autoimmune response^{104,105}. The mechanisms proposed for regulation of adaptive immunity by TGF β are the SMAD-mediated inhibition of T cell proliferation and the suppression of a T_H1 cell phenotype¹⁰⁶. Recent work has also shown that TGF β mediates the immunosuppressive differentiation of T cells⁸⁹. Treatment of naive T cells with TGF β induces the expression of the transcription factor forkhead box protein P3 (FOXP3), which drives the phenotypical conversion of a naive T cell to a T_{Reg} cell^{107,108}. Interestingly, T_{Reg} cell-induced suppression of the adaptive immune response is also mediated through the expression of TGF β ^{109–111}. Addition of IL-6 to the TGF β treatment of naive T cells induces a completely different cellular phenotype to that of T_{Reg} cells¹¹². IL-6 and TGF β induce T_H17 cell differentiation through the suppression of FOXP3 and the activation of the transcription factor retinoic acid receptor-related orphan receptor γ (ROR γ)¹¹³. Various pro-tumorigenic and antitumorigenic functions of T_H17 cells are primarily mediated through the expression of IL-17 in these cells. Interestingly, the function of IL-17 is also context dependent: IL-17 functions as a tumour promoter by inducing angiogenesis and tumour cell proliferation in immune-deficient hosts, but functions as a tumour suppressor in hosts with intact immune systems by enhancing anti-tumour immune functions. In addition to the secretion of IL-17, the presence of T_H17 cells has been associated with

T helper 1 cell

(T_H1 cell). A primary effector cell of the adaptive immune system, which is classically defined as antitumorigenic.

1D11

An antibody which is specifically directed towards the three transforming growth factor- β ligands and used to sequester these ligands, thus inhibiting TGF β signalling.

Losartan

An angiotensin II receptor antagonist that has been shown to have the off-target effects of downregulating expression of type I transforming growth factor- β receptor (*Tgfr1*) and *Tgfr2*.

increased tumorigenesis that is mediated by neutrophil recruitment and differentiation to an N2 phenotype, and by the induction of IL-6 secretion¹¹⁴.

Despite the numerous pro-tumorigenic functions that are induced by TGF β stimulation of adaptive immune cells, work using TGF β signalling-deficient T cells supports an antitumorigenic function for TGF β signalling in T cells. Thus, as in epithelial cells and fibroblasts, TGF β signalling in T cells seems to have a contextual dependence that drives its dichotomous functions in tumour progression. Attenuation of TGF β signalling in CD8⁺ T cells, through the expression of a dominant-negative TGFBR2, results in uncontrolled cellular proliferation¹⁵². Seminal work from the Letterio laboratory¹¹⁵ has shown that genetic deletion of *Smad4* in T cells results in the spontaneous development of gastrointestinal tumours. The abundant expression of T_H2 cell cytokines has been shown to mediate this phenotype. Just as TGFBR2-deficient fibroblasts are capable of inducing epithelial hyperplasia, so are T cells that are deficient in TGF β signalling. Counter-intuitively, analysis of SMAD4-deficient T cells shows a significant enrichment of T_H17 cells¹¹⁶. Tumours that arise in mice with *Smad4*-null T cells are characterized by an increased expression of IL-6 and IL-11. This increased expression seems to be sufficient to induce T_H17 cell conversion in T cells despite the previously established requirement for TGF β in this process. However, these studies only examined the canonical TGF β pathway. The requirement for the non-canonical pathway in the induction of T_H17 cells has yet to be addressed.

Although the individual components of both the innate and the adaptive immune system undoubtedly have an important role in tumour progression, it is important to think systemically when considering therapeutic intervention; all cells involved in an immune response interact with and respond to the other components; granulocytes secrete chemokines to promote recruitment of phagocytes and antigen-presenting cells; dendritic cells clear abnormal cells and activate T cells; and so on. Taking a macroscopic view in the context of TGF β signalling, it can be concluded that TGF β signalling in the immune system primarily functions as an antagonist that suppresses the recognition and the clearance of tumour cells. It is this feature that has driven current efforts to use anti-TGF β therapeutic regimens to activate T cell activity and thus to improve current chemotherapeutics.

Conclusions and therapeutic perspectives

Epithelia that lack the ability to respond to TGF β have various developmental defects, but the loss of TGF β signalling in selected organs is not, in itself, tumorigenic. However, introduction of oncogenic manipulations causes these cells to be more aggressive and metastatic than their TGF β -responsive counterparts^{45,46,49,53}. Although many of the pro-tumorigenic effects of TGF β have been ascribed to the ability of the protein to promote an EMT in epithelial cells, the recently identified effects of this cytokine on the stroma have proved to be an equally important pro-metastatic mechanism. Given

the important effects seen in tumour progression following manipulation of TGF β responsiveness, efficacious intervention in the TGF β pathway remains a highly sought-after goal for many researchers. As a result of evidence that supports the idea that loss of TGFBR in epithelial cells has a profound effect on the stroma, as well as showing that increased ligand expression by the tumour is associated with poor patient prognosis, the tumour microenvironmental effects of this pathway are increasingly relevant. Rakesh Jain has long been a proponent of the stromal normalization hypothesis, which states that inhibition and reversal of the stromal changes that occur during tumour formation would slow tumour progression and would aid therapeutic intervention¹¹⁷. The use of VEGF inhibitors and relaxin, which is a pregnancy-related hormone shown to induce collagen remodelling, has given credence to this hypothesis; TGF β inhibitors have recently been added to this list^{118,119}. The use of conventional chemotherapeutics with the specific TGF β -targeted antibody 1D11 as well as nonspecific inhibitors such as losartan (which is an angiotensin II receptor inhibitor that has been shown to reduce active TGF β levels in tumours), significantly increases the delivery and the efficacy of chemotherapeutic regimens. The use of these inhibitors in mouse models promoted vessel stability, decreased collagen deposition and significantly improved the efficacy of doxorubicin and doxil treatments¹²⁰.

Numerous studies have shown that systemically intervening in the TGF β pathway during cancer progression elicits antitumorigenic effects on epithelial cell phenotypes and through microenvironmental changes (TABLE 2). Several Phase I and Phase II clinical trials of TGF β -neutralizing antibodies and of small-molecule inhibitors to both ligand and receptor have so far been shown to be safe and efficacious; many trials are still accruing patients^{121–124}. For a complete and comprehensive examination of therapeutic treatments targeting the TGF β pathway in malignant and non-malignant disease progression, see REF. 125.

Interestingly, studies show that the effects of TGF β inhibition are efficacious even in the treatment of tumours that lack the central mediator of canonical TGF β signalling, SMAD4 (REF. 126). Such data highlight the importance not only of the microenvironment but also of microenvironmental TGF β signalling on tumour progression. TGF β -inhibitory treatment regimens tested in numerous mouse models of different cancer types, as well as in early-stage clinical trials, have shown benefits; increased activity of antitumoural adaptive immune cells is thought to be responsible^{127,128}. However, inhibition of the TGF β signalling pathway in specific components of the microenvironment has once again introduced controversial findings into the field of TGF β research: suppression of TGF β signalling in fibroblasts, in an effort to reduce the desmoplastic response in these cells, promoted tumour progression through the augmentation of inflammatory cell infiltration^{81,88}. Despite this, clinical trials involving TGF β inhibition, either through genetic abrogation of TGF β sensitivity (such as by

Table 2 | Identified microenvironmental effects of TGF β inhibitors in preclinical tumour models

Drug name	Cancers tested	Microenvironmental effects	Refs
TGFβ ligand trap			
P144	Melanoma	Increased dendritic cell, natural killer cell and T cell activity	144
1D11	Lung, breast and glioblastoma	<ul style="list-style-type: none"> Increased natural killer cell infiltration and CD8⁺ T cell-mediated tumour cell killing Decreased blood vessel area 	119,145, 146
TGFβ antisense			
AP12009	Pancreatic	Increased immune cell-mediated cytotoxicity	123
Receptor inhibitors			
GW788388	Oesophageal squamous cell carcinoma	<ul style="list-style-type: none"> Decreased fibroblast activation Decreased vascular development 	147
LY2109761	Hepatocellular carcinoma and glioblastoma	<ul style="list-style-type: none"> Decreased angiogenesis Increased pericyte coverage 	148,149
LY364947	Glioblastoma	Increased vascular permeability	146
LY3022859	Breast, pancreas and colon	<ul style="list-style-type: none"> Increased natural killer cell and cytotoxic T cell-mediated tumour cell killing Decreased T_{Reg} cell presence Decreased MDSC infiltration 	126
SB431542	Breast	Increased dendritic cell maturation	98
SD-208	Breast cancer bone metastasis and glioma	<ul style="list-style-type: none"> Decreased osteoclast differentiation Increased immune cell-mediated cytotoxicity 	127
SX-007	Glioma	Increased presence of CD3 ⁺ T cells	150
SM-16	Lung	<ul style="list-style-type: none"> Increased number of activated CD8⁺ T cells Increased presence of CD4⁺ T cells 	151

MDSC, myeloid-derived suppressor cells; TGF β , transforming growth factor- β ; T_{Reg}, regulatory T.

the adoptive transfer of T cells expressing dominant-negative forms of the TGFBR or through the use of inhibitory antibodies, have progressed (ClinicalTrials.gov identifiers: [NCT00368082](#), [NCT01582269](#) and [NCT00844064](#)). Although modest and mixed results are consistently obtained from these trials, this is probably due to varying stromal compositions and to the pleiotropic effects of TGF β .

Studies that elucidate the function of TGF β signalling in specific cellular components of the stroma and that establish an efficacious means of specifically targeting the pathways in these cells are imperative in the resolution of the contextual dependence of TGF β on the tumour microenvironment. Indeed, it is probable that, as with epithelial cell TGF β signalling, there is a contextual dependence that controls the tumour-promoting or tumour-suppressing actions of TGF β in the tumour microenvironment. As cellular components and their

interactions drive this microenvironmental context, future studies will need to address not only signalling but also the stromal milieu in the observed response. Without a thorough understanding of the contextual dependence of TGF β signalling in the stroma it is probable that the implementation of systemic therapeutics that target the TGF β pathway will yield inconclusive results because of tumour heterogeneity. Thus, novel methods of delivery that facilitate specific targeting to a cellular type or evasion of pathway activation are being investigated, including the adoptive transfer of T cells that have an abrogated TGF β response, thus circumventing the antitumorigenic response to TGF β ¹²⁹. This highly specific manipulation should provide beneficial effects on tumour progression, without introducing controversial results by affecting TGF β signalling in other stromal and epithelial cells.

- Siegel, P. M. & Massague, J. Cytostatic and apoptotic actions of TGF β in homeostasis and cancer. *Nature Rev. Cancer* **3**, 807–821 (2003).
- Massague, J. G1 cell-cycle control and cancer. *Nature* **432**, 298–306 (2004).
- Gorsch, S. M., Memoli, V. A., Stukel, T. A., Gold, L. I. & Arrick, B. A. Immunohistochemical staining for transforming growth factor β 1 associates with disease progression in human breast cancer. *Cancer Res.* **52**, 6949–6952 (1992).
- Hasegawa, Y. *et al.* Transforming growth factor- β 1 level correlates with angiogenesis, tumor progression, and prognosis in patients with non-small cell lung carcinoma. *Cancer* **91**, 964–971 (2001).
- Miettinen, P. J., Ebner, R., Lopez, A. R. & Derynck, R. TGF- β induced transdifferentiation of mammary epithelial cells to mesenchymal cells: involvement of type I receptors. *J. Cell Biol.* **127**, 2021–2036 (1994).
- Caulin, C., Scholl, F. G., Frontelo, P., Gamallo, C. & Quintanilla, M. Chronic exposure of cultured transformed mouse epidermal cells to transforming growth factor- β 1 induces an epithelial-mesenchymal transdifferentiation and a spindle tumoral phenotype. *Cell Growth Differ.* **6**, 1027–1035 (1995).
- Walker, R. A., Dearing, S. J. & Gallacher, B. Relationship of transforming growth factor β 1 to extracellular matrix and stromal infiltrates in invasive breast carcinoma. *Br. J. Cancer* **69**, 1160–1165 (1994).
- Wikstrom, P., Stattin, P., Franck-Lissbrant, I., Damber, J. E. & Bergh, A. Transforming growth factor β 1 is associated with angiogenesis, metastasis, and poor clinical outcome in prostate cancer. *Prostate* **37**, 19–29 (1998).
- Hazebag, S., Gorter, A., Kenter, G. G., van den Broek, L. & Fleuren, G. Transforming growth factor- β 1 induces tumor stroma and reduces tumor infiltrate in cervical cancer. *Hum. Pathol.* **33**, 1193–1199 (2002).
- Bierie, B. & Moses, H. L. TGF- β and cancer. *Cytokine Growth Factor Rev.* **17**, 29–40 (2006).
- Massague, J. TGF β signalling in context. *Nature Rev. Mol. Cell Biol.* **13**, 616–630 (2012).

12. Rifkin, D. B. Latent transforming growth factor- β (TGF- β) binding proteins: orchestrators of TGF- β availability. *J. Biol. Chem.* **280**, 7409–7412 (2005).
13. Shi, Y. & Massague, J. Mechanisms of TGF- β signaling from cell membrane to the nucleus. *Cell* **113**, 685–700 (2003).
14. Stenvers, K. L. *et al.* Heart and liver defects and reduced transforming growth factor β 2 sensitivity in transforming growth factor β type III receptor-deficient embryos. *Mol. Cell. Biol.* **23**, 4371–4385 (2003).
15. Moustakas, A. *et al.* The transforming growth factor β receptors types I, II, and III form hetero-oligomeric complexes in the presence of ligand. *J. Biol. Chem.* **268**, 22215–22218 (1993).
16. Derynck, R. & Zhang, Y. E. Smad-dependent and Smad-independent pathways in TGF- β family signalling. *Nature* **425**, 577–584 (2003).
17. Feng, X. H. & Derynck, R. Specificity and versatility in TGF- β signaling through Smads. *Annu. Rev. Cell Dev. Biol.* **21**, 659–693 (2005).
18. Schmierer, B. & Hill, C. S. Kinetic analysis of Smad nucleocytoplasmic shuttling reveals a mechanism for transforming growth factor β -dependent nuclear accumulation of Smads. *Mol. Cell. Biol.* **25**, 9845–9858 (2005).
19. Moustakas, A. & Heldin, C. H. Non-Smad TGF- β signals. *J. Cell Sci.* **118**, 3573–3584 (2005).
20. Massague, J. & Gomis, R. R. The logic of TGF β signaling. *FEBS Lett.* **580**, 2811–2820 (2006).
21. Ewen, M. E., Oliver, C. J., Sluss, H. K., Miller, S. J. & Peeper, D. S. p53-dependent repression of CDK4 translation in TGF- β -induced G1 cell-cycle arrest. *Genes Dev.* **9**, 204–217 (1995).
22. Polyak, K. *et al.* p27Kip1, a cyclin-Cdk inhibitor, links transforming growth factor- β and contact inhibition to cell cycle arrest. *Genes Dev.* **8**, 9–22 (1994).
23. Hannon, G. J. & Beach, D. p15INK4B is a potential effector of TGF- β -induced cell cycle arrest. *Nature* **371**, 257–261 (1994).
24. Pardoll, K. & Moustakas, A. Actions of TGF- β as tumor suppressor and pro-metastatic factor in human cancer. *Biochim. Biophys. Acta* **1775**, 21–62 (2007).
25. Bottlinger, E. P. *et al.* Expression of a dominant-negative mutant TGF- β type II receptor in transgenic mice reveals essential roles for TGF- β in regulation of growth and differentiation in the exocrine pancreas. *EMBO J.* **16**, 2621–2633 (1997).
26. Amendt, C., Schirmacher, P., Weber, H. & Blessing, M. Expression of a dominant negative type II TGF- β receptor in mouse skin results in an increase in carcinoma incidence and an acceleration of carcinoma development. *Oncogene* **17**, 25–34 (1998).
27. Huber, M. A., Kraut, N. & Beug, H. Molecular requirements for epithelial-mesenchymal transition during tumor progression. *Curr. Opin. Cell Biol.* **17**, 548–558 (2005).
28. Dumont, N., Bakin, A. V. & Arteaga, C. L. Autocrine transforming growth factor- β signaling mediates Smad-independent motility in human cancer cells. *J. Biol. Chem.* **278**, 3275–3285 (2003).
29. Giampieri, S. *et al.* Localized and reversible TGF β signalling switches breast cancer cells from cohesive to single cell motility. *Nature Cell Biol.* **11**, 1287–1296 (2009).
- This paper shows the identification of TGF β signalling as a mediator of single-cell invasion and of vascular metastasis, with the cells that are unable to respond to TGF β showing a cohesive migratory phenotype.**
30. Muraoka-Cook, R. S. *et al.* Activated type I TGF β receptor kinase enhances the survival of mammary epithelial cells and accelerates tumor progression. *Oncogene* **25**, 3408–3423 (2006).
31. Safina, A., Vandette, E. & Bakin, A. V. ALK5 promotes tumor angiogenesis by upregulating matrix metalloproteinase-9 in tumor cells. *Oncogene* **26**, 2407–2422 (2007).
32. Schniewind, B. *et al.* Dissecting the role of TGF- β type I receptor/ALK5 in pancreatic ductal adenocarcinoma: Smad activation is crucial for both the tumor suppressive and prometastatic function. *Oncogene* **26**, 4850–4862 (2007).
33. Esquela-Kerscher, A. & Slack, F. J. Oncomirs — microRNAs with a role in cancer. *Nature Rev. Cancer* **6**, 259–269 (2006).
34. Yang, P. *et al.* TGF- β -miR-34a-CCL22 signaling-induced Treg cell recruitment promotes venous metastases of HBV-positive hepatocellular carcinoma. *Cancer Cell* **22**, 291–303 (2012).
35. Davis, B. N., Hilyard, A. C., Lagna, G. & Hata, A. SMAD proteins control DROSHA-mediated microRNA maturation. *Nature* **454**, 56–61 (2008).
- This is the first established link between SMAD activity and the maturation of pri-miRNA through DROSHA, specifically following BMP treatment.**
36. Wang, J., Wang, Y., Ma, Y., Lan, Y. & Yang, X. Transforming growth factor β -regulated microRNA-29a promotes angiogenesis through targeting the phosphatase and tensin homolog in endothelium. *J. Biol. Chem.* **288**, 10418–10426 (2013).
37. Liu, Y. *et al.* MicroRNA-494 is required for the accumulation and functions of tumor-expanded myeloid-derived suppressor cells via targeting of PTEN. *J. Immunol.* **188**, 5500–5510 (2012).
38. Wang, S. E. *et al.* Transforming growth factor β engages TACE and ErbB3 to activate phosphatidylinositol-3 kinase/Akt in ErbB2-overexpressing breast cancer and desensitizes cells to trastuzumab. *Mol. Cell. Biol.* **28**, 5605–5620 (2008).
39. Sartor, M. A. *et al.* ConceptGen: a gene set enrichment and gene set relation mapping tool. *Bioinformatics* **26**, 456–463 (2010).
40. Hills, C. E., Willars, G. B. & Brunskill, N. J. Proinsulin C-peptide antagonizes the profibrotic effects of TGF- β 1 via up-regulation of retinoic acid and HGF-related signaling pathways. *Mol. Endocrinol.* **24**, 822–831 (2010).
41. Maupin, K. A. *et al.* Glycogene expression alterations associated with pancreatic cancer epithelial-mesenchymal transition in complementary model systems. *PLoS ONE* **5**, e13002 (2010).
42. Coulouarn, C., Factor, V. M. & Thorgerisson, S. S. Transforming growth factor- β gene expression signature in mouse hepatocytes predicts clinical outcome in human cancer. *Hepatology* **47**, 2059–2067 (2008).
43. Massague, J. TGF β in cancer. *Cell* **134**, 215–230 (2008).
44. Mima, K. *et al.* Epithelial-mesenchymal transition expression profiles as a prognostic factor for disease-free survival in hepatocellular carcinoma: clinical significance of transforming growth factor- β signaling. *Oncol. Lett.* **5**, 149–154 (2013).
45. Forrester, E. *et al.* Effect of conditional knockout of the type II TGF- β receptor gene in mammary epithelia on mammary gland development and polyomavirus middle T antigen induced tumor formation and metastasis. *Cancer Res.* **65**, 2296–2302 (2005).
46. Lu, S. L. *et al.* Loss of transforming growth factor- β type II receptor promotes metastatic head-and-neck squamous cell carcinoma. *Genes Dev.* **20**, 1331–1342 (2006).
47. Paiva, C. E. *et al.* Absence of TGF- β RII predicts bone and lung metastasis and is associated with poor prognosis in stage III breast tumors. *Cancer Biomark* **11**, 209–217 (2012).
- This paper is a histochemical study of TGF β signalling in breast cancer and shows a correlation between decreased signalling in the tumour epithelium and increased tumour metastasis to the lungs and the bone. This article also indicates that low TGFBR2 levels indicate poor disease-free and overall survival.**
48. Levy, L. & Hill, C. S. Alterations in components of the TGF- β superfamily signaling pathways in human cancer. *Cytokine Growth Factor Rev.* **17**, 41–58 (2006).
49. Malkoski, S. P. *et al.* Loss of transforming growth factor β type II receptor increases aggressive tumor behavior and reduces survival in lung adenocarcinoma and squamous cell carcinoma. *Clin. Cancer Res.* **18**, 2173–2183 (2012).
50. Brier, B. & Moses, H. L. Tumour microenvironment: TGF β : the molecular Jekyll and Hyde of cancer. *Nature Rev. Cancer* **6**, 506–520 (2006).
51. Yang, L. *et al.* Abrogation of TGF β signaling in mammary carcinomas recruits Gr-1⁺CD11b⁺ myeloid cells that promote metastasis. *Cancer Cell* **13**, 23–35 (2008).
52. Ijichi, H. *et al.* Inhibiting Cxcr2 disrupts tumor-stromal interactions and improves survival in a mouse model of pancreatic ductal adenocarcinoma. *J. Clin. Invest.* **121**, 4106–4117 (2011).
53. Lin, S. *et al.* Attenuation of TGF- β signaling suppresses premature senescence in a p21-dependent manner and promotes oncogenic Ras-mediated metastatic transformation in human mammary epithelial cells. *Mol. Biol. Cell* **23**, 1569–1581 (2012).
54. Gewin, L. *et al.* TGF- β receptor deletion in the renal collecting system exacerbates fibrosis. *J. Am. Soc. Nephrol.* **21**, 1334–1343 (2010).
55. Brier, B. *et al.* Abrogation of TGF- β signaling enhances chemokine production and correlates with prognosis in human breast cancer. *J. Clin. Invest.* **119**, 1571–1582 (2009).
- This paper is the first to establish of a poor clinical outcome related to the genetic changes, particularly the altered chemokine expression, that occur following ablation of TGF β signalling in the tumour epithelium.**
56. Finak, G. *et al.* Stromal gene expression predicts clinical outcome in breast cancer. *Nature Med.* **14**, 518–527 (2008).
- This is the first identification of a link between stromal gene expression changes and the outcome of patients with breast cancer.**
57. Knudsen, E. S. *et al.* Progression of ductal carcinoma *in situ* to invasive breast cancer is associated with gene expression programs of EMT and myoepithelia. *Breast Cancer Res. Treat.* **133**, 1009–1024 (2012).
58. Vogelstein, B. *et al.* Genetic alterations during colorectal-tumor development. *N. Engl. J. Med.* **319**, 525–532 (1988).
59. Hanahan, D. & Coussens, L. M. Accessories to the crime: functions of cells recruited to the tumor microenvironment. *Cancer Cell* **21**, 309–322 (2012).
60. Nombela-Arrieta, C., Ritz, J. & Silbstein, L. E. The elusive nature and function of mesenchymal stem cells. *Nature Rev. Mol. Cell Biol.* **12**, 126–131 (2011).
61. Watabe, T. & Miyazono, K. Roles of TGF- β family signaling in stem cell renewal and differentiation. *Cell Res.* **19**, 103–115 (2009).
62. Cui, B. G. & Karnoub, A. E. Mesenchymal stem cells in tumor development: emerging roles and concepts. *Cell Adh. Migr.* **6**, 220–230 (2012).
63. Tang, Y. *et al.* TGF- β 1-induced migration of bone mesenchymal stem cells couples bone resorption with formation. *Nature Med.* **15**, 757–765 (2009).
64. Zhao, L. & Hantash, B. M. TGF- β 1 regulates differentiation of bone marrow mesenchymal stem cells. *Vitam. Horm.* **87**, 127–141 (2011).
65. Kurpinski, K. *et al.* Transforming growth factor- β and notch signaling mediate stem cell differentiation into smooth muscle cells. *Stem Cells* **28**, 734–742 (2010).
66. Wan, M. *et al.* Injury-activated transforming growth factor β controls mobilization of mesenchymal stem cells for tissue remodeling. *Stem Cells* **30**, 2498–2511 (2012).
67. Jung, Y. *et al.* Recruitment of mesenchymal stem cells into prostate tumours promotes metastasis. *Nature Commun.* **4**, 1795 (2013).
68. Quante, M. *et al.* Bone marrow-derived myofibroblasts contribute to the mesenchymal stem cell niche and promote tumor growth. *Cancer Cell* **19**, 257–272 (2011).
69. Karnoub, A. E. *et al.* Mesenchymal stem cells within tumour stroma promote breast cancer metastasis. *Nature* **449**, 557–563 (2007).
70. Spaeth, E. L. *et al.* Mesenchymal stem cell transition to tumor-associated fibroblasts contributes to fibrovascular network expansion and tumor progression. *PLoS ONE* **4**, e4992 (2009).
71. Shanguan, L. *et al.* Inhibition of TGF- β /Smad signaling by BAMBI blocks differentiation of human mesenchymal stem cells to carcinoma-associated fibroblasts and abolishes their protumor effects. *Stem Cells* **30**, 2810–2819 (2012).
72. Verona, E. V. *et al.* Transforming growth factor- β signaling in prostate stromal cells supports prostate carcinoma growth by up-regulating stromal genes related to tissue remodeling. *Cancer Res.* **67**, 5737–5746 (2007).
73. Border, W. A. & Noble, N. A. Transforming growth factor β in tissue fibrosis. *N. Engl. J. Med.* **331**, 1286–1292 (1994).
74. Leask, A. & Abraham, D. J. TGF- β signaling and the fibrotic response. *FASEB J.* **18**, 816–827 (2004).
75. Boyd, N. F. *et al.* Mammographic density and the risk and detection of breast cancer. *N. Engl. J. Med.* **356**, 227–236 (2007).
76. Desmouliere, A., Geinoz, A., Gabbiani, F. & Gabbiani, G. Transforming growth factor- β 1 induces α -smooth muscle actin expression in granulation tissue myofibroblasts and in quiescent and growing cultured fibroblasts. *J. Cell Biol.* **122**, 103–111 (1993).
77. Sime, P. J., Xing, Z., Graham, F. L., Csaky, K. G. & Gaudie, J. Adenovector-mediated gene transfer of active transforming growth factor- β 1 induces prolonged severe fibrosis in rat lung. *J. Clin. Invest.* **100**, 768–776 (1997).

78. Sonnylal, S. *et al.* Postnatal induction of transforming growth factor β signaling in fibroblasts of mice recapitulates clinical, histologic, and biochemical features of scleroderma. *Arthritis Rheum.* **56**, 334–344 (2007).
79. Navab, R. *et al.* Prognostic gene-expression signature of carcinoma-associated fibroblasts in non-small cell lung cancer. *Proc. Natl Acad. Sci. USA* **108**, 7160–7165 (2011).
80. Hawinkels, L. J. *et al.* Interaction with colon cancer cells hyperactivates TGF- β signaling in cancer-associated fibroblasts. *Oncogene* <http://dx.doi.org/10.1038/nc.2012.536> (2012).
81. Calon, A. *et al.* Dependency of colorectal cancer on a TGF- β -driven program in stromal cells for metastasis initiation. *Cancer Cell* **22**, 571–584 (2012).
This paper establishes TGF β -responsive stromal signalling, especially from fibroblasts, as a driver of tumour progression and metastasis. This stromal programme could be blocked with the addition of TGF β inhibitors to slow tumour growth.
82. Bhowmick, N. A. *et al.* TGF- β signaling in fibroblasts modulates the oncogenic potential of adjacent epithelia. *Science* **303**, 848–851 (2004).
This is the first report showing that genetic mutations in stromal fibroblast can result in the development of carcinoma formation from the adjacent epithelium.
83. Cheng, N. *et al.* Loss of TGF- β type II receptor in fibroblasts promotes mammary carcinoma growth and invasion through upregulation of TGF- α , MSP- and HGF-mediated signaling networks. *Oncogene* **24**, 5053–5068 (2005).
84. Franco, O. E. *et al.* Altered TGF- β signaling in a subpopulation of human stromal cells promotes prostatic carcinogenesis. *Cancer Res.* **71**, 1272–1281 (2011).
85. Meng, W. *et al.* Downregulation of TGF- β receptor types II and III in oral squamous cell carcinoma and oral carcinoma-associated fibroblasts. *BMC Cancer* **11**, 88 (2011).
86. Xu, B. J. *et al.* Quantitative analysis of the secretome of TGF- β signaling-deficient mammary fibroblasts. *Proteomics* **10**, 2458–2470 (2010).
87. Bacman, D. *et al.* TGF- β receptor 2 downregulation in tumour-associated stroma worsens prognosis and high-grade tumours show more tumour-associated macrophages and lower TGF- β 1 expression in colon carcinoma: a retrospective study. *BMC Cancer* **7**, 156 (2007).
88. Achyut, B. R. *et al.* Inflammation-mediated genetic and epigenetic alterations drive cancer development in the neighboring epithelium upon stromal abrogation of TGF- β signaling. *PLoS Genet.* **9**, e1003251 (2013).
This is an interesting paper showing that inflammatory infiltrates mediate the pro-tumorigenic functions of fibroblasts that lack TGF β signalling, particularly through the induction of epigenetic silencing of p15 and p16 in epithelial cells.
89. Flavell, R. A., Sanjabi, S., Wrzesinski, S. H. & Licona-Limon, P. The polarization of immune cells in the tumour environment by TGF β . *Nature Rev. Immunol.* **10**, 554–567 (2010).
90. Gigante, M., Gesualdo, L. & Ranieri, E. TGF- β : a master switch in tumor immunity. *Curr. Pharm. Des.* **18**, 4126–4134 (2012).
91. Gong, D. *et al.* TGF β signaling plays a critical role in promoting alternative macrophage activation. *BMC Immunol.* **13**, 31 (2012).
92. Murray, P. J. & Wynn, T. A. Protective and pathogenic functions of macrophage subsets. *Nature Rev. Immunol.* **11**, 723–737 (2011).
93. Fridlender, Z. G. & Albelda, S. M. Tumor-associated neutrophils: friend or foe? *Carcinogenesis* **33**, 949–955 (2012).
94. Fridlender, Z. G. *et al.* Polarization of tumor-associated neutrophil phenotype by TGF- β : “N1” versus “N2” TAN. *Cancer Cell* **16**, 183–194 (2009).
This is the first paper to establish an N2 neutrophil phenotype and to show that this phenotype is driven by TGF β signalling. These neutrophils promote tumour progression through the inhibition of antitumorigenic CD8 $^{+}$ T cell activity.
95. Marcoc, J. P. *et al.* TGF- β is responsible for NK cell immaturity during ontogeny and increased susceptibility to infection during mouse infancy. *Nature Immunol.* **13**, 843–850 (2012).
This paper identifies TGF β as a factor that inhibits NK cell maturation and shows that loss of TGF β signalling is a mark of stage F NK cell development.
96. Leavy, O. Maturation and function of NK cells. *Nature Rev. Immunol.* **12**, 150 (2012).
97. Laouar, Y., Sutterwala, F. S., Gorelik, L. & Flavell, R. A. Transforming growth factor- β controls T helper type 1 cell development through regulation of natural killer cell interferon- γ . *Nature Immunol.* **6**, 600–607 (2005).
98. Tanaka, H. *et al.* Transforming growth factor β signaling inhibitor, SB-431542, induces maturation of dendritic cells and enhances anti-tumor activity. *Oncol. Rep.* **24**, 1637–1643 (2010).
99. Novitskiy, S. V. *et al.* Deletion of TGF- β signaling in myeloid cells enhances their anti-tumorigenic properties. *J. Leukoc. Biol.* **92**, 641–651 (2012).
100. Perrot, I. *et al.* Dendritic cells infiltrating human non-small cell lung cancer are blocked at immature stage. *J. Immunol.* **178**, 2763–2769 (2007).
101. Roncarolo, M. G., Levings, M. K. & Traversari, C. Differentiation of T regulatory cells by immature dendritic cells. *J. Exp. Med.* **193**, F5–9 (2001).
102. Shull, M. M. *et al.* Targeted disruption of the mouse transforming growth factor- β 1 gene results in multifocal inflammatory disease. *Nature* **359**, 693–699 (1992).
103. Kulkarni, A. B. *et al.* Transforming growth factor β 1 null mutation in mice causes excessive inflammatory response and early death. *Proc. Natl Acad. Sci. USA* **90**, 770–774 (1993).
104. Marie, J. C., Liggitt, D. & Rudensky, A. Y. Cellular mechanisms of fatal early-onset autoimmunity in mice with the T cell-specific targeting of transforming growth factor- β receptor. *Immunity* **25**, 441–454 (2006).
105. Bommireddy, R. *et al.* Self-antigen recognition by TGF β 1-deficient T cells causes their activation and systemic inflammation. *Lab Invest.* **86**, 1008–1019 (2006).
106. Li, M. O. & Flavell, R. A. TGF- β : a master of all T cell trades. *Cell* **134**, 392–404 (2008).
107. Chen, W. *et al.* Conversion of peripheral CD4 $^{+}$ CD25 $^{-}$ naive T cells to CD4 $^{+}$ CD25 $^{+}$ regulatory T cells by TGF- β induction of transcription factor Foxp3. *J. Exp. Med.* **198**, 1875–1886 (2003).
108. Tone, Y. *et al.* Smad3 and NFAT cooperate to induce Foxp3 expression through its enhancer. *Nature Immunol.* **9**, 194–202 (2008).
109. Chen, Y., Kuchroo, V. K., Inobe, J., Hafler, D. A. & Weiner, H. L. Regulatory T cell clones induced by oral tolerance: suppression of autoimmune encephalomyelitis. *Science* **265**, 1237–1240 (1994).
110. Vignali, D. A., Collison, L. W. & Workman, C. J. How regulatory T cells work. *Nature Rev. Immunol.* **8**, 523–532 (2008).
111. von Boehmer, H. & Daniel, C. Therapeutic opportunities for manipulating T_{reg} cells in autoimmunity and cancer. *Nature Rev. Drug Discov.* **12**, 51–63 (2013).
112. Mangan, P. R. *et al.* Transforming growth factor- β induces development of the T_H17 lineage. *Nature* **441**, 231–234 (2006).
This paper identifies TGF β as a primary factor in the induced differentiation of T_H17 cell commitment in a subpopulation of T cells — a process that is independent of T_{reg} cells.
113. Ivanov, I. I. *et al.* The orphan nuclear receptor ROR γ directs the differentiation program of proinflammatory IL-17 $^{+}$ T helper cells. *Cell* **126**, 1121–1133 (2006).
114. Whiteside, T. L. What are regulatory T cells (Treg) regulating in cancer and why? *Semin. Cancer Biol.* **22**, 327–334 (2012).
115. Kim, B. G. *et al.* Smad4 signalling in T cells is required for suppression of gastrointestinal cancer. *Nature* **441**, 1015–1019 (2006).
This seminal paper identifies of TGF β superfamily signalling in T cells as an inhibitor of tumour progression due to the suppression of the expression of T_H1 cytokines.
116. Ghoreschi, K. *et al.* Generation of pathogenic Th17 cells in the absence of TGF- β signalling. *Nature* **467**, 967–971 (2010).
117. Jain, R. K. Normalizing tumor microenvironment to treat cancer: bench to bedside to biomarkers. *J. Clin. Oncol.* **31**, 2205–2218 (2013).
118. Huang, Y. *et al.* Vascular normalizing doses of antiangiogenic treatment reprogram the immunosuppressive tumor microenvironment and enhance immunotherapy. *Proc. Natl Acad. Sci. USA* **109**, 17561–17566 (2012).
119. Liu, J. *et al.* TGF- β blockade improves the distribution and efficacy of therapeutics in breast carcinoma by normalizing the tumor stroma. *Proc. Natl Acad. Sci. USA* **109**, 16618–16623 (2012).
This is an interesting paper showing that TGF β inhibition-normalized breast tumour stroma enhances chemotherapeutic efficacy.
120. Diop-Frimpong, B., Chauhan, V. P., Krane, S., Boucher, Y. & Jain, R. K. Losartan inhibits collagen I synthesis and improves the distribution and efficacy of nanotherapeutics in tumors. *Proc. Natl Acad. Sci. USA* **108**, 2909–2914 (2011).
121. Mead, A. L., Wong, T. T., Cordeiro, M. F., Anderson, I. K. & Khaw, P. T. Evaluation of anti-TGF- β 2 antibody as a new postoperative anti-scarring agent in glaucoma surgery. *Invest. Ophthalmol. Vis. Sci.* **44**, 3394–3401 (2003).
122. Bogdahn, U. *et al.* Targeted therapy for high-grade glioma with the TGF- β 2 inhibitor trabedersen: results of a randomized and controlled phase IIb study. *Neuro Oncol.* **13**, 132–142 (2011).
123. Schlingensiepen, K. H. *et al.* Transforming growth factor- β 2 gene silencing with trabedersen in pancreatic cancer. *Cancer Sci.* **102**, 1193–1200 (2011).
124. Roldan Urgoiti, G. B., Singh, A. D. & Easaw, J. C. Extended adjuvant temozolomide for treatment of newly diagnosed glioblastoma multiforme. *J. Neurooncol.* **108**, 173–177 (2012).
125. Akhurst, R. J. & Hata, A. Targeting the TGF β signalling pathway in disease. *Nature Rev. Drug Discov.* **11**, 790–811 (2012).
126. Zhong, Z. *et al.* Anti-transforming growth factor β receptor II antibody has therapeutic efficacy against primary tumor growth and metastasis through multi-effects on cancer, stroma, and immune cells. *Clin. Cancer Res.* **16**, 1191–1205 (2010).
This paper shows that anti-TGF β therapeutics slow tumour progression by modulating the composition of immune cell infiltrates in favour of an antitumorigenic phenotype.
127. Uhl, M. *et al.* SD-208, a novel transforming growth factor β receptor I kinase inhibitor, inhibits growth and invasiveness and enhances immunogenicity of murine and human glioma cells *in vitro* and *in vivo*. *Cancer Res.* **64**, 7954–7961 (2004).
128. Kim, S. *et al.* Systemic blockade of transforming growth factor- β signaling augments the efficacy of immunogene therapy. *Cancer Res.* **68**, 10247–10256 (2008).
129. Foster, A. E. *et al.* Antitumor activity of EBV-specific T lymphocytes transduced with a dominant negative TGF- β receptor. *J. Immunother.* **31**, 500–505 (2008).
130. Azad, N., Zahnow, C. A., Rudin, C. M. & Baylin, S. B. The future of epigenetic therapy in solid tumours: lessons from the past. *Nature Rev. Clin. Oncol.* **10**, 256–266 (2013).
131. Jones, P. A. & Baylin, S. B. The fundamental role of epigenetic events in cancer. *Nature Rev. Genet.* **3**, 415–428 (2002).
132. Zhang, Q. *et al.* TGF- β regulates DNA methyltransferase expression in prostate cancer, correlates with aggressive capabilities, and predicts disease recurrence. *PLoS ONE* **6**, e25168 (2011).
133. Ammanamanchi, S. & Brattain, M. G. Restoration of transforming growth factor- β signaling through receptor RI induction by histone deacetylase activity inhibition in breast cancer cells. *J. Biol. Chem.* **279**, 32620–32625 (2004).
134. Kang, S. H. *et al.* Transcriptional repression of the transforming growth factor- β type I receptor gene by DNA methylation results in the development of TGF- β resistance in human gastric cancer. *Oncogene* **18**, 7280–7286 (1999).
135. Osada, H. *et al.* Heterogeneous transforming growth factor (TGF)- β unresponsiveness and loss of TGF- β receptor type II expression caused by histone deacetylation in lung cancer cell lines. *Cancer Res.* **61**, 8331–8339 (2001).
136. Khin, S. S. *et al.* Epigenetic alteration by DNA promoter hypermethylation of genes related to transforming growth factor- β (TGF- β) signaling in cancer. *Cancers* **3**, 982–993 (2011).
137. Yeh, K. T. *et al.* Aberrant TGF β /SMAD4 signaling contributes to epigenetic silencing of a putative tumor suppressor, RunX1T1 in ovarian cancer. *Epigenetics* **6**, 727–739 (2011).

138. Papageorgis, P. *et al.* Smad signaling is required to maintain epigenetic silencing during breast cancer progression. *Cancer Res.* **70**, 968–978 (2010).
139. Pan, X., Chen, Z., Huang, R., Yao, Y. & Ma, G. Transforming growth factor β 1 induces the expression of collagen type I by DNA methylation in cardiac fibroblasts. *PLoS ONE* **8**, e60335 (2013).
140. Hu, B., Gharaee-Kermani, M., Wu, Z. & Phan, S. H. Epigenetic regulation of myofibroblast differentiation by DNA methylation. *Am. J. Pathol.* **177**, 21–28 (2010).
141. Ross, S. *et al.* Smads orchestrate specific histone modifications and chromatin remodeling to activate transcription. *EMBO J.* **25**, 4490–4502 (2006).
142. Sun, G. *et al.* Epigenetic histone methylation modulates fibrotic gene expression. *J. Am. Soc. Nephrol.* **21**, 2069–2080 (2010).
143. Glenisson, W., Castronovo, V. & Waltregny, D. Histone deacetylase 4 is required for TGF β 1-induced myofibroblastic differentiation. *Biochim. Biophys. Acta* **1773**, 1572–1582 (2007).
144. Diaz-Valdes, N. *et al.* Induction of monocyte chemoattractant protein-1 and interleukin-10 by TGF β 1 in melanoma enhances tumor infiltration and immunosuppression. *Cancer Res.* **71**, 812–821 (2011).
145. Terabe, M. *et al.* Synergistic enhancement of CD8 $^{+}$ T cell-mediated tumor vaccine efficacy by an anti-transforming growth factor- β monoclonal antibody. *Clin. Cancer Res.* **15**, 6560–6569 (2009).
146. Hardee, M. E. *et al.* Resistance of glioblastoma-initiating cells to radiation mediated by the tumor microenvironment can be abolished by inhibiting transforming growth factor- β . *Cancer Res.* **72**, 4119–4129 (2012).
147. Noma, K. *et al.* The essential role of fibroblasts in esophageal squamous cell carcinoma-induced angiogenesis. *Gastroenterology* **134**, 1981–1993 (2008).
148. Mazzocca, A., Fransvea, E., Lavezzari, G., Antonaci, S. & Giannelli, G. Inhibition of transforming growth factor β receptor I kinase blocks hepatocellular carcinoma growth through neo-angiogenesis regulation. *Hepatology* **50**, 1140–1151 (2009).
149. Zhang, M. *et al.* Trimodal glioblastoma treatment consisting of concurrent radiotherapy, temozolomide, and the novel TGF- β receptor I kinase inhibitor LY2109761. *Neoplasia* **13**, 537–549 (2011).
150. Tran, T. T. *et al.* Inhibiting TGF- β signaling restores immune surveillance in the SMA-560 glioma model. *Neuro Oncol.* **9**, 259–270 (2007).
151. Garrison, K. *et al.* The small molecule TGF- β signaling inhibitor SM16 synergizes with agonistic OX40 antibody to suppress established mammary tumors and reduce spontaneous metastasis. *Cancer Immunol. Immunother.* **61**, 511–521 (2012).
152. Lucas, P. J., Kim, S. J., Melby, S. J. & Gress, R. E. Disruption of T. cell homeostasis in mice expressing a T. cell-specific dominant negative transforming growth factor β II receptor. *J. Exp. Med.* **191**, 1187–1196 (2000).

Acknowledgements

This work is supported by US National Institutes of Health (NIH) Grants CA085492 and CA102162. The authors would also like to thank all of the members of the Moses Laboratory for their thoughtful discussions. In addition, the authors would like to apologize to any investigators whose work could not be included owing to space limitations.

Competing interests statement

The authors declare no competing financial interests.

DATABASES

ClinicalTrials.gov: <http://www.clinicaltrials.gov>
NCT00368082 | NCT01582269 | NCT00844064

ALL LINKS ARE ACTIVE IN THE ONLINE PDF