


Radio detection of high energy neutrinos in the Greenland icecap

Arthur Adriaens



Department of Physics and Astronomy

Promotor: Prof. dr. Dirk Ryckbosch Dirk.Ryckbosch@ugent.be

Accompanist: Bob Oeyen Bob.Oeyen@ugent.be

Master's dissertation submitted in partial fulfilment of the requirements for the degree of master in Physics and Astronomy

CONTENTS

1	Neutrinos	2
1.1	Discovery	2
1.2	Standard model	2
1.3	Outside sources	2
1.3.1	Cosmic neutrinos	2
1.3.2	Oscillations	3
1.3.3	Majorana	3
2	Radio detection	4

CHAPTER

1

NEUTRINOS

1.1 Discovery

1.2 Standard model

1.3 Outside sources

1.3.1 Cosmic neutrinos

To estimate the temperature of the neutrinos who decoupled at the start of the universe, we can take a look at conservation of entropy [1] (...) The entropy before and after decoupling are:

$$s(a_1) = \frac{2\pi^2}{45} \left(2 + \frac{7}{8}(2 + 2 + 3 + 3)\right) T_1^3 \quad (1.1)$$

$$= \frac{2\pi^2}{45} \frac{86}{8} T_1^3 \quad (1.2)$$

$$s(a_2) = \frac{2\pi^2}{45} (2T_\gamma^3 + \frac{7}{8}(6)T_\nu^3) \quad (1.3)$$

$$(1.4)$$

Conservation of entropy:

$$s(a_1)a_1^3 = s(a_2)a_2^3 \quad (1.5)$$

$$\frac{86}{8}(T_1 a_1)^3 = \left(2 \left(\frac{T_\gamma}{T_\nu} \right)^3 + \frac{42}{8} \right) (T_\nu a_2)^3 \quad (1.6)$$

$$\frac{86}{8} = 2 \left(\frac{T_\gamma}{T_\nu} \right)^3 + \frac{42}{8} \quad (1.7)$$

$$\frac{44}{16} = \left(\frac{T_\gamma}{T_\nu} \right)^3 \quad (1.8)$$

$$\left(\frac{T_\gamma}{T_\nu} \right) = \left(\frac{11}{4} \right)^{1/3} \quad (1.9)$$

i.e

$$T_\nu = \left(\frac{4}{11} \right)^{1/3} T_\gamma \quad (1.10)$$

Φ

1.3.2 Oscillations

1.3.3 Majorana

CHAPTER

2

RADIO DETECTION

BIBLIOGRAPHY

- [1] Scott Dodelson. Modern Cosmology. Academic Press, Amsterdam, 2003.