


Tutorial 02

Measure & Probability Space

Xujie Zhao

6. & 9. May 2025



About Tutorial

- **Tutorial:** In-Person, 2h by two time periods per week
 - 12:00-14:00 Tue, 03.11.018 (Group B33)
 - 10:00-12:00 Fri, 01.06.011 (Group E24)

- The slides are not guaranteed to be accurate! (Written by myself)

- **Materials of this tutorial:** Please scan the QR-Code 

- **PART I.** Recap of the Lecture
- **PART II.** Exercises from the Exercise Sheet
- **PART III.** Additional Exercises (if time permits)



I. Recap

- **Measure and measure space**
 - ① - Definition
 - ② - Properties of measures
- **Probability measure and probability space**
 - ③ - Definition
 - ④ - Inclusion-exclusion principle (Siebformel)
- **Cumulative distribution function (cdf)**
 - ⑤ - Definition
 - ⑥ - Properties (*self-read on script 1.22*)
- **Discrete probability space**
 - ⑦ - Definition
 - ⑧ - Classical urn models

II. Exercise Sheet

Exercise 1. True or false? Justify your answers either by a proof or a counterexample.

- (i) For a probability space $(\Omega, \mathcal{A}, P)$ we have $P(\Omega \setminus A) = 1 - P(A)$ for all $A \in \mathcal{A}$.
- (ii) Let $(\Omega, \mathcal{A}, \mu)$ be a measure space, $A, B \in \mathcal{A}$ such that $\mu(A \cap B) < \infty$. Then $\mu(B \setminus A) = \mu(B) - \mu(A \cap B)$.
- (iii) For a probability space $(\Omega, \mathcal{A}, P)$ and a sequence $(A_i)_{i \in \mathbb{N}}$ in $\mathcal{A}$, we have

$$P \left(\bigcup_{i \in \mathbb{N}} A_i \right) = \sup_{i \in \mathbb{N}} P(A_i).$$

Let $S \subseteq \mathbb{R}$. A real number $u \in \mathbb{R}$ is called an *upper bound* of S if $x \leq u$ for all $x \in S$. An upper bound s of S is called *supremum* or *least upper bound* if for all upper bounds u of S in $\mathbb{R}$ we have $s \leq u$. We then write $\sup(S) = s$. When $S = \{x_i \mid i \in \mathbb{N}\} \subseteq \mathbb{R}$ is at most countable, we also write $\sup_{i \in \mathbb{N}} x_i = s$. *Suprema and infima* can be defined for subsets of any partially ordered set.

II. Exercise Sheet

(iv) Let $(\Omega, \mathcal{A}, P)$ be a probability space and $A, B, C \in \mathcal{A}$. Then

$$P(A \cup B \cup C) = P(A) + P(B) + P(C) - P(A \cap B) - P(A \cap C) - P(B \cap C) + P(A \cap B \cap C)$$

(v) The triple consisting of $\mathbb{N}$, $\mathcal{P}(\mathbb{N})$, and the set function

$$\mu : \mathcal{P}(\mathbb{N}) \rightarrow [0, \infty], A \mapsto \begin{cases} 0 & A \text{ finite} \\ \infty & A \text{ infinite} \end{cases}$$

form a measure space.

(vi) We consider the two-time fair coin flip, where we interpret the two sides to count as 0 or 1. We are only interested in the sum of the two outcomes. Then the cdf is given by

$$F(x) = \begin{cases} 0 & x < 0, \\ \frac{1}{4} & x \in [0, 1), \\ \frac{3}{4} & x \in [1, 2), \\ 1 & x \geq 2. \end{cases}$$

II. Exercise Sheet



Exercise 2. Let $\Omega = \{\square, \begin{smallmatrix} \square \\ \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet \\ \square \end{smallmatrix}, \begin{smallmatrix} \bullet & \bullet \\ \square & \square \end{smallmatrix}, \begin{smallmatrix} \bullet & \bullet \\ \bullet & \bullet \end{smallmatrix}, \begin{smallmatrix} \bullet & \bullet \\ \bullet & \bullet \end{smallmatrix}\}$, representing the rolling of a fair die. Assume a uniform distribution over Ω and let $p \in [0, 1]$ be the probability of obtaining at least one $\begin{smallmatrix} \bullet \\ \square \end{smallmatrix}$ when throwing six fair dice at once. Compute p (using rigorous mathematical notation, i.e., setting up the probability spaces properly).

II. Exercise Sheet



Exercise 3 (*Inclusion-exclusion formula*). Let $(\Omega, \mathcal{A}, P)$ be a probability space, $(A_i)_{i \in [n]} \in \mathcal{A}^{[n]}$ for a given $n \in \mathbb{N}$. Then

$$P(A_1 \cup \dots \cup A_n) = \sum_{k=1}^n (-1)^{k-1} \sum_{1 \leq i_1 < \dots < i_k \leq n} P(A_{i_1} \cap \dots \cap A_{i_k})$$

In particular, it holds that

$$P(A_1 \cup A_2) = P(A_1) + P(A_2) - P(A_1 \cap A_2) .$$

[Hint: Use complete induction.]

II. Exercise Sheet

Exercise 4. The DWT tutor Angelika supervises an exercise group with n sheets being submitted anonymously. After grading, Angelika randomly redistributes the sheets. What is the probability that no one gets their original sheet back? How does this probability behave as $n \rightarrow \infty$?

[Hint: Use the inclusion-exclusion formula and that $\exp(x) = \sum_{k=0}^{\infty} \frac{x^k}{k!}$]

III. Additional Exercises

Exercise 5. After a long party night, Anna and Bob are the only ones left at the Marienplatz subway station, where a train departs every 10 minutes. Unfortunately, the trains are so overcrowded that only one person can board at a time. Each person on the platform has an equal probability of successfully boarding. In addition, two new people arrive at the station during the time before the next train arrives. Determine the probability that Anna

- (a) can board the n -th train,
(b) can board no later than the n -th train,
where $n \in \mathbb{N}$.

(a) Let event A_n be "Anna boarded on the n -th subway".
(Everytime subway leaves, # people waiting + 1)

$\hookrightarrow$ When n -th subway comes, there will be $n+1$ people waiting.
To board on the n -th subway, Anna has to miss first $(n-1)$ subways.

$$P(A_1) = \frac{1}{2} \quad P(A_2) = (1 - \frac{1}{2}) \times \frac{1}{3} = \frac{1}{6} \quad \dots$$

$$P(A_n) = (1 - \frac{1}{2})(1 - \frac{1}{3}) \dots (1 - \frac{1}{n}) \cdot \frac{1}{n+1} = \frac{1}{2} \times \frac{2}{3} \times \frac{3}{4} \dots \times \frac{n-1}{n} \cdot \frac{1}{n+1} = \frac{1}{n(n+1)}$$

(b) Let event B be "no later than ..."

$$B = \bigcup_{i=1}^n A_i, \quad P(B) = P\left(\bigcup_{i=1}^n A_i\right)$$

$$= \sum_{i=1}^n P(A_i)$$

$$= \frac{1}{1 \times 2} + \frac{1}{2 \times 3} + \dots + \frac{1}{n(n+1)}$$

$$= 1 - \frac{1}{2} + \frac{1}{2} - \frac{1}{3} + \dots + \frac{1}{n} - \frac{1}{n+1}$$

$$= \frac{n+1-1}{n+1} = \frac{n}{n+1}$$

Materials

