

Sequences of real and complex numbers

July 28, 2024

This chapter, as the title suggests, is dedicated to the study of sequences of real and complex numbers. The first sections contain preliminary notions, conventional notations and a brief overview on primary notions necessary to make this book as self-contained as possible.

1 Real numbers

Let X, Y be two nonempty sets. A *triple* $\rho = (X, Y; \mathcal{R})$, where $\mathcal{R} \subseteq X \times Y$, is called a *binary relation*. The set X is called the *domain* of the relation ρ , Y is the *codomain* or the *range* of ρ while \mathcal{R} represents the *graph* of relation ρ . If $(x, y) \in \mathcal{R}$ we say that x is ρ related to y (or x is in relation ρ with y) and we write $x\rho y$.

The notion of binary relation, frequently used in many branches of mathematics and computer sciences, is in fact a generalization of the notion of function. A *function* $f = (X, Y; \mathcal{R})$ is a relation with the property that for every $x \in X$ there exists a unique $y \in Y$ such that xfy (xfy is denoted usually by $y = f(x)$).

Definition 1 A binary relation $\rho = (X, Y; \mathcal{R})$ is called:

- reflexive, if $x\rho x$ for all $x \in X$;
- symmetric, if $x\rho y \implies y\rho x$;
- antisymmetric, if $x\rho y$ and $y\rho x \implies x = y$;
- transitive, if $x\rho y$ and $y\rho z \implies x\rho z$;

- equivalence, if it is reflexive, symmetric and transitive;
- order relation, if it is reflexive, antisymmetric and transitive;
- totally order relation, if it an order relation and for all $x, y \in X$ we have xpy or ypx .

Example 1.1 Let $\mathcal{P}(X)$ be the collection of all subsets of a given set X . Then the equality relation, " $=$ ", and the inclusion relation, " \subseteq ", are an equivalence respectively an order relation on $\mathcal{P}(X)$.

A set X endowed with an order relation is called an *ordered set*. An order relation is usually denoted by " \leq ".

Definition 2 Let (X, \leq) be an ordered set and A a nonempty subset of X . An element $x \in X$ is called an upper bound (a lower bound) of A if

$$a \leq x \quad (x \leq a) \quad \text{for every } a \in A.$$

A set possessing a lower and an upper bound is called a bounded set.

The numbers a and b are called an lower (upper) bound of the the sequence $(a_n)_{n \geq 1}$

Definition 3 Let A be a bounded subset of an ordered set (X, \leq) . The greatest lower bound (the smallest upper bound) of A (if there exist) is called the infimum (supremum) of A and are denoted by $\inf A$ ($\sup A$).

If $\inf A$ or $\sup A$ belong to A , they are called the minimum respectively the maximum of A and are denoted by $\min A$ and $\max A$.

Example 1.2 Consider the set $A = \{1, \frac{1}{2}, \frac{1}{3}, \dots\}$. Then $\inf A = 0$ and $\sup A = 1$.

Definition 4 Let $f : X \rightarrow Y$ be a function, $A \subseteq X$, $B \subseteq Y$. The sets

$$\begin{aligned} f(A) &= \{f(x) | x \in A\} \\ f^{-1}(B) &= \{x \in X | f(x) \in B\} \end{aligned}$$

are called the image of A , respectively the counterimage of A through the function f . The set $f(X)$ is called the range of f .

Definition 5 A set \mathbb{R} endowed with the algebraic operations " $+$: $\mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$ ", " \cdot : $\mathbb{R} \times \mathbb{R} \rightarrow \mathbb{R}$," and a totally order relation " \leq " satisfying the following properties:

- a) $(\mathbb{R}, +, \cdot)$ is a commutative field;
- b)
 - If $x, y \in \mathbb{R}$, $x \leq y$, then $x + z \leq y + z$ for every $z \in \mathbb{R}$.
 - If $x, y \in \mathbb{R}$, $x \leq y$, then $xz \leq yz$ for every $z \geq 0$.
- c) Every nonempty and upper bounded subset of \mathbb{R} has a supremum;

is called a set of real numbers.

Remark 1.1 1) The first two conditions represent the algebraic part of Definition 5. The last condition, called the axiom of upper bound of Cantor-Dedekind, is the starting point for the basic results of mathematical analysis.

- 2) Some example of sets satisfying the conditions of Definition 5 are the set of infinite decimal fractions and the set of points of an ordered axe. One can prove that every two sets satisfying the conditions of Definition 5 are isomorphic.

If the set \mathbb{R} is given, then one defines the set \mathbb{N} of *natural numbers*, the set \mathbb{Z} of *integers* and the set \mathbb{Q} of *rational numbers* by:

$$\begin{aligned}\mathbb{N} &= \{0, 1, 2, \dots\} \\ \mathbb{Z} &= \{x \in \mathbb{R} | x \in \mathbb{N} \text{ or } -x \in \mathbb{N}\} \\ \mathbb{Q} &= \{xy^{-1} | x, y \in \mathbb{Z}, y \neq 0\}.\end{aligned}$$

One defines also the sets

$$\begin{aligned}\mathbb{N}^* &= \mathbb{N} \setminus \{0\} \\ \mathbb{R}_+ &= \{x \in \mathbb{R} | 0 \leq x\} \\ \mathbb{R}_+^* &= \{x \in \mathbb{R} | 0 < x\} \\ \mathbb{R}_- &= \{x \in \mathbb{R} | x \leq 0\} \\ \mathbb{R}_-^* &= \{x \in \mathbb{R} | x < 0\},\end{aligned}$$

where $x < y$ means $x \leq y$ and $x \neq y$.

The following characterizations of the supremum and the infimum of a subset of \mathbb{R} hold.

Proposition 1.1 *Let A be a nonempty subset of \mathbb{R} . Then $a = \sup A$ if and only if the following conditions are fulfilled:*

1. $x \leq a$ for every $x \in A$;
2. for every $\varepsilon > 0$, there exists $x_\varepsilon \in A$ such that $a - \varepsilon < x_\varepsilon$.

Proposition 1.2 *Let A be a nonempty subset of \mathbb{R} . Then $a = \inf A$ if and only if the following conditions are fulfilled:*

1. $a \leq x$ for every $x \in A$;
2. for every $\varepsilon > 0$, there exists $x_\varepsilon \in A$ such that $x_\varepsilon < a + \varepsilon$.

The proof of the previous propositions follows easily from the definition of the supremum and infimum of a set.

Proposition 1.3 *Every nonempty lower bounded subset of \mathbb{R} possesses an infimum.*

Proposition 1.4 *(Archimedes's axiom)*

For every $x, y \in \mathbb{R}_+^$ there exists $n \in \mathbb{N}$ such that $nx > y$.*

Proposition 1.5 *For every $x \in \mathbb{R}$ there exists a unique $n \in \mathbb{Z}$ such that $n \leq x < n + 1$.*

The number n from Proposition 1.5 is called the *integer part* of x and is denoted by $[x]$.

The *absolute value* of a real number x is defined by

$$|x| = \begin{cases} x, & \text{if } x \geq 0 \\ -x, & \text{if } x < 0. \end{cases}$$

For $a, b \in \mathbb{R}$, $a < b$ one defines the following bounded intervals:

$$\begin{aligned}(a, b) &= \{x \in \mathbb{R} | a < x < b\} \\ [a, b] &= \{x \in \mathbb{R} | a \leq x \leq b\} \\ [a, b) &= \{x \in \mathbb{R} | a \leq x < b\} \\ (a, b] &= \{x \in \mathbb{R} | a < x \leq b\}\end{aligned}$$

The sets

$$\begin{aligned}(a, +\infty) &= \{x \in \mathbb{R} | a < x\} \\ [a, +\infty) &= \{x \in \mathbb{R} | a \leq x\} \\ (-\infty, a) &= \{x \in \mathbb{R} | x < a\} \\ (-\infty, a] &= \{x \in \mathbb{R} | x \leq a\} \\ (-\infty, +\infty) &= \mathbb{R},\end{aligned}$$

are called *unbounded intervals*.

Definition 6 A set $V \subseteq \mathbb{R}$ is called a neighborhood of $x \in \mathbb{R}$ if there exists $\varepsilon > 0$ such that $(x - \varepsilon, x + \varepsilon) \subseteq V$.

By $\mathcal{V}(x)$ we denote the collection of all neighborhoods of $x \in \mathbb{R}$.

Definition 7 Let A be a subset of \mathbb{R} . An element $a \in A$ is called an interior point of A if there exists $V \in \mathcal{V}(a)$ such that $V \subseteq A$.

Definition 8 An element $x \in \mathbb{R}$ is called an cluster point (accumulation point) of a set $A \subseteq \mathbb{R}$ if for every $V \in \mathcal{V}(x)$ we have $V \cap (A \setminus \{x\}) \neq \emptyset$.

We denote by A' the set of all cluster points of A and by $\text{int } A$ the interior of the set A . By $\overline{\mathbb{R}}$ we denote the set defined by $\overline{\mathbb{R}} = \mathbb{R} \cup \{-\infty, +\infty\}$, where $-\infty$ and $+\infty$ are two elements satisfying the following properties of order:

$$-\infty < +\infty, \quad -\infty < x, \quad x < +\infty, \quad \text{for every } x \in \mathbb{R}.$$

We extend the operations " + " and " · " from \mathbb{R} to $\overline{\mathbb{R}}$ setting by definition:

$$\begin{aligned} x + (+\infty) &= (+\infty) + x = +\infty, & x \in \mathbb{R} \cup \{+\infty\} \\ x + (-\infty) &= (-\infty) + x = -\infty, & x \in \mathbb{R} \cup \{-\infty\} \\ x \cdot (+\infty) &= (+\infty) \cdot x = +\infty, & x \in \mathbb{R}_+^* \cup \{+\infty\} \\ x \cdot (-\infty) &= (-\infty) \cdot x = -\infty, & x \in \mathbb{R}_+^* \cup \{-\infty\} \\ \frac{x}{+\infty} &= \frac{x}{-\infty} = 0, & x \in \mathbb{R}. \end{aligned}$$

Definition 9 A set $V \subseteq \mathbb{R}$ is called a neighbourhood of $+\infty$ (or $-\infty$) if there exists $a \in \mathbb{R}$ with the property $(a, +\infty) \subseteq V$ (or $(-\infty, a) \subseteq V$).

The families of all neighbourhood of $+\infty$ and $-\infty$ are denoted by $\mathcal{V}(+\infty)$ and $\mathcal{V}(-\infty)$.

2 Sequences of real numbers

Definition 10 A function $f : \mathbb{N}^* \rightarrow \mathbb{R}$ is called sequence of real numbers.

Usually a sequence is denoted by $(a_n)_{n \geq 1}$ or (a_n) , where $a_n = f(n)$, $n \in \mathbb{N}^*$.

Definition 11 A sequence $(a_n)_{n \geq 1}$ is called monotonically increasing (decreasing) if

$$a_n \leq a_{n+1} \quad (a_n \geq a_{n+1}) \text{ for all } n \geq 1.$$

Definition 12 A sequence $(a_n)_{n \geq 1}$ is said to be bounded if there exists $a, b \in \mathbb{R}$ with the property

$$a \leq a_n \leq b \text{ for all } n \geq 1.$$

A sequence which is not bounded is said to be unbounded.

Definition 13 An element $a \in \overline{\mathbb{R}}$ is called the limit of a sequence $(a_n)_{n \geq 1}$ if for every $V \in \mathcal{V}(a)$ there exists $n_V \in \mathbb{N}^*$ such that $a_n \in V$ for every $n \geq n_V$.

If a sequence has a limit, then it is unique. We denote usually $a = \lim_{n \rightarrow \infty} a_n$ ($a = \lim a_n$) or $a_n \rightarrow a$.

A sequence having a finite limit is said to be *convergent*. If a sequence is not convergent it is called *divergent*. The following theorem is a characterization of convergent sequences.

Theorem 1 *A sequence $(a_n)_{n \geq 1}$ is convergent if and only if there exists $a \in \mathbb{R}$ such that for all $\varepsilon > 0$, there exists $n_\varepsilon \in \mathbb{N}^*$ with the property that for all $n \geq n_\varepsilon$ we have $|a_n - a| < \varepsilon$.*

Proof. Necessity Suppose that there exists $\lim_{n \rightarrow \infty} a_n = a$, $a \in \mathbb{R}$ and take $\varepsilon > 0$. Let $V = (a - \varepsilon, a + \varepsilon)$ be a neighbourhood of a . Then

$$\exists n_\varepsilon \in \mathbb{N} \text{ such that } a_n \in V, \forall n \geq n_\varepsilon \iff |a_n - a| < \varepsilon, \forall n \geq n_\varepsilon.$$

Sufficiency Let V be a neighbourhood of a . There exists $\varepsilon > 0$ such that $(a - \varepsilon, a + \varepsilon) \subseteq V$. It follows $a_n \in V$ for all $n \geq n_\varepsilon$, hence $\lim_{n \rightarrow \infty} a_n = a$. ■

The next result, known as Weierstrass theorem, is frequently used in the study of the convergence of a sequence.

Theorem 2 *Every monotonic and bounded sequence of real numbers is convergent.*

Proof. Suppose that $(a_n)_{n \geq 1}$ is an increasing and upper bounded sequence. Then the set $A = \{a_n | n \geq 1\}$ is upper bounded, hence there exists $a = \sup A$, $a \in \mathbb{R}$. Then for all $\varepsilon > 0$, $\exists n_\varepsilon \in \mathbb{N}$ such that $a - \varepsilon < a_{n_\varepsilon}$. Since $(a_n)_{n \geq 1}$ is monotonically increasing it follows $0 < a - a_n < \varepsilon$ for all $n \geq n_\varepsilon$. Hence $(a_n)_{n \geq 1}$ is convergent to a . ■

Theorem 3 *(Principle of nested intervals) Let $(I_n)_{n \geq 1}$, $I_n = [a_n, b_n]$, be a sequence of closed nested intervals, $I_{n+1} \subseteq I_n$, $\forall n \geq 1$. Then $\bigcap_{n \geq 1} I_n \neq \emptyset$. If $\lim_{n \rightarrow \infty} (b_n - a_n) = 0$, then $\bigcap_{n \geq 1} I_n$ is a singleton*

Proof. Define $A = \{a_n | n \in \mathbb{N}^*\}$, $B = \{b_n | n \in \mathbb{N}^*\}$. The relation $I_{n+1} \subseteq I_n$, $n \geq 1$, implies that $(a_n)_{n \geq 1}$ is increasing and $(b_n)_{n \geq 1}$ is decreasing. Since b_1

is an upper bound of A and a_1 is a lower bound of B there exists $\sup A = a$
 $\inf B = b$, $a, b \in \mathbb{R}$. By the relation

$$a_n \leq a \leq b \leq b_n, \quad n \geq 1,$$

it follows $[a, b] \subseteq \cap_{n \geq 1} I_n$. If $b_n - a_n \rightarrow 0$, clearly $a = b$. ■

Definition 14 Let $(a_n)_{n \geq 1}$ be a sequence and $(n_k)_{k \geq 1}$ a strictly increasing sequence of positive integers. Then $(a_{n_k})_{k \geq 1}$ is called a subsequence of $(a_n)_{n \geq 1}$.

Example 2.1 $(a_{2n})_{n \geq 1}, (a_{2n-1})_{n \geq 1}$ are subsequences of the sequence $(a_n)_{n \geq 1}$.

Corollary 2.1 (Cesàro lemma) A bounded sequence of real numbers contains at least a convergent subsequence.

Proof. Let $(x_n)_{n \geq 1}$ be a bounded sequence in \mathbb{R} . There exists $a, b \in \mathbb{R}$ such that $a \leq x_n \leq b$, $\forall n \geq 1$. Let $c = \frac{a+b}{2}$. From the intervals $[a, c]$, $[c, b]$ we choose the interval which contains an infinite number of terms of $(x_n)_{n \geq 1}$ and we denote it by $I_1 = [a_1, b_1]$. The set $\{n \in \mathbb{N}^* | x_n \in [a_1, b_1]\}$ is infinite and $b_1 - a_1 = \frac{b-a}{2}$. Let $c_1 = \frac{a_1+b_1}{2}$. From the intervals $[a_1, c_1]$, $[c_1, b_1]$ we choose the interval which contains an infinite number of terms of $(x_n)_{n \geq 1}$ and we denote it by $I_2 = [a_2, b_2]$. We have $b_2 - a_2 = \frac{b-a}{2^2}$.

Denoting $I_0 = [a, b]$ one obtains a sequence $I_n = [a_n, b_n]$, $n \geq 0$, with the property that $I_{n+1} \subseteq I_n$ and $b_n - a_n = \frac{b-a}{2^n}$, $n \geq 0$. Taking account of Theorem 3 it follows $\cap_{n \geq 0} I_n = \{x\}$, $x \in [a, b]$. Now we choose from every interval I_k a term $x_{n_k} \in I_k$, $k \geq 0$. We have $|x_{n_k} - x| \leq \frac{b-a}{2^k}$, $k \geq 0$, hence $\lim_{k \rightarrow \infty} x_{n_k} = x$. ■

Cesaro lemma is also called as Bolzano-Weierstrass Theorem.

Definition 15 A sequence $(a_n)_{n \geq 1}$ is said to be fundamental or Cauchy sequence if for every $\varepsilon > 0$ there exists $n_\varepsilon \in \mathbb{N}^*$ such that for all $m, n \geq n_\varepsilon$ we have $|a_m - a_n| < \varepsilon$.

The above definition is equivalent to the next one:

A sequence $(a_n)_{n \geq 1}$ is *fundamental* if for every $\varepsilon > 0$, there exists $n_\varepsilon \in \mathbb{N}^*$ such that $\forall n \geq n_\varepsilon, \forall p \in \mathbb{N} |a_{n+p} - a_n| < \varepsilon$.

Remark 2.1 *The notion of Cauchy sequence was introduced to obtain a characterization convergent sequences without using their limit.*

Theorem 4 *A sequence of real numbers is convergent if and only if it is a Cauchy sequence.*

Proof. Necessity Suppose that $(a_n)_{n \geq 1}$ is convergent and let $\lim_{n \rightarrow \infty} a_n = a$, $a \in \mathbb{R}$.

Let $\varepsilon > 0$ be a fixed number. Then there exists $n_\varepsilon \in \mathbb{N}^*$ such that for all $n \geq n_\varepsilon$ $|a_n - a| < \frac{\varepsilon}{2}$. Now, for all $m, n \geq n_\varepsilon$ we have

$$|a_m - a_n| \leq |a_m - a| + |a - a_n| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon,$$

therefore $(a_n)_{n \geq 1}$ is a Cauchy sequence.

Sufficiency Suppose $(a_n)_{n \geq 1}$ is a Cauchy sequence.

- 1) We prove first that $(a_n)_{n \geq 1}$ is bounded. For $\varepsilon = 1$ in Definition 15 follows that $\exists n_1 \in \mathbb{N}$ such that $|a_n - a_{n_1}| < 1$, $n \geq n_1$. Then

$$|a_n| \leq |a_n - a_{n_1}| + |a_{n_1}| < 1 + |a_{n_1}|, \quad \forall n \geq n_1,$$

hence $(a_n)_{n \geq 1}$ is a bounded sequence.

- 2) We prove that $(a_n)_{n \geq 1}$ is a convergent sequence. From the boundedness of $(a_n)_{n \geq 1}$ it follows that there exists a convergent subsequence $(a_{n_k})_{k \geq 1}$, $\lim_{n \rightarrow \infty} a_n = a$. Let $\varepsilon > 0$ be a positive number. By the convergence of $(a_{n_k})_{k \geq 1}$ we find $k_\varepsilon \in \mathbb{N}$ with the property

$$(2.1) \quad |a_{n_k} - a| < \frac{\varepsilon}{2}, \quad \forall k \geq k_\varepsilon.$$

Since $(a_n)_{n \geq 1}$ is a Cauchy sequence it follows that there exists $n_\varepsilon \in \mathbb{N}^*$ with the property

$$(2.2) \quad |a_m - a_n| < \frac{\varepsilon}{2}, \quad \forall m, n \geq n_\varepsilon.$$

Now take $N_\varepsilon = \max\{n_\varepsilon, n_{k_\varepsilon}\}$. For every $n \geq N_\varepsilon$ we have in view of the relations (2.1) and (2.2)

$$|a_n - a| \leq |a_n - a_{n_{k_\varepsilon}}| + |a_{n_{k_\varepsilon}} - a| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon,$$

hence $(a_n)_{n \geq 1}$ is a convergent sequence.

■

Example 2.2 Prove that $(a_n)_{n \geq 1}$,

$$a_n = \frac{\sin 1}{1^2} + \frac{\sin 2}{2^2} + \cdots + \frac{\sin n}{n^2}$$

is a convergent sequence.

Solution. We prove that $(a_n)_{n \geq 1}$ is a Cauchy sequence. For every $n, p \in \mathbb{N}^*$ we have:

$$\begin{aligned} |a_{n+p} - a_n| &= \left| \frac{\sin(n+1)}{(n+1)^2} + \frac{\sin(n+2)}{(n+2)^2} + \cdots + \frac{\sin(n+p)}{(n+p)^2} \right| \\ &\leq \left| \frac{\sin(n+1)}{(n+1)^2} \right| + \left| \frac{\sin(n+2)}{(n+2)^2} \right| + \cdots + \left| \frac{\sin(n+p)}{(n+p)^2} \right| \\ &\leq \frac{1}{(n+1)^2} + \frac{1}{(n+2)^2} + \cdots + \frac{1}{(n+p)^2} \\ &\leq \frac{1}{n(n+1)} + \frac{1}{(n+1)(n+2)} + \cdots + \frac{1}{(n+p-1)(n+p)} \\ &= \left(\frac{1}{n} - \frac{1}{n+1} \right) + \left(\frac{1}{n+1} - \frac{1}{n+2} \right) + \cdots + \left(\frac{1}{n+p-1} - \frac{1}{n+p} \right) \\ &= \frac{1}{n} - \frac{1}{n+p} < \frac{1}{n}. \end{aligned}$$

For every $\varepsilon > 0$, $\exists n_\varepsilon \in \mathbb{N}^*$ such that $\frac{1}{n} < \varepsilon$, for all $n \geq n_\varepsilon$. Thus

$$|a_{n+p} - a_n| \leq \varepsilon, \quad \forall n \geq n_\varepsilon, p \geq 1.$$

Since $(a_n)_{n \geq 1}$ is a fundamental sequence it is convergent. ■

Example 2.3 Prove that $\lim_{n \rightarrow \infty} \left(1 + \frac{1}{2} + \cdots + \frac{1}{n} \right) = \infty$.

Solution. Define $(a_n)_{n \geq 1}$, by

$$a_n = 1 + \frac{1}{2} + \cdots + \frac{1}{n}.$$

Since $(a_n)_{n \geq 1}$ is monotonically increasing there exists $\lim a_n \in \overline{\mathbb{R}}$. We prove that $(a_n)_{n \geq 1}$ is not a Cauchy sequence. For every $n \geq 1$ we have

$$\begin{aligned} |a_{2n} - a_n| &= \frac{1}{n+1} + \frac{1}{n+2} + \cdots + \frac{1}{2n} \\ &\geq \underbrace{\frac{1}{2n} + \frac{1}{2n} + \cdots + \frac{1}{2n}}_{n \text{ times}} = \frac{1}{2} \end{aligned}$$

Since $(a_n)_{n \geq 1}$ is divergent and its limit exists, we conclude that $\lim_{n \rightarrow \infty} a_n = +\infty$. ■

3 The upper limit and the lower limit of a sequence

Definition 16 Let $(a_n)_{n \geq 1}$ be a sequence of real numbers. We call $a \in \overline{\mathbb{R}}$ a limit point of the sequence $(a_n)_{n \geq 1}$ if every neighbourhood of a contains infinitely many terms of $(a_n)_{n \geq 1}$.

For a sequence $(a_n)_{n \geq 1}$ we denote by $L(a_n)$ the set of its limit points.

Example 3.1 a) For $(a_n)_{n \geq 1}$, $a_n = (-1)^n$, $L(a_n) = \{-1, 1\}$.

b) The sequence $(a_n)_{n \geq 1}$, $a_n = \sin \frac{n\pi}{2}$, has $L(a_n) = \{-1, 0, 1\}$.

Theorem 5 The element $a \in \overline{\mathbb{R}}$ is a limit point of $(a_n)_{n \geq 1}$ if and only if there exists a subsequence $(a_{n_k})_{k \geq 1}$ of $(a_n)_{n \geq 1}$ with $\lim_{k \rightarrow \infty} a_{n_k} = a$.

Proof. Necessity Suppose that $a \in L(a_n)$ and $a \in \mathbb{R}$. Let $V_k = (a - \frac{1}{k}, a + \frac{1}{k})$, $k \in \mathbb{N}^*$, be a sequence of neighbourhood of a . Every V_k contains infinitely many terms of (a_n) . Let $a_{n_1} \in V_1$, $a_{n_2} \in V_2, \dots$, with $n_1 < n_2 < \dots$. We have

$$|a_{n_k} - a| < \frac{1}{k}, \quad k \geq 1,$$

hence $\lim_{k \rightarrow \infty} a_{n_k} = a$. Analogously follows the existence of $(a_{n_k})_{k \geq 1}$ for $a \in \{-\infty, +\infty\}$.

Sufficiency

Suppose that $(a_{n_k})_{k \geq 1}$ is a subsequence of $(a_n)_{n \geq 1}$ with $\lim_{k \rightarrow \infty} a_{n_k} = a$. Then every neighbourhood of a contains infinitely many elements of $(a_n)_{n \geq 1}$, thus $a \in L(a_n)$. ■

Definition 17 Let $(a_n)_{n \geq 1}$ be a sequence of real numbers. Then $\sup L(a_n)$ is called the upper limit of $(a_n)_{n \geq 1}$ and $\inf L(a_n)$ is called the lower limit of $(a_n)_{n \geq 1}$.

We denote the upper limit and the lower limit of $(a_n)_{n \geq 1}$ by

$$\overline{\lim} a_n, \quad \text{respectively} \quad \underline{\lim} a_n,$$

or by

$$\limsup a_n, \quad \text{respectively} \quad \liminf a_n,$$

Example 3.2 a) For the sequence $(a_n)_{n \geq 1}$, defined by

$$a_n := 1, 1, \frac{1}{2}, 1, \frac{1}{2}, \frac{1}{3}, 1, \frac{1}{2}, \frac{1}{3}, \frac{1}{4}, \dots,$$

we get $L(a_n) = \{1, \frac{1}{2}, \frac{1}{3}, \dots\} \cup \{0\}$ therefore $\overline{\lim} a_n = 1$ and $\underline{\lim} a_n = 0$.

b) It can be proved that the sequence $(a_n)_{n \geq 1}$, $a_n = \sin n$, has $L(a_n) = [-1, 1]$, $\overline{\lim} a_n = 1$ and $\underline{\lim} a_n = -1$.

Theorem 6 A sequence $(a_n)_{n \geq 1}$ has a limit if and only if

$$\overline{\lim} a_n = \underline{\lim} a_n.$$

4 Some remarkable sequences

Proposition 4.1 (the number e) The sequence $(e_n)_{n \geq 1}$,

$$e_n = \left(1 + \frac{1}{n}\right)^n,$$

is monotonically increasing and bounded above. By definition

$$e = \lim_{n \rightarrow \infty} \left(1 + \frac{1}{n}\right)^n.$$

Proof. Define $(e'_n)_{n \geq 1}$, $e'_n = \left(1 + \frac{1}{n}\right)^{n+1}$. We prove that $(e'_n)_{n \geq 1}$ is monotonically decreasing. We have

$$\begin{aligned} \frac{e'_n}{e'_{n+1}} &= \left(\frac{n+1}{n}\right)^{n+1} \left(\frac{n+1}{n+2}\right)^{n+2} = \left(\frac{(n+1)^2}{n^2+2n}\right)^{n+1} \cdot \frac{n+1}{n+2} \\ &= \left(1 + \frac{1}{n^2+2n}\right)^{n+1} \cdot \frac{n+1}{n+2} > \left(1 + (n+1)\frac{1}{n^2+2n}\right) \cdot \frac{n+1}{n+2} \\ &= \frac{n(n+2)^2+1}{n(n+2)^2} > 1, \quad \forall n \geq 1, \end{aligned}$$

taking account of Bernoulli's inequality:

$$(1+t)^n > 1+nt, \quad t \in (-1, \infty) \setminus \{0\}, n \in \mathbb{N}^*.$$

We prove that $(e_n)_{n \geq 1}$ is an increasing sequence.

$$\begin{aligned} \frac{e_{n+1}}{e_n} &= \left(\frac{n+2}{n+1}\right)^{n+1} \left(\frac{n}{n+1}\right)^n = \left(\frac{(n+2)}{n+1}\right)^{n+1} \frac{n+1}{n} \\ &= \left(\frac{n^2+2n}{(n+1)^2}\right)^{n+1} \frac{n+1}{n} = \left(1 - \frac{1}{(n+1)^2}\right)^{n+1} \frac{n+1}{n} \\ &> \left(1 - (n+1)\frac{1}{(n+1)^2}\right) \frac{n+1}{n} = 1, \quad \forall n \geq 1, \end{aligned}$$

By the relation

$$e_1 < e_2 < \cdots < e_{n+1} < \cdots < e'_{n+1} < e'_n < \cdots < e'_2 < e'_1$$

follows that (e'_n) , (e_n) are convergent sequences and by

$$0 < e'_n - e_n = \left(1 + \frac{1}{n}\right)^n \frac{1}{n} < \frac{e_n}{n} \leq \frac{e_1}{n}, \quad n \geq 1,$$

one gets

$$\lim e_n = \lim e'_n := e.$$

■

Corollary 4.1 *The following inequality holds*

$$\left(1 + \frac{1}{n}\right)^n < e < \left(1 + \frac{1}{n}\right)^{n+1}, \quad \forall n \in \mathbb{N}^*.$$

Remark 4.1 *The sequence $(E_n)_{n \geq 1}$,*

$$E_n = 1 + \frac{1}{1!} + \frac{1}{2!} + \cdots + \frac{1}{n!}$$

is monotonically increasing and $\lim_{n \rightarrow \infty} E_n = e$ ($e \notin \mathbb{Q}$, $e \simeq 2,71828\dots$).

Proposition 4.2 *(Euler's constant) The sequence $(\gamma_n)_{n \geq 1}$,*

$$\gamma_n = 1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \ln n$$

is monotonically decreasing and bounded below. We denote

$$\lim_{n \rightarrow \infty} \gamma_n := \gamma$$

(γ is called Euler's constant and $\gamma = 0,577\dots$).

Proof. From Corollary 4.1 follows

$$(4.1) \quad \frac{1}{n+1} < \ln(n+1) - \ln n < \frac{1}{n}, \quad \forall n \geq 1.$$

Summing the previous inequalities one gets

$$(4.2) \quad \sum_{k=1}^n \frac{1}{k+1} < \ln(n+1) < \sum_{k=1}^n \frac{1}{k}, \quad \forall n \geq 1.$$

We have

$$\gamma_{n+1} - \gamma_n = \frac{1}{n+1} - \ln(n+1) + \ln n$$

and taking account of (4.1) follows $\gamma_{n+1} < \gamma_n$, $n \geq 1$.

Now, in view of inequality (4.2) we get

$$1 + \frac{1}{2} + \frac{1}{3} + \cdots + \frac{1}{n} - \ln n > \ln(n+1) - \ln n > 0, \quad n \geq 1.$$

■

The following theorems are often used for finding the limit of a sequence.

Theorem 7 (Sandwich Theorem) Let $(a_n)_{n \geq 1}$, $(b_n)_{n \geq 1}$, $(c_n)_{n \geq 1}$ be sequences of real numbers with the property that

$$a_n \leq b_n \leq c_n \text{ for all } n \geq n_0.$$

If

$$\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} c_n = l, l \in \overline{\mathbb{R}},$$

then $\lim_{n \rightarrow \infty} b_n = l$.

Theorem 8 (Stolz-Cesaro I) Let $(a_n)_{n \geq 1}$, $(b_n)_{n \geq 1}$ be two sequences of real numbers satisfying the conditions:

- 1) $(b_n)_{n \geq 1}$ is strictly monotone and unbounded;
- 2) there exists $\lim_{n \rightarrow \infty} \frac{a_{n+1} - a_n}{b_{n+1} - b_n} = l, l \in \overline{\mathbb{R}};$

Then $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = l$.

Theorem 9 (Stolz-Cesaro II) Let $(a_n)_{n \geq 1}$, $(b_n)_{n \geq 1}$ be two sequences of real numbers satisfying the conditions:

- 1) $\lim_{n \rightarrow \infty} a_n = \lim_{n \rightarrow \infty} b_n = 0;$
- 2) $(b_n)_{n \geq 1}$ is strictly monotone;
- 3) there exists $\lim_{n \rightarrow \infty} \frac{a_{n+1} - a_n}{b_{n+1} - b_n} = l, l \in \overline{\mathbb{R}};$

Then $\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = l$.

Proof. We prove only Theorem 9. Suppose that $(b_n)_{n \geq 1}$ is monotonically increasing and $l \in \mathbb{R}$. Let ε be a positive number. There exists $n_\varepsilon \in \mathbb{N}$ such that for all $n \geq n_\varepsilon$

$$\left| \frac{a_{n+1} - a_n}{b_{n+1} - b_n} - l \right| < \varepsilon$$

or

$$(b_{n+1} - b_n)(l - \varepsilon) < a_{n+1} - a_n < (b_{n+1} - b_n)(l + \varepsilon), \quad n \geq n_\varepsilon.$$

Adding the previous relations from n to $n + p - 1$ follows

$$\begin{aligned} (l - \varepsilon) \sum_{k=n}^{n+p-1} (b_{k+1} - b_k) &< \sum_{k=n}^{n+p-1} (a_{k+1} - a_k) < (l + \varepsilon) \sum_{k=n}^{n+p-1} (b_{k+1} - b_k) \\ (l - \varepsilon)(b_{n+p} - b_n) &< a_{n+p} - a_n < (l + \varepsilon)(b_{n+p} - b_n) \\ -\varepsilon &< \frac{a_{n+p} - a_n}{b_{n+p} - b_n} - l < \varepsilon, \end{aligned}$$

for all $n \geq n_\varepsilon, p \geq 1$. Letting $p \rightarrow \infty$ in the previous relation one gets

$$-\varepsilon \leq \frac{a_n}{b_n} - l \leq \varepsilon, \quad n \geq n_\varepsilon,$$

hence

$$\lim_{n \rightarrow \infty} \frac{a_n}{b_n} = l.$$

■

Corollary 4.2 *Let $(a_n)_{n \geq 1}$, be a sequence of positive numbers with the property that there exists $\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n} = l, l \in \mathbb{R}$. Then $\lim_{n \rightarrow \infty} \sqrt[n]{a_n} = l$.*

Proof.

$$\begin{aligned} \lim_{n \rightarrow \infty} \sqrt[n]{a_n} &= \lim_{n \rightarrow \infty} e^{\ln \sqrt[n]{a_n}} = e^{\lim_{n \rightarrow \infty} \frac{\ln a_n}{n}} \\ &= e^{\lim_{n \rightarrow \infty} \frac{\ln a_{n+1} - \ln a_n}{n+1-n}} = e^{\lim_{n \rightarrow \infty} \frac{a_{n+1}}{a_n}} = e^{\ln l} = l. \end{aligned}$$

■

5 Sequences of complex numbers

A sequence of complex numbers $(z_n)_{n \geq 1}$ has the form

$$z_n = a_n + ib_n, \quad n \geq 1,$$

where $(a_n)_{n \geq 1}$ and $(b_n)_{n \geq 1}$ are sequences of real numbers, i.e., $a_n = \Re z_n$, $b_n = \Im z_n$. Many notions and theorems carry over from real to complex sequences.

Definition 18 *A sequence $(z_n)_{n \geq 1}$ of complex numbers is said to be convergent to $z \in \mathbb{C}$ if for every $\varepsilon > 0$ there exists $n_\varepsilon \in \mathbb{N}^*$ such that $|z_n - z| < \varepsilon$ for every $n \geq n_\varepsilon$.*

Definition 19 *A sequence $(z_n)_{n \geq 1}$ of complex numbers is said to be fundamental (or Cauchy sequence) if for every $\varepsilon > 0$ there exists $n_\varepsilon \in \mathbb{N}^*$ such that for all $m, n \geq n_\varepsilon$ we have $|z_m - z_n| < \varepsilon$.*

Theorem 10 *A sequence $(z_n)_{n \geq 1}$ of complex numbers converges to $z \in \mathbb{C}$ if and only if the real sequences $(\Re z_n)_{n \geq 1}$ and $(\Im z_n)_{n \geq 1}$ converge to $\Re z$ and $\Im z$ respectively.*

Proof. Consider $z_n = a_n + ib_n$, $n \geq 1$, $z = a + bi$, $a, b, a_n, b_n \in \mathbb{R}$.

Necessity

Suppose that $z_n \rightarrow z$ and let $\varepsilon > 0$ be given. Then $\exists n_\varepsilon \in \mathbb{N}^*$ such that

$$|z_n - z| < \varepsilon \quad \forall n \geq n_\varepsilon.$$

We have

$$\begin{aligned} |a_n - a| &< |z_n - z| < \varepsilon, \\ |b_n - b| &< |z_n - z| < \varepsilon, \quad \forall n \geq n_\varepsilon, \end{aligned}$$

thus $a_n \rightarrow a$, $b_n \rightarrow b$.

Sufficiency Suppose $a_n \rightarrow a$, $b_n \rightarrow b$ and prove that $z_n \rightarrow z$. Let $\varepsilon > 0$ be given. Then there exist $n'_\varepsilon, n''_\varepsilon \in \mathbb{N}^*$ such that

$$\begin{aligned} |a_n - a| &< \frac{\varepsilon}{2}, \quad \forall n \geq n'_\varepsilon, \\ |b_n - b| &< \frac{\varepsilon}{2}, \quad \forall n \geq n''_\varepsilon. \end{aligned}$$

Taking $n_\varepsilon = \max\{n'_\varepsilon, n''_\varepsilon\}$ it follows

$$|z_n - z| < |a_n - a| + |b_n - b| < \frac{\varepsilon}{2} + \frac{\varepsilon}{2} = \varepsilon, \quad \forall n \geq n_\varepsilon,$$

hence $\lim_{n \rightarrow \infty} z_n = z$. ■

As for real sequences we can prove.

Theorem 11 *A sequence of complex numbers is convergent if and only if it is a Cauchy sequence.*

Since the complex field \mathbb{C} is not an ordered field, all notions and proprieties where the order is involved do not make sense for complex sequences or they need modifications. The sandwich theorem does not hold; there is no notion of monotone sequence, upper and lower limit. Nevertheless the notion of boundedness, subsequence and limit points do hold in the complex case too.