

Nonlinear Programming

- Let f be a real valued function defined on $\Omega \subseteq \mathbb{R}^n$.

Nonlinear Programming

- Let f be a real valued function defined on $\Omega \subseteq \mathbb{R}^n$.
- The problem is to minimize or maximize $f(\mathbf{x})$ subject to $\mathbf{x} \in \Omega$, where f need not be a linear function.

Nonlinear Programming

- Let f be a real valued function defined on $\Omega \subseteq \mathbb{R}^n$.
- The problem is to minimize or maximize $f(\mathbf{x})$ subject to $\mathbf{x} \in \Omega$, where f need not be a linear function.
- **Definition 1:** A point (or an element) $\mathbf{x}^* \in \Omega$ is called a local minimum (maximum) of f if there exists an $\epsilon > 0$, such that $\mathbf{x} \in \Omega$ and $\|\mathbf{x} - \mathbf{x}^*\| < \epsilon$ implies $f(\mathbf{x}^*) \leq f(\mathbf{x})$ ($f(\mathbf{x}^*) \geq f(\mathbf{x})$) .

Nonlinear Programming

- Let f be a real valued function defined on $\Omega \subseteq \mathbb{R}^n$.
- The problem is to minimize or maximize $f(\mathbf{x})$ subject to $\mathbf{x} \in \Omega$, where f need not be a linear function.
- **Definition 1:** A point (or an element) $\mathbf{x}^* \in \Omega$ is called a local minimum (maximum) of f if there exists an $\epsilon > 0$, such that $\mathbf{x} \in \Omega$ and $\|\mathbf{x} - \mathbf{x}^*\| < \epsilon$ implies $f(\mathbf{x}^*) \leq f(\mathbf{x})$ ($f(\mathbf{x}^*) \geq f(\mathbf{x})$) .
- **Definition 2:** A point $\mathbf{x}^* \in \Omega$ is called a global minimum (maximum) of f if $f(\mathbf{x}^*) \leq f(\mathbf{x})$ ($f(\mathbf{x}^*) \geq f(\mathbf{x})$) for all $\mathbf{x} \in \Omega$.

Nonlinear Programming

- Let f be a real valued function defined on $\Omega \subseteq \mathbb{R}^n$.
- The problem is to minimize or maximize $f(\mathbf{x})$ subject to $\mathbf{x} \in \Omega$, where f need not be a linear function.
- **Definition 1:** A point (or an element) $\mathbf{x}^* \in \Omega$ is called a local minimum (maximum) of f if there exists an $\epsilon > 0$, such that $\mathbf{x} \in \Omega$ and $\|\mathbf{x} - \mathbf{x}^*\| < \epsilon$ implies $f(\mathbf{x}^*) \leq f(\mathbf{x})$ ($f(\mathbf{x}^*) \geq f(\mathbf{x})$) .
- **Definition 2:** A point $\mathbf{x}^* \in \Omega$ is called a global minimum (maximum) of f if $f(\mathbf{x}^*) \leq f(\mathbf{x})$ ($f(\mathbf{x}^*) \geq f(\mathbf{x})$) for all $\mathbf{x} \in \Omega$.

- **Definition 3:** A vector $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is said to be a **feasible direction** at $\mathbf{x}^* \in \Omega$, if there exists a $c > 0$ such that for all t , $0 \leq t \leq c$, $\mathbf{x}^* + t\mathbf{d} \in \Omega$.

- **Definition 3:** A vector $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is said to be a **feasible direction** at $\mathbf{x}^* \in \Omega$, if there exists a $c > 0$ such that for all t , $0 \leq t \leq c$, $\mathbf{x}^* + t\mathbf{d} \in \Omega$.
- **Example 1:** Let $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.

- **Definition 3:** A vector $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is said to be a **feasible direction** at $\mathbf{x}^* \in \Omega$, if there exists a $c > 0$ such that for all t , $0 \leq t \leq c$, $\mathbf{x}^* + t\mathbf{d} \in \Omega$.
- **Example 1:** Let $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- At $[0, 0]^T$ if $\mathbf{d} = [d_1, d_2]^T$ is a feasible direction then $d_1 \geq 0$ and $d_2 \geq 0$.

- **Definition 3:** A vector $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is said to be a **feasible direction** at $\mathbf{x}^* \in \Omega$, if there exists a $c > 0$ such that for all t , $0 \leq t \leq c$, $\mathbf{x}^* + t\mathbf{d} \in \Omega$.
- **Example 1:** Let $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- At $[0, 0]^T$ if $\mathbf{d} = [d_1, d_2]^T$ is a feasible direction then $d_1 \geq 0$ and $d_2 \geq 0$.
- At $[0, \frac{1}{2}]^T$ if \mathbf{d} is a feasible direction then $d_1 \geq 0$ but d_2 can be any real number.

- **Definition 3:** A vector $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is said to be a **feasible direction** at $\mathbf{x}^* \in \Omega$, if there exists a $c > 0$ such that for all t , $0 \leq t \leq c$, $\mathbf{x}^* + t\mathbf{d} \in \Omega$.
- **Example 1:** Let $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- At $[0, 0]^T$ if $\mathbf{d} = [d_1, d_2]^T$ is a feasible direction then $d_1 \geq 0$ and $d_2 \geq 0$.
- At $[0, \frac{1}{2}]^T$ if \mathbf{d} is a feasible direction then $d_1 \geq 0$ but d_2 can be any real number.
- At $[\frac{1}{2}, 0]^T$ if \mathbf{d} is a feasible direction then $d_2 \geq 0$ but d_1 can be any real number.

- **Definition 3:** A vector $\mathbf{d} \in \mathbb{R}^n$, $\mathbf{d} \neq \mathbf{0}$ is said to be a **feasible direction** at $\mathbf{x}^* \in \Omega$, if there exists a $c > 0$ such that for all t , $0 \leq t \leq c$, $\mathbf{x}^* + t\mathbf{d} \in \Omega$.
- **Example 1:** Let $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- At $[0, 0]^T$ if $\mathbf{d} = [d_1, d_2]^T$ is a feasible direction then $d_1 \geq 0$ and $d_2 \geq 0$.
- At $[0, \frac{1}{2}]^T$ if \mathbf{d} is a feasible direction then $d_1 \geq 0$ but d_2 can be any real number.
- At $[\frac{1}{2}, 0]^T$ if \mathbf{d} is a feasible direction then $d_2 \geq 0$ but d_1 can be any real number.
- At $[\frac{1}{2}, \frac{1}{2}]^T$ any $\mathbf{d} \in \mathbb{R}^2$ will be a feasible direction.

First order necessary conditions for a local minimum

- **Remark 1:** If \mathbf{x}^* is an interior point of Ω then any $\mathbf{d} \in \mathbb{R}^n, \mathbf{d} \neq \mathbf{0}$ is a feasible direction at \mathbf{x}^* .

First order necessary conditions for a local minimum

- **Remark 1:** If \mathbf{x}^* is an interior point of Ω then any $\mathbf{d} \in \mathbb{R}^n, \mathbf{d} \neq \mathbf{0}$ is a feasible direction at \mathbf{x}^* .
- **Theorem 1:** Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function (that is, the first order partial derivatives of f exists and are continuous as functions from Ω to \mathbb{R}). If \mathbf{x}^* is a local minimum point then for any feasible direction \mathbf{d} at \mathbf{x}^* , $\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0$,

First order necessary conditions for a local minimum

- **Remark 1:** If \mathbf{x}^* is an interior point of Ω then any $\mathbf{d} \in \mathbb{R}^n, \mathbf{d} \neq \mathbf{0}$ is a feasible direction at \mathbf{x}^* .
- **Theorem 1:** Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function (that is, the first order partial derivatives of f exists and are continuous as functions from Ω to \mathbb{R}). If \mathbf{x}^* is a local minimum point then for any feasible direction \mathbf{d} at \mathbf{x}^* , $\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0$, where $(\nabla f(\mathbf{x}^*))_i = (\frac{\partial f}{\partial x_i})(\mathbf{x}^*)$ (the gradient vector of f at \mathbf{x}^* is written as a row vector), and $\mathbf{d} \in \mathbb{R}^n$ is a column vector.

First order necessary conditions for a local minimum

- **Remark 1:** If \mathbf{x}^* is an interior point of Ω then any $\mathbf{d} \in \mathbb{R}^n, \mathbf{d} \neq \mathbf{0}$ is a feasible direction at \mathbf{x}^* .
- **Theorem 1:** Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function (that is, the first order partial derivatives of f exists and are continuous as functions from Ω to \mathbb{R}). If \mathbf{x}^* is a local minimum point then for any feasible direction \mathbf{d} at \mathbf{x}^* , $\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0$,
where $(\nabla f(\mathbf{x}^*))_i = (\frac{\partial f}{\partial x_i})(\mathbf{x}^*)$ (the gradient vector of f at \mathbf{x}^* is written as a row vector), and $\mathbf{d} \in \mathbb{R}^n$ is a column vector.
- **Theorem 2 :** Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function. Let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum point of f then $\nabla f(\mathbf{x}^*) = \mathbf{0}$

- **Example 1:** Consider the following problem:

- **Example 1:** Consider the following problem:

$$\text{Minimize } f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$$

- **Example 1:** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$

subject to $x_1 \geq 0, x_2 \geq 0$, hence

$\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.

- **Example 1:** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$

subject to $x_1 \geq 0, x_2 \geq 0$, hence

$\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.

- $\nabla f(\mathbf{x}) = (3x_1^2 - 2x_1 x_2, -x_1^2 + 4x_2) = [0, 0]$ has two solutions $x_1 = 0, x_2 = 0$ and $x_1 = 6, x_2 = 9$.

- **Example 1:** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$
subject to $x_1 \geq 0, x_2 \geq 0$, hence
 $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- $\nabla f(\mathbf{x}) = (3x_1^2 - 2x_1 x_2, -x_1^2 + 4x_2) = [0, 0]$ has two solutions $x_1 = 0, x_2 = 0$ and $x_1 = 6, x_2 = 9$.
- So $[6, 9]^T$ and $[0, 0]^T$ satisfies the **first order necessary conditions** for a local minimum.

- **Example 1:** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$

subject to $x_1 \geq 0, x_2 \geq 0$, hence

$\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.

- $\nabla f(\mathbf{x}) = (3x_1^2 - 2x_1 x_2, -x_1^2 + 4x_2) = [0, 0]$ has two solutions $x_1 = 0, x_2 = 0$ and $x_1 = 6, x_2 = 9$.
- So $[6, 9]^T$ and $[0, 0]^T$ satisfies the **first order necessary conditions** for a local minimum.
- $[0, 0]^T$ is also a local minimum of f .

- **Example 1:** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$

subject to $x_1 \geq 0, x_2 \geq 0$, hence

$\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.

- $\nabla f(\mathbf{x}) = (3x_1^2 - 2x_1 x_2, -x_1^2 + 4x_2) = [0, 0]$ has two solutions $x_1 = 0, x_2 = 0$ and $x_1 = 6, x_2 = 9$.
- So $[6, 9]^T$ and $[0, 0]^T$ satisfies the **first order necessary conditions** for a local minimum.
- $[0, 0]^T$ is also a local minimum of f .
- **Exercise:** Will $[0, 0]^T$ be a local minimum point of f given in **Example 1**, if $\Omega = \mathbb{R}^2$?

- **Example 2 :** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$

- **Example 2 :** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$
subject to $x_1 \geq 0, x_2 \geq 0$.

- **Example 2 :** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$
subject to $x_1 \geq 0, x_2 \geq 0$.

- Hence $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$

- **Example 2 :** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$
subject to $x_1 \geq 0, x_2 \geq 0$.

- Hence $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$
- Note that f is a continuously differentiable function

- **Example 2 :** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$
subject to $x_1 \geq 0, x_2 \geq 0$.
- Hence $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$
- Note that f is a continuously differentiable function
- Check that $\mathbf{x}^* = [\frac{1}{2}, 0]^T$, satisfies the **first order necessary conditions** for a locally minimum

- **Example 2 :** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$
subject to $x_1 \geq 0, x_2 \geq 0$.
- Hence $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$
- Note that f is a continuously differentiable function
- Check that $\mathbf{x}^* = [\frac{1}{2}, 0]^T$, satisfies the **first order necessary conditions** for a locally minimum
- Check that f has a global minimum at $x_1 = \frac{1}{2}, x_2 = 0$

Second order necessary conditions for a point to be a local minimum

- **Theorem 3:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function (that is all the second order partial derivatives of f (given by $\frac{\partial^2 f}{\partial x_j \partial x_i}$) exists and are continuous as functions from Ω to \mathbb{R}).

Second order necessary conditions for a point to be a local minimum

- **Theorem 3:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function (that is all the second order partial derivatives of f (given by $\frac{\partial^2 f}{\partial x_j \partial x_i}$) exists and are continuous as functions from Ω to \mathbb{R}).
If \mathbf{x}^* is a local minimum of f then for any feasible direction \mathbf{d} at \mathbf{x}^*

Second order necessary conditions for a point to be a local minimum

- **Theorem 3:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function (that is all the second order partial derivatives of f (given by $\frac{\partial^2 f}{\partial x_j \partial x_i}$) exists and are continuous as functions from Ω to \mathbb{R}).
If \mathbf{x}^* is a local minimum of f then for any feasible direction \mathbf{d} at \mathbf{x}^*
 1. $\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0$.

Second order necessary conditions for a point to be a local minimum

- **Theorem 3:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function (that is all the second order partial derivatives of f (given by $\frac{\partial^2 f}{\partial x_j \partial x_i}$) exists and are continuous as functions from Ω to \mathbb{R}).
If \mathbf{x}^* is a local minimum of f then for any feasible direction \mathbf{d} at \mathbf{x}^*
 1. $\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0$.
 2. If $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$, then $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*)\mathbf{d} \geq 0$.

Second order necessary conditions for a point to be a local minimum

- **Theorem 3:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function (that is all the second order partial derivatives of f (given by $\frac{\partial^2 f}{\partial x_j \partial x_i}$) exists and are continuous as functions from Ω to \mathbb{R}).

If \mathbf{x}^* is a local minimum of f then for any feasible direction \mathbf{d} at \mathbf{x}^*

1. $\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0$.
2. If $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$, then $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*)\mathbf{d} \geq 0$.

Note: The matrix $\nabla^2 f$ (also denoted by H) is called the Hessian matrix of f ,

Second order necessary conditions for a point to be a local minimum

- **Theorem 3:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function (that is all the second order partial derivatives of f (given by $\frac{\partial^2 f}{\partial x_j \partial x_i}$) exists and are continuous as functions from Ω to \mathbb{R}).

If \mathbf{x}^* is a local minimum of f then for any feasible direction \mathbf{d} at \mathbf{x}^*

1. $\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0$.
2. If $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$, then $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*)\mathbf{d} \geq 0$.

Note: The matrix $\nabla^2 f$ (also denoted by H) is called the Hessian matrix of f ,

$$(\nabla^2 f(\mathbf{x}^*))_{ij} = \frac{\partial^2 f}{\partial x_j \partial x_i}(\mathbf{x}^*) = \left[\frac{\partial^2 f}{\partial x_j \partial x_i} \right]_{\mathbf{x}^*}.$$

- **Theorem 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function and let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum of f then

- **Theorem 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function and let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum of f then
 1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$.

- **Theorem 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function and let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum of f then
 1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$.
 2. $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} \geq 0$ for all vectors $\mathbf{d} \in \mathbb{R}^n$ (or $\nabla^2 f(\mathbf{x}^*)$ is positive semidefinite).

- **Theorem 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function and let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum of f then
 1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$.
 2. $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} \geq 0$ for all vectors $\mathbf{d} \in \mathbb{R}^n$ (or $\nabla^2 f(\mathbf{x}^*)$ is positive semidefinite).
- **Example 1:** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$

- **Theorem 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function and let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum of f then
 1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$.
 2. $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} \geq 0$ for all vectors $\mathbf{d} \in \mathbb{R}^n$ (or $\nabla^2 f(\mathbf{x}^*)$ is positive semidefinite).
- **Example 1:** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$
subject to $x_1 \geq 0, x_2 \geq 0$,

- **Theorem 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function and let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum of f then
 1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$.
 2. $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} \geq 0$ for all vectors $\mathbf{d} \in \mathbb{R}^n$ (or $\nabla^2 f(\mathbf{x}^*)$ is positive semidefinite).
- **Example 1:** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$
subject to $x_1 \geq 0, x_2 \geq 0$,
hence $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.

- Theorem 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function and let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum of f then
 - $\nabla f(\mathbf{x}^*) = \mathbf{0}$.
 - $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} \geq 0$ for all vectors $\mathbf{d} \in \mathbb{R}^n$ (or $\nabla^2 f(\mathbf{x}^*)$ is positive semidefinite).
- Example 1:** Consider the following problem:
 Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$
 subject to $x_1 \geq 0, x_2 \geq 0$,
 hence $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- $\nabla f(\mathbf{x}) = (3x_1^2 - 2x_1 x_2, -x_1^2 + 4x_2) = [0, 0]$ has two solutions

- Theorem 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function and let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* is a local minimum of f then
 - $\nabla f(\mathbf{x}^*) = \mathbf{0}$.
 - $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} \geq 0$ for all vectors $\mathbf{d} \in \mathbb{R}^n$ (or $\nabla^2 f(\mathbf{x}^*)$ is positive semidefinite).
- Example 1:** Consider the following problem:
 Minimize $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$
 subject to $x_1 \geq 0, x_2 \geq 0$,
 hence $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- $\nabla f(\mathbf{x}) = (3x_1^2 - 2x_1 x_2, -x_1^2 + 4x_2) = [0, 0]$ has two solutions
- $x_1 = 0, x_2 = 0$ and $x_1 = 6, x_2 = 9$.

- $[6, 9]^T$ is an interior point of the feasible region Ω hence any $\mathbf{d} \in \mathbb{R}^2$ is a feasible direction at $[6, 9]^T$.

- $[6, 9]^T$ is an interior point of the feasible region Ω hence any $\mathbf{d} \in \mathbb{R}^2$ is a feasible direction at $[6, 9]^T$.
- $[6, 9]^T$ satisfies the **first order necessary conditions** for a local minimum.

- $[6, 9]^T$ is an interior point of the feasible region Ω hence any $\mathbf{d} \in \mathbb{R}^2$ is a feasible direction at $[6, 9]^T$.
- $[6, 9]^T$ satisfies the **first order necessary conditions** for a local minimum.
- $\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 6x_1 - 2x_2 & -2x_1 \\ -2x_1 & 4 \end{pmatrix}$.

- $[6, 9]^T$ is an interior point of the feasible region Ω hence any $\mathbf{d} \in \mathbb{R}^2$ is a feasible direction at $[6, 9]^T$.
- $[6, 9]^T$ satisfies the **first order necessary conditions** for a local minimum.
- $\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 6x_1 - 2x_2 & -2x_1 \\ -2x_1 & 4 \end{pmatrix}$.

- At $\mathbf{x}^* = [6, 9]^T$, $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 18 & -12 \\ -12 & 4 \end{pmatrix}$.

- At $\mathbf{x}^* = [6, 9]^T$, $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 18 & -12 \\ -12 & 4 \end{pmatrix}$.
- But $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} = -2$ for $\mathbf{d} = [1, 1]^T$.

- At $\mathbf{x}^* = [6, 9]^T$, $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 18 & -12 \\ -12 & 4 \end{pmatrix}$.
- But $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} = -2$ for $\mathbf{d} = [1, 1]^T$.
- Hence $\mathbf{x}^* = [6, 9]^T$ since it does not satisfy the **second order necessary conditions for a local minimum** is not a local minimum point of f .

- At $\mathbf{x}^* = [6, 9]^T$, $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 18 & -12 \\ -12 & 4 \end{pmatrix}$.
- But $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} = -2$ for $\mathbf{d} = [1, 1]^T$.
- Hence $\mathbf{x}^* = [6, 9]^T$ since it does not satisfy the **second order necessary conditions for a local minimum** is not a local minimum point of f .
- Hence the first order necessary conditions are **necessary** but **not sufficient** for a point to be a local minimum.

- At $\mathbf{x}^* = [0, 0]^T$, $\mathbf{d} \neq \mathbf{0}$ is a feasible direction if and only if $d_1 \geq 0$ and $d_2 \geq 0$.

- At $\mathbf{x}^* = [0, 0]^T$, $\mathbf{d} \neq \mathbf{0}$ is a feasible direction if and only if $d_1 \geq 0$ and $d_2 \geq 0$.
Since $\nabla f(\mathbf{x}^*) = \mathbf{0}$, $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$ for all \mathbf{d} .

- At $\mathbf{x}^* = [0, 0]^T$, $\mathbf{d} \neq \mathbf{0}$ is a feasible direction if and only if $d_1 \geq 0$ and $d_2 \geq 0$.

Since $\nabla f(\mathbf{x}^*) = \mathbf{0}$, $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$ for all \mathbf{d} .

Since $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}$,

- At $\mathbf{x}^* = [0, 0]^T$, $\mathbf{d} \neq \mathbf{0}$ is a feasible direction if and only if $d_1 \geq 0$ and $d_2 \geq 0$.

Since $\nabla f(\mathbf{x}^*) = \mathbf{0}$, $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$ for all \mathbf{d} .

Since $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}$, $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} = 4d_2^2 \geq 0$ for all \mathbf{d} .

- At $\mathbf{x}^* = [0, 0]^T$, $\mathbf{d} \neq \mathbf{0}$ is a feasible direction if and only if $d_1 \geq 0$ and $d_2 \geq 0$.

Since $\nabla f(\mathbf{x}^*) = \mathbf{0}$, $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$ for all \mathbf{d} .

Since $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}$, $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} = 4d_2^2 \geq 0$ for all \mathbf{d} .

- $[0, 0]^T$ satisfies both the **first and second order necessary conditions** for a local minimum.

- At $\mathbf{x}^* = [0, 0]^T$, $\mathbf{d} \neq \mathbf{0}$ is a feasible direction if and only if $d_1 \geq 0$ and $d_2 \geq 0$.

Since $\nabla f(\mathbf{x}^*) = \mathbf{0}$, $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$ for all \mathbf{d} .

Since $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}$, $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} = 4d_2^2 \geq 0$ for all \mathbf{d} .

- $[0, 0]^T$ satisfies both the **first and second order necessary conditions** for a local minimum.
- If $\Omega = \mathbb{R}^2$ in Example 1 then $[0, 0]^T$ although satisfies both the **first and second order necessary conditions** is not a local minimum.

- At $\mathbf{x}^* = [0, 0]^T$, $\mathbf{d} \neq \mathbf{0}$ is a feasible direction if and only if $d_1 \geq 0$ and $d_2 \geq 0$.

Since $\nabla f(\mathbf{x}^*) = \mathbf{0}$, $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$ for all \mathbf{d} .

Since $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}$, $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} = 4d_2^2 \geq 0$ for all \mathbf{d} .

- $[0, 0]^T$ satisfies both the **first and second order necessary conditions** for a local minimum.
- If $\Omega = \mathbb{R}^2$ in Example 1 then $[0, 0]^T$ although satisfies both the **first and second order necessary conditions** is not a local minimum.
- For $\mathbf{d} = [-1, 0]^T$ which is a feasible direction at $[0, 0]^T$, $f(\mathbf{x}^* + t\mathbf{d}) = f(\mathbf{0} + t\mathbf{d}) < f(\mathbf{0})$, for all $t > 0$,

- At $\mathbf{x}^* = [0, 0]^T$, $\mathbf{d} \neq \mathbf{0}$ is a feasible direction if and only if $d_1 \geq 0$ and $d_2 \geq 0$.

Since $\nabla f(\mathbf{x}^*) = \mathbf{0}$, $\nabla f(\mathbf{x}^*)\mathbf{d} = 0$ for all \mathbf{d} .

Since $\nabla^2 f(\mathbf{x}^*) = \begin{pmatrix} 0 & 0 \\ 0 & 4 \end{pmatrix}$, $\mathbf{d}^T \nabla^2 f(\mathbf{x}^*) \mathbf{d} = 4d_2^2 \geq 0$ for all \mathbf{d} .

- $[0, 0]^T$ satisfies both the **first and second order necessary conditions** for a local minimum.
- If $\Omega = \mathbb{R}^2$ in Example 1 then $[0, 0]^T$ although satisfies both the **first and second order necessary conditions** is not a local minimum.
- For $\mathbf{d} = [-1, 0]^T$ which is a feasible direction at $[0, 0]^T$, $f(\mathbf{x}^* + t\mathbf{d}) = f(\mathbf{0} + t\mathbf{d}) < f(\mathbf{0})$, for all $t > 0$, so $[0, 0]^T$ is not a local minimum of f .

- **Example 2 :** Consider the following problem:

$$\text{Minimize } f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$$

- **Example 2 :** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$

subject to $x_1 \geq 0, x_2 \geq 0$.

- **Example 2 :** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$
subject to $x_1 \geq 0, x_2 \geq 0$.
- Note that f is a twice continuously differentiable function.

- **Example 2 :** Consider the following problem:

Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$

subject to $x_1 \geq 0, x_2 \geq 0$.

- Note that f is a twice continuously differentiable function.
- Check that $\mathbf{x}^* = [\frac{1}{2}, 0]^T$, **satisfies the first and the second order necessary conditions** for \mathbf{x}^* to be locally minimum.

- **Example 2 :** Consider the following problem:

$$\text{Minimize } f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$$

subject to $x_1 \geq 0, x_2 \geq 0$.

- Note that f is a twice continuously differentiable function.
- Check that $\mathbf{x}^* = [\frac{1}{2}, 0]^T$, **satisfies the first and the second order necessary conditions** for \mathbf{x}^* to be locally minimum.
- $\mathbf{x}^* = [\frac{1}{2}, 0]^T$, is the global minimum of f .

- **Example 2 :** Consider the following problem:
Minimize $f(x_1, x_2) = x_1^2 - x_1 + x_2 + x_1 x_2$
subject to $x_1 \geq 0, x_2 \geq 0$.
- Note that f is a twice continuously differentiable function.
- Check that $\mathbf{x}^* = [\frac{1}{2}, 0]^T$, **satisfies the first and the second order necessary conditions** for \mathbf{x}^* to be locally minimum.
- $\mathbf{x}^* = [\frac{1}{2}, 0]^T$, is the global minimum of f .
- **Definition:** A real symmetric matrix A is said to be positive semidefinite (negative semidefinite) if $\mathbf{x}^T A \mathbf{x} \geq 0$ ($\mathbf{x}^T A \mathbf{x} \leq 0$) for all $\mathbf{x} \in \mathbb{R}^n$.

- **Definition:** A real symmetric matrix A is said to be positive definite (negative definite) if $\mathbf{x}^T A \mathbf{x} > 0$ ($\mathbf{x}^T A \mathbf{x} < 0$) for all **nonzero** vectors $\mathbf{x} \in \mathbb{R}^n$.

- **Definition:** A real symmetric matrix A is said to be positive definite (negative definite) if $\mathbf{x}^T A \mathbf{x} > 0$ ($\mathbf{x}^T A \mathbf{x} < 0$) for all **nonzero** vectors $\mathbf{x} \in \mathbb{R}^n$.
- **Theorem:** If A is a symmetric, $n \times n$, real matrix then the following statements are equivalent:

- **Definition:** A real symmetric matrix A is said to be positive definite (negative definite) if $\mathbf{x}^T A \mathbf{x} > 0$ ($\mathbf{x}^T A \mathbf{x} < 0$) for all **nonzero** vectors $\mathbf{x} \in \mathbb{R}^n$.
- **Theorem:** If A is a symmetric, $n \times n$, real matrix then the following statements are equivalent:
- A is positive semidefinite.

- **Definition:** A real symmetric matrix A is said to be positive definite (negative definite) if $\mathbf{x}^T A \mathbf{x} > 0$ ($\mathbf{x}^T A \mathbf{x} < 0$) for all **nonzero** vectors $\mathbf{x} \in \mathbb{R}^n$.
- **Theorem:** If A is a symmetric, $n \times n$, real matrix then the following statements are equivalent:
 - A is positive semidefinite.
 - All eigenvalues of A are nonnegative.

- **Definition:** A real symmetric matrix A is said to be positive definite (negative definite) if $\mathbf{x}^T A \mathbf{x} > 0$ ($\mathbf{x}^T A \mathbf{x} < 0$) for all **nonzero** vectors $\mathbf{x} \in \mathbb{R}^n$.
- **Theorem:** If A is a symmetric, $n \times n$, real matrix then the following statements are equivalent:
 - A is positive semidefinite.
 - All eigenvalues of A are nonnegative.
 - All principal minors of A are nonnegative.

- **Definition:** A real symmetric matrix A is said to be positive definite (negative definite) if $\mathbf{x}^T A \mathbf{x} > 0$ ($\mathbf{x}^T A \mathbf{x} < 0$) for all **nonzero** vectors $\mathbf{x} \in \mathbb{R}^n$.
- **Theorem:** If A is a symmetric, $n \times n$, real matrix then the following statements are equivalent:
 - A is positive semidefinite.
 - All eigenvalues of A are nonnegative.
 - All principal minors of A are nonnegative.
- **Definition:** $\lambda \in \mathbb{C}$ is called an eigenvalue of an $n \times n$ matrix A if there exists an $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ (that is atleast one component of \mathbf{x} is nonzero) such that $A\mathbf{x} = \lambda\mathbf{x}$.

- **Definition:** A real symmetric matrix A is said to be positive definite (negative definite) if $\mathbf{x}^T A \mathbf{x} > 0$ ($\mathbf{x}^T A \mathbf{x} < 0$) for all **nonzero** vectors $\mathbf{x} \in \mathbb{R}^n$.
- **Theorem:** If A is a symmetric, $n \times n$, real matrix then the following statements are equivalent:
 - A is positive semidefinite.
 - All eigenvalues of A are nonnegative.
 - All principal minors of A are nonnegative.
- **Definition:** $\lambda \in \mathbb{C}$ is called an eigenvalue of an $n \times n$ matrix A if there exists an $\mathbf{x} \in \mathbb{C}^n$, $\mathbf{x} \neq \mathbf{0}$ (that is at least one component of \mathbf{x} is nonzero) such that $A\mathbf{x} = \lambda\mathbf{x}$.
- For example the $\mathbf{0}$ matrix has all n eigenvalues equal to 0, the identity matrix I_n has all n eigenvalues equal to 1, and for an **upper triangular matrix**, the diagonal entries are its eigenvalues.

- **Definition:** If A is an $n \times n$ matrix and $\alpha \subseteq \{1, \dots, n\}$, $\beta \subseteq \{1, \dots, n\}$ then $A[\alpha, \beta]$ is the (sub)matrix obtained from A , by deleting all the rows of A which do not belong to α , and by deleting all the columns of A which do not belong to β .

- **Definition:** If A is an $n \times n$ matrix and $\alpha \subseteq \{1, \dots, n\}$, $\beta \subseteq \{1, \dots, n\}$ then $A[\alpha, \beta]$ is the (sub)matrix obtained from A , by deleting all the rows of A which do not belong to α , and by deleting all the columns of A which do not belong to β .
- If $\alpha = \beta$, then $A[\alpha, \alpha]$ is called a **principal submatrix** of A and $\det A[\alpha, \alpha]$ is called a **principal minor** of A .

- **Definition:** If A is an $n \times n$ matrix and $\alpha \subseteq \{1, \dots, n\}$, $\beta \subseteq \{1, \dots, n\}$ then $A[\alpha, \beta]$ is the (sub)matrix obtained from A , by deleting all the rows of A which do not belong to α , and by deleting all the columns of A which do not belong to β .
- If $\alpha = \beta$, then $A[\alpha, \alpha]$ is called a **principal submatrix** of A and $\det A[\alpha, \alpha]$ is called a **principal minor** of A .
- If $\alpha = \beta = \{i\}$, where $i \in \{1, \dots, n\}$, then $A[\alpha, \alpha] = [a_{ii}]$, and $\det A[\alpha, \alpha] = a_{ii}$, the i th diagonal entry.

- **Definition:** If A is an $n \times n$ matrix and $\alpha \subseteq \{1, \dots, n\}$, $\beta \subseteq \{1, \dots, n\}$ then $A[\alpha, \beta]$ is the (sub)matrix obtained from A , by deleting all the rows of A which do not belong to α , and by deleting all the columns of A which do not belong to β .
- If $\alpha = \beta$, then $A[\alpha, \alpha]$ is called a **principal submatrix** of A and $\det A[\alpha, \alpha]$ is called a **principal minor** of A .
- If $\alpha = \beta = \{i\}$, where $i \in \{1, \dots, n\}$, then $A[\alpha, \alpha] = [a_{ii}]$, and $\det A[\alpha, \alpha] = a_{ii}$, the i th diagonal entry.
- If $\alpha = \beta = \{i, j\}$, where $i, j \in \{1, \dots, n\}$ and $i < j$, then

- **Definition:** If A is an $n \times n$ matrix and $\alpha \subseteq \{1, \dots, n\}$, $\beta \subseteq \{1, \dots, n\}$ then $A[\alpha, \beta]$ is the (sub)matrix obtained from A , by deleting all the rows of A which do not belong to α , and by deleting all the columns of A which do not belong to β .
- If $\alpha = \beta$, then $A[\alpha, \alpha]$ is called a **principal submatrix** of A and $\det A[\alpha, \alpha]$ is called a **principal minor** of A .
- If $\alpha = \beta = \{i\}$, where $i \in \{1, \dots, n\}$, then $A[\alpha, \alpha] = [a_{ii}]$, and $\det A[\alpha, \alpha] = a_{ii}$, the i th diagonal entry.
- If $\alpha = \beta = \{i, j\}$, where $i, j \in \{1, \dots, n\}$ and $i < j$, then
$$A[\alpha, \alpha] = \begin{bmatrix} a_{ii} & a_{ij} \\ a_{ji} & a_{jj} \end{bmatrix}.$$

- **Definition:** If A is an $n \times n$ matrix and $\alpha \subseteq \{1, \dots, n\}$, $\beta \subseteq \{1, \dots, n\}$ then $A[\alpha, \beta]$ is the (sub)matrix obtained from A , by deleting all the rows of A which do not belong to α , and by deleting all the columns of A which do not belong to β .
- If $\alpha = \beta$, then $A[\alpha, \alpha]$ is called a **principal submatrix** of A and $\det A[\alpha, \alpha]$ is called a **principal minor** of A .
- If $\alpha = \beta = \{i\}$, where $i \in \{1, \dots, n\}$, then $A[\alpha, \alpha] = [a_{ii}]$, and $\det A[\alpha, \alpha] = a_{ii}$, the i th diagonal entry.
- If $\alpha = \beta = \{i, j\}$, where $i, j \in \{1, \dots, n\}$ and $i < j$, then

$$A[\alpha, \alpha] = \begin{bmatrix} a_{ii} & a_{ij} \\ a_{ji} & a_{jj} \end{bmatrix}.$$
 If $\alpha = \{1, \dots, n\}$, then $A[\alpha, \alpha] = A$, and $\det A[\alpha, \alpha] = \det(A)$.

- **Definition:** If A is an $n \times n$ matrix and $\alpha \subseteq \{1, \dots, n\}$, $\beta \subseteq \{1, \dots, n\}$ then $A[\alpha, \beta]$ is the (sub)matrix obtained from A , by deleting all the rows of A which do not belong to α , and by deleting all the columns of A which do not belong to β .
- If $\alpha = \beta$, then $A[\alpha, \alpha]$ is called a **principal submatrix** of A and $\det A[\alpha, \alpha]$ is called a **principal minor** of A .
- If $\alpha = \beta = \{i\}$, where $i \in \{1, \dots, n\}$, then $A[\alpha, \alpha] = [a_{ii}]$, and $\det A[\alpha, \alpha] = a_{ii}$, the i th diagonal entry.
- If $\alpha = \beta = \{i, j\}$, where $i, j \in \{1, \dots, n\}$ and $i < j$, then

$$A[\alpha, \alpha] = \begin{bmatrix} a_{ii} & a_{ij} \\ a_{ji} & a_{jj} \end{bmatrix}.$$
 If $\alpha = \{1, \dots, n\}$, then $A[\alpha, \alpha] = A$, and $\det A[\alpha, \alpha] = \det(A)$.
- **Remark:** A nonsingular (nonzero determinant) **positive semidefinite matrix** is **positive definite**.

- **Sufficient conditions for a local minima :**

Theorem 4 : Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function.

- **Sufficient conditions for a local minima :**

Theorem 4 : Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function.

Let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* satisfies the following conditions

- **Sufficient conditions for a local minima :**

Theorem 4 : Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function.

Let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* satisfies the following conditions

1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$

- **Sufficient conditions for a local minima :**

Theorem 4 : Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function.

Let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* satisfies the following conditions

1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$
2. $\nabla^2 f(\mathbf{x}^*)$ is positive definite,

- **Sufficient conditions for a local minima :**

Theorem 4 : Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function.

Let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* satisfies the following conditions

1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$

2. $\nabla^2 f(\mathbf{x}^*)$ is positive definite,

then \mathbf{x}^* is a local minimum point of f .

- **Sufficient conditions for a local minima :**

Theorem 4 : Let $f : \Omega \rightarrow \mathbb{R}$ be a twice continuously differentiable function.

Let \mathbf{x}^* be an interior point of Ω . If \mathbf{x}^* satisfies the following conditions

1. $\nabla f(\mathbf{x}^*) = \mathbf{0}$

2. $\nabla^2 f(\mathbf{x}^*)$ is positive definite,

then \mathbf{x}^* is a local minimum point of f .

- **Remark:** Since maximizing f is same as minimizing $-f$, all the previous theorems have corresponding analogues for a maximization problem with some obvious changes.

- **Definition 4:** A real valued function f defined on a convex set $\Omega \subseteq \mathbb{R}^n$ is said to be a **convex function** on Ω if for all $\mathbf{x}, \mathbf{y} \in \Omega$ and all $0 \leq \alpha \leq 1$,
$$f(\alpha \mathbf{x} + (1 - \alpha) \mathbf{y}) \leq \alpha f(\mathbf{x}) + (1 - \alpha) f(\mathbf{y}).$$

- **Definition 4:** A real valued function f defined on a convex set $\Omega \subseteq \mathbb{R}^n$ is said to be a **convex function** on Ω if for all $\mathbf{x}, \mathbf{y} \in \Omega$ and all $0 \leq \alpha \leq 1$,
$$f(\alpha \mathbf{x} + (1 - \alpha) \mathbf{y}) \leq \alpha f(\mathbf{x}) + (1 - \alpha) f(\mathbf{y}).$$
- **Definition 5:** An $f : \Omega \rightarrow \mathbb{R}$ is said to be a **concave function** if $-f$ is a **convex function**.

- **Definition 4:** A real valued function f defined on a convex set $\Omega \subseteq \mathbb{R}^n$ is said to be a **convex function** on Ω if for all $\mathbf{x}, \mathbf{y} \in \Omega$ and all $0 \leq \alpha \leq 1$,
$$f(\alpha \mathbf{x} + (1 - \alpha) \mathbf{y}) \leq \alpha f(\mathbf{x}) + (1 - \alpha) f(\mathbf{y}).$$
- **Definition 5:** An $f : \Omega \rightarrow \mathbb{R}$ is said to be a **concave function** if $-f$ is a **convex function**.
- **Theorem 1 :** If f is a convex function on Ω (a convex set), then the set $S = \{\mathbf{x} \in \Omega : f(\mathbf{x}) \leq c\}$ is a convex set (for all real c).

- **Definition 4:** A real valued function f defined on a convex set $\Omega \subseteq \mathbb{R}^n$ is said to be a **convex function** on Ω if for all $\mathbf{x}, \mathbf{y} \in \Omega$ and all $0 \leq \alpha \leq 1$,
$$f(\alpha \mathbf{x} + (1 - \alpha) \mathbf{y}) \leq \alpha f(\mathbf{x}) + (1 - \alpha) f(\mathbf{y}).$$
- **Definition 5:** An $f : \Omega \rightarrow \mathbb{R}$ is said to be a **concave function** if $-f$ is a **convex function**.
- **Theorem 1 :** If f is a convex function on Ω (a convex set), then the set $S = \{\mathbf{x} \in \Omega : f(\mathbf{x}) \leq c\}$ is a convex set (for all real c).
- **Theorem 2:** Let f be a continuously differentiable function defined on a convex set, $\Omega \subseteq \mathbb{R}^n$, then f is convex on Ω if and only if

- **Definition 4:** A real valued function f defined on a convex set $\Omega \subseteq \mathbb{R}^n$ is said to be a **convex function** on Ω if for all $\mathbf{x}, \mathbf{y} \in \Omega$ and all $0 \leq \alpha \leq 1$,

$$f(\alpha \mathbf{x} + (1 - \alpha) \mathbf{y}) \leq \alpha f(\mathbf{x}) + (1 - \alpha) f(\mathbf{y}).$$
- **Definition 5:** An $f : \Omega \rightarrow \mathbb{R}$ is said to be a **concave function** if $-f$ is a **convex function**.
- **Theorem 1 :** If f is a convex function on Ω (a convex set), then the set $S = \{\mathbf{x} \in \Omega : f(\mathbf{x}) \leq c\}$ is a convex set (for all real c).
- **Theorem 2:** Let f be a continuously differentiable function defined on a convex set, $\Omega \subseteq \mathbb{R}^n$, then f is convex on Ω if and only if

$$f(\mathbf{y}) \geq f(\mathbf{x}) + \nabla f(\mathbf{x})(\mathbf{y} - \mathbf{x}) \text{ for all } \mathbf{x}, \mathbf{y} \in \Omega.$$

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has atleast one interior point), then f is a **convex function** on Ω if and only if

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has atleast one interior point), then f is a **convex function** on Ω if and only if
for all $\mathbf{x} \in \Omega$,

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has atleast one interior point), then f is a **convex function** on Ω if and only if
for all $\mathbf{x} \in \Omega$, $\nabla^2 f(\mathbf{x})$ is positive semidefinite.

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has at least one interior point), then f is a **convex function** on Ω if and only if
for all $\mathbf{x} \in \Omega$, $\nabla^2 f(\mathbf{x})$ is positive semidefinite.
- **Revisiting Example 1** : Let $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$ be defined on

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has at least one interior point), then f is a **convex function** on Ω if and only if
for all $\mathbf{x} \in \Omega$, $\nabla^2 f(\mathbf{x})$ is positive semidefinite.
- **Revisiting Example 1** : Let $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$ be defined on
 $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has at least one interior point), then f is a **convex function** on Ω if and only if
for all $\mathbf{x} \in \Omega$, $\nabla^2 f(\mathbf{x})$ is positive semidefinite.
- **Revisiting Example 1** : Let $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$ be defined on
 $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- Since $\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 6x_1 - 2x_2 & -2x_1 \\ -2x_1 & 4 \end{pmatrix}$,

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has at least one interior point), then f is a **convex function** on Ω if and only if
for all $\mathbf{x} \in \Omega$, $\nabla^2 f(\mathbf{x})$ is positive semidefinite.
- **Revisiting Example 1** : Let $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$ be defined on
 $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- Since $\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 6x_1 - 2x_2 & -2x_1 \\ -2x_1 & 4 \end{pmatrix}$,
- At $x_1 = 1, x_2 = 3$,

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has at least one interior point), then f is a **convex function** on Ω if and only if
for all $\mathbf{x} \in \Omega$, $\nabla^2 f(\mathbf{x})$ is positive semidefinite.
- **Revisiting Example 1** : Let $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$ be defined on
 $\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- Since $\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 6x_1 - 2x_2 & -2x_1 \\ -2x_1 & 4 \end{pmatrix}$,
- At $x_1 = 1, x_2 = 3$, $\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 0 & -2 \\ -2 & 4 \end{pmatrix}$
is clearly not positive semidefinite.

- **Theorem 3** : Let f be a twice continuously differentiable function on a convex set Ω (where Ω has at least one interior point), then f is a **convex function** on Ω if and only if

for all $\mathbf{x} \in \Omega$, $\nabla^2 f(\mathbf{x})$ is positive semidefinite.
- **Revisiting Example 1** : Let $f(x_1, x_2) = x_1^3 - x_1^2 x_2 + 2x_2^2$ be defined on

$\Omega = \{[x_1, x_2]^T : x_1 \geq 0, x_2 \geq 0\}$.
- Since $\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 6x_1 - 2x_2 & -2x_1 \\ -2x_1 & 4 \end{pmatrix}$,
- At $x_1 = 1, x_2 = 3$, $\nabla^2 f(\mathbf{x}) = \begin{pmatrix} 0 & -2 \\ -2 & 4 \end{pmatrix}$

is clearly not positive semidefinite.

Hence f is not a convex function on Ω .

- **Remark :** Since minimizing f is same as maximizing $-f$, all the previous theorems for **minimizing** a **convex function** have corresponding analogues for **maximizing** a **concave function**.

- **Remark :** Since minimizing f is same as maximizing $-f$, all the previous theorems for **minimizing** a **convex function** have corresponding analogues for **maximizing** a **concave function**.

Theorem 4: Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function.

- **Remark :** Since minimizing f is same as maximizing $-f$, all the previous theorems for **minimizing a convex function** have corresponding analogues for **maximizing a concave function**.

Theorem 4: Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function. If f is a convex function on Ω , then \mathbf{x}^* is a local (global) minimum of f if and only if

- **Remark :** Since minimizing f is same as maximizing $-f$, all the previous theorems for **minimizing a convex function** have corresponding analogues for **maximizing a concave function**.

Theorem 4: Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function. If f is a convex function on Ω , then \mathbf{x}^* is a local (global) minimum of f if and only if for all feasible direction \mathbf{d} at \mathbf{x}^*

$$\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0.$$

- **Remark :** Since minimizing f is same as maximizing $-f$, all the previous theorems for **minimizing a convex function** have corresponding analogues for **maximizing a concave function**.

Theorem 4: Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function. If f is a convex function on Ω , then \mathbf{x}^* is a local (global) minimum of f if and only if for all feasible direction \mathbf{d} at \mathbf{x}^*

$$\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0.$$

- **Corollary 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function. If f is a convex function on Ω and \mathbf{x}^* an interior point of Ω , then \mathbf{x}^* is a local (global) minimum for f if and only if
- $$\nabla f(\mathbf{x}^*) = \mathbf{0}.$$

- **Remark :** Since minimizing f is same as maximizing $-f$, all the previous theorems for **minimizing a convex function** have corresponding analogues for **maximizing a concave function**.

Theorem 4: Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function. If f is a convex function on Ω , then \mathbf{x}^* is a local (global) minimum of f if and only if for all feasible direction \mathbf{d} at \mathbf{x}^*

$$\nabla f(\mathbf{x}^*)\mathbf{d} \geq 0.$$

- **Corollary 4:** Let $f : \Omega \rightarrow \mathbb{R}$ be a continuously differentiable function. If f is a convex function on Ω and \mathbf{x}^* an interior point of Ω , then \mathbf{x}^* is a local (global) minimum for f if and only if
$$\nabla f(\mathbf{x}^*) = \mathbf{0}.$$
- The above result is **not** necessarily true if f is **not convex** as you have already seen in **Example 1**.