



December 5, 2022

Knotted loops & 4-dimensional spaces






Knots and links

- A knot is a embedded circle in 3-dimensional space






Knots and links

- A knot is a embedded circle in 3-dimensional space







Knots and links


- A knot is a embedded circle in 3-dimensional space




- A link is a collection of knots



Satellite operations




A pattern P




A knot K


Satellite operations



A pattern P




A knot K




The satellite $P(K)$


Satellite operations





A pattern P




A knot K




The satellite $P(K)$




Satellite operations



A pattern Q





A link L



The satellite $Q(L)$




Isotopy of knots

- Two knots/links are **isotopic** if one can be deformed to the other



Isotopy of knots




- Two knots/links are **isotopic** if one can be deformed to the other



Three pictures of the **trivial knot**, or **unknot**

Isotopy of knots




- Two knots/links are **isotopic** if one can be deformed to the other



Three pictures of the **trivial knot**, or **unknot**

Isotopy of knots

- Two knots/links are **isotopic** if one can be deformed to the other




Three pictures of the **trivial knot**, or **unknot**

Slice knots and links

- A knot is **trivial** precisely if it bounds a disc in 3-dimensional space



Slice knots and links

- A knot is **trivial** precisely if it bounds a disc in 3-dimensional space



Slice knots and links



- A knot is **trivial** precisely if it bounds a disc in 3-dimensional space



- A knot is **slice** if it bounds a disc in \mathbb{R}^4_+ , the upper half-space in 4-dimensional space



Slice knots and links

- A knot is **trivial** precisely if it bounds a disc in 3-dimensional space





- A knot is **slice** if it bounds a disc in \mathbb{R}^4_+ , the upper half-space in 4-dimensional space
- A link is **slice** if it bounds a collection of discs in \mathbb{R}^4_+



A slice knot





A slice knot





A slice knot





A slice knot





A slice knot






A slice knot




A slice knot




A slice knot





A slice knot



A slice knot



A slice knot




Satellite knots and sliceness

Theorem [Cochran-Davis-R. 2014]

For infinitely many patterns P , and all knots K , if $P(K)$ is slice, then K is slice.

E.g.




Satellite knots and sliceness

Theorem [Cochran-Davis-R. 2014]

For infinitely many patterns P , and all knots K , if $P(K)$ is slice, then K is slice.

E.g.




- More generally, with respect to the notion of **concordance**, these patterns are **injective** functions

Satellite knots and sliceness

Theorem [Cochran-Davis-R. 2014]

For infinitely many patterns P , and all knots K , if $P(K)$ is slice, then K is slice.

E.g.




- More generally, with respect to the notion of **concordance**, these patterns are **injective** functions
- Further work on **iterated** satellites, **bijection** of patterns, ...
[R. 2013, R. 2015, Cochran-R. 2016, Feller-Park-R. 2019, Davis-Park-R. 2021]

4-dimensional spaces



Manifolds

- Manifolds are spaces which are locally like Euclidean space






Manifolds

- Manifolds are spaces which are locally like Euclidean space






Manifolds

- Manifolds are spaces which are locally like Euclidean space



Manifolds


- Manifolds are spaces which are locally like Euclidean space



- The universe appears to be a 3-dimensional manifold
- Space-time forms a 4-dimensional manifold
- The n -dimensional sphere is given by $\{(x_1, x_2, \dots, x_{n+1}) \mid x_1^2 + x_2^2 + \dots + x_{n+1}^2 = 1\}$




Classifying manifolds

- Dimension 1: the circle






Classifying manifolds

- Dimension 1: the circle
- Dimension 2: classified by genus



Classifying manifolds

- Dimension 1: the circle
- Dimension 2: classified by genus



The distinct circles in a manifold M are recorded in the fundamental group $\pi_1(M)$

Classifying manifolds

- Dimension 1: the circle
- Dimension 2: classified by genus
- Dimension 3: classified, decompose into geometric pieces [Perelman 2003]

Classifying manifolds


- Dimension 1: the circle
- Dimension 2: classified by genus
- Dimension 3: classified, decompose into geometric pieces [Perelman 2003]
- Dimensions 5 & higher: surgery theory

Classifying manifolds

- Dimension 1: the circle
- Dimension 2: classified by genus
- Dimension 3: classified, decompose into geometric pieces [Perelman 2003]
- What about dimension four?
- Dimensions 5 & higher: surgery theory


Surgery to classify surfaces

- Idea: modify the surface in a controlled manner to reduce complexity
 - find geometrically dual pairs of circles, detecting genus




Surgery to classify surfaces

- Idea: modify the surface in a controlled manner to reduce complexity
 - find geometrically dual pairs of circles, detecting genus




Surgery to classify surfaces

- Idea: modify the surface in a controlled manner to reduce complexity
 - find **geometrically dual** pairs of circles, detecting genus




Surgery to classify surfaces

- Idea: modify the surface in a controlled manner to reduce complexity
 - find geometrically dual pairs of circles, detecting genus




- the dual circle ensures that the result is less complex





Surgery to classify surfaces

- Idea: modify the surface in a controlled manner to reduce complexity
 - find **geometrically dual** pairs of circles, detecting genus



- the dual circle ensures that the result is less complex



Surgery for 4-manifolds

- For 4-manifolds, we need to find and control *circles* as well as *spheres*
 - In general, we must control spheres up to the middle dimension
- The group $\pi_2(M)$ records the types of spheres in a manifold M

Surgery for 4-manifolds

- For 4-manifolds, we need to find and control *circles* as well as *spheres*
 - In general, we must control spheres up to the middle dimension
- The group $\pi_2(M)$ records the types of spheres in a manifold M

Goal for surgery in 4-manifolds

Let M be a 4-manifold. Try to represent every algebraically dual pair in $\pi_2(M)$ by a *geometrically dual* pair of spheres in M .

Sphere embedding theorem

Let M be a 4-manifold. Every algebraically dual pair in $\pi_2(M)$ can be represented by a geometrically dual pair of spheres in M .


For $\pi_1(M)$ trivial: Freedman 1982

For $\pi_1(M)$ good: Freedman-Quinn 1990 + Powell-R.-Teichner 2018


For general $\pi_1(M)$: open

- Good groups include all finite groups, abelian groups, solvable groups, elementary abelian groups, groups of subexponential growth, ...
- Consequences include: 4-dimensional Poincaré conjecture, classification of simply connected 4-manifolds, topological transversality, ...

Explaining 4-manifolds



- 2013+2016:
 - semester programmes in Bonn, Germany
 - lectures by Freedman and others
 - notes developed into accessible textbook
 - published 2021, Oxford University Press



Sphere embedding theorem

Theorem [Casson 1970s, Freedman 1982, Freedman-Quinn 1990]

M a connected 4-manifold, $\pi_1(M)$ good, Σ a simply connected 2-manifold.

$F: \Sigma \longrightarrow M$ such that

- the algebraic self-intersection number of F vanishes
- &
- F has a framed algebraically dual sphere.

Then F is homotopic, relative to $\partial\Sigma$, to an embedding, with a geometrically dual sphere.

Sphere embedding theorem

Theorem [Casson 1970s, Freedman 1982, Freedman-Quinn 1990]

M a connected 4-manifold, $\pi_1(M)$ good, Σ a simply connected 2-manifold.

$F: \Sigma \longrightarrow M$ such that

- the algebraic self-intersection number of F vanishes
- &
- F has a framed algebraically dual sphere.

Then F is homotopic, relative to $\partial\Sigma$, to an embedding, with a geometrically dual sphere.

$\pi_1(M) \neq 1$, Powell-R.-Teichner 2018

~~Sphere~~ embedding theorem Surface

Theorem [Casson 1970s, Freedman 1982, Freedman-Quinn 1990,
Powell-R.-Teichner 2018, Stong 1994, Kasprowski-Powell-R.-Teichner 2022]

M a connected 4-manifold, $\pi_1(M)$ good, Σ a simply connected 2-manifold.

$F: \Sigma \longrightarrow M$ such that

- the algebraic self-intersection number of F vanishes
- F has a framed algebraically dual sphere.

$\begin{matrix} \uparrow & \uparrow \\ \partial\Sigma \hookrightarrow \partial M \end{matrix}$

Then F is homotopic, relative to $\partial\Sigma$, to an embedding, with a geometrically dual sphere

~~Sphere~~ embedding theorem Surface

Theorem [Casson 1970s, Freedman 1982, Freedman-Quinn 1990,
Powell-R.-Teichner 2018, Stong 1994, Kasprowski-Powell-R.-Teichner 2022]

M a connected 4-manifold, $\pi_1(M)$ good, Σ a ~~simply~~ connected 2-manifold.

$F: \Sigma \longrightarrow M$ such that

- the algebraic self-intersection number of F vanishes
- F has a ~~framed~~ algebraically dual sphere.

$\begin{matrix} \uparrow & \uparrow \\ \partial\Sigma \hookrightarrow \partial M & \end{matrix}$

Then F is homotopic, relative to $\partial\Sigma$, to an embedding, with a ~~geometrically dual~~ sphere

~~Sphere~~ embedding theorem Surface

Theorem [Casson 1970s, Freedman 1982, Freedman-Quinn 1990,
Powell-R.-Teichner 2018, Stong 1994, Kasprowski-Powell-R.-Teichner 2022]

M a connected 4-manifold, $\pi_1(M)$ good, Σ a ~~simply~~ connected 2-manifold.

$F: \Sigma \longrightarrow M$ such that


- the algebraic self-intersection number of F vanishes
- F has a ~~framed~~ algebraically dual sphere.

$\begin{array}{ccc} \uparrow & \uparrow \\ \partial\Sigma \hookrightarrow \partial M & & \end{array}$

Then F is homotopic, relative to $\partial\Sigma$, to an embedding, with a ~~geometrically dual~~ sphere if and only if the Kervaire-Milnor invariant $km(F) \in \mathbb{Z}/2$ vanishes.


The Kervaire-Milnor invariant

- Assume that F has trivial algebraic self-intersection number



The Kervaire-Milnor invariant


- Assume that F has trivial algebraic self-intersection number



- Then the self-intersections are in isolated double points, paired by Whitney discs

The Kervaire-Milnor invariant


- Assume that F has trivial algebraic self-intersection number



- Then the self-intersections are in isolated double points, paired by Whitney discs
- $km(F, \{W\}) = \sum |\text{Int}(W) \pitchfork F| \bmod 2$

The Kervaire-Milnor invariant

- Assume that F has trivial algebraic self-intersection number



- Then the self-intersections are in isolated double points, paired by Whitney discs
- $km(F, \{W\}) = \sum |\text{Int}(W) \pitchfork F| \bmod 2$
- When is this independent of $\{W\}$?

Other directions

- More classification results for 4-manifolds [Kasprowski-Powell-R. 2022+]
- The effect of Gluck twisting on general 4-manifolds [Kasprowski-Powell-R. 2022]
- Construction of pairs of smooth 4-manifolds that are homotopy equivalent but not homeomorphic [Kasprowski-Powell-R. 2022+]
- Study of the mapping class group of the universal exotic smooth structure on \mathbb{R}^4 [Gompf-Orson-R. 2022+]
- Finding embedded spheres in knot traces (no algebraically dual sphere) [Feller-Miller-Nagel-Orson-Powell-R. 2021]
- Slicing knots in general 4-manifolds [Kjuchukova-Miller-R.-Sakalli 2021, Kasprowski-Powell-R.-Teichner 2022, Marengon-Miller-R.-Stipsicz 2022]

Links & 4-manifolds

Links and 4-manifolds

Theorem [Casson-Freedman 1984]


Surgery works for all 4-manifolds precisely if every member of a certain family of links is slice (with free fundamental group).

Links and 4-manifolds

Theorem [Casson-Freedman 1984]

Surgery works for all 4-manifolds precisely if every member of a certain family of links is slice (with free fundamental group).

E.g.




Slice?

Links and 4-manifolds


Theorem [Casson-Freedman 1984]

Surgery works for all 4-manifolds precisely if every member of a certain family of links is slice (with free fundamental group).

E.g.



Slice?




Not slice!

Links and 4-manifolds


Theorem [Casson-Freedman 1984]

Surgery works for all 4-manifolds precisely if every member of a certain family of links is slice (with free fundamental group).

E.g.



Slice?



Not slice!

Key open question for classifying 4-manifolds: decide if these links are slice.

Questions?