Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 7092\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 7887\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 032\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387 \end{array}$

Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 7092\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 7887\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 032\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387 \end{array}$

Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 702\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 7887\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 032\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387 \end{array}$

Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 7092\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 988\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 7857\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 032\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387\\ \end{array}$

Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6186\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 7092\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 7887\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 032\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387\\ \end{array}$

Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 7092\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 7887\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 932\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387\\ \end{array}$

Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 7092\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 7887\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 032\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387 \end{array}$

Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 702\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 191\ 502\ 7587\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6388\\ 032\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387 \end{array}$

Graham's number

an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:


 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 7092\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 7887\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 939\ 40\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 932\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387 \end{array}$

Graham's number


an enormous number in Ramsey theory whose first digit might remain unknown for all time but whose last 500 digits can be easily computed and are shown below:

 $\begin{array}{c} \dots 024\ 259\ 5069\ 506\ 473\ 8395\ 657\ 479\ 1365\ 193\ 517\ 9833\ 453\ 536\ 2521\\ 430\ 035\ 4012\ 602\ 677\ 1622\ 672\ 160\ 4198\ 106\ 522\ 6316\ 935\ 518\ 8780\\ 388\ 144\ 8314\ 065\ 252\ 6168\ 785\ 095\ 5526\ 460\ 510\ 7117\ 200\ 099\ 702\\ 912\ 495\ 4437\ 888\ 749\ 6062\ 882\ 911\ 7250\ 630\ 013\ 0362\ 293\ 491\ 6080\\ 254\ 594\ 6149\ 457\ 887\ 1427\ 832\ 350\ 8292\ 421\ 020\ 9182\ 589\ 675\ 3560\\ 430\ 869\ 9380\ 168\ 924\ 9889\ 268\ 099\ 5101\ 690\ 559\ 1995\ 119\ 502\ 785\\ 178\ 308\ 3701\ 834\ 023\ 6474\ 548\ 882\ 2221\ 615\ 732\ 2801\ 013\ 297\ 4509\\ 273\ 445\ 9450\ 434\ 330\ 0901\ 096\ 928\ 0253\ 527\ 518\ 3328\ 988\ 446\ 1508\\ 940\ 424\ 8265\ 018\ 193\ 8515\ 625\ 357\ 9639\ 961\ 899\ 3967\ 905\ 496\ 6380\\ 032\ 223\ 4872\ 396\ 701\ 8485\ 186\ 439\ 0591\ 045\ 756\ 2726\ 246\ 419\ 5387 \end{array}$


The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:



The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:



The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:



The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:

$$G = 3 \underbrace{\uparrow \cdots \cdots \uparrow 3}_{3 \underbrace{\uparrow \cdots \cdots \uparrow 3}}$$

$$2 \uparrow \uparrow 4 = 2^{2^2} = 65536$$

$$2 \uparrow \uparrow \uparrow 4 = 2 \uparrow \uparrow (2 \uparrow \uparrow (2 \uparrow \uparrow 2))$$

$$2 \uparrow \uparrow \uparrow \uparrow 4 = 2 \uparrow \uparrow \uparrow (2 \uparrow \uparrow \uparrow (2 \uparrow \uparrow \uparrow 2))$$

$$3 \uparrow \cdots \uparrow 3$$

$$3 \uparrow \cdots \uparrow 3$$

$$3 \uparrow \cdots \uparrow 3$$

The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:

$$G = 3 \underbrace{\uparrow \cdot \cdots \cdot \uparrow 3}_{3 \underbrace{\uparrow \cdot \cdots \cdot \uparrow 3}}$$

$$2 \uparrow \uparrow 4 = 2^{2^2} = 65536$$

$$2 \uparrow \uparrow \uparrow 4 = 2 \uparrow \uparrow (2 \uparrow \uparrow (2 \uparrow \uparrow 2))$$


$$2 \uparrow \uparrow \uparrow \uparrow 4 = 2 \uparrow \uparrow \uparrow (2 \uparrow \uparrow \uparrow (2 \uparrow \uparrow \uparrow 2))$$

$$3 \uparrow \cdot \cdots \cdot \uparrow 3$$

$$3 \uparrow \cdot \cdots \cdot \uparrow 3$$

$$3 \uparrow \cdot \cdots \cdot \uparrow 3$$

The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:



The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:

$$G = 3 \underbrace{\uparrow \cdots \cdots \uparrow 3}_{3 \underbrace{\uparrow \cdots \cdots \uparrow 3}}$$

$$2 \uparrow \uparrow 4 = 2^{2^2} = 65536$$

$$2 \uparrow \uparrow \uparrow 4 = 2 \uparrow \uparrow (2 \uparrow \uparrow (2 \uparrow \uparrow 2))$$

$$2 \uparrow \uparrow \uparrow \uparrow 4 = 2 \uparrow \uparrow \uparrow (2 \uparrow \uparrow \uparrow (2 \uparrow \uparrow \uparrow 2))$$

$$3 \uparrow \cdots \uparrow 3$$

$$3 \uparrow \cdots \uparrow 3$$

$$3 \uparrow \cdots \uparrow 3$$

The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:

The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:

$$G = 3 \underbrace{\uparrow \cdots \cdots \uparrow 3}_{3 \underbrace{\uparrow \cdots \cdots \uparrow 3}}$$

$$2 \uparrow \uparrow 4 = 2^{2^2} = 65536$$

$$2 \uparrow \uparrow \uparrow 4 = 2 \uparrow \uparrow (2 \uparrow \uparrow (2 \uparrow \uparrow 2))$$

$$2 \uparrow \uparrow \uparrow \uparrow 4 = 2 \uparrow \uparrow \uparrow (2 \uparrow \uparrow \uparrow (2 \uparrow \uparrow \uparrow 2))$$

$$3 \uparrow \cdots \uparrow 3$$

$$3 \uparrow \cdots \uparrow 3$$

$$3 \uparrow \cdots \uparrow 3$$

The depicted coloring of the edges connecting any two vertices of the cube is **monophilic** because it contains a singly-colored plane region spanned by four vertices. But not all colorings of the cube are monophilic, and similarly for the four-, five-, ..., and 13-dimensional cube. In contrast, any coloring of the G-dimensional cube is monophilic. The definition of G uses **hyperoperators**:

