

# Non-regular languages

(Pumping Lemma)

Non-regular languages

$$\{a^n b^n : n \geq 0\}$$

$$\{vv^R : v \in \{a,b\}^*\}$$

Regular languages

$$a^* b$$

$$b^* c + a$$

$$b + c(a + b)^*$$


*etc...*

How can we prove that a language  $L$  is not regular?

Prove that there is no DFA or NFA or RE that accepts  $L$





**Difficulty:** this is not easy to prove  
(since there is an infinite number of them)

**Solution:** use the Pumping Lemma !!!






# The Pigeonhole Principle




4 pigeons



3 pigeonholes



A pigeonhole must  
contain at least two pigeons



$n$  pigeons





.....




$m$  pigeonholes

$n > m$



.....





# The Pigeonhole Principle

$n$  pigeons


$m$  pigeonholes

$$n > m$$

There is a pigeonhole  
with at least 2 pigeons



.....






The Pigeonhole Principle

and


DFAs

Consider a DFA with 4 states



Consider the walk of a “long” string:  $aaaaab$   
(length at least 4)


A state is repeated in the walk of  $aaaaab$



The state is repeated as a result of the pigeonhole principle

Walk of  $aaaaab$

Pigeons:  
(walk states)



Are more than

Nests:  
(Automaton states)




Repeated  
state


Consider the walk of a “long” string:  $aabb$   
(length at least 4)

Due to the pigeonhole principle:

A state is repeated in the walk of  $aabb$





The state is repeated as a result of the pigeonhole principle




**In General:** If  $|w| \geq \# \text{states of DFA}$ ,  
by the pigeonhole principle,  
a state is repeated in the walk  $w$

Walk of  $w = \sigma_1 \sigma_2 \cdots \sigma_k$



$$|w| \geq \# \text{states of DFA} = m$$






# The Pumping Lemma

Take an **infinite** regular language  $L$   
(contains an infinite number of strings)

There exists a DFA that accepts  $L$



$m$   
states


Take string  $w \in L$  with  $|w| \geq m$

(number of  
states of DFA)

then, at least one state is repeated  
in the walk of  $w$

Walk in DFA of

$$w = \sigma_1 \sigma_2 \cdots \sigma_k$$




Repeated state in DFA

There could be many states repeated


Take  $q$  to be the first state repeated

One dimensional projection of walk  $w$  :




We can write  $w = xyz$

One dimensional projection of walk  $w$  :




In DFA:  $w = x y z$

contains only  
first occurrence of  $q$



Observation:  $\text{length } |xy| \leq m$  number of states of DFA




Unique States

Since, in  $xy$  no state is repeated (except  $q$ )

Observation:  $\text{length } |y| \geq 1$


Since there is at least one transition in loop






We do not care about the form of string  $z$

$z$  may actually overlap with the paths of  $x$  and  $y$



Additional string: The string  $xz$   
is accepted


Do not follow loop



Additional string:


The string  $x y y z$   
is accepted

Follow loop  
2 times



Additional string: The string  $x y y y z$   
is accepted

Follow loop  
3 times



In General:


The string

$x y^i z$

is accepted

$i = 0, 1, 2, \dots$


Follow loop  
 $i$  times





Therefore:

$$x y^i z \in L \quad i = 0, 1, 2, \dots$$




Language accepted by the DFA



In other words, we described:



The Pumping Lemma !!!



# The Pumping Lemma:

- Given a infinite regular language  $L$
- there exists an integer  $m$  (critical length)
- for any string  $w \in L$  with length  $|w| \geq m$
- we can write  $w = x y z$
- with  $|x y| \leq m$  and  $|y| \geq 1$
- such that:  $x y^i z \in L \quad i = 0, 1, 2, \dots$



In the book:

Critical length  $m$  = Pumping length  $p$

# Applications of the Pumping Lemma

## Observation:

Every language of finite size has to be regular

(we can easily construct an NFA  
that accepts every string in the language)

Therefore, every non-regular language  
has to be of infinite size

(contains an infinite number of strings)

Suppose you want to prove that  
An infinite language  $L$  is not regular

1. Assume the opposite:  $L$  is regular
2. The pumping lemma should hold for  $L$
3. Use the pumping lemma to obtain a contradiction
4. Therefore,  $L$  is not regular

# Explanation of Step 3: How to get a contradiction

---

1. Let  $m$  be the critical length for  $L$
2. Choose a particular string  $w \in L$  which satisfies the length condition  $|w| \geq m$
3. Write  $w = xyz$
4. Show that  $w' = xy^i z \notin L$  for some  $i \neq 1$
5. This gives a contradiction, since from pumping lemma  $w' = xy^i z \in L$

Note: It suffices to show that  
only one string  $w \in L$   
gives a contradiction

You don't need to obtain  
contradiction for every  $w \in L$

# Example of Pumping Lemma application

**Theorem:** The language  $L = \{a^n b^n : n \geq 0\}$   
is not regular

**Proof:** Use the Pumping Lemma

$$L = \{a^n b^n : n \geq 0\}$$

Assume for contradiction  
that  $L$  is a regular language

Since  $L$  is infinite  
we can apply the Pumping Lemma



$$L = \{a^n b^n : n \geq 0\}$$

Let  $m$  be the critical length for  $L$

Pick a string  $w$  such that:  $w \in L$

and length  $|w| \geq m$

We pick  $w = a^m b^m$

From the Pumping Lemma:

we can write  $w = a^m b^m = x y z$

with lengths  $|x y| \leq m, \quad |y| \geq 1$

$$w = xyz = a^m b^m = \overbrace{a \dots a}^m \overbrace{a \dots a b \dots b}^m$$

$x \quad y \quad z$

Thus:  $y = a^k, \quad 1 \leq k \leq m$

$$x y z = a^m b^m$$

$$y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^i z \in L$

$$i = 0, 1, 2, \dots$$

Thus:  $x y^2 z \in L$

$$x y z = a^m b^m \quad y = a^k, \quad 1 \leq k \leq m$$

From the Pumping Lemma:  $x y^2 z \in L$

$$xy^2z = \overbrace{a \dots a a \dots a a \dots a a \dots a}^{m+k} \overbrace{b \dots b}^m \in L$$

$\underbrace{\hspace{1.5cm}}_x \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{1.5cm}}_y \quad \underbrace{\hspace{3.5cm}}_z$

Thus:  $a^{m+k} b^m \in L$

$$a^{m+k}b^m \in L \qquad k \geq 1$$

---

**BUT:**  $L = \{a^n b^n : n \geq 0\}$



$$a^{m+k}b^m \notin L$$

**CONTRADICTION!!!**

Therefore: Our assumption that  $L$   
is a regular language is not true

**Conclusion:**  $L$  is not a regular language

END OF PROOF

Non-regular language  $\{a^n b^n : n \geq 0\}$

Regular languages

$$L(a^* b^*)$$