

EE5471: Assignment on Random Number Generation

Harishankar Ramachandran
EE Dept, IIT Madras

February 10, 2020

1 Reading Assignment

Chapter on Random Numbers

2 The Assignment

1. Implement the function to generate normally distributed random numbers in Python. Plot a histogram of the random numbers generated and compare to the theoretical pdf.
2. Use the Chi-squared test and determine if 100 random numbers generated above belong to a gaussian (zero mean, unit variance) distribution. Vary the bin sizes and the number of random numbers and determine the optimal test. Apply the test to random numbers generated from the above function and offset by 0.01. When can the chi-squared test distinguish between the correct random numbers and the offset ones?
3. Repeat Q2 with the KS test.
4. Consider now that the function in Q1 was the sum of two random distributions:

$$x = \begin{cases} \text{randn()} & 99.9\% \text{ of the time} \\ \text{randn()}*500 & 0.1\% \text{ of the time} \end{cases}$$

How do the two tests handle the distribution? Do they recognise that this is no longer a gaussian.

5. Consider the following function:

$$f(x,y) = u^2 + v^2 \quad (1)$$

where

$$\begin{pmatrix} u \\ v \end{pmatrix} = \begin{pmatrix} \cos \alpha & \sin \alpha \\ -\sin \alpha & \cos \alpha \end{pmatrix} \begin{pmatrix} x \\ y \end{pmatrix} - \begin{pmatrix} 0.5 \\ 0.5 \end{pmatrix} \quad (2)$$

where $\alpha = \pi \sin \left(10 \left(\sqrt{x^2 + y^2} - 0.5 \right) \right)$. We wish to compute

$$I = \int_{|f|<1} dx dy$$

- (a) Write a Python function to implement this function of two variables, x and y .
- (b) Plot a contour plot (use `contourf` to get filled contours) of the function with contour values `[0.0, 1.0]` and consider the region corresponding to $|f| < 1$.
- (c) Find a covering function that contains all the places where $|f| < 1$, and use it to estimate the integral above.