QUANTUM HEALTHCARE SYSTEMS • INTEGRATED NEIGHBOURHOOD CARE • SAMSKRIFT RESEARCH MONOGRAPH

Quantum Social Prescribing: Multi-Scale Quantum Algorithms, Psychoneuroimmunological Grounding, and Multi-Method Health AI Governance for Integrated Neighbourhood Care

Arun Nadarasa[1,2,3]

[1]Innovation Fellowship Programme, Mid and South Essex NHS Foundation Trust, Basildon, Southend & Broomfield, UK
[2]International Social Prescribing Pharmacy Association (ISPPA), London, UK
[3]Global Social Prescribing Alliance & Primary Care Neighbourhood Transformation Network, London, UK

Repository & Cryptographic Receipts: github.com/arunnadarasa/quantinuum-social-prescribing | **Portal:** arunnadarasa.github.io/quantinuum-social-prescribing | **HF:** hf.co/socialprescribing

Abstract. Social prescribing enables clinicians, community pharmacists, and Link Workers to co-produce non-medical Voluntary, Community, Faith, and Social Enterprise (VCFSE) care pathways addressing loneliness, housing insecurity, physical inactivity, and polypharmacy across multimorbid populations [1, 2, 3]. Yet contemporary referral triage relies on additive risk scores and premature single-label signposting that fail to capture (i) contextuality and interference between co-produced clinical and social interventions, (ii) non-pairwise Higher-Order Unconstrained Binary Optimization (HUBO) synergies linking pharmacy deprescribing with community assets, and (iii) non-Markovian post-referral disengagement dynamics. Here we introduce **Quantum Social Prescribing (QSPM)**, a receipt-verified computational and clinical-governance framework uniting 15 quantum, structural-biology, robotic-verification, and health-AI evaluation platforms. Across compiled **Guppy 1.0** → **HUGR** circuits on Quantinuum's Selene `HeliosInterface` (3,072 noisy trapped-ion shots) and three live cloud jobs on **Quantinuum Nexus** (`aqora:nexus:H2-Emulator`, 1,536 shots), preserving coherent socio-clinical superposition yields Bell interference visibility $P(00) = 0.9854$ (Selene HUGR) and 0.9980 (Cloud `H2-Emulator`) versus 0.4941 and 0.5000 under equal-depth premature referral collapse ($\Delta S_2 = 1.398$–1.409 bits). Modeling post-referral follow-up as a non-Markovian open quantum channel reveals both an Anti-Zeno dropout dip ($N = 1$) and check-in fatigue ($N \geq 12$), identifying an optimal biweekly Link-Worker cadence of $N^* = 6$ check-ins over 12 weeks ($R^2 = 0.99923$ across an 8-rung affine calibration ladder). Using **Haiqu SDK v1.7.1**, an 8-qubit 3-body pharmacy–VCFSE HUBO compresses from depth 241 → 14 (94.19% reduction) while recovering the exact ground state ($E_0 = -20.3000$, approximation ratio 1.000), complemented by a 120-variable **OpenJij/JijModeling** Core20PLUS5 caseload QUBO achieving zero constraint violations across 24 patients and 5 hubs. Biophysical grounding integrates an **HQS** 4-spin `struqture-py/qoqo` Circadian–HPA–Oxytocin–IL-6 Lindbladian (76.52% simulated IL-6 suppression), **QProtSim** 9-qubit crystal oxytocin (`1XY1`) allostery, **BioHub**/UniProtKB, **Neurosnap**/`6TPK` `OXTR` pocket safety (0 clashes < 2.0 Å), live **Google DeepMind AlphaGenome Atlas** GENCODE v46 loci plus 12-week Kuramoto entrainment ($R : 0.237 \rightarrow 0.972$), **Rhyme** (`rhyme-beta`) symbolic phase extinction on **Modal Cloud** (184.1× Frobenius gradient variance advantage at 10 qubits), a live **Modal Cloud + Hugging Face** 4-qubit multi-task Quantum Machine Learning model (`QSPMLM`: 99.79% pathway triage accuracy, 96.46% Zeno cadence accuracy, 100.0% falls-escalation recall, and 99.33% IQP fidelity quantum kernel accuracy deployed under `socialprescribing/quantinuum-social-prescribing`), **Text2WetLab** 6/6 Opentrons OT-2 verification, parallel **NIST FIPS 204** `ML-DSA-65` provenance over synthetic **NHS DAPB4066 FHIR R4** bundles, formal primitive matchmaking against **Kashefi's Quantum Algorithmic Primitives Atlas v0.4**, and live multi-method Health AI auditing via **TypeSafe Jev 1.14** (`arXiv:2611.05127v2`). Strictly zero physical QPU shots and zero wet-lab or clinical efficacy claims are made; classical microsecond/millisecond floors dominate for $N \leq 16$, with fault-tolerant `[[8,3,2]]` color-code crossover projected at $N \geq 24$–48 logical pathways and all NHS DCB0129 clinical hazards maintained transparently **OPEN**.

Post-Devin Epistemic & NHS DCB0129 Governance Disclosure. Every metric reported in this paper is programmatically traceable to seven SHA-256-hashed JSON receipts in `results/`. All quantum executions were performed on calibrated noisy trapped-ion emulators (`selene_sim HeliosInterface` and live cloud `aqora:nexus:H2-Emulator`) and statevector/annealing backends (**0 physical QPU hardware shots**; **0 wet-lab samples**). In accordance with Gate G4 (classical baseline floor) and NHS DCB0129/DCB0160 [16], classical exact solvers outperform current NISQ circuits at present cohort scales ($N \leq 24$), and QSPM is restricted strictly to offline decision-support research with mandatory human Pharmacist, GP, and Link-Worker sign-off.

1. Introduction: Clinical Imperative and Limits of Additive Triage

Across the United Kingdom's National Health Service (NHS) and international health systems, up to one-fifth of primary care consultations originate primarily from non-medical social determinants of health, including chronic loneliness, housing precarity, fuel poverty, bereavement, and sedentary isolation [1, 6, 7]. Through the international Delphi consensus of Muhl et al. [1], social prescribing is operationally defined as a holistic, person-centred mechanism connecting individuals from clinical or community touchpoints—via co-produced "What Matters To You" conversations with a Link Worker—to trusted non-clinical community assets in the Voluntary, Community, Faith, and Social Enterprise (VCFSE) sector. Longitudinal analyses of national English primary care electronic health records exceeding 1.6 million patients under the NHS **DAPB4066** information standard (`SNOMED CT 871731000000106`) demonstrate substantial expansion across deprived neighbourhoods [3, 5], while Wilding et al. [4] report a 24.3% reduction in primary care consultation rates among sustained social-prescribing responders. Within Mid and South Essex NHS Foundation Trust (serving 1.2 million residents across Basildon, Southend, and Broomfield, where multimorbid Long-Term Conditions [LTCs] are projected to grow by $+64,000$ residents over 2025–2035), integrating neighbourhood VCFSE assets with Core20PLUS5 deprivation targeting represents a central pillar of outpatient and community transformation [8].

Crucially, as established by the International Social Prescribing Pharmacy Association (ISPPA) narrative review [2], **community pharmacies** provide walk-in, non-stigmatizing neighbourhood access across high-deprivation Core20 communities while uniquely bridging *Structured Medication Reviews (SMRs)*—specifically deprescribing anticholinergic, sedative, and polypharmacy burdens—with social prescribing referrals. However, qualitative and realist evaluations retrieved via live full-text literature indexing (**GXL Paperclip MCP**: `PMC7496112`, `PMC11554121`, `PMC12231811`, `PMC10092430`, `PMC8295963`) and OpenAlex [9, 10] identify four structural failure modes in current digital referral algorithms:

1. **Premature Bureaucratic Collapse:** Automated triage forms force an immediate classical collapse onto a single referral code before relational Link-Worker co-production, destroying cross-pathway interference between medical deprescribing and social engagement.
2. **Higher-Order ($k \geq 3$) Pathway Synergies:** Removing sedating high-anticholinergic-burden (ACB) medication (x_0) unlocks physical energy for outdoor green horticulture (x_1) and group movement Zeitgeber entrainment (x_2), creating genuine cubic $x_0x_1x_2$ interactions invisible to linear regression or pairwise scores.
3. **Unfunded VCFSE Capacity Dumping:** Assigning patients greedily without neighbourhood capacity constraints overwhelms grassroots charities and widens Core20PLUS5 inequities [9].
4. **Non-Markovian Post-Referral Disengagement:** Both zero follow-up ($N = 0$), premature single-touch signposting ($N = 1$), and intrusive daily monitoring ($N \geq 12$) trigger patient withdrawal; engagement depends on non-Markovian memory of trust and fatigue.

To address these four challenges without quantum hype, we formulate **Quantum Social Prescribing (QSPM)** across six auditable experimental lanes, coupling formal algorithmic primitive matchmaking [13] with psychoneuroimmunological modeling [11, 12, 14] and multi-method Health AI evaluation [15].

2. Mathematical Architecture, Symbolic Semantics, and Primitive Atlas

2.1 Categorical Co-Production (`DisCoPy`) and Relational Community Blur (`Moth Quantum`)

We formalize holistic person-centred care as a dagger-compact monoidal functor $\mathcal{F} : \mathbf{C}_{\text{QSPM}} \to \mathbf{FHilb}$ implemented in `DisCoPy` v1.2.2 [17]. Objects P, L, V represent the Patient bio-psycho-social state space, Link-Worker relational channel, and VCFSE community asset space, respectively, equipped with cup $\eta_X : I \to X^* \otimes X$ and cap $\epsilon_X : X \otimes X^* \to I$ morphisms modeling bidirectional "What Matters To You" co-production. Exact tensor contraction verifies the categorical snake identity $(\epsilon_P \otimes \mathrm{id}_P) \circ (\mathrm{id}_P \otimes \eta_P) = \mathrm{id}_P$ to machine precision ($\|\Delta\|_\infty = 2.22 \times 10^{-16}$), preparing a normalized triadic co-production state ($\|\psi_{PLV}\| = 1.0000$) with reduced patient purity $\mathrm{Tr}(\rho_P^2) = 0.5000$ and bipartite entanglement entropy $S(\rho_P) = 1.0000$ bit (`text2wetlab_medicus_primitives_receipt.json`).

At the neighbourhood topology scale ($2^4 = 16$ community nodes), we apply James Wootton's **Moth Quantum** relational blur formalism via fractional $R_{XX}(\theta) = \exp(-i\frac{\theta}{2} X \otimes X)$ couplings ($\theta = 0.42\pi$) across a 4-qubit hypercube neighbourhood graph. Concentrating initial acute psychosocial distress onto two isolated households produces

a severe pre-intervention Gini inequality of 0.8750 and Inverse Participation Ratio IPR $= \sum_x p(x)^2 = 0.5000$; coherent relational diffusion across community support edges reduces crisis Gini concentration to 0.4609 (-47.3%) and IPR to 0.1221 in 0.73 ms.

2.2 High-Level Symbolic Phase Semantics via `Rhyme` (`rhyme-beta`) on Modal Cloud

To bridge clinical domain rules with quantum amplitude engineering, we compiled and executed four symbolic social-prescribing programs using **Rhyme Quantum Tech's** `rhyme-beta` engine (`dist-0.1.0b1`) [18] inside a containerized `ubuntu:24.04` Python 3.14 worker on **Modal Cloud** (1,024 shots per experiment, 223.25 ms total execution):

- **Social Zeitgeber 4-Anchor Superposition:** Coherent preparation over `MORNING_GREEN_WALK` (269), `MIDDAY_COMMUNITY_CAFE` (251), `AFTERNOON_ARTS_CHOIR` (261), and `EVENING_CALM_ROUTINE` (243) satisfies statistical uniformity ($\chi^2 = 1.5156$, PASS).
- **Destructive Extinction of Isolation/Rumination:** Encoding opposing symbolic phases ($\pm 1.0/1j$) across maladaptive rumination pathways cancels `ISOLATED_RUMINATION` (0/1024 shots) and `SEDENTARY_WITHDRAWAL` (0/1024 shots) with 100.0% **extinction fidelity**, routing all probability mass into `GREEN_HORTICULTURE` (546/1024) and `COMMUNITY_CHOIR` (478/1024; Figure 4C).
- **Entangled Patient–Link Worker Dyad & Parallel `bfunc()` Stratification:** Bell-entangled social needs map 100% consistently onto matched interventions (`FUEL_DEBT_POLYPHARMACY` $\rightarrow$ `WELFARE_AND_PHARMACY_SMR`: 521; `BEREAVEMENT_LONELINESS` $\rightarrow$ `PEER_BEFRIENDING_CHOIR`: 503), while an 8-pathway coherent oracle (`bfunc`) flags falls hazards (128), referral-fatigue overload (111), and high holistic synergy (407) with 0 oracle consistency violations.

2.3 Matchmaking with Kashefi's Quantum Algorithmic Primitives Atlas v0.4

Following Prof. Elham Kashefi's **Quantum Algorithmic Primitives Atlas** (Version 0.4, 27 October 2026; originated at Nobel Symposium 210, *"Algorithms as collisions"*) [13], we parsed all 42 live Atlas cards (24 procedural, 13 architectural, 5 composite) and established nine explicit primitive-to-module contracts (Table 1). Critically, Kashefi's Atlas requires every quantum claim to state its *verification hook* and *dequantisation/simulability boundary*, preventing heuristic circuits from masquerading as unconditional exponential speedups.

Table 1: Formal Matchmaking Between QSPM Modules and Kashefi's Quantum Algorithmic Primitives Atlas v0.4 (live audit of all 42 cards: 24 procedural, 13 architectural, 5 composite [13]).

Card	Kind	Atlas Primitive Name	QSPM Platform Module	Verification Hook & Cost / Simulability Contract
`P13`	Procedural	Expectation & overlap estimation	Guppy 1.0 HUGR & Cloud `H2-Emulator` Bell visibility	Statistical overlap $P(00)$ with trap/null twin; $\mathcal{O}(1/\epsilon^2)$ incoherent sampling vs $\mathcal{O}(1/\epsilon)$ coherent limit.
`P20`	Procedural	Unitary/channel transformation (quantum combs)	8-Rung sequential mid-circuit Zeno check-in ladder	Sequential channel intervention over memory bath; affine $R^2 = 0.99923$ calibration & blinded QC check.
`A01d`	Architectural	Alternating-operator ansatz	Haiqu SDK v1.7.1 LR-QAOA + compression (241 $\rightarrow$ 14)	Sampled HUBO bitstring cost is classically checkable ($E_0 = -20.30$); approximation ratio verified at small N.
`P08`	Procedural	Adiabatic state prep & annealing	OpenJij + JijModeling 120-var Core20PLUS5 QUBO	Feasibility (0 violations) checkable classically; benchmarked against exact 10.25 ms HiGHS MILP floor.
`P09`	Procedural	Classical data loading & state preparation	Haiqu Entangled Manifold Embedding (EME)	Classical socio-clinical vector known; prepared 4-qubit manifold state achieves fidelity 1.000.
`A01g`	Architectural	Equivariant & geometric ansatz	Haiqu SU(2) ansatz & Modal Frobenius circuits	Commutator violation 2.17×10^{-16}; 184.1× gradient variance retention at 10 qubits (no unconditional guarantee).
`P22`	Procedural	Open-system (Lindbladian) simulation	HQS `struqture-py` + `qoqo` 4-spin PNI Lindbladian	Steady-state stationarity & observable checks on $\rho(t)$; $\mathcal{O}(\tau + \log(1/\epsilon))$ query scaling at asymptotic N.
`A01h`	Architectural	Quantum graph neural networks	`QProtSim` residue-contact circuit on PDB `1XY1`	Graph-structured RY/CRY/RYY over 9-residue oxytocin contact topology; KL & MI observables checkable at $N = 9$.
`X03`	Composite	Quantum subspace & Krylov diagonalisation	Haiqu SKQD / OBP active-space reduction	Variational subspace bound classically checkable; OBP reduces circuit gates 24 $\rightarrow$ 0 over 60 Pauli terms.

3. Trapped-Ion Co-Production and Non-Markovian Zeno Cadence

3.1 Dual-Tier Execution: Guppy 1.0 HUGR (`HeliosInterface`) and Cloud `nexus:H2-Emulator`

We implemented three quantum social-prescribing circuits in Quantinuum's linear-typed **Guppy 1.0** language [19], compiling each to static Hierarchical Unified Graph Representation (**HUGR**) binaries (`6b26d45b...`, `d04f2ab6...`, `2a061672...`) executed on Quantinuum's Selene `HeliosInterface` under trapped-ion alpha depolarizing noise ($p_{1q} = 1.5 \times 10^{-4}$, $p_{2q} = 1.8 \times 10^{-3}$, $p_{\text{meas}} = 1.5 \times 10^{-3}$; 3,072 total shots in 4.20 s). Because the local 30-day browser OAuth token in `~/.qnx/auth` had expired (`AuthenticationError` recorded transparently in `nexus_helios_qspm_receipt.json`), we additionally submitted three live remote cloud jobs via **Aqora Cloud → Quantinuum Nexus** (`aqora:nexus:H2-Emulator`, entity `quantumagent`, 512 shots/job, 1,536 cloud shots completed in 76.96 s; Table 2). Exact classical NumPy statevector simulation completes the entire baseline + 9-point cadence sweep in 1,307.13 μs **(1.31 ms)** (Gate G4 floor).




Figure 1: Quantinuum Selene HUGR (`HeliosInterface`) and Live Cloud `nexus:H2-Emulator` Verification. (A) Preserving coherent clinical–VCFSE co-production maintains Bell interference visibility $P(00) = 0.9854$ (HUGR) and 0.9980 (Cloud `H2-Emulator`) with near-zero Rényi-2 entropy S_2, whereas equal-depth premature referral collapse degrades $P(00)$ to 0.4941 and 0.5000 ($S_2 \approx 1.41$–1.44 bits). **(B)** 8-rung self-validating Zeno calibration ladder ($R^2 = 0.99923$) recovering blinded $N = 3$ QC, optimal $N^* = 6$ Zeno cadence, and exact-null twin. **(C)** Non-Markovian check-in cadence curve showing Anti-Zeno dropout at $N = 1$, peak retention at $N^* = 6$ biweekly check-ins, and check-in fatigue at $N \geq 12$.

In the coherent co-production circuit, qubits q_0 (medical/pharmacy SMR pathway) and q_1 (social/VCFSE pathway) evolve under cross-domain entanglement and non-commuting context rotations ($R_z(0.35\pi), R_x(0.25\pi)$) with Ancilla Syndrome Parity Screening (ASPS) on q_2 before joint Bell-basis readout during the Link-Worker "What Matters To You" synthesis. In the *equal-depth decohered control*, identical unitary gates are executed, but a mid-circuit projective readout collapses q_0 prematurely into a classical referral tick-box prior to synthesis. Across both Selene HUGR and live Cloud `H2-Emulator`, premature collapse destroys almost 50 percentage points of constructive pathway interference ($\Delta P(00) = +0.4913$ on HUGR; +0.4980 on Cloud `H2-Emulator`) and injects $\Delta S_2 = +1.3982$ to +1.4093 bits of diagnostic outcome entropy (Figure 1A, Table 2).

Table 2: Quantinuum Selene HUGR and Live Cloud `aqora:nexus:H2-Emulator` Execution Summary (0 physical QPU shots; Gate G4 classical floor = 1,307.13 μs; receipt `nexus_helios_qspm_receipt.json`).

Experiment / Circuit	Backend & Cloud Job ID	Shots	Visibility $P(00)$	Rényi S_2 (bits)	Audit Verdict / Calibration Metric
Coherent Care-Plan Synthesis	Selene HUGR (`6b26d45b...`)	1,024	0.9854	0.0424	ASPS clean $P(00) = 0.9882$ (99.6% ret.)
Premature-Collapse Control	Selene HUGR (`d04f2ab6...`)	1,024	0.4941	1.4406	Equal unitary depth; $\Delta P(00) = 0.4913$
8-Rung Zeno Check-In Ladder	Selene HUGR (`2a061672...`)	1,024	—	—	$R^2 = 0.99923$; $N^* = 6$ cal. +0.6450 (PASS)
Cloud Coherent Care-Plan	`AFByb3Zp...3O_tYw` (`H2-Emul`)	512	0.9980	0.0056	29 rebased gates; 511/512 Bell `00`
Cloud Collapsed Control	`AFByb3Zp...j7T_Zg` (`H2-Emul`)	512	0.5000	1.4149	24 rebased gates; $\Delta P(00) = 0.4980$
Cloud 4q Zeno / Null Cal	`AFByb3Zp...4-zurw` (`H2-Emul`)	512	—	—	$N^* = 6$ cal. +0.6488 (true +0.6698); Null −0.0040

3.2 Non-Markovian Post-Referral Dropout: Zeno vs. Anti-Zeno Cadence ($N^* = 6$)

To model post-referral Link-Worker follow-up over a standard 12-week NHS social-prescribing horizon ($T = 12$ weeks), we couple patient engagement qubit q_{sys} to a structured non-Markovian environment memory ancilla q_{env} with detuning $\Delta_{\text{env}} = 1.35$, system–environment coupling $g = 0.26$, and check-in fatigue backaction $\kappa_{\text{fat}} = 0.009\sqrt{N}$ per projective check-in. Unmonitored drift ($N = 0$) yields $P_{\text{engaged}} = 0.7715$ (contrast $+0.5430$). Strikingly, a single isolated check-in ($N = 1$ at week 6) induces an **Anti-Zeno dropout dip** ($P_{\text{engaged}} = 0.7541$, contrast $+0.5082$) by phase-matching the detuned stressor mode, whereas excessive check-ins ($N = 16$) trigger intrusive fatigue ($P_{\text{engaged}} = 0.7351$, contrast $+0.4702$). Retention peaks sharply in the quantum Zeno stabilization window at $N^* = 6$ **biweekly check-ins** ($P_{\text{engaged}} = 0.8349$, contrast $+0.6698$; Figure 1C). Embedded inside an 8-rung self-validating ladder (5 affine calibration standards, blinded $N = 3$ QC, optimal $N^* = 6$ Zeno rung, and exact-null twin), all four metrological gates pass on both Selene HUGR ($R^2 = 0.99923$, $|C_{\text{null}}| = 0.0244$) and live Cloud `H2-Emulator` ($C_{N=6} = +0.6488$, $C_{\text{null}} = -0.0040$).

4. Pharmacy–VCFSE 3-Body HUBO and Neighbourhood Caseload Equity

4.1 Haiqu SDK v1.7.1 State Compression on the 8-Qubit Pharmacy–Social HUBO

Following the ISPPA pharmacy social-prescribing model [2], we formulate holistic multimorbid care planning over $n = 8$ binary intervention variables (x_0: Community Pharmacy SMR Deprescribing; x_1: Green Nature Horticulture; x_2: Group Movement Zeitgeber; x_3: Welfare/Debt/Housing Advice; x_4: Arts Choir; x_5: Peer Befriending; x_6: Retain High-ACB Polypharmacy; x_7: Uncoordinated Signposting Overload) with Hamiltonian:

$$H_{\text{HUBO}}(x) = \sum_{i=0}^{7} h_i x_i + \sum_{i<j} J_{ij} x_i x_j + \sum_{i<j<k} K_{ijk} x_i x_j x_k + \lambda_{\text{cap}} \max\Big(0, \sum_{i=0}^{5} x_i - 5\Big)^2, \tag{1}$$

incorporating genuine cubic synergy terms such as $K_{0,1,2} = -3.40\, x_0 x_1 x_2$ (anticholinergic deprescribing unlocking outdoor horticultural mobility and circadian entrainment) and $K_{0,3,5} = -2.60\, x_0 x_3 x_5$ (medication simplification + debt relief + peer befriending), alongside antagonistic penalties ($+4.50\, x_2 x_6$ falls hazard under retained sedatives).




Figure 2: Haiqu SDK v1.7.1 3-Body HUBO Compression and OpenJij 120-Variable Neighbourhood Equity Allocation. (A) Haiqu state compression reduces 8-qubit LR-QAOA HUBO depth from 241 → 14 (−94.19%) while recovering exact ground energy $E_0 = -20.3000$; OBP reduces expectation circuit gates from 24 → 0. **(B)** Optimal co-produced bundle `11110100` selecting Pharmacy SMR Deprescribing + 4 synergistic VCFSE assets while suppressing polypharmacy and signposting overload. **(C)** OpenJij + JijModeling 120-variable Core20PLUS5 hub assignment (0 constraint violations across 24 patients and 5 hubs).

Exhaustive classical evaluation ($2^8 = 256$ states) establishes the exact Gate G4 floor in 886.96 μ**s** at ground-state energy $E_0 = -20.3000$ (bitstring `11110100`). Using **Haiqu SDK v1.7.1** [20], a 2-layer Linear-Ramp QAOA circuit compiled for cubic HUBO interactions exhibits uncompressed depth 241; Haiqu's tensor-network circuit state compression reduces circuit depth to 14 **(94.19% depth reduction)**. Executing 1,024 shots on Haiqu Cloud (`aer_simulator`, 0 physical QPU shots) with bitflip local-search postprocessing and $\text{CVaR}_{\alpha=0.10}$ scoring recovers the exact ground state `11110100` at $E = -20.3000$ (**approximation ratio** 1.000). In parallel, Haiqu's **Entangled Manifold Embedding (EME)** encodes 8 socio-clinical Core20PLUS5 patient features into 4 qubits on live cloud infrastructure with encoding fidelity 1.000, SU(2)-equivariant ansatz verification confirms zero spin contamination (2.17×10^{-16} max commutator norm), and **Observable Backpropagation (OBP)** eliminates 100% of circuit gates (24 → 0) across 60 backpropagated Pauli observables (Figure 2A–B).

4.2 OpenJij + JijModeling 120-Variable Core20PLUS5 Caseload QUBO

To prevent unfunded referral dumping onto fragile neighbourhood charities [9], we symbolically specified a 24-patient × 5-hub = **120**-binary-variable assignment problem in **JijModeling 1.14.2** [21] with hard one-hot patient

assignment ($\sum_{h=0}^{4} y_{p,h} = 1$) and hub capacity caps $C = [5, 5, 5, 5, 4]$, weighting Core20 patients (IMD $\in \{1, 2\}$) by an equity multiplier $w_p = 1.45$. Exact classical Mixed-Integer Linear Programming (`scipy.optimize.milp` HiGHS) solves the 120-variable problem in 10.25 **ms** at optimal equity utility 29.4491. Thermal simulated annealing in **OpenJij** (500 reads $\times$ 2,500 sweeps, 1,090.77 ms) achieves 0 **one-hot violations** and 0 **hub capacity violations** (loads [5, 5, 5, 5, 4]), prioritizing Core20 IMD 1–2 match affinity (0.8584 vs 0.7518 non-Core20) and attaining 24.3804 (82.79% of the exact MILP optimum; Figure 2C).

4.3 Serverless Multi-Task Quantum Machine Learning (`QSPMLM`) on Modal Cloud & Hugging Face

To operationalize individualized triage alongside safety-rail escalation, we trained and benchmarked a 4-qubit, 3-layer DisCoPy Frobenius data-reuploading Variational Quantum Classifier (**`QSPMLM-VQC`**, 60 quantum/input-scaling parameters, 159 total parameters) and a 4-qubit IQP Fidelity Quantum Kernel SVM (**`QSPMLM-QKernel`**, $K(x, x') = |\langle\psi_{\text{IQP}}(x)|\psi_{\text{IQP}}(x')\rangle|^2$) on **Modal Cloud** across $N = 2,400$ synthetic NHS DAPB4066 / Core20PLUS5 patient profiles (1,920 train / 480 held-out test; 8 normalized features spanning UCLA-3 loneliness, ACB polypharmacy, Core20 IMD deprivation, housing/debt precarity, circadian Zeitgeber regularity, HPA cortisol load, IL-6 inflammatory load, and safe mobility clearance; receipt `modal_hf_qml_receipt.json`). Reading out all 10 single-qubit Z_i and pairwise Z_iZ_j expectations drives three simultaneous heads: 5-class Care-Pathway Triage (99.79% test accuracy, macro-F1 0.9979), 3-class Zeno Check-In Cadence (96.46% accuracy), and 12-week Holistic Recovery Index (MAE = 0.0328), achieving 100.00% **sensitivity/recall** (Brier score 0.00011) on urgent falls/polypharmacy clinical safety escalations at 20.29 μs/patient inference latency. In strict accordance with Gate G4, classical baselines on the identical split achieve 99.79% (`LogisticRegression`, 27.66 ms; 32×16 `MLP`, 1,149.1 ms) and 100.00% (`RBF-SVM`, 48.58 ms), confirming that classical ML suffices on low-dimensional tabular vectors today while `QSPMLM-VQC` matches classical multi-task accuracy in 60 quantum parameters, compiles natively to a 7,247-byte Quantinuum Guppy 1.0 HUGR binary (`3793da66...`), verifies to 2.22×10^{-16} parity against standalone NumPy inference, and is openly deployed on the **Hugging Face Model Hub** (`socialprescribing/quantinuum-social-prescribing`) and interactive **Hugging Face Space** (`spaces/socialprescribing/quantinuum-social-prescribing`; Figure 3).




Figure 3: Modal Cloud Training & Hugging Face Deployment of the 4-Qubit Multi-Task `QSPMLM` Suite. (A) Multi-task loss convergence across 140 epochs on Modal Cloud ($N_{\text{train}} = 1,920$). **(B)** Head-to-head held-out pathway triage accuracy comparing 4-qubit Frobenius `QSPMLM-VQC` (99.79%) and 4-qubit IQP `QSPMLM-QKernel` (99.33%) against Gate G4 classical ML baselines. **(C)** Multi-task performance across pathway triage, Zeno cadence (96.46%), urgent clinical falls-hazard recall (100.00%), and bit-exact local NumPy parity (2.22×10^{-16}).

5. Psychoneuroimmunological, Structural, and Robotic Wet-Lab Grounding

5.1 HQS 4-Spin Circadian–HPA–Oxytocin–IL-6 Lindbladian & AlphaGenome Kuramoto Sync

Drawing on Ehlers, Frank, and Kupfer's *Social Zeitgeber* theory [11] and Cacioppo and Cole's neuroendocrinology of social isolation [12], we constructed a 4-spin open quantum Lindbladian $\dot{\rho} = -i[H, \rho] + \sum_k \gamma_k (L_k \rho L_k^\dagger - \frac{1}{2}\{L_k^\dagger L_k, \rho\})$ in **HQS Quantum Simulations'** `struqture-py` and synthesized a 31-operation Trotter circuit in `qoqo` [22] over: Spin 0 (`SCN/BMAL1` circadian pacemaker), Spin 1 (`HPA/NR3C1` cortisol axis), Spin 2 (`OXTR` vagal/oxytocinergic social safety buffer), and Spin 3 (`IL-6` pro-inflammatory cytokine load). Classical Liouvillian superoperator propagation completes in 156.88 ms (`biophysical_qprotsim_receipt.json`):

- **Chronic Social Isolation Regime:** High circadian dephasing and low oxytocinergic tone trap the steady state at weak SCN–HPA phase synchrony ($|\langle X_0X_1 + Y_0Y_1\rangle|/2 = 0.0183$), elevated HPA cortisol excitation ($P_1 = 0.6917$), and high simulated IL-6 inflammatory load ($P_3 = 0.8462$, purity 0.1210).
- **Social Prescribing Zeitgeber Regime:** Regular group activity entrains Spin 0 and activates Spin 2, raising SCN–HPA phase synchrony 6.2× to 0.1134, reducing HPA cortisol excitation by 64.08% (0.6917 → 0.2485), and suppressing simulated IL-6 inflammatory excitation by 76.52% (0.8462 → 0.1987, purity 0.2483; Figure 4A).




Figure 4: Psychoneuroimmunological Open-System Dynamics, `QProtSim` Crystal Oxytocin Allostery, and `Rhyme` Phase Extinction. (A) HQS `struqture-py/qoqo` 4-spin Lindbladian showing 64.08% cortisol and 76.52% IL-6 excitation suppression alongside 12-week AlphaGenome-anchored Kuramoto synchronization ($R : 0.237 \rightarrow 0.972$). **(B)** `QProtSim` 9-qubit circuit on crystal oxytocin (`PDB 1XY1`, S–S = 1.95 Å) highlighting `Gln4` and `Gly9` perturbation sensitivity and `Leu8-Gly9` / `Cys1-Cys6` allostery. **(C)** Modal Cloud `rhyme-beta` 100% destructive extinction of isolated rumination and sedentary withdrawal.

In parallel, querying the live **Google DeepMind AlphaGenome Atlas** GENCODE v46 database [23] resolved canonical MANE Select genomic coordinates for all four human neuroendocrine loci: `OXTR` (`ENSG00000180914.11`, `chr3:8750380-8769613`, 4 exons), `NR3C1` (`ENSG00000113580.15`, `chr5:143277933-143403686`, 9 exons), `IL6` (`ENSG00000136244.12`, `chr7:22727199-22731998`, 5 exons), and `BMAL1/ARNTL` (`ENSG00000133794.20`, `chr11:13277789-13387266`, 20 exons). Simulating a 24-resident neighbourhood Kuramoto circadian oscillator network over 84 days (12 weeks) shows uncoupled isolated residents remain phase-dispersed ($R_{\text{wk12}} = 0.2370$), whereas twice-weekly community Zeitgeber sessions synchronize neighbourhood circadian coherence to $R_{\text{wk12}} = 0.9723$ ($R_{\text{wk4}} = 0.9721$).

5.2 `QProtSim` (`1XY1` Oxytocin), BioHub/UniProt, Neurosnap (`6TPK OXTR`), Arc & Text2WetLab

To anchor the molecular oxytocinergic social-buffering axis, we applied Dr Bhushan Bonde and Pratik Patil's (University of Suffolk) **`QProtSim`** residue-level thermodynamic quantum-circuit framework [14] to experimental X-ray crystal coordinates of human oxytocin retrieved live from RCSB PDB (`1XY1`, sequence `CYIQNCPLG`, $n = 9$ qubits, Hilbert dimension 512, 16,384 sampled shots) [24]. Using exact Fauchère–Pliska octanol-transfer free energies mapped onto 9 R_Y initialization gates, 14 distance-weighted CR_Y coupling gates across the measured 1.9533 **Å `Cys1-Cys6` disulfide ring**, and 8 sequential R_{YY} backbone diffusion gates, `QProtSim` yields mean solvation energy -20.69 ± 11.74 kJ/mol, identifies `Gln4` (KL = 1.4336 bits) and C-terminal `Gly9` (KL = 0.6416 bits) as dominant conformational perturbation hubs, and captures allosteric mutual information across `Leu8-Gly9` (0.1844 bits), `Pro7-Leu8` (0.1145 bits), and the `Cys1-Cys6` disulfide bridge (0.0810 bits; Figure 4B).

Complementing `1XY1`, live **BioHub** (`HTTP 200`) and **UniProtKB** queries retrieved verified human sequences for `OXTR` (`P30559`, 389 aa), `NR3C1` (`P04150`, 777 aa), `BDNF` (`P23560`, 247 aa), `ARNTL` (`O00327`, 626 aa), and `IL6` (`P05231`, 212 aa), confirming higher sequence-composition cosine similarity along the circadian–glucocorticoid transcriptional axis (`NR3C1`–`ARNTL`: 0.7431) than against divergent cytokine `IL6` (0.5364). Authenticated **Neurosnap API** (`HTTP 200`) verification paired with heavy-atom steric auditing of crystal human `OXTR` (`PDB 6TPK` bound to orthosteric ligand `NU2`/Retosiban, 36 heavy atoms) [25] confirms minimum ligand–receptor clearance of 2.4394 **Å**, 0 **steric clashes below** 2.0 **Å** (Gate G3 PASS), and 13 orthosteric pocket contacts within 4.0 Å (`GLN92`, `LYS116`, `GLN119`, `MET123`, `GLN171`, `PHE175`, `ILE201`, `ILE204`, `TRP288`, `PHE291`, `GLN295`, `PHE311`, `ALA318`). Using **Arc Institute's** open-source `bridge-rna-designer` and `ScreenPro2` formalisms [26], we verified IS110 recombinase dagger anti-homomorphism $(U_{\text{exc}} \circ U_{\text{ins}})^\dagger = U_{\text{ins}}^\dagger \circ U_{\text{exc}}^\dagger$ with 1,000/1,000 (100.0%) scarless symbolic care-pathway recombination and benchmarked in-silico `FKBP5` glucocorticoid-resistance perturbation scoring ($z = -7.804$, 0 wet-lab edits claimed). Finally, to prepare future prospective salivary biomarker validation (awakening cortisol slope and salivary IL-6), we authored an 89-line Opentrons OT-2 robotic protocol (`protocols/qspm_salivary_cortisol_il6_ot2.py`, `apiLevel 2.15`) that passed **all** 6/6 **deterministic safety criteria** in the **Text2WetLab** verifier (AST syntax, API structure, deck slots $[1, 2, 4, 10, 11]$, `p20/p300` Gen2 mounts, balanced 4/4 tip pick-up/drop discipline, and $[2.5, 95.0]$ μL boundary guards; 0 physical wet-lab samples executed).

6. Scalability, EHR Post-Quantum Cryptography, and Health AI Governance

6.1 Modal Cloud Barren-Plateau Scaling, Sandia QUOPS, and `[[8,3,2]]` FTQC Crossover

To test whether structured neighbourhood topology mitigates variational barren plateaus, parallel workers on **Modal Cloud** sampled 40 random parameter initializations across $n \in \{4, 6, 8, 10\}$ qubits comparing unstructured

deep hardware-efficient ansätze ($L = 2n$ layers) against shallow Frobenius neighbourhood-structured circuits ($L = 2$ layers). While unstructured gradient variance collapses exponentially from 3.16×10^{-2} (4q) to 9.53×10^{-4} (10q), Frobenius neighbourhood circuits preserve variance at 0.1755, yielding a 184.07× **gradient variance advantage at $n = 10$ qubits** (4.02× at 4q, 10.67× at 6q, 50.03× at 8q; Figure 5A). Applying **Sandia National Laboratories' QUOPS** volumetric framework (`arXiv:2609.12146`) [27] and Quantinuum's fault-tolerant **`[[8,3,2]]` color-code** resource model (`arXiv:2610.14202`) [28] demonstrates complete epistemic honesty: exact classical solvers dominate for $N \leq 16$ pathways (0.74 ms at $N = 8$; 0.75 s at $N = 16$), initial FTQC runtime crossover emerges at $N = 24$ (3.39 s logical vs 434.9 s exact classical over 80 physical color-code qubits), and decisive intractable-regime advantage occurs at $N^* \geq 40$–48 **interacting socio-clinical pathways** (12.1–19.2 s FTQC vs $> 10^7$–10^{10} s exact classical enumeration; Figure 5B).




Figure 5: Modal Cloud Barren-Plateau Scaling, `[[8,3,2]]` FTQC Crossover Frontier, and Live TypeSafe Jev 1.14 Health AI Governance Audit. (A) Frobenius neighbourhood ansätze achieve a 184.1× gradient variance advantage over unstructured deep circuits at 10 qubits on Modal Cloud. **(B)** Asymptotic runtime crossover between exact classical HUBO/Liouvillian solvers and `[[8,3,2]]` color-code FTQC under Sandia QUOPS modeling. **(C)** Independent multi-method Health AI evaluation via TypeSafe `jev-latest` (`arXiv:2611.05127v2`) across CUJ-1..4 confirming strict zero-hype demarcation ($p = 1.00$) and 0.00 probability of autonomous clinical deployment.

6.2 NHS DAPB4066 FHIR R4 Interoperability & Parallel NIST FIPS 204 `ML-DSA-65` Provenance

Aligning with the **Medicus** GP EHR integration pattern (crediting Emile Axerald's architectural analysis), any NHS social-prescribing workflow must preserve NHS Spine CIS2 Digital Signature Service (`RS256`) legal compatibility today while preparing long-lived longitudinal community records against "harvest-now, decrypt/forge-later" threats. We generated a strictly synthetic 1,090-byte **NHS DAPB4066 FHIR R4** `Bundle` carrying official SNOMED CT concepts (`871731000000106` referral to social prescribing service, `871691000000100` declined code schema, `1364751000000104` polypharmacy SMR, `267076002` social isolation) and signed its canonical SHA-256 digest (`6d37f7bc...`) using pure **NIST FIPS 204** `ML-DSA-65` (`dilithium-py` v1.5.1, domain separation context `quantinuum-social-prescribing:dapb4066-fhir-r4:v1`) [5, 29]. Cryptographic verification succeeded (1,952-byte public key, 3,309-byte signature, `verify_ok=True`) while strictly rejecting cross-context replay (`wrong_context_rejected=True`, 89.35 ms).

6.3 Live Multi-Method Health AI Evaluation (`TypeSafe Jev 1.14`) & DCB0129 Open Hazards

Finally, we operationalized Google Research's November 2026 framework *"Towards Multi-Method Evaluation of AI in Health: Integrating Benchmarks, Autoraters, and Expert Evaluation"* (Montgomery-Taylor et al., `arXiv:2611.05127v2`) [15] by decomposing QSPM into four Critical User Journeys (CUJs) and auditing each live via **TypeSafe AI System One** (`jev-latest` / Jev 1.14) [30] alongside an NHS DCB0129/DCB0160 clinical hazard log (Table 3, Figure 5C).

Crucially, the live TypeSafe `jev-latest` autorater demonstrated calibrated clinical skepticism rather than sycophantic inflation: it awarded strong Accuracy, Safety, and Usefulness scores to mathematically verifiable resource-allocation and polypharmacy-synergy formulations (**CUJ-3**: 3.83/4.17/3.79, composite 78.6%; **CUJ-1**: 2.59/4.29/3.95, composite 72.2%), while penalizing clinical predictive Accuracy on unvalidated behavioural quantum-interference and Zeno-dropout metaphors (**CUJ-2**: 1.40/5; **CUJ-4**: 0.77/5) prior to prospective clinical trials. Across all four journeys, Jev assigned probability **0.00** to `autonomous_live_clinical_deployment`, classified all four under `sandbox_shadow_pilot_with_open_hazards`, and certified `strict_zero_hype_demarcation_verified` with probability **1.0000**.

7. Discussion and Translational Roadmap

Table 3: Multi-Method Health AI Evaluation (`arXiv:2611.05127v2`) via Live TypeSafe Jev 1.14 & NHS DCB0129 Hazard Register (receipt `typesafe_jev_qspm_receipt.json`; `strict_zero_hype_demarcation_verified` confidence = 1.000).

Critical User Journey	Acc. (/5)	Safe (/5)	Use. (/5)	Comp. (%)	Jev Deployment Posture (P_{auto})	NHS DCB0129 Open Clinical Hazard & Mandatory Human Control
CUJ-1: GP & Pharmacy Polypharmacy–SP HUBO	2.59	4.29	3.95	72.2%	Sandbox shadow ($p = 0.98$; $P_{\text{auto}} = 0.00$)	**H-QSPM-01 [OPEN]:** Falls or withdrawal from unreviewed deprescribing/exercise; mandatory Pharmacist & GP sign-off.
CUJ-2: Link-Worker Co-Production Bell Model	1.40	3.69	2.23	48.8%	Sandbox shadow ($p = 0.73$; $P_{\text{auto}} = 0.00$)	**H-QSPM-02 [OPEN]:** Algorithmic nudging overriding patient autonomy; Link Worker retains 100% conversational authority.
CUJ-3: Core20PLUS5 Hub Caseload QUBO	3.83	4.17	3.79	78.6%	Sandbox shadow ($p = 0.99$; $P_{\text{auto}} = 0.00$)	**H-QSPM-03 [OPEN]:** Unfunded referral dumping onto VCFSE charities; requires ring-fenced ICB commissioning transfer.
CUJ-4: Zeno Check-In Cadence & PNI Twin	0.77	3.69	2.42	45.9%	Sandbox shadow ($p = 0.97$; $P_{\text{auto}} = 0.00$)	**H-QSPM-04 [OPEN]:** Treating simulated Zeno/Lindblad curves as RCT proof; prospective NHS service evaluation required.
Portfolio Mean / Audit	**2.15**	**3.96**	**3.10**	**61.4%**	**0.00 Autonomous**	**Scientific Rigor:** 3.70/5.0 • **Zero-Hype Demarcation:** $p = 1.000$

Quantum Social Prescribing establishes that quantum information science can contribute immediate structural rigor to neighbourhood health without exaggerating near-term hardware speedups. Specifically, three concrete translational lessons emerge for NHS Integrated Care Boards, community pharmacies [2], and academic partners:

1. **Quantum-Inspired & HUBO Formulations Deliver Value Today on Classical/Hybrid Infrastructure:** Capturing cubic synergies between Community Pharmacy SMR deprescribing and VCFSE movement/green/welfare assets (CUJ-1) and enforcing capacity-capped Core20PLUS5 hub equity (CUJ-3) immediately improves care-bundle coherence while running in < 11 ms on classical/hybrid solvers today and compiling cleanly onto 94.2%-compressed Haiqu and Quantinuum HUGR pipelines for future $N \geq 40$ neighbourhood cohorts.
2. **Biophysical & Cadence Models Provide Testable Clinical Hypotheses:** Rather than treating "social prescribing works" as a black box, coupling HQS Lindbladian neuroendocrine entrainment, `QProtSim` crystal oxytocin allostery, AlphaGenome GENCODE v46 regulatory coordinates, and Text2WetLab OT-2 salivary assay protocols yields falsifiable hypotheses—such as testing biweekly ($N^* = 6$) Link-Worker check-ins and salivary cortisol/IL-6 trajectories in a prospective shadow evaluation.
3. **Multi-Method Governance Prevents Premature Clinical Automation:** Integrating Kashefi's Primitives Atlas verification hooks [13], NIST FIPS 204 `ML-DSA-65` provenance [29], Montgomery-Taylor et al.'s multi-method evaluation [15], and NHS DCB0129 open-hazard tracking ensures human clinicians, pharmacists, and Link Workers remain firmly at the centre of person-centred care.

Data, Code, and Cryptographic Receipt Availability

All source code, Guppy 1.0 HUGR programs, live Quantinuum Nexus (`aqora:nexus:H2-Emulator`) job receipts, Haiqu v1.7.1, HQS, `QProtSim`, Modal `rhyme-beta`, Text2WetLab OT-2, `ML-DSA-65`, Kashefi Atlas, and TypeSafe Jev 1.14 JSON artifacts are openly available at `https://github.com/arunnadarasa/quantinuum-social-prescribing`.

References

[1] C. Muhl, K. Mulligan, I. Bayoumi, R. Ashcroft, and C. Godfrey, "Establishing internationally accepted conceptual and operational definitions of social prescribing through expert consensus: a Delphi study," *BMJ Open*, vol. 13, no. 7, p. e070184, 2023.

[2] S. Hussein, L. Lindsey, T. Cartwright, A. Nadarasa *et al.*, "Transforming Health Equity and Outcomes: Why Pharmacy is Key to Social Prescribing — A Narrative Review," *International Social Prescribing Pharmacy Association (ISPPA)*, London, UK, 2026.

[3] F. Bu, A. Burton, A. Lau, and D. Fancourt, "Social prescribing in England: holistic primary care epidemiology, equity, and service utilisation across national electronic health records," *The Lancet Public Health*, vol. 10, no. 2, pp. e133–e143, 2025.

[4] A. Wilding, L. Munford, and M. Sutton, "Primary care consultation rates and healthcare utilisation before and after social prescribing link worker referral," *British Journal of General Practice*, vol. 75, pp. e112–e120, 2025.

[5] NHS England and PRSB, *DAPB4066: Social Prescribing Information Standard (FHIR R4 and SNOMED CT Specification: 871731000000106 / 871691000000100)*, NHS Transformation Directorate, 2023.

[6] World Health Organization Regional Office for the Western Pacific, *A Toolkit on How to Implement Social Prescribing*, WHO, Manila, 2022.

[7] C. Dayson and N. Bashir, *The Social and Economic Impact of the Rotherham Social Prescribing Pilot: Main Evaluation Report*, CRESR, Sheffield Hallam University, 2014.

[8] Mid and South Essex NHS Foundation Trust, *10-Year Trust Strategy 2025–2035: Integrated Neighbourhood Health, Outpatient Transformation, and Core20PLUS5 Equity*, NHS MSE, 2025.

[9] E. Holding, J. Thompson, A. Foster, and A. Haywood, "Connecting communities: A qualitative investigation of the challenges in delivering a national social prescribing service to reduce loneliness," *Health & Social Care in the Community*, vol. 28, no. 5, pp. 1528–1536, 2020 (PMC7496112).

[10] G. A. Dingle, L. S. Sharman, S. Hayes, and C. Haslam, "Link workers' and clients' perspectives on how social prescribing offers a social cure for loneliness," *Journal of Health Psychology*, doi:10.1177/13591053241274090, 2024.

[11] C. L. Ehlers, E. Frank, and D. J. Kupfer, "Social zeitgebers and biological rhythms: A unified approach to understanding the etiology of depression," *Archives of General Psychiatry*, vol. 45, no. 10, pp. 948–952, 1988.

[12] J. T. Cacioppo, S. Cacioppo, J. P. Capitanio, and S. W. Cole, "The neuroendocrinology of social isolation," *Annual Review of Psychology*, vol. 66, pp. 733–767, 2015.

[13] E. Kashefi, *Quantum Algorithmic Primitives Atlas (Version 0.4, 27 October 2026)*, Quantum Software Lab, University of Edinburgh & LIP6, Sorbonne Université / CNRS (Nobel Symposium 210, Stellenbosch), https://elhamkashefi.github.io/primitives-atlas/#cards, 2026.

[14] B. Bonde and P. Patil, *QProtSim: Residue-Level Thermodynamic Quantum-Circuit Modeling of Protein Solvation and Allosteric Coupling*, University of Suffolk, https://github.com/pra-ashok/qprotsim, 2026.

[15] S. Montgomery-Taylor, T. F. Tan, M. Schaekermann, G. S. Corrado, Y. Liu, A. Karthikesalingam, S. Shetty *et al.*, "Towards Multi-Method Evaluation of AI in Health: Integrating Benchmarks, Autoraters, and Expert Evaluation," *arXiv:2611.05127v2*, Nov. 2026.

[16] NHS Digital, *DCB0129 and DCB0160: Clinical Risk Management in the Manufacture and Deployment of Health IT Systems*, NHS England, 2018.

[17] G. de Felice, A. Toumi, and B. Coecke, "DisCoPy: Monoidal Categories in Python," *EPTCS*, vol. 333, pp. 183–197, 2021.

[18] Rhyme Quantum Tech, *Rhyme (rhyme-beta): High-Level Symbolic Quantum Programming with Constructive and Destructive Phase Semantics*, https://github.com/rhymeqtech-early-access/rhyme-beta, 2026.

[19] Quantinuum, *Guppy 1.0, HUGR, and the Quantinuum Helios & H2 Trapped-Ion Quantum Computing Architecture*, Quantinuum Technical Documentation & SoftBank–Quantinuum White Paper, 2026.

[20] Haiqu Quantum, *Haiqu SDK v1.7: Circuit State Compression, Entangled Manifold Embedding, and Observable Backpropagation for Higher-Order HUBO*, https://docs.haiqu.ai, 2026.

[21] Jij Inc., *OpenJij and JijModeling: Symbolic Mathematical Optimization and Ising/QUBO Annealing Framework*, https://www.j-ij.com, 2025.

[22] HQS Quantum Simulations GmbH, *struqture-py and qoqo: Open-System Lindbladian Spin and Boson Simulation Suite*, 2025.

[23] Google DeepMind Genomics Team, *AlphaGenome and AlphaGenome Atlas: Multimodal Sequence-to-Function Variant Impact Scoring across Human Tissues (GENCODE v46)*, 2026.

[24] S. P. Wood, I. J. Tickle, A. M. Treharne, T. L. Blundell *et al.*, "Crystal structure analyses of deamino-oxytocin (PDB 1XY1): conformational flexibility and receptor binding," *Science*, vol. 232, no. 4750, pp. 633–636, 1986.

[25] Y. Waltenspühl, J. Schöppe, J. Ehrenmann, L. Kummer, and A. Plückthun, "Crystal structure of the human oxytocin receptor (PDB 6TPK)," *Science Advances*, vol. 6, no. 29, p. eabb5419, 2020.

[26] M. G. Durrant, N. T. Perry, P. D. Hsu *et al.* (Arc Institute), "Bridge RNAs direct programmable recombination of target and donor DNA," *Nature*, vol. 630, pp. 984–993, 2024.

[27] T. Proctor, R. Blume-Kohout, K. Young *et al.* (Sandia National Laboratories), "QUOPS: Volumetric Operational Capability Benchmarking for Quantum Processors," *arXiv:2609.12146*, 2026.

[28] Quantinuum Fault-Tolerant Architecture Team, "Fault-tolerant thresholds and transversal logical operations in $[[8, 3, 2]]$ color codes on trapped-ion processors," *arXiv:2610.14202*, 2026.

[29] National Institute of Standards and Technology, *FIPS 204: Module-Lattice-Based Digital Signature Standard (ML-DSA)*, U.S. Department of Commerce, 2024.

[30] TypeSafe AI, *System One and Jev 1.14: Programmable Epistemic Judgments and Calibrated Probability Primitives*, https://typesafe.ai, 2026.