THAPAR INSTITUTE OF ENGINEERING & TECHNOLOGY, PATIALA

UCS321 – EST Project

Forecasting Air Quality Index around IndianOil Refinery Zones using ARIMA and LSTM Models

Technical Project Report — Chennai Corridor (CPCL Manali Refinery Zone)

Submitted by: Group 2W1

Name	Roll No.
Arush Saini	1025230031
Jiya Wadhwa	1025230094
Darsh Marwaha	1025230032
Tahoor	1025230126
Arshita Doomra	1025230123

Course: UCS321 – EST Project

Institute: Thapar Institute of Engineering & Technology, Patiala

Date: November 2026

Abstract

This project evaluates classical state-space and deep learning sequence models for forecasting daily Air Quality Index (AQI) around the Chennai Petroleum Corporation Limited (CPCL) Manali refinery corridor, a subsidiary of Indian Oil Corporation Limited (IOCL). Using daily Central Pollution Control Board (CPCB) ambient monitoring data for Chennai from 2016-01-01 to 2020-06-30 (1,643 calendar days, excluding a 2015 monitoring outage), the dataset is partitioned chronologically into Train (1,096 days), Validation (243 days), Test (182 days), and an untouched COVID-19 lockdown Stress window (122 days). Strict leakage controls govern 15 anomaly-corrected days, gap-adaptive imputation, upper-bound winsorization, and train-only scaling. Across 1-day-ahead forecasting on the Test split, ARIMAX(2,1,1) with three annual Fourier harmonics and lagged PM2.5 achieved the lowest error (RMSE 25.97, MAE 18.45, R^2 0.6657, CPCB category accuracy 76.92%), outperforming a hybrid ARIMAX-LSTM residual model (RMSE 26.81, MAE 19.91), a tuned CNN-LSTM (RMSE 27.63, MAE 21.21), Pure ARIMA (RMSE 30.58, MAE 19.23), Univariate LSTM (RMSE 31.73, MAE 22.87), and naive persistence (RMSE 32.10, MAE 19.56). In 7-day-ahead evaluation across 175 origins, a validation-selected Epoch-2 Direct Multi-Step LSTM beats persistence only at horizons 5–7, while recursive ARIMA-Fourier leads horizons 2–6. During the lockdown stress window (mean AQI 78.81), Pure ARIMA achieved the lowest RMSE (13.51) followed by persistence (14.45), whereas exogenous and deep models degraded. Split-conformal intervals attained 96.70% empirical test coverage.

Table of Contents

Abstract 2

Table of Contents 3

List of Figures 4

List of Tables 4

1. Introduction 5

2. Background 6

3. Dataset 8

4. Methodology 9

5. Data Preprocessing 12

6. Exploratory Data Analysis & Visualization 14

7. ARIMA / ARIMAX Modelling 17

8. LSTM / Deep Learning Modelling 20

9. Results & Comparison 22

10. Discussion 31

11. Limitations & Future Scope 32

12. Conclusion 33

References 33

Appendix A: Environment and Reproducibility 34

Appendix B: Top-10 Classical Grid Search & Optuna Best Hyperparameters 35

List of Figures

Figure 4.1: End-to-end AQI forecasting pipeline illustrating sequential preprocessing, exploratory diagnostics, parallel ARIMA/LSTM tracks, and evaluation.
Figure 5.1: Time series overlay comparing raw vs imputed values across short and medium monitoring gaps.
Figure 5.2: Pollutant feature distributions showing upper-bound winsorization at the Training 99.5th percentile.
Figure 5.3: Final cleaned daily Chennai AQI series (2016–2020) partitioned into Train, Validation, Test, and Stress windows.
Figure 6.1: Spearman correlation matrix across AQI and criteria pollutant concentrations on the Training split.
Figure 6.2: STL decomposition of Chennai daily AQI separating long-term trend, annual seasonal cycle, and residual spikes.
Figure 6.3: ACF and PACF correlograms of raw (d=0) and first-differenced (d=1) Chennai daily AQI.
Figure 7.1: Four-panel residual diagnostic suite for ARIMAX(2,1,1)+Fourier(K=3)+PM2.5 confirming white-noise residuals with heavy positive tails.
Figure 7.2: ARIMAX 1-day walk-forward forecast vs actual AQI on the Test partition with 95% parametric prediction intervals.
Figure 8.1: Stage 2 deep learning architecture tournament comparing CNN-LSTM, GRU, BiLSTM, and Stacked LSTM across 5 seeds.
Figure 9.1: Master Test period benchmark comparing RMSE and MAE across classical, deep learning, hybrid, and baseline models.
Figure 9.2: 2x2 factorial comparison showing Test RMSE improvements from exogenous pollutant inputs across both paradigms.
Figure 9.3: Pairwise Diebold–Mariano HLN-corrected two-sided p-value heatmap across benchmark models.
Figure 9.4: Operational alert Precision, Recall, and F1 scores at AQI > 100 and AQI > 200 thresholds.
Figure 9.5: CPCB qualitative health category confusion matrices for ARIMAX, CNN-LSTM, and Hybrid on the Test split.
Figure 9.6: RMSE and MAE degradation between normal Test days (bottom 90%) and spike days (top 10%).
Figure 9.7: Test period forecast trajectory overlay comparing ARIMAX, CNN-LSTM, and Hybrid with zoomed Diwali 2019 inset.
Figure 9.8: Multi-horizon RMSE growth from h=1 to h=7 days on the 175 aligned Test origins.
Figure 9.9: Comparison of Split-Conformal, Parametric Gaussian, and MC-Dropout prediction intervals over a 30-day Test window.
Figure 9.10: COVID lockdown stress window trajectories comparing Pure ARIMA, ARIMAX, CNN-LSTM, Hybrid, and Persistence against actual AQI.

List of Tables

Table 3.1: Pre-COVID Candidate Series Comparison (2015-01-01 to 2020-02-29)
Table 4.1: Authoritative Dataset Partitions and Summary Statistics (FINAL_NUMBERS.md)
Table 5.1: Upper-Bound-Only Winsorization Thresholds and Clipping Counts (Train 99.5th Percentile)
Table 6.1: Stationarity Test Diagnostics on Training AQI (2016–2018, N=1,096)
Table 7.1: Validation Comparison Across Differencing Regimes (d=0 vs d=1) and Model Families
Table 9.1: Master 1-Day-Ahead Benchmark Results on Test Split (2019-09-01 to 2020-02-29, N=182 Days)
Table 9.2: 2x2 Factorial Grid: Algorithmic Paradigm vs Information Scope (Test RMSE)
Table 9.3: HLN-Corrected Diebold–Mariano Pairwise Hypothesis Tests (Test Period, T=182)
Table 9.4: Operational Alert Evaluation on Test Split (N=182 Days)
Table 9.5: Top 5 Worst-Forecast Test Days Audit (FINAL_NUMBERS.md)
Table 9.6: Multi-Horizon Forecast Benchmark Across 175 Identical Test Origins (h = 1 to 7 Days)
Table 9.7: 95% Prediction Interval Calibration Benchmark (Test N=182, Stress N=122)
Table 9.8: Validation (N=243) and COVID Lockdown Stress Window (N=122) Metrics with Stress Mean Bias
Table B.1: Top 10 ARIMA/ARIMAX Configurations Ranked by Validation RMSE (outputs/results/p5_grid_search.csv)
Table B.2: Optuna Best Hyperparameters (Trial #10, CNN-LSTM, Feature Set CORE, Lookback L=7)

1. Introduction

1.1 Air Quality Monitoring in Industrial Refinery Zones

Urban and industrial air quality management relies on timely quantification of criteria pollutants regulated under national ambient standards. Petroleum refining complexes handle high-temperature catalytic cracking, sulfur recovery, combustion in process heaters and boilers, and hydrocarbon storage, which collectively emit particulate matter (PM2.5, PM10), nitrogen oxides (NOx), sulfur dioxide (SO2), carbon monoxide (CO), and volatile organic compounds (VOCs). In coastal industrial corridors such as north Chennai, emissions from refining, thermal power generation, port freight, and urban traffic interact with sea-breeze circulations and seasonal monsoons to shape daily ambient pollutant concentrations.

Short- to medium-term forecasting of the composite Air Quality Index (AQI) provides environmental engineering teams with advance visibility into deteriorating dispersion conditions. Rather than reacting only after regulatory limits are exceeded, accurate 1-day and 7-day ambient forecasts support proactive operational planning, ambient risk communication, and schedule coordination around meteorological stagnation episodes.

1.2 Problem Statement and Objectives

This engineering design project addresses the following problem specification:

> *"Indian Oil Corporation seeks to forecast the Air Quality Index (AQI) around its refinery zones. Using historical pollutant levels, an ARIMA and LSTM-based forecasting model is needed to predict AQI values and support proactive emission control."*

To fulfill this specification rigorously, five engineering objectives were established:

- **Objective 1 (Data Integrity & Site Selection):** Audit multi-station and city-level CPCB monitoring archives around the Chennai Petroleum Corporation Limited (CPCL) Manali refinery zone and construct a continuous, leakage-free daily modeling series.
- **Objective 2 (Domain-Faithful Preprocessing):** Preserve authentic high-pollution physical events (such as festival firecrackers and winter inversions) while isolating verified telemetry sensor malfunctions, applying gap-adaptive imputation and train-only scaling.
- **Objective 3 (Classical & Deep Learning Modeling):** Develop and tune both classical state-space models (ARIMA, ARIMA with annual Fourier harmonics, and exogenous

ARIMAX) and recurrent neural networks (Stacked LSTM, GRU, BiLSTM, CNN-LSTM, direct 7-day LSTM, and an ARIMAX+LSTM hybrid).

- **Objective 4 (Rigorous Benchmarking & Uncertainty Quantification):** Compare all models across identical chronological partitions using regression accuracy, CPCB category classification, Harvey–Leybourne–Newbold corrected Diebold–Mariano tests, regulatory alert metrics, and calibrated prediction intervals.
- **Objective 5 (Structural-Break Stress Testing):** Evaluate model robustness without retraining during the 2020 COVID-19 national lockdown structural break (1 March to 30 June 2020).

1.3 Scope of the Report

This report documents the complete seven-phase engineering pipeline. Section 2 presents conceptual background on CPCB AQI, time-series modeling, and statistical evaluation. Section 3 details dataset provenance and the formal selection of Chennai city-level daily data. Section 4 outlines the end-to-end architecture and chronological leakage controls. Sections 5 and 6 report preprocessing and exploratory diagnostics. Sections 7 and 8 detail classical ARIMA/ARIMAX and deep learning model development. Section 9 synthesizes unified benchmark results, statistical tests, multi-step errors, intervals, and the lockdown stress test. Sections 10 through 12 examine modeling considerations, limitations, future scope, and final conclusions.

2. Background

2.1 CPCB National Air Quality Index and Categories

The Central Pollution Control Board (CPCB) introduced India's National Air Quality Index in 2014 to translate complex multivariate pollutant concentrations into a single public-health communication scale ranging from 0 to 500. Under CPCB methodology, sub-indices are computed for up to eight pollutants (PM10, PM2.5, NO2, SO2, CO, O3, NH3, and Pb) using piecewise-linear breakpoint functions based on 24-hour averages (8-hour averages for CO and O3). The daily composite AQI is assigned as the maximum sub-index across monitored species, provided at least three pollutants including PM2.5 or PM10 are valid. The index maps onto six qualitative health categories: Good (0–50), Satisfactory (51–100), Moderate (101–200), Poor (201–300), Very Poor (301–400), and Severe (401–500).

2.2 Classical Time Series: ARIMA, ARIMAX, and Fourier Seasonality

Autoregressive Integrated Moving Average (ARIMA(p, d, q)) models represent a univariate series after d orders of non-seasonal differencing as a linear combination of p lagged observations and q lagged forecast innovations. When external covariates such as lagged fine particulate matter PM2.5(t−1) are introduced into the observation equation, the formulation extends to ARIMAX. For daily environmental series exhibiting annual seasonality (period m = 365.25 days), classical seasonal ARIMA (SARIMA) becomes computationally intractable because seasonal differencing creates a 365-dimensional state-space transition matrix. An efficient alternative represents annual periodicity through K pairs of deterministic Fourier sine and cosine harmonics entered as exogenous regressors, capturing smooth seasonal cycles with only 2K parameters.

2.3 Recurrent Neural Networks: LSTM, GRU, and CNN-LSTM

Long Short-Term Memory (LSTM) networks address vanishing gradients in sequence modeling by regulating information flow across time steps via forget, input, and output gates paired with an additive cell state. Gated Recurrent Units (GRU) simplify this architecture by merging cell and hidden states through reset and update gates, reducing parameter counts on smaller datasets. A hybrid CNN-LSTM prepends a 1-D temporal convolutional layer before recurrent layers to extract local multi-pollutant motifs (such as multi-day stagnation ramps) prior to sequence aggregation. Training deep networks with Huber loss stabilizes gradients when occasional extreme pollution spikes introduce heavy right-tail residuals.

2.4 Forecasting Evaluation, Statistical Testing, and Conformal Intervals

Point forecast accuracy is evaluated using scale-dependent Root Mean Squared Error (RMSE, penalizing large spike errors) and Mean Absolute Error (MAE), relative Mean Absolute Percentage Error (MAPE), and Coefficient of Determination (R^2). Category classification uses overall CPCB accuracy, Macro-F1 (accounting for rare-class imbalance), and Cohen's Quadratic Weighted Kappa (QWK), which penalizes multi-category misclassifications quadratically. Predictive accuracy differences between two competing models are assessed formally via the Diebold–Mariano (DM) test with the Harvey–Leybourne–Newbold (HLN) small-sample correction, supplemented by Bonferroni adjustment when conducting multiple pairwise tests. Finally, prediction uncertainty is quantified via Monte Carlo (MC) dropout for epistemic parameter spread and split-conformal prediction, which calibrates distribution-free residual quantiles on held-out validation errors.

3. Dataset

3.1 Data Source and Granularity Selection

The empirical dataset originates from the public Kaggle archive 'Air Quality Data in India (2015–2020)' compiled by Vopani from official CPCB Continuous Ambient Air Quality Monitoring Stations (CAAQMS). Although the archive provides both hourly and daily files (city_hour.csv, station_hour.csv, city_day.csv, station_day.csv, and stations.csv), this project strictly utilizes daily aggregated records. Daily granularity aligns directly with 24-hour CPCB regulatory reporting intervals, suppresses sub-daily rush-hour boundary-layer volatility, stabilizes linear state-space estimation, and yields tractable sequence lengths for deep neural training.

3.2 Location Rationale and Formal Series Selection Rule

Chennai was selected as the primary geographic target because Indian Oil Corporation's subsidiary, Chennai Petroleum Corporation Limited (CPCL), operates its major 10.5 MMTPA Manali Refinery complex in north Chennai. Within Chennai, CPCB metadata lists four monitoring stations: TN001 (Alandur Bus Depot), TN002 (Manali Village), TN003 (Manali CPCB), and TN004 (Velachery Residential Area). Two stations (TN002 and TN003) sit inside the Manali industrial corridor.

To prevent subjective selection, an explicit quantitative data-quality rule was enforced prior to modeling over the pre-lockdown audit window (2015-01-01 to 2020-02-29, 1,886 calendar days): a Manali station would be selected if and only if it contained (1) at least 3 years of historical depth (>= 1,095 days), (2) <= 20% missing AQI, and (3) no continuous missing gap exceeding 30 days. Otherwise, the Chennai city-level daily series in city_day.csv would be used. Table 3.1 summarizes the audit outcome.

Table 3.1: Pre-COVID Candidate Series Comparison (2015-01-01 to 2020-02-29)

Candidate Series	Span (Days)	Valid Days	Missing AQI (%)	Max Gap (Days)	Gaps >30d	Usable Pollutants
Chennai City-Level (Selected)	**1,886**	**1,761**	**6.63%**	**82**	**1**	**10**
Manali CPCB (Station TN003)	1,886	1,296	31.28%	411	2	7
Manali Village (Station TN002)	256	218	14.84%	26	0	11
Alandur Bus Depot (Station TN001)	1,886	1,541	18.29%	149	2	9

Station TN002 contained only 256 days (beginning June 2019), failing the 3-year historical depth criterion. Station TN003 suffered 31.28% missingness and two extended outages (411

days in 2016–2017 and 136 days in 2015), failing both the missingness and gap-length rules. Consequently, the Chennai city-level series (city_day.csv) was formally selected. Furthermore, calendar year 2015 was excluded entirely because it contained an 82-day monitoring outage (September–November 2015). Starting the study on 2016-01-01 through 2020-06-30 yields 1,643 continuous calendar days with near-zero post-2015 outages.

3.3 Pollutant Screening and Feature Sets

Twelve candidate pollutants in the pre-lockdown Chennai series were screened against a 30% missingness ceiling. PM10 (87.86% missing) and Xylene (100.00% missing) were excluded. Among the ten remaining species, NH3 exhibited 24.50% missingness (including a 411-day early gap) and near-zero Pearson correlation with AQI ($r = 0.0075$), leaving nine consistently recorded features (ALL9: PM2.5, NO, NO2, NOx, CO, SO2, O3, Benzene, Toluene). Because PM2.5 (6.20% missing, $r = 0.6741$ in full-span screening), CO (1.33% missing, $r = 0.3515$), O3 (2.60% missing, $r = 0.2059$), NO2 (1.91% missing), and SO2 (1.91% missing) directly represent primary combustion and refinery-relevant species, these five formed the primary CORE feature set for multivariate deep learning.

4. Methodology

4.1 Overall System Architecture

The forecasting framework follows a strict chronological pipeline progressing from data ingestion and auditing through domain-aware preprocessing, exploratory diagnostics, parallel classical and deep learning modeling tracks, unified multi-metric benchmarking, and structural-break stress testing. Figure 4.1 illustrates the complete end-to-end workflow.




Figure 4.1: End-to-end AQI forecasting pipeline illustrating sequential preprocessing, exploratory diagnostics, parallel ARIMA/LSTM tracks, and evaluation.

4.2 Chronological Partitioning and Stress-Test Design

To eliminate temporal data leakage and preserve real-world forecasting deployment conditions, random cross-validation shuffling was strictly prohibited. The 1,643-day series

was partitioned into four non-overlapping chronological blocks summarized in Table 4.1. The primary modeling series terminates on 2020-02-29, isolating the nationwide COVID-19 lockdown period (2020-03-01 to 2020-06-30) as an untouched out-of-distribution stress test.

Table 4.1: Authoritative Dataset Partitions and Summary Statistics (FINAL_NUMBERS.md)

Split Name	Start Date	End Date	Days (N)	Mean AQI	Std AQI	Primary Function
Training	2016-01-01	2018-12-31	1,096	112.86	46.91	Model fitting, parameter & scaler calibration
Validation	2019-01-01	2019-08-31	243	103.42	36.72	HPO, early stopping, conformal calibration
Test	**2019-09-01**	**2020-02-29**	**182**	**95.73**	**45.03**	**Held-out benchmarking & DM statistical tests**
Stress (Lockdown)	**2020-03-01**	**2020-06-30**	**122**	**78.81**	**20.38**	**Out-of-distribution structural break testing**
Full Series	2016-01-01	2020-06-30	1,643	107.04	44.92	Complete continuous audited series

4.3 Strict Leakage Controls

Four strict temporal isolation safeguards governed every modeling phase:

- **Train-Only Parameter Fitting:** Seasonal imputation climatologies, 99.5th-percentile upper winsorization thresholds, and MinMaxScaler(0, 1) bounds were fitted exclusively on the 2016–2018 Training partition and applied downstream without refitting on Validation, Test, or Stress data.
- **Target-Timestamp Window Partitioning:** In sequence window generation (src/sequences.py), sliding input windows [t−L+1, ..., t] were partitioned strictly by their target timestamp (t+1 or t+1..t+7), allowing lookback context across split boundaries while preventing any future information leak.
- **Validation-Only Selection:** Grid searches, architecture tournaments, Optuna trials, conformal calibration quantiles, and Direct 7-Day LSTM training epochs were selected strictly on Validation errors before ever evaluating the Test split.
- **Identical Multi-Step Origins:** All multi-horizon models (Direct 7-Day LSTM, Recursive ARIMA-Fourier, Persistence) were evaluated on the identical set of 175 Test origins (2019-09-01 to 2020-02-22) where all seven forward horizons lie inside the Test partition.

5. Data Preprocessing

5.1 Ordered Preprocessing Pipeline

Data preparation was implemented reproducibly in notebooks/03_preprocessing.ipynb across eleven deterministic steps: (1) loading city_day.csv, filtering Chennai, and clipping dates to 2016-01-01 through 2020-06-30 (1,643 days); (2) reindexing to a continuous daily DatetimeIndex; (3) physical bounds validation (confirming 0 negative values or CPCB >500 violations across all columns); (4) telemetry sensor anomaly removal; (5) split-wise missingness audit; (6) gap-adaptive imputation; (7) strict calendar-attributed spike inspection and upper-only winsorization; (8) cyclical calendar feature encoding; (9) chronological split partitioning; (10) train-isolated MinMaxScaler(0, 1) fitting and export; and (11) persistence of clean unscaled (for ARIMA) and scaled (for LSTM) CSV splits.

5.2 Sensor Anomaly Removal: Station TN004 Telemetry Malfunction

Exploratory cross-station verification uncovered an isolated telemetry anomaly in August–September 2016 where Station TN004 (Velachery) logged extreme CO spikes (up to 17–30 mg/m³ at station level, city-aggregated CO 2.08–11.03 mg/m³) driving reported city AQI to 215–449 while contemporaneous fine particulate matter PM2.5 remained in the Good range (12–23 ug/m³) and Station TN001 reported AQI near 64. Cross-checking station_day.csv verified that city AQI on 19–22 September 2016 (AQI 372, 437, 449, 434) and 11 preceding days between 2016-08-15 and 2016-09-18 (AQI 215–266) was driven by the same TN004 CO sensor fault. Consequently, AQI and CO for all 15 confirmed anomaly days were set to NaN, flagged with anomaly_removed=1, and re-imputed. This reduced the maximum target AQI from the artificial 449.00 down to the genuine Diwali 2017 peak of 431.00.

5.3 Missingness Audit and Gap-Adaptive Imputation

After anomaly removal, missingness was confined strictly to the Training split (2016–2018), where AQI was missing on 4.65% of days (51 days across 14 gaps, max gap length 17 days), PM2.5 on 3.01% (33 days, max gap 17 days), CO on 3.38% (37 days, max gap 11 days), O3 on 3.47%, NO2 and SO2 on 3.01% each, Benzene on 4.56%, and Toluene on 4.38%. Validation, Test, and Stress splits all had 0.00% missingness across AQI and all CORE/ALL9 pollutants. Short gaps (<= 3 consecutive days) were filled by linear interpolation; medium gaps (4–30 days) were imputed using level-adjusted Training day-of-year (DOY) median climatologies. As shown in Figure 5.1, zero missing values remained post-imputation while preserving natural seasonal trajectories.




Figure 5.1: Time series overlay comparing raw vs imputed values across short and medium monitoring gaps.

5.4 Authentic Spike Preservation and Upper-Only Winsorization

Target AQI spikes were explicitly NOT clipped or removed as outliers because extreme values represent genuine public-health hazards (including Diwali 2017 at AQI 431.0 and Bhogi 2017 at AQI 360.0). For pollutant inputs, upper-bound-only winsorization was applied using the Training 99.5th percentile threshold (Table 5.1). Lower clipping was intentionally omitted so clean monsoon scavenging readings and the 2020 lockdown drop remained unattenuated. Figure 5.2 illustrates the resulting upper-bound feature distributions, and Figure 5.3 displays the final cleaned AQI series across all four chronological partitions.

Table 5.1: Upper-Bound-Only Winsorization Thresholds and Clipping Counts (Train 99.5th Percentile)

Pollutant	Train 99.5th Percentile	Total Clipped Days (of 1,643)	Clipped Share (%)
PM2.5	182.114 ug/m³	7	0.43%
NO	26.575 ug/m³	13	0.79%
NO2	40.212 ug/m³	8	0.49%
NOx	40.788 ppb	18	1.10%
CO	3.033 mg/m³	6	0.37%
SO2	28.769 ug/m³	7	0.43%
O3	72.189 ug/m³	11	0.67%
Benzene	12.315 ug/m³	8	0.49%
Toluene	18.258 ug/m³	7	0.43%




Figure 5.2: Pollutant feature distributions showing upper-bound winsorization at the Training 99.5th percentile.




Figure 5.3: Final cleaned daily Chennai AQI series (2016–2020) partitioned into Train, Validation, Test, and Stress windows.

6. Exploratory Data Analysis & Visualization

6.1 Long-Term Trend, Seasonality, and Category Distribution

Exploratory analysis on the cleaned series reveals a multi-year downward drift in ambient AQI prior to the 2020 lockdown: Training (2016–2018) averaged 112.86 ± 46.91 (median 99.0, skewness 1.64), Validation (Jan–Aug 2019) averaged 103.42 ± 36.72, and Test (Sep 2019–Feb 2020) averaged 95.73 ± 45.03 (Table 4.1). Calendar year 2016 exhibited substantially higher mean (128.71) and volatility (std 45.16) than 2017–2019 (annual means 102.94–105.49), reflecting an ordinary least-squares trend slope of -7.616 AQI/year across annual means.

Seasonally, northeast winter (Nov–Jan) records the highest ambient levels (mean 126.67 ± 49.83) due to lower mixing heights and festival emissions, whereas the southwest monsoon (Jun–Aug) exhibits wet scavenging lows (96.30 ± 32.47). Conversely, day-of-week averages remain relatively flat (Mon 106.49 to Sat 116.13, Sun 112.77; cross-day std ~3.34 AQI),

indicating the absence of a strong weekend effect. Regarding CPCB health bands, over 93% of days fall within Satisfactory (51–100) or Moderate (101–200), while Good (<=50), Very Poor (301–400), and Severe (>400) together account for under 1.5% of observations, creating strong class imbalance for multi-class evaluation.

6.2 Pollutant Correlations and STL Decomposition

Within the Training partition, PM2.5 exhibits the strongest contemporaneous correlation with AQI (Pearson r = 0.766, Spearman rho = 0.731; Figure 6.1) and the strongest next-day predictive cross-correlation (r = 0.729 against AQI(t+1)), followed by O3 (0.286) and CO (0.103). By contrast, NO2 (0.016) and SO2 (0.046) exhibit weak direct linear cross-correlation with next-day AQI. Lagged cross-correlations decay steadily beyond lags 3 to 7, directly motivating a compact lookback window for sequence models. Additive STL decomposition (Figure 6.2) quantifies annual seasonal strength at Fs = 0.5672 (m = 365), whereas weekly seasonal strength is Fs = 0.0079 (m = 7), justifying the omission of weekly seasonal differencing.




Figure 6.1: Spearman correlation matrix across AQI and criteria pollutant concentrations on the Training split.




Figure 6.2: STL decomposition of Chennai daily AQI separating long-term trend, annual seasonal cycle, and residual spikes.

6.3 Stationarity Diagnostics and ACF / PACF Profile

Augmented Dickey–Fuller (ADF) and Kwiatkowski–Phillips–Schmidt–Shin (KPSS) tests were applied to Training AQI (Table 6.1). Raw AQI yields conflicting evidence: ADF rejects a unit root (stat = -5.6682, p = 9.05e-7), whereas KPSS rejects level stationarity (stat = 1.1417, p = 0.01) due to the multi-year trend shift. After first differencing (d=1), both tests agree on stationarity (ADF p = 1.99e-27; KPSS stat = 0.0856, p = 0.10).

Table 6.1: Stationarity Test Diagnostics on Training AQI (2016–2018, N=1,096)

Series	ADF Statistic	ADF p-Value	KPSS Statistic	KPSS p-Value	Combined Verdict
Raw AQI (d = 0)	-5.6682	9.05e-07 (Reject H0)	1.1417	0.010 (Reject H0)	Trend-nonstationary (conflicting)
First-Differenced AQI (d = 1)	**-14.8172**	**1.99e-27 (Reject H0)**	**0.0856**	**0.100 (Fail to reject)**	**Strictly Stationary**

The ACF and PACF profiles (Figure 6.3) show a sharp PACF cutoff after lag 1 (0.678 on raw AQI), suggesting low-order AR dynamics. Upon differencing (d=1), negative ACF spikes appear at lags 1 and 2 (-0.152, -0.170) alongside negative PACF across lags 1–9—an over-differencing signature—coupled with weak daily lag-365 autocorrelation (r = 0.0816 just

above the +/-0.0592 Bartlett band). Accordingly, both d=0 (stationary with deterministic Fourier harmonics) and d=1 were systematically tested during Phase 5 grid search.



Figure 6.3: ACF and PACF correlograms of raw (d=0) and first-differenced (d=1) Chennai daily AQI.

7. ARIMA / ARIMAX Modelling

7.1 Baselines and Systematic Grid Search

Three non-parametric baselines anchored model evaluation: Naive Daily Persistence (AQI_hat(t+1) = AQI(t)), 7-Day Moving Average, and 365-Day Seasonal Naive. Across 80+ classical configurations in outputs/results/p5_grid_search.csv, three incremental families were tested using orders p, q in {0..3} and d in {0, 1}: M1 (Pure ARIMA), M2 (ARIMA + annual Fourier pairs K in {1, 2, 3}), and M3 (ARIMAX + Fourier + lagged exogenous PM2.5(t−1)). All models were ranked strictly by Validation RMSE (Table 7.1).

Table 7.1: Validation Comparison Across Differencing Regimes (d=0 vs d=1) and Model Families

Model Family & Specification	Differencing	Fourier K	Exog PM2.5	AIC	Val RMSE	Val MAE

Model Family & Specification	Differencing	Fourier K	Exog PM2.5	AIC	Val RMSE	Val MAE
M1 Best d=0: ARIMA(2,0,2)	d = 0	0	No	10,865.37	30.441	20.524
M1 Best d=1: ARIMA(1,1,1)	d = 1	0	No	10,858.23	30.381	20.454
M2 Best d=1: ARIMA(1,1,1)+F3	d = 1	3	No	10,853.91	30.124	19.966
M3 Best d=0: ARIMAX(1,0,0)+F3+PM2.5	d = 0	3	Yes	10,601.14	30.128	18.164
M3 Best d=1: ARIMAX(2,1,1)+F3+PM2.5	**d = 1**	**3**	**Yes**	**10,533.66**	**28.907**	**17.333**

Critical AIC Caveat across Differencing Orders: Raw Akaike Information Criterion (AIC) values are mathematically non-comparable between d=0 and d=1 models because differencing alters both the effective sample size (N−d) and the probability Jacobian of the target variable. Therefore, model selection between d=0 and d=1 relied strictly on out-of-sample Validation RMSE and MAE rather than AIC. Comparing best candidates shows that d=1 outperforms d=0 (Val RMSE 28.907 vs 30.128, Val MAE 17.333 vs 18.164), consistent with first-differencing tracking multi-day local level drift more responsively than a global constant mean.

7.2 Exogenous Fourier Harmonics, PM2.5, and Training-Window Sensitivity

Incorporating K=3 annual Fourier harmonic pairs (6 sinusoidal terms at period 365.25 days) reduced validation error smoothly relative to Pure ARIMA, and adding 1-day-lagged PM2.5(t−1) yielded a further sharp improvement (Val RMSE 30.124 -> 28.907). To verify whether including 2016 (which exhibited higher mean AQI 128.71) helped or hurt 2019 calibration, a training-window sensitivity check compared the full 2016–2018 window (N=1,096 days) against a truncated 2017–2018 window (N=730 days). Retaining 2016 improved Validation RMSE from 29.920 to 28.907 (Val MAE 18.665 -> 17.333), consistent with needing three complete annual cycles to calibrate K=3 seasonal harmonics.

7.3 Residual Diagnostics, Walk-Forward Evaluation, and Multi-Step Results

Residual diagnostics on the fitted ARIMAX(2,1,1)+F3+PM2.5 model (Figure 7.1) yield a near-zero mean (-0.273, std 29.248) with positive skewness (1.384) and high kurtosis (10.868) caused by unmodeled episodic spikes. Ljung–Box independence tests do not reject white noise at lag 10 (Q = 10.61, p = 0.3889) or lag 20 (Q = 17.70, p = 0.6073). Evaluated via fixed-parameter walk-forward forecasting (Kalman filtering without refitting) over the 182-day Test set (Figure 7.2), ARIMAX achieved Test RMSE 25.97, MAE 18.45, R^2 0.6657, and CPCB accuracy 76.92% (versus Pure ARIMA Test RMSE 30.58 and Persistence 32.10). For

7-day recursive forecasting without future measured PM2.5, ARIMA-Fourier was unrolled across the 175 Test origins, achieving RMSE 32.25 (h=1), 42.83 (h=2), 46.05 (h=3), 47.83 (h=4), 49.33 (h=5), 50.06 (h=6), and 50.35 (h=7), outperforming Persistence (32.72 to 61.16) at every horizon.




Figure 7.1: Four-panel residual diagnostic suite for ARIMAX(2,1,1)+Fourier(K=3)+PM2.5 confirming white-noise residuals with heavy positive tails.




Figure 7.2: ARIMAX 1-day walk-forward forecast vs actual AQI on the Test partition with 95% parametric prediction intervals.

8. LSTM / Deep Learning Modelling

8.1 Sequence Construction and Staged Exploration (Stages 1–3)

Deep learning experiments were executed on CPU across 5 deterministic random seeds (42, 123, 456, 789, 1011) using Train-fitted MinMaxScaler(0, 1) inputs (src/train_dl.py). Sliding windows of shape (L, D) were strictly partitioned by target date. A three-stage search avoided unguided over-parameterization:

- **Stage 1 — Feature Set x Lookback Factorial Grid (80 runs):** Using a baseline 2-layer Stacked LSTM (64 units, dropout 0.2, Adam lr=1e-3, MSE), four feature sets (Univariate U, Parsimonious P, CORE C, ALL9 A) were crossed with four lookbacks L in {7, 14, 30, 60}. Compact L=7 with CORE5 (D=8 including AQI and sin/cos DOY) achieved the lowest 5-seed Validation RMSE of 27.375 +/- 0.513 (vs CORE5 L=14 at 27.906, Parsimonious L=7 at 28.177, ALL9 L=7 at 28.195, and Univariate L=7 at 29.765).
- **Stage 2 — Architecture Tournament (40 runs):** On the top two Stage 1 configurations (C, L=7 and C, L=14), four recurrent topologies were compared across 5 seeds (Figure 8.1). CNN-LSTM (Conv1D followed by LSTM layers) won with Validation RMSE 26.241 +/- 1.043 on CORE5 (L=7), outperforming Stacked GRU (27.148 at L=14, 27.207 at L=7), Stacked LSTM (27.375), and BiLSTM (27.427).
- **Stage 3 — Optuna Bayesian HPO (60 trials):** Tree-structured Parzen Estimator (TPE) optimization with Median Pruning searched layer units, dropout, recurrent dropout, learning rate, batch size, L2 regularization, and loss function (MSE vs Huber). Trial #10 achieved 2-seed Val RMSE 24.850 using Huber loss, batch size 16, lr = 0.00438, units (80, 48), dropout 0.15, and recurrent dropout 0.20 (learning rate accounted for 67.9% of HPO importance). Five-seed confirmation yielded Val RMSE 25.496 +/- 0.994 at a median of 18 epochs.




Figure 8.1: Stage 2 deep learning architecture tournament comparing CNN-LSTM, GRU, BiLSTM, and Stacked LSTM across 5 seeds.

8.2 Final Retraining, MC Dropout, 7-Day Direct LSTM, and Hybrid Model

In Stage 4, the winning CNN-LSTM configuration was retrained on Train+Validation (1,339 days) across 5 seeds for 18 epochs, forming a 5-seed ensemble that achieved Test RMSE 27.63, MAE 21.21, R^2 0.6214, and CPCB accuracy 68.68%. A standalone Univariate Stacked LSTM (L=7) benchmarked separately yielded Test RMSE 31.73 and MAE 22.87. Using 100 stochastic forward passes per seed (500 total), MC Dropout produced narrow 95% bands (mean width 31.69 AQI) that undercovered Test observations at 37.91%, reflecting epistemic parameter uncertainty only rather than full observation noise.

In Stage 5, a Direct Multi-Step CNN-LSTM with a Dense(7) output head was trained to emit h=1..7 simultaneously. Crucially, to prevent any test-set leakage, the retraining epoch count was determined strictly by evaluating Validation RMSE across all 7 horizons epoch-by-epoch across 5 seeds. Validation error reached its minimum at Epoch 2 (mean Val RMSE = 33.88) before rising rapidly due to the level shift between 2016–2018 (mean 112.86) and 2019 (mean 103.42). Retrained on Train+Val for 2 epochs across 5 seeds (src/retrain_7day.py), the Direct 7-Day LSTM achieved RMSE 40.39 (h=1), 47.30 (h=2), 51.46 (h=3), 54.42 (h=4), 56.08 (h=5), 54.48 (h=6), and 53.04 (h=7). Finally, in Stage 6, a Residual Hybrid model (trained on L=7 lagged ARIMAX errors plus pollutants) improved standalone DL Test RMSE from 27.63 down to 26.81 (MAE 19.91, R^2 0.6435).

9. Results & Comparison

9.1 Master Test Benchmark and 2x2 Paradigm Grid

Table 9.1 consolidates 1-day-ahead performance across all eight evaluated models on the 182-day Test period (visualized in Figure 9.1). ARIMAX(2,1,1)+F3+PM2.5 achieved the strongest score on every metric except MAPE (where Pure ARIMA's slightly smaller low-end predictions gave 19.33% vs 20.25%), attaining Test RMSE 25.97, MAE 18.45, R^2 0.6657, CPCB accuracy 76.92%, and QWK 0.6471.

Table 9.1: Master 1-Day-Ahead Benchmark Results on Test Split (2019-09-01 to 2020-02-29, N=182 Days)

Model Family	Specific Model	RMSE	MAE	MAPE (%)	R^2	CPCB Acc	Macro F1	QWK
Classical Champion	**ARIMAX(2,1,1)+F3+PM2.5**	**25.97**	**18.45**	**20.25**	**0.6657**	**76.92%**	**0.3664**	**0.6471**
Ensemble Hybrid	Hybrid ARIMAX + Residual LSTM	26.81	19.91	22.58	0.6435	74.73%	0.3644	0.6296
DL Champion	CNN-LSTM (L=7, CORE)	27.63	21.21	23.92	0.6214	68.68%	0.3167	0.5384
Classical	Pure ARIMA(2,1,1)	30.58	19.23	19.33	0.5362	74.18%	0.3473	0.5587
Deep Learning	Univariate Stacked LSTM (L=7)	31.73	22.87	25.43	0.5008	70.88%	0.3205	0.5226
Baseline	Naive Daily Persistence (182d)	32.10	19.56	19.40	0.4889	73.63%	0.3396	0.5951
Baseline	7-Day Moving Average	42.44	26.49	27.19	0.1067	64.84%	0.2465	0.3405
Baseline	Seasonal Naive (365-Day Lag)	71.61	53.86	61.80	-1.5426	36.81%	0.1179	-0.0554




Figure 9.1: Master Test period benchmark comparing RMSE and MAE across classical, deep learning, hybrid, and baseline models.

Disentangling algorithmic paradigm from information scope in a balanced 2x2 factorial grid (Table 9.2, Figure 9.2) reveals consistent benefits from exogenous pollutant data: adding PM2.5 to classical ARIMA reduces Test RMSE by 15.1% (30.58 -> 25.97), while adding CORE pollutants to LSTM reduces Test RMSE by 12.9% (31.73 -> 27.63). Across both univariate and multivariate scopes, classical state-space models achieve lower Test RMSE than their deep learning counterparts (-3.6% univariate; -6.0% multivariate).

Table 9.2: 2x2 Factorial Grid: Algorithmic Paradigm vs Information Scope (Test RMSE)

Algorithmic Paradigm	Univariate (AQI Only)	Multivariate (AQI + Pollutants)	Multivariate Gain
Classical (ARIMA Family)	**30.58 (Pure ARIMA)**	**25.97 (ARIMAX+F3+PM2.5)**	**-4.61 (-15.1%)**
Deep Learning (LSTM Family)	31.73 (Univariate LSTM)	27.63 (CNN-LSTM, CORE)	-4.10 (-12.9%)
Classical Advantage	-1.15 (-3.6%)	-1.66 (-6.0%)	—




Figure 9.2: 2x2 factorial comparison showing Test RMSE improvements from exogenous pollutant inputs across both paradigms.

9.2 Diebold–Mariano Statistical Hypothesis Tests

Pairwise Diebold–Mariano tests with HLN small-sample correction under squared-error loss (h=1, T=182) are summarized in Table 9.3 and Figure 9.3. At an unadjusted alpha = 0.05 significance level, ARIMAX significantly outperforms Persistence (p = 0.0132), Pure ARIMA (p = 0.0297), CNN-LSTM (p = 0.0127), and Hybrid (p = 0.0001), whereas CNN-

LSTM vs Persistence does not reach significance (p = 0.0758). However, when applying a conservative Bonferroni correction across the family of 10 pairwise tests (alpha_adj = 0.05 / 10 = 0.005), only the ARIMAX vs Hybrid comparison remains statistically significant (p = 0.0001).

Table 9.3: HLN-Corrected Diebold–Mariano Pairwise Hypothesis Tests (Test Period, T=182)

Comparison (M1 vs M2)	DM Stat	p-Value	Sig. at Unadjusted alpha=0.05?	Sig. at Bonferroni alpha=0.005?
ARIMAX vs Persistence	-2.504	0.0132	Yes	No
ARIMAX vs Pure ARIMA	-2.192	0.0297	Yes	No
ARIMAX vs CNN-LSTM	-2.519	0.0127	Yes	No
ARIMAX vs Hybrid	**-3.935**	**0.0001**	**Yes**	**Yes (p < 0.005)**
CNN-LSTM vs Persistence	-1.786	0.0758	No	No
Pure ARIMA vs Persistence	-0.861	0.3903	No	No
Hybrid vs Persistence	-2.126	0.0349	Yes	No
Hybrid vs CNN-LSTM	-1.284	0.2010	No	No
Hybrid vs Pure ARIMA	-1.751	0.0818	No	No
CNN-LSTM vs Pure ARIMA	-1.488	0.1385	No	No




Figure 9.3: Pairwise Diebold–Mariano HLN-corrected two-sided p-value heatmap across benchmark models.

9.3 Operational Alerts and Category Classification

Binary alert metrics were evaluated at CPCB action thresholds AQI > 100 (Moderate+, 52 actual Test days) and AQI > 200 (Poor+, 10 actual Test days), as displayed in Table 9.4 and Figure 9.4, alongside category confusion matrices in Figure 9.5. At AQI > 100, Hybrid achieved 86.54% recall and 0.7031 F1, while ARIMAX achieved the highest precision (64.52%) with 76.92% recall and 0.7018 F1, clearly surpassing Persistence (F1 0.6476). At AQI > 200, ARIMAX and Hybrid tied at 85.71% precision, 60.00% recall, and 0.7059 F1, outperforming CNN-LSTM (recall 40.00%, F1 0.5333), though the 10-day sample (5.5% frequency) limits rare-event statistical power.

Table 9.4: Operational Alert Evaluation on Test Split (N=182 Days)

Threshold & Metric	ARIMAX	Hybrid	CNN-LSTM	Pure ARIMA	Persistence

Threshold & Metric	ARIMAX	Hybrid	CNN-LSTM	Pure ARIMA	Persistence
AQI > 100 Precision (52 actual days)	64.52%	59.21%	53.01%	61.11%	64.15%
AQI > 100 Recall	76.92%	86.54%	84.62%	63.46%	65.38%
AQI > 100 F1 Score	**0.7018**	**0.7031**	**0.6519**	**0.6226**	**0.6476**
AQI > 200 Precision (10 actual days)	85.71%	85.71%	80.00%	100.00%	70.00%
AQI > 200 Recall	60.00%	60.00%	40.00%	50.00%	70.00%
AQI > 200 F1 Score	**0.7059**	**0.7059**	**0.5333**	**0.6667**	**0.7000**




Figure 9.4: Operational alert Precision, Recall, and F1 scores at AQI > 100 and AQI > 200 thresholds.




Figure 9.5: CPCB qualitative health category confusion matrices for ARIMAX, CNN-LSTM, and Hybrid on the Test split.

9.4 Stratified Spike Error Analysis and Worst-Day Calendar Audit

Decomposing Test errors between normal days (bottom 90%, actual AQI < 134.9, N=163) and spike days (top 10%, AQI >= 134.9, N=19) highlights systematic negative bias on episodic peaks (Figure 9.6). While ARIMAX achieved 18.78 RMSE (+9.56 bias) on normal days, all models degraded sharply on top-10% spikes (mean negative bias ranging from -27.6 to -55.5 AQI). Table 9.5 audits the top 5 worst-forecast days: exactly 2 of the 5 worst days correspond to calendar-verified festival events (Diwali on 2019-10-27 and Bhogi on 2020-01-

14), while the remaining 3 days carry 'No matched event'. Figure 9.7 illustrates the Diwali 2019 under-prediction across all models.

Table 9.5: Top 5 Worst-Forecast Test Days Audit (FINAL_NUMBERS.md)

Date	Actual AQI	ARIMAX Error	CNN-LSTM Error	Calendar Match	Context
2019-10-27	**306.0**	**-140.81**	**-144.74**	**Diwali 2019**	**Verified festival pyrotechnics spike**
2020-01-14	**230.0**	**-92.22**	**-85.63**	**Bhogi 2020**	**Verified harvest bonfire smoke event**
2019-11-04	222.0	-90.43	-83.72	No matched event	Sudden post-monsoon jump from 91 to 222 AQI
2019-11-12	77.0	+72.90	+66.91	No matched event	Over-prediction after rapid ambient drop
2019-11-07	273.0	-67.96	-75.09	No matched event	Unmodeled severe pollution buildup




Figure 9.6: RMSE and MAE degradation between normal Test days (bottom 90%) and spike days (top 10%).




Figure 9.7: Test period forecast trajectory overlay comparing ARIMAX, CNN-LSTM, and Hybrid with zoomed Diwali 2019 inset.

9.5 Multi-Horizon Forecasting (h = 1 to 7 Days)

Evaluated across the 175 aligned Test origins (Table 9.6, Figure 9.8), the validation-selected Epoch-2 Direct Multi-Step LSTM beats Persistence only at horizons 5, 6, and 7 (56.08 vs 58.42; 54.48 vs 61.38; 53.04 vs 61.16), whereas both Persistence and recursive ARIMA-Fourier outperform it at h=1..4. Recursive ARIMA-Fourier achieves the lowest RMSE across every horizon (32.25 to 50.35), consistent with the hypothesis that deterministic annual Fourier harmonics constrain multi-day trajectory dispersion.

Table 9.6: Multi-Horizon Forecast Benchmark Across 175 Identical Test Origins (h = 1 to 7 Days)

Horizon	Persist RMSE	ARIMA-F3 RMSE	LSTM-7d RMSE	Persist MAE	ARIMA-F3 MAE	LSTM-7d MAE	LSTM Beats Persist?
h = 1	32.72	32.25	40.39	20.15	22.37	31.96	No
h = 2	47.14	42.83	47.30	28.99	31.64	37.45	No
h = 3	51.01	46.05	51.46	32.01	35.01	41.85	No
h = 4	53.46	47.83	54.42	33.79	36.72	44.91	No
h = 5	**58.42**	**49.33**	**56.08**	**36.80**	**38.07**	**46.92**	**Yes (56.08 < 58.42)**
h = 6	**61.38**	**50.06**	**54.48**	**38.05**	**38.87**	**44.46**	**Yes (54.48 < 61.38)**
h = 7	**61.16**	**50.35**	**53.04**	**37.53**	**39.34**	**42.76**	**Yes (53.04 < 61.16)**




Figure 9.8: Multi-horizon RMSE growth from h=1 to h=7 days on the 175 aligned Test origins.

9.6 Uncertainty Quantification: Prediction Intervals

Split-conformal calibration using finite-sample Validation residual quantiles (q = 63.87 for ARIMAX, q = 57.71 for LSTM) achieved 96.70% and 96.15% empirical coverage on Test (Table 9.7, Figure 9.9), and 98.36% on Stress, whereas MC Dropout covered only 37.91% on Test. Note that split-conformal theoretical marginal guarantees assume data exchangeability, which time series with level drift do not strictly satisfy; empirical coverage here therefore serves as an observed calibration check rather than a formal guarantee.

Table 9.7: 95% Prediction Interval Calibration Benchmark (Test N=182, Stress N=122)

Method	Interval Type	Val Quantile q	Mean Width (AQI)	Test Coverage	Stress Coverage
ARIMAX Split-Conformal	**Distribution-free**	**63.87**	**127.74**	**96.70%**	**98.36%**
LSTM Split-Conformal	**Distribution-free**	**57.71**	**115.43**	**96.15%**	**98.36%**
ARIMAX Parametric	Gaussian (+/-1.96 sigma)	N/A	114.39	96.15%	97.54%
LSTM MC-Dropout	Epistemic parameter	N/A	31.69	37.91%	N/A




Figure 9.9: Comparison of Split-Conformal, Parametric Gaussian, and MC-Dropout prediction intervals over a 30-day Test window.

9.7 COVID-19 Lockdown Stress Test (2020-03-01 to 2020-06-30)

During the 122-day COVID-19 national lockdown structural break (mean AQI 78.81 vs pre-lockdown full-series mean 109.30 and Test mean 95.73), Pure ARIMA achieved the lowest stress RMSE (13.51), followed by persistence (14.45); models using exogenous or seasonal terms (ARIMAX 17.74, multivariate CNN-LSTM 20.28) performed worst, with the LSTMs over-predicting (mean bias +7.84 / +10.55) and ARIMAX under-predicting (−8.32). Table 9.8 and Figure 9.10 summarize Validation vs Stress performance alongside mean forecast bias.

Table 9.8: Validation (N=243) and COVID Lockdown Stress Window (N=122) Metrics with Stress Mean Bias

Model Name	Val RMSE	Val MAE	Stress RMSE	Stress MAE	Stress R^2	Stress Acc	Stress Bias
Pure ARIMA(2,1,1)	**30.49**	**20.45**	**13.51**	**9.06**	**0.5567**	**86.89%**	**-1.15**
Naive Persistence	**34.27**	**20.85**	**14.45**	**9.41**	**0.4928**	**87.70%**	**-0.31**
7-Day MA	34.50	24.27	14.65	10.22	0.4793	86.89%	-0.82
Hybrid ARIMAX+LSTM	—	—	15.62	9.46	0.4080	86.89%	-3.19
Univariate LSTM	29.61	20.03	15.98	12.77	0.3801	86.07%	+7.84
ARIMAX(2,1,1)+F3+PM2.5	28.91	17.33	17.74	12.37	0.2361	80.33%	-8.32
CNN-LSTM (Multivariate)	25.63	17.03	20.28	15.40	0.0019	81.97%	+10.55
Seasonal Naive (365d)	46.33	34.76	37.93	28.82	-2.4913	62.30%	+22.70




Figure 9.10: COVID lockdown stress window trajectories comparing Pure ARIMA, ARIMAX, CNN-LSTM, Hybrid, and Persistence against actual AQI.

10. Discussion

10.1 Why ARIMAX Outperformed CNN-LSTM on This Dataset

A central empirical outcome of this study is that classical ARIMAX(2,1,1)+F3+PM2.5 outperformed the tuned CNN-LSTM on the held-out Test split (RMSE 25.97 vs 27.63; MAE 18.45 vs 21.21) despite extensive architecture search and Bayesian optimization. Three technical considerations are consistent with this outcome. First, daily series length (~1,096 training samples, 1,339 Train+Val) sits within the small-data regime where parsimonious linear state-space models with fewer than a dozen free parameters are less prone to variance than multi-layer neural networks with thousands of weights. Second, CPCB daily AQI behaves largely as a locally persistent autoregressive process anchored linearly to yesterday's PM2.5; first-differencing (d=1) tracks downward annual drift (112.86 -> 103.42 -> 95.73) naturally without over-anchoring to 2016–2018 mean levels. Third, validation selection across 180+ deep learning configurations introduced optimistic validation selection bias: while CNN-LSTM beat ARIMAX on Validation (25.63 vs 28.91), part of that advantage may reflect overfitting to the 243-day Validation regime rather than transferable generalization to the Test regime.

10.2 Festivals as the Primary Failure Mode and Regime-Shift Sensitivity

Every evaluated model relies solely on lagged ambient sensors and smooth seasonal harmonics. Because Diwali pyrotechnics and Bhogi bonfires inject sharp, single-day emission impulses disconnected from preceding-day PM2.5, all models exhibit large negative errors on the top-5 worst days (e.g., -140.78 on Diwali 2019). Without explicit calendar event flags, historical sensor lags cannot anticipate cultural discontinuities. During the 2020 COVID

lockdown stress test, a possible reason Pure ARIMA (RMSE 13.51, bias -1.15) and Persistence (14.45, bias -0.31) remained accurate is that pure univariate differencing adapts immediately to the preceding day's new low-emission baseline, whereas models relying on pre-lockdown seasonal priors or multivariate weights exhibited larger biases (-8.32 for ARIMAX; +7.84 and +10.55 for LSTMs).

10.3 Potential Operational Use

Day-ahead ARIMAX(2,1,1)+F3+PM2.5 forecasts, coupled with split-conformal prediction intervals, can serve as an informational early-warning input for industrial environmental monitoring. Operational decision thresholds demonstrate measured utility on the test set: at AQI > 100 (Moderate+), the model achieved 64.5% precision and 76.9% recall (F1 = 0.702 across 52 actual alert days); at AQI > 200 (Poor+), it achieved 85.7% precision and 60.0% recall (F1 = 0.706 across 10 actual alert days, though rare-event sample size is limited). Festival days represent a known failure mode where all models exhibit large negative forecast bias; addressing this requires explicit calendar and event indicator features in future work. Finally, because the underlying CPCB monitoring represents city-level ambient air quality rather than fence-line perimeter monitoring, these forecasts reflect regional atmospheric conditions and do not isolate the CPCL Manali refinery's specific operational emissions contribution.

11. Limitations & Future Scope

11.1 Project Limitations

- **City-Level Ambient vs Fence-Line Resolution:** Station-level gaps at Manali (TN002 depth, TN003 31.28% missingness) necessitated city-level Chennai data, which blends refinery emissions with urban vehicular, port, and domestic signals.
- **Absence of Meteorological Predictors:** Planetary boundary layer height, wind speed and direction, relative humidity, and atmospheric stability fields were absent from the CPCB dataset, limiting non-festival stagnation prediction.
- **Severe-Spike Sparsity and Non-Exchangeability:** Only 10 Test days exceeded AQI 200 (5.5%), restricting statistical confidence for extreme alerts, while distribution drift prevents time-series observations from strictly satisfying conformal exchangeability assumptions.

11.2 Future Scope

- **On-Site Refinery CEMS and Fence-Line Integration:** Combine Continuous Emission Monitoring System (CEMS) stack logs from CPCL Manali with dedicated perimeter micro-sensors to isolate refinery-attributable ambient increments.
- **Coupling with Numerical Weather Prediction (NWP):** Incorporate forecasted wind vectors and boundary-layer depth to improve stagnation forecasting on non-festival spike days.
- **Calendar Event Indicators and Adaptive Updating:** Add explicit festival binary/lead-time indicators (Diwali, Bhogi) and online rolling recalibration to mitigate structural regime shifts.

12. Conclusion

This study built a reproducible, leakage-free pipeline for forecasting daily Chennai ambient AQI around Indian Oil Corporation's CPCL Manali refinery zone across classical state-space and deep learning paradigms. Across 182 held-out Test days, ARIMAX(2,1,1) combined with three annual Fourier harmonics and 1-day lagged PM2.5 emerged as the top day-ahead model (RMSE 25.97, MAE 18.45, R^2 0.6657, CPCB accuracy 76.92%), outperforming a Hybrid ARIMAX+LSTM model (26.81), CNN-LSTM (27.63), Pure ARIMA (30.58), Univariate LSTM (31.73), and naive persistence (32.10), with exogenous PM2.5 improving accuracy by 12.9%–15.1% across both paradigms.

Rigorous evaluation also established important empirical boundaries: under Bonferroni correction across 10 Diebold–Mariano tests, only ARIMAX vs Hybrid remained statistically significant ($p = 0.0001$); in 7-day-ahead forecasting, validation-selected Epoch-2 Direct LSTM outperformed persistence only at horizons 5–7 while recursive ARIMA-Fourier led all horizons; and during the 2020 COVID-19 lockdown, univariate Pure ARIMA (RMSE 13.51) and persistence (14.45) proved most resilient to sudden emission drops. Together with split-conformal uncertainty intervals (96.70% Test coverage) and clear city-level scope caveats, the ARIMAX framework offers a transparent baseline for proactive environmental monitoring.

References

[1] Vopani (R. Rao), "Air Quality Data in India (2015–2020)," Kaggle. [Online]. Available: https://www.kaggle.com/datasets/rohanrao/air-quality-data-in-india.

[2] Central Pollution Control Board (CPCB), National Air Quality Index, Ministry of Environment, Forest and Climate Change, Government of India, New Delhi, 2014.

[3] G. E. P. Box, G. M. Jenkins, G. C. Reinsel, and G. M. Ljung, Time Series Analysis: Forecasting and Control, 5th ed. Hoboken, NJ, USA: Wiley, 2015.

[4] R. J. Hyndman and G. Athanasopoulos, Forecasting: Principles and Practice, 3rd ed. Melbourne, Australia: OTexts. [Online]. Available: https://otexts.com/fpp3/.

[5] S. Hochreiter and J. Schmidhuber, "Long short-term memory," Neural Computation, vol. 9, no. 8, 1997.

[6] K. Cho, B. van Merrienboer, C. Gulcehre, D. Bahdanau, F. Bougares, H. Schwenk, and Y. Bengio, "Learning phrase representations using RNN encoder–decoder for statistical machine translation," in Proc. Conf. Empirical Methods in Natural Language Processing (EMNLP), 2014.

[7] R. B. Cleveland, W. S. Cleveland, J. E. McRae, and I. Terpenning, "STL: A seasonal-trend decomposition procedure based on loess," Journal of Official Statistics, vol. 6, no. 1, 1990.

[8] D. A. Dickey and W. A. Fuller, "Distribution of the estimators for autoregressive time series with a unit root," Journal of the American Statistical Association, vol. 74, no. 366a, 1979.

[9] D. Kwiatkowski, P. C. B. Phillips, P. Schmidt, and Y. Shin, "Testing the null hypothesis of stationarity against the alternative of a unit root," Journal of Econometrics, vol. 54, 1992.

[10] G. M. Ljung and G. E. P. Box, "On a measure of lack of fit in time series models," Biometrika, vol. 65, no. 2, 1978.

[11] F. X. Diebold and R. S. Mariano, "Comparing predictive accuracy," Journal of Business & Economic Statistics, vol. 13, no. 3, 1995.

[12] D. Harvey, S. Leybourne, and P. Newbold, "Testing the equality of prediction mean squared errors," International Journal of Forecasting, vol. 13, no. 2, 1997.

[13] Y. Gal and Z. Ghahramani, "Dropout as a Bayesian approximation: Representing model uncertainty in deep learning," in Proc. Int. Conf. Machine Learning (ICML), 2016.

[14] A. N. Angelopoulos and S. Bates, "A gentle introduction to conformal prediction and distribution-free uncertainty quantification," arXiv:2107.07511, 2021.

[15] T. Akiba, S. Sano, T. Yanase, T. Ohta, and M. Koyama, "Optuna: A next-generation hyperparameter optimization framework," in Proc. ACM SIGKDD Int. Conf. Knowledge Discovery & Data Mining (KDD), 2019.

[16] J. Cohen, "Weighted kappa: Nominal scale agreement provision for scaled disagreement or partial credit," Psychological Bulletin, vol. 70, no. 4, 1968.

Appendix A: Environment and Reproducibility

All experiments were executed deterministically on Windows 11 using Python 3.11.0 (64-bit) in a dedicated virtual environment (.venv) with Jupyter kernel ucs321-aqi. Deep learning models trained on native CPU (TensorFlow 2.16.1) with fixed random seeds [42, 123, 456, 789, 1011]. Key pinned packages in requirements.txt include numpy==1.26.4, pandas==2.2.2, scipy==1.13.1, scikit-learn==1.5.0, statsmodels==0.14.2, pmdarima==2.0.4, tensorflow==2.16.1, optuna==3.6.1, matplotlib==3.8.4, and seaborn==0.13.2.

Exact End-to-End Execution Order:

1. notebooks/02_data_audit.ipynb — Missingness audit and series selection.

2. notebooks/03_preprocessing.ipynb — TN004 anomaly removal, imputation, upper winsorization, chronological splits, and train-only scaling.

3. notebooks/04_eda.ipynb — Trend, seasonality, STL, correlation, and ADF/KPSS diagnostics.

4. notebooks/05_arima.ipynb — Baselines, 80+ ARIMA/ARIMAX grid search, walk-forward Test/Stress and 175-origin multi-step evaluation.

5. python src/train_dl.py -> python src/retrain_7day.py -> python src/train_univariate_lstm.py -> notebooks/06_lstm.ipynb — Deep learning pipeline.

6. python src/run_phase7.py -> python src/draw_flow_diagram.py -> notebooks/07_evaluation.ipynb -> python src/check_consistency.py — Final benchmark and verification.

Appendix B: Top-10 Classical Grid Search & Optuna Best Hyperparameters

Table B.1: Top 10 ARIMA/ARIMAX Configurations Ranked by Validation RMSE (outputs/results/p5_grid_search.csv)

Rank	Model Family	Order	K	PM2.5	AIC	BIC	Val RMSE	Val MAE
1	**M3_ARIMAX_F3_PM25**	**(2,1,1)**	**3**	**True**	**10533.66**	**10588.65**	**28.907**	**17.333**
2	M3_ARIMAX_F3_PM25	(1,1,2)	3	True	10527.14	10582.12	29.278	17.690
3	M2_ARIMAX_F3	(1,1,1)	3	False	10853.91	10898.90	30.124	19.966
4	M3_ARIMAX_F3_PM25	(1,0,0)	3	True	10601.14	10651.14	30.128	18.164
5	M3_ARIMAX_F3_PM25	(1,1,1)	3	True	10553.69	10603.68	30.142	18.206
6	M3_ARIMAX_F1_PM25	(1,1,1)	1	True	10550.26	10580.25	30.153	18.038
7	M2_ARIMAX_F2	(1,1,1)	2	False	10850.33	10885.32	30.165	20.014
8	M3_ARIMAX_F2_PM25	(1,1,1)	2	True	10553.31	10593.30	30.215	18.113
9	M2_ARIMAX_F3	(2,1,1)	3	False	10853.57	10903.55	30.244	20.017
10	M2_ARIMAX_F1	(1,1,1)	1	False	10847.23	10872.22	30.254	20.173

Table B.2: Optuna Best Hyperparameters (Trial #10, CNN-LSTM, Feature Set CORE, Lookback L=7)

Hyperparameter	Selected Value	Role in Architecture
units_l1 / units_l2	**80 / 48**	**Conv1D / LSTM hidden filter and state dimensions**
dropout / recurrent_dropout	0.15 / 0.20	Feedforward & recurrent regularization (also used in MC Dropout)
learning_rate	**0.00438**	**Adam step size (67.9% HPO parameter importance)**
batch_size	16	Mini-batch size (15.3% HPO importance)
l2_reg	1.11e-06	Weight decay regularization penalty
loss_fn	huber	Robust loss reducing extreme-spike gradient distortion
median_epochs (1-day / 7-day)	18 / 2	Validation-selected training epochs for Train+Val retraining