

Lab 3

Sophie Lindberg, Arvid Edenheim

5/13/2019

1 - Normal model, mixture of normal model with semi-conjugate prior

1a) - Normal model

The code for the Gibbs implementation can be seen below.

```
# draw mu
drawMu <- function(prevMu, prevSigma) {
  tauSq <- 1/( (n/prevSigma) + (1/tau0^2) )
  w <- (n/prevSigma)/((n/prevSigma) + (1/tau0^2))
  mu <- w*dataMean + (1-w)*mu0
  draw <- rnorm(1, mu, sqrt(tauSq))
  return (draw)
}

#inv chi square
invChiSquare <- function(v, s) {
  return(v*s / rchisq(1,v))
}

# draw sigma
drawSigma <- function(mu) {
  sum <- 0
  for (i in 1:n) {
    sum <- sum + (data[i,1] - mu)^2
  }
  s <- (v0*sigma0 + sum)/(n+v0)
  return(invChiSquare(vn, s))
}

mu <- c()
sigma2 <- c()

currMu <- 32
currSigma <- sigma0
for (i in 1:iter) {
  if(i %% 2 == 0) {
    currMu <- drawMu(currMu, currSigma)
  } else {
    currSigma <- drawSigma(currMu)
  }
  mu <- c(mu, currMu)
  sigma2 <- c(sigma2, currSigma)
}
```

By plotting the trajectories of the sampled Markov chains it can be seen that σ and μ converges around 39 and 32 respectively.

1b) - Mixture normal model

The results of the Gibbs sampling data augmentation algorithm given in NormalMixtureModel.R resulted can be seen below.






Figure 1: Analyzing the convergence



Both μ and σ converged after a few samples.





Figure 2: Graphical comparison

1c) - Graphical comparison

The figure below shows a comparison between the samplers. The blue line represents the normal density from exercise a, and the green line represents the mixture of normals density from exercise b. Both of the models have limitations when it comes to fit the data.

Including Plots

You can also embed plots, for example:



Note that the `echo = FALSE` parameter was added to the code chunk to prevent printing of the R code that generated the plot.