

Spontaneous emission in an exponential model

A. D. Kammogne¹ and L. C. Fai¹

¹*Mesoscopic and Multilayers Structures Laboratory, Faculty of Science,
Department of Physics, University of Dschang, Cameroon*

(Dated: December 11, 2024)

The phenomenon of spontaneous emission can lead to the creation of an imaginary coupling and a shift. To explore this, we utilized the renormalized first Nikitin model, revealing an exponential detuning variation with a phase and an imaginary coupling along with the shift. By employing the time-dependent Schrödinger equation, we investigated the behavior of our system. Our findings indicate that the imaginary coupling provides specific information, while the shift generates allowed and forbidden zones in the energy diagram of the real part of the energy. In the diagram of the imaginary part of the energy, time dictates order or chaos in the system and identifies the information transmission zone. Notably, the first Nikitin model exhibits similarities to the Rabi model in the short-time approximation. Our theoretical conclusions are consistent with numerical solutions.

INTRODUCTION

The concept of spontaneous emission underpins many physical phenomena, such as non-Hermiticity [1–10], dissipation [11–20] between quantum systems, the production of lasers [21–30], and various other natural occurrences. Non-Hermiticity has become a focal topic of scientific discussion worldwide and has garnered significant attention in the field of quantum mechanics. In some cases, spontaneous emission and non-Hermiticity are interrelated. For instance, [31] demonstrated that excessive spontaneous emission in a laser oscillator increases noise in non-Hermitian optical systems. Furthermore, [32] confirmed spontaneous emission at exceptional points through photoluminescence measurements, while [33] developed a general theory of spontaneous emission, highlighting its applicability to dispersive media.

Several authors suggest that the chirality of non-Hermitian models, such as the SSH (Su-Schrieffer-Heeger) model [34], generates a complex Berry phase, explaining the presence of gapless edge modes. Yuto Ashida et al. [35] explored the topology of non-Hermitian systems, introducing the concept of band topology in complex spectra and defining their classifications. Another study [36] showed that spontaneous decay induces quantum phase transitions in non-Hermitian systems, revealing a connection between dissipation and non-Hermitian properties in quantum systems. Non-Hermitian systems are characterized by complex eigenenergies comprising real and imaginary components. This notion can be counterintuitive because quantum mechanics dictates that a system must be Hermitian, with real eigenenergies, to measure observables.

In studying the interaction between a two-level system and its environment, we propose a new scenario where spontaneous emission generates imaginary coupling, as illustrated in Fig. 1 accompanied by a static shift. In this context, we consider two quantum states, denoted as $|1\rangle$ and $|2\rangle$, represented respectively by positive detuning $\Omega(t)$ and negative detuning $-\Omega(t)$. These states are coupled via an imaginary term $i\Delta$ and the static shift.




FIG. 1. Spontaneous emission between two diabatic states during their interaction. In this process, the states $|1\rangle$ and $|2\rangle$ are associated with the detuning $\Omega(t)$ and $-\Omega(t)$. Injection of the magnetic field is at the origin of the creation of spontaneous emission characterized by an imaginary coupling followed by a shift. This reflects the non-hermitian character of our Hamiltonian

To investigate the dynamics of this scenario, we selected the exponential Nikitin model, characterized by exponentially varying detuning, imaginary coupling, and a static shift term. The imaginary term primarily drives the non-Hermitian nature of our Hamiltonian. While quantum mechanics generally permits the measurement of observables only when the spectrum is real, this work examines how complex spectra emerge when coupling becomes complex. The Nikitin model has been extensively explored in previous works, such as those by Nesbitt et al. [37] and Vitanov [38], who highlighted its similarity to the Demkov-Kunike model under strong coupling. The Nikitin model also describes decay processes [39] and has applications in interferometry [40]. Furthermore, collision dynamics between two states have been described using an exponential model by V. Sidis et al. [41].

In this work, we present a novel perspective on sponta-

neous emission, incorporating imaginary coupling with a shift in the off-diagonal Hamiltonian terms. Spontaneous emission plays a crucial role in Bose-Einstein condensates [42–53], radio-active process [54–63] and ultra-cold trapped gas [64–74]. Our aim is to construct a theoretical framework to better understand this model's behavior and elucidate the role of imaginary coupling.

This paper is organized as follows: in the first section, we present the exponential Nikitin model. In the second section, we solve the time-dependent Schrödinger equation to analyze the model's dynamics. In the third section, we discuss the energy spectrum. Finally, we summarize our findings and propose future research directions.

I. EXPONENTIAL MODEL

To describe the interaction between diabatic states $|1\rangle$ and $|2\rangle$ during spontaneous emission, we select the first exponential Nikitin model due to its exponentially varying parameters.

Our model is described by the Hamiltonian:

$$H(t) = \Omega(t)\sigma_z + \delta\sigma_x, \quad (1.1)$$

where $\Omega(t)$ and δ represent, respectively, the detuning and the Rabi frequency. Each parameter plays a crucial role in describing the model. The detuning $\Omega(t)$ determines the direction of the magnetic field via the Pauli matrix σ_z , while the Rabi frequency measures the coupling strength in the transverse direction σ_x . These quantities are defined as follows:

$$\Omega(t) = \frac{1}{2}(A \exp(\alpha t + \beta) + \epsilon), \quad (1.2)$$

$$\delta = \frac{1}{2}(i\Delta + \epsilon). \quad (1.3)$$

As shown, the detuning $\Omega(t)$ varies exponentially, as observed in [39], but with the additional feature of a phase β . Here, A represents the amplitude, responsible for the signal strength; α is the sweep velocity induced by an external field; and β is the phase, which controls oscillations' increase or decrease. The coupling δ includes an imaginary component due to the effects of spontaneous emission, accompanied by a shift ϵ . At this stage, the shift acts as the real coupling responsible for the interaction between the levels. The Pauli matrices σ_ω with $\omega = x, y, z$ are defined as the generators of the $SU(2)$ Lie algebra describing the dynamics of the two-level system.

Compared to the work of [39], which studied the Nikitin model with real detuning and coupling in the context of decay, this work focuses on spontaneous emission with imaginary coupling. The combination of imaginary coupling and the shift introduces nonlinearity to our system.

II. METHODS

Among the various methods available to study the dynamics of this model—such as the Bloch picture [75, 76] with the Lindblad equation [77], the Feynman Path integral [78], the Schrödinger picture with the time-dependent Schrödinger [39], and so on—we chose the Schrödinger picture. This method provides a wealth of information, including probability amplitudes, the propagator, and the transition probabilities.

A. Schrödinger picture

The dynamics of our model are determined using the time-dependent Schrödinger equation, which establishes the relationship between probabilities and their evolution through the matrix elements of the propagator. The non-Hermitian nature of our system arises from the effects of spontaneous emission, introducing an imaginary coupling. The Schrödinger equation is expressed as:

$$i\frac{d}{dt}C(t) = H(t)C(t). \quad (2.1)$$

Using an appropriate gauge transformation $C_1(t) = \psi_1(t) \exp\left(-\frac{i}{2}\int_{t_0}^t \Omega(t_1) dt_1\right)$, and a change of variables $x(t) = \exp(\alpha t + \beta)$, the differential equation transforms into:

$$x^2 \frac{d^2\psi_1(x)}{dx^2} + x(a - bx) \frac{d\psi_1(x)}{dx} + c^2\psi_1(x) = 0. \quad (2.2)$$

As observed, the differential equation now depends on the variable x incorporating the phase β . The parameters a, b and c are defined as follows:

$$a = 1 + i\frac{\epsilon}{\alpha}, \quad (2.3)$$

$$b = -i\frac{A}{\alpha}, \quad (2.4)$$

$$c = \frac{i\Delta + \epsilon}{2\alpha}. \quad (2.5)$$

At this stage, these parameters acquire new roles: a introduces a shift to the system, b represents the signal strength, and c captures the contribution of spontaneous emission, including the imaginary coupling. Since equation (2.2) is not directly solvable, we apply the ansatz transformation $\psi_1(x) = x^\mu z(x)$. This reformulates the differential equation into the confluent hypergeometric equation:

$$x^2 \frac{d^2z(x)}{dx^2} + (\gamma - bx) \frac{dz(x)}{dx} - \mu bz(x) = 0, \quad (2.6)$$




FIG. 2. Variation of the different populations (ground and excited) of the first Nikitin model (Exp1) versus ϵ . We have considered $A = 2.0/\alpha, \beta = 1.5/\alpha$. we remark that the shift and the imaginary coupling are responsible for the loss of information when their values are high in cases (a₁) and (a₂). The exact results (Blue line) agree with the numerical solutions (red line).

where the parameters γ, b , and μ are given by:

$$\mu_{1,2} = \frac{1 - a \mp \sqrt{(1 - a)^2 - 4c^2}}{2}, \quad (2.7)$$

$$\gamma = 2\mu + a. \quad (2.8)$$

III. RESULTS

A. Probabilities amplitudes

The solution to equation (2.6) includes two linearly independent functions, $M(\mu, \gamma, z)$ and $U(\mu, \gamma, z)$. Using the ansatz transformation, the probability amplitudes




FIG. 3. Variation of the different populations (ground state and excited state) of the first Nikitin model (Exp1) versus the coupling. We have considered $A = 2.0, t = 5, \alpha = 1$. Exact results (Blue line) agrees with the numerical solutions (red line).

can be expressed as:

$$C_1(x, x_0) = x^\lambda \exp\left(-\frac{b}{2}x\right) (a_+(x_0) U_1(x) + a_-(x_0) V_1(x)), \quad (3.1)$$

$$C_2(x, x_0) = x^\lambda \exp\left(-\frac{b}{2}x\right) (a_+(x_0) U_2(x) + a_-(x_0) V_2(x)). \quad (3.2)$$

Here,

$$a_\pm(x_0) = b_\pm x_0^{-\lambda} \exp\left(\frac{b}{2}x_0\right), \quad (3.3)$$

$$U_1(x) = x^\mu M(\mu, \gamma, z), \quad (3.4)$$

$$V_1(x) = x^\mu U(\mu, \gamma, z), \quad (3.5)$$

$$U_2(x) = -\frac{i\mu}{c} x^\mu \left(M(\mu, \gamma, z) + \frac{x}{\gamma} M(\mu + 1, \gamma + 1, z) \right), \quad (3.6)$$

$$V_2(x) = -\frac{i\mu}{c}x^\mu (U(\mu, \gamma, z) - xU(\mu+1, \gamma+1, z)), \quad (3.7)$$

The parameter λ is defined as:

$$\lambda = \frac{i\epsilon}{2\alpha}. \quad (3.8)$$




FIG. 4. Evolution of the different populations at the ground state and excited state of the first Nikitin model (Exp1) versus the shift ϵ for different values of the decay rates. We have considered $A = 2.0$, $t = 5.0$, $\Delta = 0.5$, $\alpha = 1$. The theoretical results (Blue line) agree with the numerical solutions (red line).

The functions $M(\mu, \gamma, x)$ and $U(\mu, \gamma, x)$ represent Kummer's confluent hypergeometric functions. Their properties, as described in [79], allow us to derive expressions for the derivatives:

$$\frac{d}{dz}M(\mu, \gamma, x) = \frac{\mu}{\gamma}M(\mu+1, \gamma+1, x), \quad (3.9)$$

$$\frac{d}{dz}U(\mu, \gamma, x) = -\mu U(\mu+1, \gamma+1, x). \quad (3.10)$$

The constants $b_{\pm}x_0^{-\lambda}$ are determined using [39].

B. Propagator and probabilities

To study the system's evolution between an initial time t_0 and an arbitrary time t , we use the propagator $C(x) = U(x, x_0)C(x_0)$. The components of the propagator are given by:

$$U_{k1}(x, x_0) = \frac{\omega_{k1}(x, x_0)}{\omega_{12}(x, x_0)} \exp\left(-\frac{b}{2}x\right) \exp\left(\frac{b}{2}x_0\right), \quad (3.11)$$

$$U_{k2}(x, x_0) = -\frac{\omega_{k2}(x, x_0)}{\omega_{12}(x, x_0)} \exp\left(-\frac{b}{2}x\right) \exp\left(\frac{b}{2}x_0\right). \quad (3.12)$$

The matrices elements are:

$$\omega_{kk'}(x, x_0) = V_k(x_0)U_{k'}(x) - U_k(x_0)V_{k'}(x), \quad (3.13)$$

$$\omega_{12}(x_0, x_0) = U_2(x_0)V_1(x_0) - U_1(x_0)V_2(x_0). \quad (3.14)$$

Using the Wronskian of the confluent hypergeometric function:

$$W = \frac{\Gamma(\gamma)}{\Gamma(\mu)}x^{-\gamma} \exp(x), \quad (3.15)$$

we calculate the transition parameter between levels 1 to level 2

$$\omega_{12}(x_0, x_0) = \frac{i\Gamma(\gamma)}{c\Gamma(\mu)}x_0^{2\mu-\gamma+1} \exp(x_0), \quad (3.16)$$

responsible for the transition between the ground and the excited state. The propagator element is expressed by

$$U_{12}(x, x_0) = ic\frac{\Gamma(\mu)}{\Gamma(\gamma)}(F(x, x_0) - F(x_0, x)) \exp(\vartheta(x, x_0)), \quad (3.17)$$

$$\begin{aligned} U_{22}(x, x_0) = & \mu b^{\gamma-1} \frac{\Gamma(\mu)}{\Gamma(\gamma)}(F(x, x_0) - F(x_0, x)) \\ & + \frac{bx}{\gamma} M(\mu+1, \gamma+1, x_0) U(\mu, \gamma, x) \\ & + bx_0 M(\mu+1, \gamma+1, x) U(\mu, \gamma, x_0) \exp(\vartheta(x, x_0)). \end{aligned} \quad (3.18)$$

The two matrices of the propagator are necessary to describe the transition or the survival of the particle. we easily determine the survival probability given by:

$$P_{22}(x, x_0) = \text{Re}(U_{22}(x, x_0)) + \text{Im}(U_{22}(x, x_0)). \quad (3.19)$$

The probability of transition is given by:

$$P_{12}(x, x_0) = \text{Re}(U_{12}(x, x_0)) + \text{Im}(U_{12}(x, x_0)). \quad (3.20)$$

C. Interpretation of the results

Figures 2, 3, and 4, describe the evolution of the ground and excited-state populations during spontaneous emission.

In Fig. 2, we observe the evolution of the first Nikitin model for the ground and excited-state populations as a function of time when the coupling and the shift vary. Regarding the excited-state population, when $\Delta > 0$ and $\epsilon > 0$ the system demonstrates an increase in oscillations with a weak amplitude, as shown in panel (a_1). Conversely, when $\Delta < 0$ and $\epsilon < 0$, as depicted in panel (a_3), the system also oscillates, but its amplitude doubles. This indicates that the imaginary coupling and the shift enhance information transfer at low values. When $\Delta = 0$ and $\epsilon \neq 0$ in panel (a_5), oscillations occur with very low amplitude due to the absence of coupling. In the ground-state population, oscillations decrease whether the imaginary coupling increases (a_2) or decreases (a_4), implying that this parameter does not significantly influence population evolution. However, when $\Delta = 0$ and $\epsilon \neq 0$, the oscillations increase, and transmission is maximal when the imaginary part of the coupling vanishes.

In Fig. 3, we plot the evolution of the different populations against the coupling when the phase β and the shift ϵ vary. In both the ground and excited states, the two populations exhibit exponential decay due to the absence of oscillations. This decay is less pronounced in panels (a_1), (a_2), (a_3), (a_5). However, in panels (a_4) and (a_6), the decay is more pronounced due to the high values of the shift in the ground-state population. In this case, the population exceeds one, an effect attributed to the high phase. Therefore, the shift and phase are responsible for the rapid decay of the population in this scenario.

Figure 4 illustrates the variation of the previous states versus the shift ϵ as the phase varies. In panels (c_2), (c_4) and (c_6) the population decreases exponentially. In contrast, panels (c_1), (c_3) and (c_5) exhibit slight oscillations due to the high value of the phase β in the excited-state population. This implies that the phase β is critical for the potential transmission of information in the ground-state population.

IV. ENERGIES OF THE NIKITIN MODEL

To study the energies of the Nikitin model, we analyze the eigenvalues of our system, which yield:

$$E_{\pm} = \pm (i\Delta + \epsilon) \csc(2\vartheta(t)), \quad (4.1)$$

where

$$\tan(2\vartheta(t)) = -\frac{i\Delta + \epsilon}{\Omega(t)}. \quad (4.2)$$

The eigenenergy is complex due to the imaginary coupling term, which highlights the non-Hermitian nature of




FIG. 5. Permitted and forbidden zone on the energy diagram of the real part. We have considered $\alpha = -15.0/A$, $t = 7.0/A$. We use Mathematica to plot these figures and we choose the Rainbow color as color function. The shift plays a great function in the creation of the barrier to control quantum information.

this model. The eigenenergy can be separated into real and imaginary parts:

$$\text{Re}(E_{\pm}) = \frac{1}{2} |Z^{1/2}(t)| \cos \phi, \quad (4.3)$$

$$\text{Im}(E_{\pm}) = \frac{1}{2} |Z^{1/2}(t)| \sin \phi, \quad (4.4)$$




FIG. 6. (Color online) The imaginary part of the energy against the coupling and the phase. In this figure, we have considered $A/\epsilon = 20/\epsilon$, $t = 15/\epsilon$, $\alpha = 0.5/\epsilon$. The sweep velocity α and the time t are responsible for the order or the chaos in the system for high values. Remind that the imaginary part of the energy describes the loss of quantum information.

with

$$\tan(2\phi) = \frac{2\epsilon\Delta}{\Omega^2(t) + \epsilon^2 - \Delta^2(t)}, \quad (4.5)$$

$$Z^{1/2}(t) = \frac{2\epsilon\Delta}{\Omega^2(t) + \epsilon^2 - \Delta^2(t)}. \quad (4.6)$$

Figures 5 and 6 provide graphical representations of the real and imaginary parts of the energy. The real part indicates energy gain, while the imaginary part reflects information loss.

In Fig.5, the color bar $Re(E_-)$ and $Re(E_+)$ illustrates the evolution of eigenenergies for the ground and excited states. We observe a transmission or forbidden zone for specific information. When $\epsilon \leq 2$, the system creates a barrier (white color) that prevents information transmission near zero on the coupling axis (panels (a₁) and (a₂)). When $\epsilon > 2$, the system generates a barrier (blue color) that allows information transmission in the same zone. Both phenomena occur over a long phase duration.

Figure 6 shows the imaginary part of the energy as a function of the phase and coupling. The color bar depicts the variation in imaginary coupling for the ground (a₁) and excited states (a₂). The transmission of information is localized (between $-30/\alpha$ and $-10/\alpha$) due to information loss. The sweep velocity and amplitude influence order or chaos in the system for high values. The imaginary part describes the system's quantum information loss.

V. SIMILARITY WITH THE RABI MODEL

The first exponential Nikitin model shows similarities with the Rabi model when time approaches $-\infty$ in an exponential function. Under these conditions, the detuning becomes $\Omega(t) = \epsilon$, while the coupling remains a complex function $i\Delta + \epsilon$. To study the dynamics of this model, we again use the time-dependent Schrödinger equation. After applying appropriate gauge transformations, the differential equation for the system becomes:




FIG. 7. (Color online) a) Interferogram of the population versus the time and the shift, we remark the presence of many waves around the crossing region at $\epsilon = 0$, we can assimilate the white line at $\epsilon = 0$ as a mirror that reflects any waves in the opposite direction. b) Similarity between the exact results and the numerical solutions against time of the excited-state population. We consider $\Delta = 0.2$

$$\frac{d^2\psi_{1,2}(t)}{dt^2} \mp i\epsilon \frac{d\psi_{1,2}(t)}{dt} + \frac{1}{4}(i\Delta + \epsilon)^2\psi_{1,2}(t) = 0. \quad (5.1)$$

The initial conditions of the Rabi problem $\psi_1(0) = 1$ and $\psi_2(0) = 0$, help solve this differential equation and determine the survival and transition probabilities. The survival probability is given by:

$$P_{Rabi}(t) = \frac{4\epsilon^2 + 3(i\Delta + \epsilon)^2}{4(\epsilon^2 + (i\Delta + \epsilon)^2)} \sin^2 \left(\frac{\sqrt{\epsilon^2 + (i\Delta + \epsilon)^2}}{2} t \right). \quad (5.2)$$

In Fig.7, we plot the variation of the population as a function of time. In frame (a), the interferogram of the population shows the presence of multiple waves around a mirror at $\epsilon = 0$, which reflects them in the opposite direction. These waves are generated by the small values of the coupling, leading us to conclude that the coupling is responsible for the interferometry observed in the system. Interferometry allows the extraction of significant information about the system.

CONCLUSION

We have presented the first exponential Nikitin model under the influence of spontaneous emission. This emission is accounted for by the presence of an imaginary coupling combined with a shift. The inclusion of the imaginary coupling introduces nonlinearity into the system's evolution and contributes to its non-Hermitian nature. To study the dynamics of this model, we determined the transition and survival probabilities as functions of time, coupling, and shift.

When the population varies with time, as shown in Fig.2, we observed that the imaginary coupling and the shift enhance the transmission of information for small values in the ground-state population. Conversely, these parameters have no significant influence on the excited-state population. When populations vary with the coupling, as illustrated in Fig.3, both ground and excited states exhibit exponential decay. Rapid decay occurs only when the phase takes a high value. Similarly, Fig.4 demonstrates exponential decay in populations, but some oscillations appear when the phase is large, indicating information transfer.

The non-Hermitian nature of our system results in complex eigenenergies, which can be separated into real and imaginary parts. The evolution of these eigenenergies indicates that energy gain is localized over a broad region, represented by the real part of the energy. Meanwhile, information loss is confined to a smaller region, represented by the imaginary part. Furthermore, the imaginary part contributes significantly to the transmission of the Rabi population, creating a "mirror" effect that reflects waves and generates oscillations.

For future work, it would be interesting to analyze spontaneous emission at different sweep velocities, utilizing both the Demkov-Kunike and the second Nikitin model.

ACKNOWLEDGMENTS

A. D. Kammogne thanked M. B. Kenmoe for interesting scientific comments and the University of Dschang

for its warm hospitality during this project.

DECLARATION

This paper has received no funding or financial support from anyone.

REFERENCES

- [1] S. Ke, D. Zhao, Q. Liu, and W. Liu, Adiabatic transfer of surface plasmons in non-Hermitian graphene waveguides, *Optical and Quantum Electronics*, 50, 1-13, (2018), <https://doi.org/10.1007/s11082-018-1661-3>.
- [2] Y. Hadad, N. Engheta, Possibility for inhibited spontaneous emission in electromagnetically open parity-time-symmetric guiding structures, *Proceedings of the National Academy of Sciences*, 117(11), 5576-5581, (2020), <https://doi.org/10.1073/pnas.1914279117>.
- [3] L. Jakóbczyk, A. Jamróz, Entanglement and nonlocality versus spontaneous emission in two-atom systems, *Physics Letters A*, 318(4-5), 318-326, (2003), <https://doi.org/10.1016/j.physleta.2003.09.044>.
- [4] W. Becker, J. K. McIver, Madey's theorems for free-electron devices, spontaneous emission, and applications, *Zeitschrift für Physik D Atoms, Molecules and Clusters*, 7, 353-372, (1988), <https://doi.org/10.1007/BF01439805>.
- [5] F. Chen, Y. Leng, Improving the capacity of quantum dense coding with both spontaneous emission and dephasing by non-Hermitian operation, *Laser Physics*, 34(6), 065202, (2024), <https://doi.org/10.1088/1555-6611/ad3aec>.
- [6] Zongping Gong, Miguel Bello, Daniel Malz, and Flore K. Kunst, Bound states and photon emission in non-Hermitian nanophotonics, *Phys. Rev. A* 106, 053517, 2022, <https://doi.org/10.1103/PhysRevA.106.053517>.
- [7] M. Khanbekyan, and J. Wiersig, Decay suppression of spontaneous emission of a single emitter in a high-Q cavity at exceptional points, *Phys. Rev. Research* 2, 023375, 2020, <https://doi.org/10.1103/PhysRevResearch.2.023375>
- [8] Wenjie Zhou, Jingfeng Liu, Juanfeng Zhu, Dmitrii Gromyko, Chengwei Qiu, Lin Wu, Exceptional points unveiling quantum limit of fluorescence rates in non-Hermitian plexcitonic single-photon sources, *APL Quantum*, 1(1), (2024), <https://doi.org/10.1063/5.0191494>.
- [9] Satoshi Sunada, Enhanced response of non-Hermitian photonic systems near exceptional points, *Phys. Rev. A* 97, 043804, (2018), <https://doi.org/10.1103/PhysRevA.97.043804>.
- [10] Yiming Huang, Yuhao Kang, and Azriel Z. Genack, Wave excitation and dynamics in non-Hermitian disordered systems, *Phys. Rev. Research* 4, 013102, (2022), <https://doi.org/10.1103/PhysRevResearch.4.013102>.
- [11] P. Ao, Quantum dynamics of a two-state system in a dissipative environment, *Physical Review B*, 43(7), 5397, (1991), <https://doi.org/10.1103/PhysRevB.43.5397>.
- [12] A. Kudlis, I. Iorsh, and I. V. Tokatly, Dissipation and spontaneous emission in quantum electrodynamical density functional theory based on optimized effective potential: A proof of concept study, *Physical Review B*, 105(5), 054317, (2022), <https://doi.org/10.1103/PhysRevB.105.054317>.
- [13] G. S. Agarwal, Quantum statistical theories of spontaneous emission and their relation to other approaches, *Quantum Optics*, 1-128, (2006), <https://doi.org/10.1007/BFb0042382>.
- [14] D. Meschede, Radiating atoms in confined space: From spontaneous emission to micromasers, *Physics Reports*, 211(5), 201-250, (1992), [https://doi.org/10.1016/0370-1573\(92\)90110-L](https://doi.org/10.1016/0370-1573(92)90110-L).
- [15] P. Ginzburg, Accelerating spontaneous emission in open resonators, *Annalen der Physik*, 528(7-8), 571-579, (2016), <https://doi.org/10.1002/andp.201500317>.
- [16] Matthew Pelton, Modified spontaneous emission in nanophotonic structures, *Nature Photonics* volume 9, pages427-435 (2015), <https://doi.org/10.1038/nphoton.2015.103>.
- [17] T. Sondergaard, B. Tromborg, General theory for spontaneous emission in active dielectric microstructures: Example of a fiber amplifier, *Phys. Rev. A* 64, 033812, (2001), <https://doi.org/10.1103/PhysRevA.64.033812>.
- [18] T. Brandes and B. Kramer, Spontaneous Emission of Phonons by Coupled Quantum Dots, *Phys. Rev. Lett.* 83, 1999, <https://doi.org/10.1103/PhysRevLett.83.3021>.
- [19] Vasili V. Klimov, M. Ducloy and V. S. Letokhov, Spontaneous emission of an atom in the presence of nanobodies, *Quantum Electronics*, Volume 31, Number 7, (2001), <https://doi.org/10.1070/QE2001v03n07ABEH002007>.
- [20] Alejandro González-Tudela, Connecting steady-states of driven-dissipative photonic lattices with spontaneous collective emission phenomena, *New Journal of Physics*, Volume 24, April (2022), <https://doi.org/10.1088/1367-2630/ac58b7>.
- [21] B. Qi, H. Z. Chen, L. Ge, P. Berini, and R. M. Ma, Parity-time symmetry synthetic lasers: Physics and devices, *Advanced Optical Materials*, 7(22), 1900694, (2019), <https://doi.org/10.1002/adom.201900694>.
- [22] Y. Yamamoto, S. Machida, and G. Björk, Microcavity semiconductor lasers with controlled spontaneous emission, *Opt Quant Electron* 24, S215-S243 (1992), <https://doi.org/10.1007/BF00625826>.

- [23] C. Henry, "Theory of spontaneous emission noise in open resonators and its application to lasers and optical amplifiers," Journal of Lightwave Technology, vol. 4, no. 3, pp. 288-297, (1986), <https://doi.org/10.1109/JLT.1986.1074715>.
- [24] Mariano A. Zimmmer, Jiming Bao, Federico Capasso, Sven Müller, Carsten Ronning, Laser action in nanowires: Observation of the transition from amplified spontaneous emission to laser oscillation, Applied Physics Letters, 93(5), (2008), <https://doi.org/10.1063/1.2965797>.
- [25] N.P. Barnes, B.M. Walsh, Amplified spontaneous emission-application to Nd:YAG lasers, IEEE Journal of Quantum Electronics, 35, 1, January (1999), <https://doi.org/10.1109/3.737626>.
- [26] A. V. Bogatskaya, E. A. Volkova and A. M. Popov, Spontaneous emission of atoms in a strong laser field, Journal of Experimental and Theoretical Physics, 125, 587-596, (2017), <https://doi.org/10.1134/S1063776117090114>.
- [27] H. Cao, J. Y. Xu, S.-H. Chang, and S. T. Ho, Transition from amplified spontaneous emission to laser action in strongly scattering media, Phys. Rev. E 61, (1985), <https://doi.org/10.1103/PhysRevE.61.1985> Phys. Rev. E 61, (1985).
- [28] F. H. Nicoll, Decrease in Spontaneous Emission at the Onset of Lasing in Semiconductors, J. Appl. Phys. 42, 2743-2746 (1971), <https://doi.org/10.1063/1.1660616>.
- [29] Weng. W. Chow, Frank Jahnke, and Christopher Gies, Light: Science and Applications volume 3, pagee201 (2014), <https://doi.org/10.1038/lسا.2014.82>.
- [30] Y. W. Huang, J.S. Wang, J. C. Moreno, and H. R. Griem, Measurement and interpretation of spontaneous line emission from a common upper level in laser-produced carbon plasmas, Erratum Phys. Rev. Lett. 65, 2747 (1990), <https://doi.org/10.1103/PhysRevLett.65.1757>.
- [31] A. E. Siegman, Excess spontaneous emission in non-Hermitian optical systems, Physical Review A, 39(3), 1253, 1989, <https://doi.org/10.1103/PhysRevA.39.1253>.
- [32] L. Ferrier, P. Bouteyre, A. Pick, S. Cueff, N. M. Dang, C. Diederichs, ..., and H. S. Nguyen, Unveiling the enhancement of spontaneous emission at exceptional points, Physical Review Letters, 129(8), 083602, 2022, <https://doi.org/10.1103/PhysRevLett.129.083602>.
- [33] A. Pick, B. Zhen, O. D. Miller, C. W. Hsu, F. Hernandez, A. W. Rodriguez, ...and S. G. Johnson, General theory of spontaneous emission near exceptional points, Optics express, 25(11), 12325-12348, 2017, <https://doi.org/10.1364/OE.25.012325>.
- [34] S. Lieu, Topological phases in the Non-Hermitian Su-Schrieffer-Heeger model, Physical Review B, 97(4), 045106, 2018, <https://doi.org/10.1103/PhysRevB.97.045106>.
- [35] Y. Ashida, Z. Gong, M. Ueda, Non-hermitian physics, Advances in Physics, 69(3), 249-435, 2020, <https://doi.org/10.1080/00018732.2021.1876991>.
- [36] T. E. Lee, and, C. K. Chan, Heralded magnetism in non-Hermitian atomic systems, Physical Review X, 4(4), 041001, 2014, <https://doi.org/10.1103/PhysRevX.4.041001>.
- [37] B. S. Nesbitt, S. F. C. O'Rourke, and D. S. F.Crothers, (1996). Analytic continuation of the Nikitin exponential model to non-zero impact parameters, Journal of Physics B: Atomic, Molecular and Optical Physics, 29(12), 2515, 1996, <https://doi.org/10.1088/0953-4075/29/12/014>.
- [38] N. V. Vitanov, Generalized Nikitin model: strong-coupling approximation, Journal of Physics B: Atomic, Molecular and Optical Physics, 27(9), 1791, 1994, <https://doi.org/10.1088/0953-4075/27/9/016>.
- [39] A. D. Kammogne, N. Issofa, and L. C. Fai, Non-resonant exponential Nikitin models with decay, Chinese Journal of Physics, 88, 690-705, 2024, <https://doi.org/10.1016/j.cjph.2023.10.013>.
- [40] A. D. Kammogne, M. B. Kenmoe, and L. C. Fai, Statistics of interferograms in three-level systems, Physics Letters A, 425, 127872, 2022, <https://doi.org/10.1016/j.physleta.2021.127872>.
- [41] V. Sidis, D. Grimbart, M. Sizun, and, M. Baer, Quantal los calculations of differential cross sections for vibrational excitation and vibronic charge transfer in H++O₂ collisions, Chemical physics letters, 163(1), 19-22, 1989, [https://doi.org/10.1016/0009-2614\(89\)80004-1](https://doi.org/10.1016/0009-2614(89)80004-1).
- [42] H. W. Li, and J. Z. Sun, Bose-Einstein condensates under a non-Hermitian spin-orbit coupling, Chinese Physics B, 30(6), 066702, 2021, <https://doi.org/10.1088/1674-1056/abd7e4>.
- [43] Radka Bach, Marek Trippenbach, and Kazimierz Rzażewski , Spontaneous emission of atoms via collisions of Bose-Einstein condensates, Phys. Rev. A 65, 063605, (2002), <https://doi.org/10.1103/PhysRevA.65.063605>.
- [44] M. R. Hush, S. S. Szigeti, A. R. R. Carvalho, and J. J. Hope, Controlling spontaneous-emission noise in measurement-based feedback cooling of a Bose-Einstein condensate, New Journal of Physics, 15(11), 113060 (2013), <https://doi.org/10.1088/1367-2630/15/11/113060>.
- [45] C. M. Savage, J. Ruostekoski, and D. F. Walls, Spontaneous photon emission stimulated by two Bose-Einstein condensates, Physical Review A, 56(3), 2046, (1997), <https://doi.org/10.1103/PhysRevA.56.2046>.
- [46] D. C. Dai, and A. P. Monkman, Observation of superfluorescence from a quantum ensemble of coherent excitons in a ZnTe crystal: Evidence for spontaneous Bose-Einstein condensation of excitons, Physical Review B-Condensed Matter and Materials Physics, 84(11), 115206, (2011), <https://doi.org/10.1103/PhysRevB.84.115206>.
- [47] D. Snoke, Spontaneous Bose coherence of excitons and polaritons, Science, 298(5597), 1368-1372, (2002), <https://doi.org/10.1126/science.1078082>.
- [48] L. W. Clark, A. Gaj, L., Feng, and C. Chin, Collective emission of matter-wave jets from driven Bose-Einstein condensates, Nature, 551(7680), 356-359, (2017), <https://doi.org/10.1038/nature24272>.
- [49] Julian Schmitt, Tobias Damm, David Dung, Christian Wahl, Frank Vewinger, Jan Klaers, and Martin Weitz, Spontaneous Symmetry Breaking and Phase Coherence of a Photon Bose-Einstein Condensate Coupled to a Reservoir, Phys. Rev. Lett. 116, 033604, (2016), <https://doi.org/10.1103/PhysRevLett.116.033604>.
- [50] J. Kasprzak, M. Richard, S. Kundermann, A. Baas, P. Jeambrun, J. M. J. Keeling, ... and L. S. Dang, Bose-Einstein condensation of exciton polaritons, Nature, 443(7110), 409-414. (2006), <https://doi.org/10.1038/nature05131>.
- [51] Hui Deng, Hartmut Haug, and Yoshihisa Yamamoto, Exciton-polariton Bose-Einstein condensation, Rev. Mod. Phys. 82, 1489, (2010), <https://doi.org/10.1103/RevModPhys.82.1489>.

- [52] J. Bloch, I. Carusotto, and M. Wouters, Non-equilibrium Bose-Einstein condensation in photonic systems, *Nature Reviews Physics*, 4(7), 470-488, (2022), <https://doi.org/10.1038/s42254-022-00464-0>.
- [53] D. J. Heinzen, Roahn Wynar, P. D. Drummond, and K. V. Kheruntsyan, Superchemistry: Dynamics of Coupled Atomic and Molecular Bose-Einstein Condensates, *Phys. Rev. Lett.* 84, 5029 (2000), <https://doi.org/10.1103/PhysRevLett.84.5029>.
- [54] M. Reisenbauer, H. Rudolph, L. Egyed, K. Hornberger, A. V. Zasedatelev, M. Abuzarli, ..., and U. Delić, Non-Hermitian dynamics and non-reciprocity of optically coupled nanoparticles, *Nature Physics*, 1-7, (2024), <https://doi.org/10.1038/s41567-024-02589-8>.
- [55] H. J. Rose and G. A. Jones, A new kind of natural radioactivity, *Nature* volume 307, pages245-247 (1984), <https://doi.org/10.1038/307245a0>.
- [56] J. Gross, and C. P. Leblond, LOCALIZATION OF RADIOACTIVE ELEMENTS:(A Review), *Canadian Medical Association Journal*, 57(2), 102, (1947), <https://doi.org/PMC1590597>.
- [57] H. J. WALKE, Spontaneous Emission of Neutrons from Radioactive Isotopes, *Nature* volume 134, page215 (1934), <https://doi.org/10.1038/134215a0>.
- [58] F. Hartmann-Boutron et D. Spanjaard, Study of radioactive impurities in solids. Part one : radiation characteristics, *J. Phys. France* 33, 285-297 (1972), <https://doi.org/10.1051/jphys:01972003302-3028500>.
- [59] J. van Brakel, The possible influence of the discovery of radio-active decay on the concept of physical probability, *Archive for history of exact sciences*, 31, 369-385, (1985), <https://doi.org/10.1007/BF00348520>.
- [60] M. Goldhaber and R. D. Hill, Leo Szilard, Radioactivity Induced by Nuclear Excitation I. Excitation by Neutrons, *Phys. Rev.* 55, 47, 1939, <https://doi.org/10.1103/PhysRev.55.47>.
- [61] V. V. Gromov, Radiation-chemical Processes in Radioactive Compounds, *Russian Chemical Reviews*, Volume 42, Number 1, (1973), <https://doi.org/10.1070/RC1973v042n01ABEH002565>.
- [62] D.N. Poenaru et M. Ivascu, ^5He radioactivity, *J. Phys. France* 45, 1099-1106 (1984), *J. Phys. France* 45, 1099-1106 (1984), <https://doi.org/10.1051/jphys:019840045070109900>.
- [63] E. Pickup and L. Voyvodic, Radioactive Nuclei Emitted in Cosmic-Ray Stars, *Phys. Rev.* 80, 1100, (1950), <https://doi.org/10.1103/PhysRev.80.1100>.
- [64] Z. Shi, L. Qin, X. Zhao., and H. Huang, Optical pattern formation in a rydberg-dressed atomic gas with non-hermitian potentials, *Photonics MDPI* 9, 11, 856, , (2022), <https://doi.org/10.3390/photonics9110856>.
- [65] Timothy P. Softley, Cold and ultracold molecules in the twenties, *Proceedings of the Royal Society A*, 479(2274), 20220806. (2023), <https://doi.org/10.1098/rspa.2022.0806>.
- [66] Paul S. Julienne and Frederick H. Mies, Collisions of ultracold trapped atoms, *JOSA B*, 6(11), 2257-2269, (1989), <https://doi.org/10.1364/JOSAB.6.002257>.
- [67] S. Ostermann, F. Piazza, and H. Ritsch, Spontaneous Crystallization of Light and Ultracold Atoms, *Phys. Rev. X* 6, 021026, (2016), <https://doi.org/10.1103/PhysRevX.6.021026>.
- [68] Paul S. Julienne, In Scattering (pp. 1043-1067), Academic Press, (2002), <https://doi.org/10.1016/B978-012613760-6/50056-5>.
- [69] Laurent Hilico, Jean-Philippe Karr, Albane Douillet, Paul Indelicato, Sebastian Wolf, and Ferdinand Schmidt Kaler, Preparing single ultra-cold antihydrogen atoms for free-fall in GBAR, *International Journal of Modern Physics: Conference Series* (Vol. 30, p. 1460269), (2014), <https://doi.org/10.1142/S2010194514602695>.
- [70] Dylan Jervis and Joseph H. Thywissen, Making an Ultracold Gas, *Quantum Gas Experiments*, pp. 5-32, (2014), <https://doi.org/10.1142/9781783264766-0002>.
- [71] I. Bloch, J. Dalibard, and S. Nascimbene , Quantum simulations with ultracold quantum gases, *Nature Physics* volume 8, pages267-276 (2012), <https://doi.org/10.1038/nphys2259>.
- [72] K. Roux, V. Helson, H. Konishi, and J. P. Brantut, New *J. Phys.* 23 043029, (2021), <https://doi.org/10.1088/1367-2630/abeb91>.
- [73] T. E. Drake, Y. Sagi, R. Paudel, J. T. Stewart, J. P. Gaebler, and D. S. Jin, Direct observation of the Fermi surface in an ultracold atomic gas, *Phys. Rev. A* 86, 031601(R), (2012), <https://doi.org/10.1103/PhysRevA.86.031601>.
- [74] L. Santos, G. V. Shlyapnikov, P. Zoller1, and M. Lewenstein, Bose-Einstein Condensation in Trapped Dipolar Gases, *Erratum Phys. Rev. Lett.* 88, 139904 (2002), <https://doi.org/10.1103/PhysRevLett.85.1791>.
- [75] K. N. Zlatanov, G. S. Vasilev, P. A. Ivanov, and N. V. Vitanov, Exact solution of the Bloch equations for the non-resonant exponential model in the presence of dephasing, *Phys. Rev. A* 92, 043404 – Published 6 October, (2015), <https://doi.org/10.1103/PhysRevA.92.043404>.
- [76] A. B. Tchapda, M. B. Kenmoe, and L. C. Fai, Landau-Zener transitions in a qubit/qutrit periodically driven in both longitudinal and transverse directions, *arXiv:1708.04184v2*, 2017 <https://doi.org/10.48550/arXiv.1708.04184>.
- [77] C. L. Cortes, M. Otten, and S. K. Gray, Non-Hermitian approach for quantum plasmonics, *The Journal of Chemical Physics*, 152(8), 2020, <https://doi.org/10.1063/1.5131762>.
- [78] F. Bagarello, and J. Feinberg, Bicoherent-state path integral quantization of a non-Hermitian Hamiltonian, *Annals of Physics*, 422, 168313, 2020, <https://doi.org/10.1016/j.aop.2020.168313>.
- [79] H. Bateman, and A. Erdélyi, Higher transcendental functions, *volume II. Bateman Manuscript Project* Mc Graw-Hill Book Company, 1953.
- [80] R. Beals, and R. Wong, Special functions: a graduate text, (Vol. 126). Cambridge University Press, 2010 .