

# The Miller-Rabin Primality Test

Aryaman Maithani

IIT Bombay

5th November 2021

# Table of Contents

- 1 Introduction
- 2 First attempt
- 3 The Miller-Rabin test

# Table of Contents

1 Introduction

2 First attempt

3 The Miller-Rabin test

# The question at hand

In this talk, we are concerned with finding an algorithm for the following problem.

# The question at hand

In this talk, we are concerned with finding an algorithm for the following problem.

## Algorithm

Input: An integer  $n > 1$ .

# The question at hand

In this talk, we are concerned with finding an algorithm for the following problem.

## Algorithm

Input: An integer  $n > 1$ .

Output:  $\text{isPrime}(n)$ .

# Naïve approach

The simplest way to do this is by

# Naïve approach

The simplest way to do this is by trial division.



# Naïve approach

The simplest way to do this is by trial division. Indeed, we simply divide  $n$  by 2, 3, 4, and so on, and see if the remainder is 0 in any case.

# Naïve approach

The simplest way to do this is by trial division. Indeed, we simply divide  $n$  by 2, 3, 4, and so on, and see if the remainder is 0 in any case. As we know, we only need to divide by numbers up to  $\sqrt{n}$ .

# Naïve approach

The simplest way to do this is by trial division. Indeed, we simply divide  $n$  by 2, 3, 4, and so on, and see if the remainder is 0 in any case. As we know, we only need to divide by numbers up to  $\sqrt{n}$ . The issue with this algorithm is that it is extremely inefficient,

# Naïve approach

The simplest way to do this is by trial division. Indeed, we simply divide  $n$  by 2, 3, 4, and so on, and see if the remainder is 0 in any case. As we know, we only need to divide by numbers up to  $\sqrt{n}$ . The issue with this algorithm is that it is extremely inefficient, requiring  $\Theta(\sqrt{n})$  operations,

# Naïve approach

The simplest way to do this is by trial division. Indeed, we simply divide  $n$  by 2, 3, 4, and so on, and see if the remainder is 0 in any case. As we know, we only need to divide by numbers up to  $\sqrt{n}$ . The issue with this algorithm is that it is extremely inefficient, requiring  $\Theta(\sqrt{n})$  operations, which is *exponential* in the *bit length*  $\log(n)$ .

# Naïve approach

The simplest way to do this is by trial division. Indeed, we simply divide  $n$  by 2, 3, 4, and so on, and see if the remainder is 0 in any case. As we know, we only need to divide by numbers up to  $\sqrt{n}$ . The issue with this algorithm is that it is extremely inefficient, requiring  $\Theta(\sqrt{n})$  operations, which is *exponential* in the *bit length*  $\log(n)$ . For example, if  $n$  has 100-decimal digits, it would take more than  $10^{33}$  years to perform  $\sqrt{n}$  divisions.

# Naïve approach

The simplest way to do this is by trial division. Indeed, we simply divide  $n$  by 2, 3, 4, and so on, and see if the remainder is 0 in any case. As we know, we only need to divide by numbers up to  $\sqrt{n}$ . The issue with this algorithm is that it is extremely inefficient, requiring  $\Theta(\sqrt{n})$  operations, which is *exponential* in the *bit length*  $\log(n)$ . For example, if  $n$  has 100-decimal digits, it would take more than  $10^{33}$  years to perform  $\sqrt{n}$  divisions.

Moreover, note that the above algorithm does *more* than what we expected from our algorithm.

# Naïve approach

The simplest way to do this is by trial division. Indeed, we simply divide  $n$  by 2, 3, 4, and so on, and see if the remainder is 0 in any case. As we know, we only need to divide by numbers up to  $\sqrt{n}$ . The issue with this algorithm is that it is extremely inefficient, requiring  $\Theta(\sqrt{n})$  operations, which is *exponential* in the *bit length*  $\log(n)$ . For example, if  $n$  has 100-decimal digits, it would take more than  $10^{33}$  years to perform  $\sqrt{n}$  divisions.

Moreover, note that the above algorithm does *more* than what we expected from our algorithm. Namely, it not only tells us that the number is prime but also produces a nontrivial factor in the case that  $n$  is composite.



# Probabilistic approach

In this talk, we describe a much faster primality testing.

# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm.

# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm. It allows for 100-decimal digits numbers to be tested in less than a second.

# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm. It allows for 100-decimal digits numbers to be tested in less than a second. Unlike the earlier algorithm, it does *not* give us a prime factor in the case that  $n$  is composite.

# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm. It allows for 100-decimal digits numbers to be tested in less than a second. Unlike the earlier algorithm, it does *not* give us a prime factor in the case that  $n$  is composite.

The catch?

# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm. It allows for 100-decimal digits numbers to be tested in less than a second. Unlike the earlier algorithm, it does *not* give us a prime factor in the case that  $n$  is composite.

The catch? This algorithm is *probabilistic*.

# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm. It allows for 100-decimal digits numbers to be tested in less than a second. Unlike the earlier algorithm, it does *not* give us a prime factor in the case that  $n$  is composite.

The catch? This algorithm is *probabilistic*. This means that the algorithm can make a mistake.

# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm. It allows for 100-decimal digits numbers to be tested in less than a second. Unlike the earlier algorithm, it does *not* give us a prime factor in the case that  $n$  is composite.

The catch? This algorithm is *probabilistic*. This means that the algorithm can make a mistake.

*However,*



# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm. It allows for 100-decimal digits numbers to be tested in less than a second. Unlike the earlier algorithm, it does *not* give us a prime factor in the case that  $n$  is composite.

The catch? This algorithm is *probabilistic*. This means that the algorithm can make a mistake.

*However*, one has control over this probability, and can make it arbitrarily small

# Probabilistic approach

In this talk, we describe a much faster primality testing. This is a polynomial time algorithm. It allows for 100-decimal digits numbers to be tested in less than a second. Unlike the earlier algorithm, it does *not* give us a prime factor in the case that  $n$  is composite.

The catch? This algorithm is *probabilistic*. This means that the algorithm can make a mistake.

*However*, one has control over this probability, and can make it arbitrarily small (but not zero).

# Some algebraic objects

For the rest of the talk, we shall assume that  $n > 1$  is an *odd* integer.

# Some algebraic objects

For the rest of the talk, we shall assume that  $n > 1$  is an *odd* integer. Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be its prime factorisation.

# Some algebraic objects

For the rest of the talk, we shall assume that  $n > 1$  is an *odd* integer. Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be its prime factorisation.

By  $\mathbb{Z}_n$ , we shall denote the ring of integers modulo  $n$ .

# Some algebraic objects

For the rest of the talk, we shall assume that  $n > 1$  is an *odd* integer. Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be its prime factorisation.

By  $\mathbb{Z}_n$ , we shall denote the ring of integers modulo  $n$ . We have a ring homomorphism

$$\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$$

# Some algebraic objects

For the rest of the talk, we shall assume that  $n > 1$  is an *odd* integer. Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be its prime factorisation.

By  $\mathbb{Z}_n$ , we shall denote the ring of integers modulo  $n$ . We have a ring homomorphism

$$\begin{aligned}\theta : \mathbb{Z}_n &\rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}} \\ [a]_n &\mapsto ([a]_{p_1^{e_1}}, \cdots, [a]_{p_r^{e_r}}).\end{aligned}$$

# Some algebraic objects

For the rest of the talk, we shall assume that  $n > 1$  is an *odd* integer. Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be its prime factorisation.

By  $\mathbb{Z}_n$ , we shall denote the ring of integers modulo  $n$ . We have a ring homomorphism

$$\begin{aligned}\theta : \mathbb{Z}_n &\rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}} \\ [a]_n &\mapsto ([a]_{p_1^{e_1}}, \cdots, [a]_{p_r^{e_r}}).\end{aligned}$$

In fact, the Chinese Remainder Theorem tells us that the above is an isomorphism.



# Some algebraic objects

For the rest of the talk, we shall assume that  $n > 1$  is an *odd* integer. Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be its prime factorisation.

By  $\mathbb{Z}_n$ , we shall denote the ring of integers modulo  $n$ . We have a ring homomorphism

$$\begin{aligned}\theta : \mathbb{Z}_n &\rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}} \\ [a]_n &\mapsto ([a]_{p_1^{e_1}}, \cdots, [a]_{p_r^{e_r}}).\end{aligned}$$

In fact, the Chinese Remainder Theorem tells us that the above is an isomorphism. This gives us a group isomorphism between the group of invertible elements of the two rings as

# Some algebraic objects

For the rest of the talk, we shall assume that  $n > 1$  is an *odd* integer. Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be its prime factorisation.

By  $\mathbb{Z}_n$ , we shall denote the ring of integers modulo  $n$ . We have a ring homomorphism

$$\begin{aligned}\theta : \mathbb{Z}_n &\rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}} \\ [a]_n &\mapsto ([a]_{p_1^{e_1}}, \cdots, [a]_{p_r^{e_r}}).\end{aligned}$$

In fact, the Chinese Remainder Theorem tells us that the above is an isomorphism. This gives us a group isomorphism between the group of invertible elements of the two rings as

$$(\mathbb{Z}_n)^* \xrightarrow{\cong} (\mathbb{Z}_{p_1^{e_1}})^* \times \cdots \times (\mathbb{Z}_{p_r^{e_r}})^*.$$

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ .

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ .

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime.

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

- 1 there is an efficient algorithm that on input  $n$  and  $\alpha \in \mathbb{Z}_n^+$ , determines if  $\alpha \in L_n$ ;



# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

- 1 there is an efficient algorithm that on input  $n$  and  $\alpha \in \mathbb{Z}_n^+$ , determines if  $\alpha \in L_n$ ;
- 2 if  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ ; and

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

- 1 there is an efficient algorithm that on input  $n$  and  $\alpha \in \mathbb{Z}_n^+$ , determines if  $\alpha \in L_n$ ;
- 2 if  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ ; and
- 3 if  $n$  is composite,  $|L_n| \leq c(n - 1)$  for some universal constant  $c < 1$ .

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

- 1 there is an efficient algorithm that on input  $n$  and  $\alpha \in \mathbb{Z}_n^+$ , determines if  $\alpha \in L_n$ ;
- 2 if  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ ; and
- 3 if  $n$  is composite,  $|L_n| \leq c(n - 1)$  for some universal constant  $c < 1$ .

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

- 1 there is an efficient algorithm that on input  $n$  and  $\alpha \in \mathbb{Z}_n^+$ , determines if  $\alpha \in L_n$ ;
- 2 if  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ ; and
- 3 if  $n$  is composite,  $|L_n| \leq c(n - 1)$  for some universal constant  $c < 1$ .

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ .

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

- 1 there is an efficient algorithm that on input  $n$  and  $\alpha \in \mathbb{Z}_n^+$ , determines if  $\alpha \in L_n$ ;
- 2 if  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ ; and
- 3 if  $n$  is composite,  $|L_n| \leq c(n - 1)$  for some universal constant  $c < 1$ .

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ ,

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

- 1 there is an efficient algorithm that on input  $n$  and  $\alpha \in \mathbb{Z}_n^+$ , determines if  $\alpha \in L_n$ ;
- 2 if  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ ; and
- 3 if  $n$  is composite,  $|L_n| \leq c(n - 1)$  for some universal constant  $c < 1$ .

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true;

# Bird's eye view of probabilistic tests

Several probabilistic primality tests, including the Miller–Rabin test, have the following general structure.

Define  $\mathbb{Z}_n^+$  to be the set of nonzero elements of  $\mathbb{Z}_n$ . Note that  $|\mathbb{Z}_n^+| = n - 1$ . Moreover,  $\mathbb{Z}_n^+ = \mathbb{Z}_n^*$  iff  $n$  is prime. Suppose also that we define a set  $L_n \subseteq \mathbb{Z}_n^+$  such that:

- 1 there is an efficient algorithm that on input  $n$  and  $\alpha \in \mathbb{Z}_n^+$ , determines if  $\alpha \in L_n$ ;
- 2 if  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ ; and
- 3 if  $n$  is composite,  $|L_n| \leq c(n - 1)$  for some universal constant  $c < 1$ .

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.



## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- 1 The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.

## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- 1 The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.
- 2 If  $n$  is prime,

## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- 1 The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.
- 2 If  $n$  is prime, then the algorithm outputs true,

## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- 1 The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.
- 2 If  $n$  is prime, then the algorithm outputs true, and it does so *correctly*.

## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- 1 The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.
- 2 If  $n$  is prime, then the algorithm outputs true, and it does so *correctly*.
- 3 If  $n$  is composite,

## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- 1 The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.
- 2 If  $n$  is prime, then the algorithm outputs true, and it does so *correctly*.
- 3 If  $n$  is composite, then the algorithm *may* output true,

## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- 1 The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.
- 2 If  $n$  is prime, then the algorithm outputs true, and it does so *correctly*.
- 3 If  $n$  is composite, then the algorithm *may* output true, with probability at most  $c^k$ .

## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- 1 The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.
- 2 If  $n$  is prime, then the algorithm outputs true, and it does so *correctly*.
- 3 If  $n$  is composite, then the algorithm *may* output true, with probability at most  $c^k$ .

In particular, note that there is a *one-sided error*.



## Algorithm

To test for primality, we set a “repetition parameter”  $k$ , and choose random elements  $\alpha_1, \dots, \alpha_k \in \mathbb{Z}_n^+$ . If  $\alpha_i \in L_n$  for all  $i \in \{1, \dots, k\}$ , then we output true; otherwise, we output false.

- ① The algorithm is efficient since we can check  $\alpha \in L_n$  efficiently.
- ② If  $n$  is prime, then the algorithm outputs true, and it does so *correctly*.
- ③ If  $n$  is composite, then the algorithm *may* output true, with probability at most  $c^k$ .

In particular, note that there is a *one-sided error*. In fancy language, this is a *Monte Carlo algorithm*.

# Table of Contents

1 Introduction

2 First attempt

3 The Miller-Rabin test

# First attempt

We now try to define a suitable candidate for  $L_n$ .

# First attempt

We now try to define a suitable candidate for  $L_n$ .

## Definition 1

$$L_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{n-1} = 1\}.$$

# First attempt

We now try to define a suitable candidate for  $L_n$ .

## Definition 1

$$L_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{n-1} = 1\}.$$

Note that we can test  $\alpha \in L_n$  efficiently, using a repeated-squaring algorithm.

# First attempt

We now try to define a suitable candidate for  $L_n$ .

## Definition 1

$$L_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{n-1} = 1\}.$$

Note that we can test  $\alpha \in L_n$  efficiently, using a repeated-squaring algorithm.

It is easy to see that  $L_n \subseteq \mathbb{Z}_n^*$ .

# First attempt

We now try to define a suitable candidate for  $L_n$ .

## Definition 1

$$L_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{n-1} = 1\}.$$

Note that we can test  $\alpha \in L_n$  efficiently, using a repeated-squaring algorithm.

It is easy to see that  $L_n \subseteq \mathbb{Z}_n^*$ . In fact,  $L_n$  is the kernel of the  $(n-1)$ -power map  $\mathbb{Z}_n^* \rightarrow \mathbb{Z}_n^*$  given by  $x \mapsto x^{n-1}$ .

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ .



# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ ,

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

## Proof sketch.

The first statement is clear.

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

## Proof sketch.

The first statement is clear. For the second, one recalls that  $L_n$  is a subgroup of  $\mathbb{Z}_n^*$ .

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

## Proof sketch.

The first statement is clear. For the second, one recalls that  $L_n$  is a subgroup of  $\mathbb{Z}_n^*$ . Thus,  $\frac{|\mathbb{Z}_n^*|}{|L_n|}$  is a positive integer.

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

## Proof sketch.

The first statement is clear. For the second, one recalls that  $L_n$  is a subgroup of  $\mathbb{Z}_n^*$ . Thus,  $\frac{|\mathbb{Z}_n^*|}{|L_n|}$  is a positive integer. Thus, if the integer is not 1, it is at least 2.



# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

## Proof sketch.

The first statement is clear. For the second, one recalls that  $L_n$  is a subgroup of  $\mathbb{Z}_n^*$ . Thus,  $\frac{|\mathbb{Z}_n^*|}{|L_n|}$  is a positive integer. Thus, if the integer is not 1, it is at least 2. Combine this with the fact that  $|\mathbb{Z}_n^*| \leq n-1$  to get the result. □

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

## Proof sketch.

The first statement is clear. For the second, one recalls that  $L_n$  is a subgroup of  $\mathbb{Z}_n^*$ . Thus,  $\frac{|\mathbb{Z}_n^*|}{|L_n|}$  is a positive integer. Thus, if the integer is not 1, it is at least 2. Combine this with the fact that  $|\mathbb{Z}_n^*| \leq n-1$  to get the result. □

However, there are infinitely many odd composite  $n$  for which  $L_n = \mathbb{Z}_n^*$

# Does it fit the bill?

## Theorem 2

If  $n$  is prime, then  $L_n = \mathbb{Z}_n^*$ . If  $n$  is composite and  $L_n \subsetneq \mathbb{Z}_n^*$ , then  $|L_n| \leq \frac{1}{2}(n-1)$ .

## Proof sketch.

The first statement is clear. For the second, one recalls that  $L_n$  is a subgroup of  $\mathbb{Z}_n^*$ . Thus,  $\frac{|\mathbb{Z}_n^*|}{|L_n|}$  is a positive integer. Thus, if the integer is not 1, it is at least 2. Combine this with the fact that  $|\mathbb{Z}_n^*| \leq n-1$  to get the result. □

However, there are infinitely many odd composite  $n$  for which  $L_n = \mathbb{Z}_n^*$  and thus, they cannot be ignored.

## Definition 3

An odd composite number  $n$  such that  $L_n = \mathbb{Z}_n^*$  is called a *Carmichael number*.

# Carmichael numbers

## Definition 3

An odd composite number  $n$  such that  $L_n = \mathbb{Z}_n^*$  is called a *Carmichael number*.

## Example

The smallest Carmichael number is  $561 = 3 \cdot 11 \cdot 17$ .

# Carmichael numbers

## Definition 3

An odd composite number  $n$  such that  $L_n = \mathbb{Z}_n^*$  is called a *Carmichael number*.

## Example

The smallest Carmichael number is  $561 = 3 \cdot 11 \cdot 17$ .

## Theorem 4

$n$  is a Carmichael number iff  $n$  is of the following form:

# Carmichael numbers

## Definition 3

An odd composite number  $n$  such that  $L_n = \mathbb{Z}_n^*$  is called a *Carmichael number*.

## Example

The smallest Carmichael number is  $561 = 3 \cdot 11 \cdot 17$ .

## Theorem 4

$n$  is a Carmichael number iff  $n$  is of the following form:

- 1  $n = p_1 \cdots p_r$  for distinct primes  $p_i$ ,

# Carmichael numbers

## Definition 3

An odd composite number  $n$  such that  $L_n = \mathbb{Z}_n^*$  is called a *Carmichael number*.

## Example

The smallest Carmichael number is  $561 = 3 \cdot 11 \cdot 17$ .

## Theorem 4

$n$  is a Carmichael number iff  $n$  is of the following form:

- ①  $n = p_1 \cdots p_r$  for distinct primes  $p_i$ ,
- ②  $r \geq 3$ ,



# Carmichael numbers

## Definition 3

An odd composite number  $n$  such that  $L_n = \mathbb{Z}_n^*$  is called a *Carmichael number*.

## Example

The smallest Carmichael number is  $561 = 3 \cdot 11 \cdot 17$ .

## Theorem 4

$n$  is a Carmichael number iff  $n$  is of the following form:

- ①  $n = p_1 \cdots p_r$  for distinct primes  $p_i$ ,
- ②  $r \geq 3$ ,
- ③  $(p_i - 1) \mid (n - 1)$  for all  $i \in \{1, \dots, r\}$ .

# Carmichael Numbers characterisation

Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number.

# Carmichael Numbers characterisation

Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

# Carmichael Numbers characterisation

Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group.

# Carmichael Numbers characterisation

Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ .

# Carmichael Numbers characterisation

Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ .

# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ ,

# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ , then  $p_i \mid n - 1$ , a contradiction.



# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ , then  $p_i \mid n - 1$ , a contradiction. Thus,  $e_i = 1$  for all  $i$ .

# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ , then  $p_i \mid n - 1$ , a contradiction. Thus,  $e_i = 1$  for all  $i$ .

Now, we must show that  $r \geq 3$ .

# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ , then  $p_i \mid n - 1$ , a contradiction. Thus,  $e_i = 1$  for all  $i$ .

Now, we must show that  $r \geq 3$ . For the sake of contradiction, assume that  $r = 2$ .

# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ , then  $p_i \mid n - 1$ , a contradiction. Thus,  $e_i = 1$  for all  $i$ .

Now, we must show that  $r \geq 3$ . For the sake of contradiction, assume that  $r = 2$ . In this case, we have  $n = p_1 p_2$  for some  $p_1 > p_2$ .

# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ , then  $p_i \mid n - 1$ , a contradiction. Thus,  $e_i = 1$  for all  $i$ .

Now, we must show that  $r \geq 3$ . For the sake of contradiction, assume that  $r = 2$ . In this case, we have  $n = p_1 p_2$  for some  $p_1 > p_2$ . We note that

$$n - 1 = p_1 p_2 - 1 = (p_1 - 1)p_2 + (p_2 - 1).$$

# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ , then  $p_i \mid n - 1$ , a contradiction. Thus,  $e_i = 1$  for all  $i$ .

Now, we must show that  $r \geq 3$ . For the sake of contradiction, assume that  $r = 2$ . In this case, we have  $n = p_1 p_2$  for some  $p_1 > p_2$ . We note that

$$n - 1 = p_1 p_2 - 1 = (p_1 - 1)p_2 + (p_2 - 1).$$

The above shows that  $p_1 - 1 \mid p_2 - 1$ ,

# Carmichael Numbers characterisation

## Proof.

Let  $n = p_1^{e_1} \cdots p_r^{e_r}$  be a Carmichael number. From earlier, we have

$$\mathbb{Z}_n^* \cong \mathbb{Z}_{p_1^{e_1}}^* \times \cdots \times \mathbb{Z}_{p_r^{e_r}}^*.$$

Since  $n - 1$  annihilates the left group, it annihilates the right group. Thus,

$$p_i^{e_i-1}(p_i - 1) \mid (n - 1)$$

for all  $i \in \{1, \dots, r\}$ . In particular,  $(p_i - 1) \mid (n - 1)$ . Moreover, if  $e_i > 1$  for some  $i$ , then  $p_i \mid n - 1$ , a contradiction. Thus,  $e_i = 1$  for all  $i$ .

Now, we must show that  $r \geq 3$ . For the sake of contradiction, assume that  $r = 2$ . In this case, we have  $n = p_1 p_2$  for some  $p_1 > p_2$ . We note that

$$n - 1 = p_1 p_2 - 1 = (p_1 - 1)p_2 + (p_2 - 1).$$

The above shows that  $p_1 - 1 \mid p_2 - 1$ , a contradiction since  $p_1 > p_2$ .

Proof (Continued).

Conversely, suppose  $n$  has the given form.



## Proof (Continued).

Conversely, suppose  $n$  has the given form. Let  $a$  be coprime to  $n$

## Proof (Continued).

Conversely, suppose  $n$  has the given form. Let  $a$  be coprime to  $n$  and hence, to each  $p_i$ .

## Proof (Continued).

Conversely, suppose  $n$  has the given form. Let  $a$  be coprime to  $n$  and hence, to each  $p_i$ . Then, by Fermat's Little Theorem, we have  $a^{p_i-1} \equiv 1 \pmod{p_i}$ .

# Carmichael Numbers characterisation

## Proof (Continued).

Conversely, suppose  $n$  has the given form. Let  $a$  be coprime to  $n$  and hence, to each  $p_i$ . Then, by Fermat's Little Theorem, we have  $a^{p_i-1} \equiv 1 \pmod{p_i}$ . Since  $n-1$  is a multiple of  $p_i-1$ , we get

## Proof (Continued).

Conversely, suppose  $n$  has the given form. Let  $a$  be coprime to  $n$  and hence, to each  $p_i$ . Then, by Fermat's Little Theorem, we have  $a^{p_i-1} \equiv 1 \pmod{p_i}$ . Since  $n-1$  is a multiple of  $p_i-1$ , we get

$$a^{n-1} \equiv 1 \pmod{p_i}$$

for all  $i \in \{1, \dots, r\}$ .

## Proof (Continued).

Conversely, suppose  $n$  has the given form. Let  $a$  be coprime to  $n$  and hence, to each  $p_i$ . Then, by Fermat's Little Theorem, we have  $a^{p_i-1} \equiv 1 \pmod{p_i}$ . Since  $n-1$  is a multiple of  $p_i-1$ , we get

$$a^{n-1} \equiv 1 \pmod{p_i}$$

for all  $i \in \{1, \dots, r\}$ . By the Chinese Remainder Theorem, we are now done. □

# Table of Contents

1 Introduction

2 First attempt

3 The Miller-Rabin test

# A Better Candidate

We now define a new set  $L'_n$  as follows.



# A Better Candidate

We now define a new set  $L'_n$  as follows.

## Definition 5

Let  $n - 1 = t2^h$

# A Better Candidate

We now define a new set  $L'_n$  as follows.

## Definition 5

Let  $n - 1 = t2^h$  where  $t$  is odd, and  $h \geq 1$ .

# A Better Candidate

We now define a new set  $L'_n$  as follows.

## Definition 5

Let  $n - 1 = t2^h$  where  $t$  is odd, and  $h \geq 1$ .

$$L'_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{t2^h} = 1 \text{ and}$$

# A Better Candidate

We now define a new set  $L'_n$  as follows.

## Definition 5

Let  $n - 1 = t2^h$  where  $t$  is odd, and  $h \geq 1$ .

$$L'_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{t2^h} = 1 \text{ and} \\ \alpha^{t2^{j+1}} = 1 \Rightarrow \alpha^{t2^j} = \pm 1 \text{ for } j = 0, \dots, h-1\}.$$

# A Better Candidate

We now define a new set  $L'_n$  as follows.

## Definition 5

Let  $n - 1 = t2^h$  where  $t$  is odd, and  $h \geq 1$ .

$$L'_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{t2^h} = 1 \text{ and} \\ \alpha^{t2^{j+1}} = 1 \Rightarrow \alpha^{t2^j} = \pm 1 \text{ for } j = 0, \dots, h-1\}.$$

The Miller-Rabin test uses this set  $L'_n$ .

# A Better Candidate

We now define a new set  $L'_n$  as follows.

## Definition 5

Let  $n - 1 = t2^h$  where  $t$  is odd, and  $h \geq 1$ .

$$L'_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{t2^h} = 1 \text{ and} \\ \alpha^{t2^{j+1}} = 1 \Rightarrow \alpha^{t2^j} = \pm 1 \text{ for } j = 0, \dots, h-1\}.$$

The Miller-Rabin test uses this set  $L'_n$ . By definition, it is clear that  $L'_n \subseteq L_n$ ,

# A Better Candidate

We now define a new set  $L'_n$  as follows.

## Definition 5

Let  $n - 1 = t2^h$  where  $t$  is odd, and  $h \geq 1$ .

$$L'_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{t2^h} = 1 \text{ and} \\ \alpha^{t2^{j+1}} = 1 \Rightarrow \alpha^{t2^j} = \pm 1 \text{ for } j = 0, \dots, h-1\}.$$

The Miller-Rabin test uses this set  $L'_n$ . By definition, it is clear that  $L'_n \subseteq L_n$ , since the **green condition** is the same from earlier.

# A Better Candidate

We now define a new set  $L'_n$  as follows.

## Definition 5

Let  $n - 1 = t2^h$  where  $t$  is odd, and  $h \geq 1$ .

$$L'_n := \{\alpha \in \mathbb{Z}_n^+ : \alpha^{t2^h} = 1 \text{ and} \\ \alpha^{t2^{j+1}} = 1 \Rightarrow \alpha^{t2^j} = \pm 1 \text{ for } j = 0, \dots, h-1\}.$$

The Miller-Rabin test uses this set  $L'_n$ . By definition, it is clear that  $L'_n \subseteq L_n$ , since the **green condition** is the same from earlier.

In fact,  $L'_n$  is precisely the set of those elements of  $L_n$  which also satisfy the **brown condition**.



# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

1  $\beta \leftarrow \alpha^t$

# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

- 1  $\beta \leftarrow \alpha^t$
- 2 if  $\beta = 1$  then return true

# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

- 1  $\beta \leftarrow \alpha^t$
- 2 if  $\beta = 1$  then return true
- 3 for  $j \leftarrow 0$  to  $h - 1$  do

# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

- 1  $\beta \leftarrow \alpha^t$
- 2 if  $\beta = 1$  then return true
- 3 for  $j \leftarrow 0$  to  $h - 1$  do
  - if  $\beta = -1$  then return true

# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

- ①  $\beta \leftarrow \alpha^t$
- ② if  $\beta = 1$  then return true
- ③ for  $j \leftarrow 0$  to  $h - 1$  do
  - if  $\beta = -1$  then return true
  - if  $\beta = 1$  then return false

# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

- ①  $\beta \leftarrow \alpha^t$
- ② if  $\beta = 1$  then return true
- ③ for  $j \leftarrow 0$  to  $h - 1$  do
  - if  $\beta = -1$  then return true
  - if  $\beta = 1$  then return false
  - $\beta \leftarrow \beta^2$

# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

- ①  $\beta \leftarrow \alpha^t$
- ② if  $\beta = 1$  then return true
- ③ for  $j \leftarrow 0$  to  $h - 1$  do
  - if  $\beta = -1$  then return true
  - if  $\beta = 1$  then return false
  - $\beta \leftarrow \beta^2$
- ④ return false



# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

- ①  $\beta \leftarrow \alpha^t$
- ② if  $\beta = 1$  then return true
- ③ for  $j \leftarrow 0$  to  $h - 1$  do
  - if  $\beta = -1$  then return true
  - if  $\beta = 1$  then return false
  - $\beta \leftarrow \beta^2$
- ④ return false

# Testing membership

Testing whether a given  $\alpha \in \mathbb{Z}_n^+$  belongs to  $L'_n$  can be done using the following algorithm:

## Algorithm (Testing membership)

- 1  $\beta \leftarrow \alpha^t$
- 2 if  $\beta = 1$  then return true
- 3 for  $j \leftarrow 0$  to  $h - 1$  do
  - if  $\beta = -1$  then return true
  - if  $\beta = 1$  then return false
  - $\beta \leftarrow \beta^2$
- 4 return false

This algorithm runs in time  $O(\text{poly}(\log(n)))$  and thus, satisfies the first criteria.

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ .

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties.

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties. This choice gives the Miller Rabin test.

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties. This choice gives the Miller Rabin test.

To put it all together, we have the test as:

## Algorithm (Miller Rabin)

- 1 input  $n$  and  $k$

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties. This choice gives the Miller Rabin test.

To put it all together, we have the test as:

## Algorithm (Miller Rabin)

- ① input  $n$  and  $k$
- ② for  $j \leftarrow 1$  to  $k$  do



# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties. This choice gives the Miller Rabin test.

To put it all together, we have the test as:

## Algorithm (Miller Rabin)

- ① input  $n$  and  $k$
- ② for  $j \leftarrow 1$  to  $k$  do
  - pick  $\alpha \in \mathbb{Z}_n^+$  randomly

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties. This choice gives the Miller Rabin test.

To put it all together, we have the test as:

## Algorithm (Miller Rabin)

- ① input  $n$  and  $k$
- ② for  $j \leftarrow 1$  to  $k$  do
  - pick  $\alpha \in \mathbb{Z}_n^+$  randomly
  - if  $\alpha \notin L'_n$  then return false

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties. This choice gives the Miller Rabin test.

To put it all together, we have the test as:

## Algorithm (Miller Rabin)

- ① input  $n$  and  $k$
- ② for  $j \leftarrow 1$  to  $k$  do
  - pick  $\alpha \in \mathbb{Z}_n^+$  randomly
  - if  $\alpha \notin L'_n$  then return false
- ③ return true

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties. This choice gives the Miller Rabin test.

To put it all together, we have the test as:

## Algorithm (Miller Rabin)

- ① input  $n$  and  $k$
- ② for  $j \leftarrow 1$  to  $k$  do
  - pick  $\alpha \in \mathbb{Z}_n^+$  randomly
  - if  $\alpha \notin L'_n$  then return false
- ③ return true

# Does it fit the bill?

## Theorem 6

If  $n$  is prime, then  $L'_n = \mathbb{Z}_n^*$ . If  $n$  is composite, then  $|L'_n| \leq \frac{1}{4}(n-1)$ .

Thus, this set  $L'_n$  does have the required properties. This choice gives the Miller Rabin test.

To put it all together, we have the test as:

## Algorithm (Miller Rabin)

- ① input  $n$  and  $k$
- ② for  $j \leftarrow 1$  to  $k$  do
  - pick  $\alpha \in \mathbb{Z}_n^+$  randomly
  - if  $\alpha \notin L'_n$  then return false
- ③ return true

Let us now prove the above theorem.

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory:

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ ,

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .



## Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

## Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ .

## Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ .

## Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map.

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map. Since  $\mathbb{Z}_n^*$  is cyclic, it follows that  $|L_n| = \gcd(\varphi(n), n-1)$ .

## Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map. Since  $\mathbb{Z}_n^*$  is cyclic, it follows that  $|L_n| = \gcd(\varphi(n), n-1)$ . We can explicitly calculate it to get



# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map. Since  $\mathbb{Z}_n^*$  is cyclic, it follows that  $|L_n| = \gcd(\varphi(n), n-1)$ . We can explicitly calculate it to get

$$|L'_n| \leq |L_n| = \gcd(\varphi(n), n-1)$$

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map. Since  $\mathbb{Z}_n^*$  is cyclic, it follows that  $|L_n| = \gcd(\varphi(n), n-1)$ . We can explicitly calculate it to get

$$|L'_n| \leq |L_n| = \gcd(\varphi(p^e), p^e - 1)$$

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map. Since  $\mathbb{Z}_n^*$  is cyclic, it follows that  $|L_n| = \gcd(\varphi(n), n-1)$ . We can explicitly calculate it to get

$$|L'_n| \leq |L_n| = \gcd(p^{e-1}(p-1), p^e - 1) =$$

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map. Since  $\mathbb{Z}_n^*$  is cyclic, it follows that  $|L_n| = \gcd(\varphi(n), n-1)$ . We can explicitly calculate it to get

$$|L'_n| \leq |L_n| = \gcd(p^{e-1}(p-1), p^e - 1) = p - 1$$

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map. Since  $\mathbb{Z}_n^*$  is cyclic, it follows that  $|L_n| = \gcd(\varphi(n), n-1)$ . We can explicitly calculate it to get

$$|L'_n| \leq |L_n| = \gcd(p^{e-1}(p-1), p^e - 1) = \frac{p^e - 1}{p^{e-1} + \dots + 1}$$

# Does it fit the bill? (Yes)

We shall frequently use the following result from group theory: If  $G$  has a cyclic group of order  $n$ , then there are exactly  $\gcd(n, m)$  many elements of  $G$  satisfying  $g^m = 1$ .

Proof.

**Case 1.**  $n$  is prime.

Note that we have  $L'_n \subseteq L_n = \mathbb{Z}_n^*$ . Thus, it suffices to prove that  $L_n \subseteq L'_n$ . But this follows because  $x^2 = 1 \Rightarrow x = \pm 1$  in a field.

**Case 2.**  $n = p^e$  for a prime  $p \geq 3$  and  $e \geq 2$ .

Recall that  $L_n$  is the kernel of the  $(n-1)$ -power map. Since  $\mathbb{Z}_n^*$  is cyclic, it follows that  $|L_n| = \gcd(\varphi(n), n-1)$ . We can explicitly calculate it to get

$$|L'_n| \leq |L_n| = \gcd(p^{e-1}(p-1), p^e - 1) = \frac{p^e - 1}{p^{e-1} + \dots + 1} \leq \frac{n-1}{4}.$$

# Does it fit the bill? (Yes)

Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

# Does it fit the bill? (Yes)

Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.



# Does it fit the bill? (Yes)

Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i 2^{h_i}$  in the usual way,

# Does it fit the bill? (Yes)

Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ ,

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ , and let  $j$  be the smallest index in  $g, \dots, h - 1$  such that  $\alpha^{t2^{j+1}} = 1$ .



# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ , and let  $j$  be the smallest index in  $g, \dots, h - 1$  such that  $\alpha^{t2^{j+1}} = 1$ . By **definition** of  $L'_n$ , we have  $\alpha^{t2^j} = -1$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ , and let  $j$  be the smallest index in  $g, \dots, h-1$  such that  $\alpha^{t2^{j+1}} = 1$ . By **definition** of  $L'_n$ , we have  $\alpha^{t2^j} = -1$ . Let  $i$  be such that  $g = h_i$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ , and let  $j$  be the smallest index in  $g, \dots, h - 1$  such that  $\alpha^{t2^{j+1}} = 1$ . By **definition** of  $L'_n$ , we have  $\alpha^{t2^j} = -1$ . Let  $i$  be such that  $g = h_i$ . Writing  $\theta(\alpha) = (\alpha_1, \dots, \alpha_r)$ ,

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ , and let  $j$  be the smallest index in  $g, \dots, h-1$  such that  $\alpha^{t2^{j+1}} = 1$ . By **definition** of  $L'_n$ , we have  $\alpha^{t2^j} = -1$ . Let  $i$  be such that  $g = h_i$ . Writing  $\theta(\alpha) = (\alpha_1, \dots, \alpha_r)$ , we have  $\alpha_i^{t2^j} = -1$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i 2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i 2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ , and let  $j$  be the smallest index in  $g, \dots, h - 1$  such that  $\alpha^{t2^{j+1}} = 1$ . By **definition** of  $L'_n$ , we have  $\alpha^{t2^j} = -1$ . Let  $i$  be such that  $g = h_i$ . Writing  $\theta(\alpha) = (\alpha_1, \dots, \alpha_r)$ , we have  $\alpha_i^{t2^j} = -1$ . Thus, the order of  $\alpha_i^t$  (in  $\mathbb{Z}_{p_i^{e_i}}^*$ ) is equal to  $2^{j+1}$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ , and let  $j$  be the smallest index in  $g, \dots, h-1$  such that  $\alpha^{t2^{j+1}} = 1$ . By **definition** of  $L'_n$ , we have  $\alpha^{t2^j} = -1$ . Let  $i$  be such that  $g = h_i$ . Writing  $\theta(\alpha) = (\alpha_1, \dots, \alpha_r)$ , we have  $\alpha_i^{t2^j} = -1$ . Thus, the order of  $\alpha_i^t$  (in  $\mathbb{Z}_{p_i^{e_i}}^*$ ) is equal to  $2^{j+1}$ . But this is a contradiction since  $2^{j+1}$  does not divide  $|\mathbb{Z}_{p_i^{e_i}}^*| = t_i2^{h_i}$ .

# Does it fit the bill? (Yes)

## Proof (continued).

**Case 3.**  $n = p_1^{e_1} \cdots p_r^{e_r}$  is the standard prime factorisation of  $n$ , with  $r > 1$ .

Let  $\theta : \mathbb{Z}_n \rightarrow \mathbb{Z}_{p_1^{e_1}} \times \cdots \times \mathbb{Z}_{p_r^{e_r}}$  be the ring isomorphism from earlier.

Write  $n - 1 = t2^h$  and  $\varphi(p_i^{e_i}) = t_i2^{h_i}$  in the usual way, and let  $g := \min\{h, h_1, \dots, h_r\}$ . Note that  $g \geq 1$ , and that each  $\mathbb{Z}_{p_i^{e_i}}^*$  is a cyclic group of order  $t_i2^{h_i}$ . Let  $\alpha \in L'_n$ .

We first show that  $\alpha^{t2^g} = 1$ . By **definition** of  $L'_n$ , we may assume  $g < h$ . Now, suppose  $\alpha^{t2^g} \neq 1$ , and let  $j$  be the smallest index in  $g, \dots, h-1$  such that  $\alpha^{t2^{j+1}} = 1$ . By **definition** of  $L'_n$ , we have  $\alpha^{t2^j} = -1$ . Let  $i$  be such that  $g = h_i$ . Writing  $\theta(\alpha) = (\alpha_1, \dots, \alpha_r)$ , we have  $\alpha_i^{t2^j} = -1$ . Thus, the order of  $\alpha_i^t$  (in  $\mathbb{Z}_{p_i^{e_i}}^*$ ) is equal to  $2^{j+1}$ . But this is a contradiction since  $2^{j+1}$  does not divide  $|\mathbb{Z}_{p_i^{e_i}}^*| = t_i2^{h_i}$ . ( $\because j \geq g = h_i$ )

# Does it fit the bill? (Yes)

Proof (continued).

For  $j = 0, \dots, h$ , define  $\rho_j$  to be the  $(t2^j)$ -power map on  $\mathbb{Z}_n^*$ .



# Does it fit the bill? (Yes)

## Proof (continued).

For  $j = 0, \dots, h$ , define  $\rho_j$  to be the  $(t2^j)$ -power map on  $\mathbb{Z}_n^*$ . From the previous claim, and the **definition** of  $L'_n$ , it follows that  $\alpha^{t2^{g-1}} = \pm 1$   $\forall \alpha \in L'_n$ .

# Does it fit the bill? (Yes)

## Proof (continued).

For  $j = 0, \dots, h$ , define  $\rho_j$  to be the  $(t2^j)$ -power map on  $\mathbb{Z}_n^*$ . From the previous claim, and the **definition** of  $L'_n$ , it follows that  $\alpha^{t2^{g-1}} = \pm 1$   $\forall \alpha \in L'_n$ . Thus,  $L'_n \subseteq \rho_{g-1}^{-1}(\{\pm 1\})$

# Does it fit the bill? (Yes)

## Proof (continued).

For  $j = 0, \dots, h$ , define  $\rho_j$  to be the  $(t2^j)$ -power map on  $\mathbb{Z}_n^*$ . From the previous claim, and the **definition** of  $L'_n$ , it follows that  $\alpha^{t2^{g-1}} = \pm 1$   $\forall \alpha \in L'_n$ . Thus,  $L'_n \subseteq \rho_{g-1}^{-1}(\{\pm 1\})$  and hence,  $|L'_n| \leq 2|\ker(\rho_{g-1})|$ .

# Does it fit the bill? (Yes)

## Proof (continued).

For  $j = 0, \dots, h$ , define  $\rho_j$  to be the  $(t2^j)$ -power map on  $\mathbb{Z}_n^*$ . From the previous claim, and the **definition** of  $L'_n$ , it follows that  $\alpha^{t2^{g-1}} = \pm 1$   $\forall \alpha \in L'_n$ . Thus,  $L'_n \subseteq \rho_{g-1}^{-1}(\{\pm 1\})$  and hence,  $|L'_n| \leq 2|\ker(\rho_{g-1})|$ . Also,

$$|\ker(\rho_j)| = \prod_{i=1}^r \gcd(t_i 2^{h_i}, t 2^j) \quad \forall j \in \{0, \dots, h\}.$$

# Does it fit the bill? (Yes)

## Proof (continued).

For  $j = 0, \dots, h$ , define  $\rho_j$  to be the  $(t2^j)$ -power map on  $\mathbb{Z}_n^*$ . From the previous claim, and the **definition** of  $L'_n$ , it follows that  $\alpha^{t2^{g-1}} = \pm 1$   $\forall \alpha \in L'_n$ . Thus,  $L'_n \subseteq \rho_{g-1}^{-1}(\{\pm 1\})$  and hence,  $|L'_n| \leq 2|\ker(\rho_{g-1})|$ . Also,

$$|\ker(\rho_j)| = \prod_{i=1}^r \gcd(t_i 2^{h_i}, t 2^j) \quad \forall j \in \{0, \dots, h\}.$$

Since  $g \leq h$  and  $g \leq h_i$  for all  $i$ , we get

# Does it fit the bill? (Yes)

## Proof (continued).

For  $j = 0, \dots, h$ , define  $\rho_j$  to be the  $(t2^j)$ -power map on  $\mathbb{Z}_n^*$ . From the previous claim, and the **definition** of  $L'_n$ , it follows that  $\alpha^{t2^{g-1}} = \pm 1$   $\forall \alpha \in L'_n$ . Thus,  $L'_n \subseteq \rho_{g-1}^{-1}(\{\pm 1\})$  and hence,  $|L'_n| \leq 2|\ker(\rho_{g-1})|$ . Also,

$$|\ker(\rho_j)| = \prod_{i=1}^r \gcd(t_i 2^{h_i}, t 2^j) \quad \forall j \in \{0, \dots, h\}.$$

Since  $g \leq h$  and  $g \leq h_i$  for all  $i$ , we get

$$2^r |\ker(\rho_{g-1})| = |\ker(\rho_g)| \leq |\ker(\rho_h)|.$$

# Does it fit the bill? (Yes)

## Proof (continued).

For  $j = 0, \dots, h$ , define  $\rho_j$  to be the  $(t2^j)$ -power map on  $\mathbb{Z}_n^*$ . From the previous claim, and the **definition** of  $L'_n$ , it follows that  $\alpha^{t2^{g-1}} = \pm 1$   $\forall \alpha \in L'_n$ . Thus,  $L'_n \subseteq \rho_{g-1}^{-1}(\{\pm 1\})$  and hence,  $|L'_n| \leq 2|\ker(\rho_{g-1})|$ . Also,

$$|\ker(\rho_j)| = \prod_{i=1}^r \gcd(t_i 2^{h_i}, t 2^j) \quad \forall j \in \{0, \dots, h\}.$$

Since  $g \leq h$  and  $g \leq h_i$  for all  $i$ , we get

$$2^r |\ker(\rho_{g-1})| = |\ker(\rho_g)| \leq |\ker(\rho_h)|.$$

Combining the **red** expressions, we get

$$|L'_n| \leq 2^{-r+1} |\ker(\rho_h)| = \frac{|L_n|}{2^{r-1}}.$$

# Does it fit the bill? (Yes)

Proof (continued).

$$|L'_n| \leq \frac{|L_n|}{2^{r-1}}$$



# Does it fit the bill? (Yes)

Proof (continued).

$$|L'_n| \leq \frac{|L_n|}{2^{r-1}}$$

If  $r \geq 3$ , then we are done

# Does it fit the bill? (Yes)

Proof (continued).

$$|L'_n| \leq \frac{|L_n|}{2^{r-1}}$$

If  $r \geq 3$ , then we are done since  $|L_n| \leq |Z_n^*| \leq n - 1$ , and  $2^{r-1} \geq 4$ .

# Does it fit the bill? (Yes)

Proof (continued).

$$|L'_n| \leq \frac{|L_n|}{2^{r-1}}$$

If  $r \geq 3$ , then we are done since  $|L_n| \leq |Z_n^*| \leq n - 1$ , and  $2^{r-1} \geq 4$ .

If  $r = 2$ , then  $n$  is not a Carmichael number and thus,

# Does it fit the bill? (Yes)

Proof (continued).

$$|L'_n| \leq \frac{|L_n|}{2^{r-1}}$$

If  $r \geq 3$ , then we are done since  $|L_n| \leq |Z_n^*| \leq n - 1$ , and  $2^{r-1} \geq 4$ .

If  $r = 2$ , then  $n$  is not a Carmichael number and thus,

$$\frac{|L_n|}{2^{r-1}} = \frac{|L_n|}{2}$$

# Does it fit the bill? (Yes)

Proof (continued).

$$|L'_n| \leq \frac{|L_n|}{2^{r-1}}$$

If  $r \geq 3$ , then we are done since  $|L_n| \leq |Z_n^*| \leq n - 1$ , and  $2^{r-1} \geq 4$ .

If  $r = 2$ , then  $n$  is not a Carmichael number and thus,

$$\frac{|L_n|}{2^{r-1}} = \frac{|L_n|}{2} \leq \frac{1}{4}(n - 1),$$

and we are again done. □