

# Extra Questions for MA 105

Aryaman Maithani

Semester: Autumn 2019

---

Notation:

$\mathbb{N} = \{1, 2, \dots\}$  denotes the set of natural numbers.

$\mathbb{Q}$  denotes the set of rational numbers.

$\mathbb{R}$  denotes the set of real numbers.

## WEEK 1

1. Let  $f$  be any bijection from  $\mathbb{N}$  to  $\mathbb{Q} \cap [0, 1]$ .  
Define the sequence  $(a_n)$  of real numbers as:  $a_n := f(n) \quad \forall n \in \mathbb{N}$ .  
Prove that  $(a_n)$  diverges or find an example of  $f$  such that  $(a_n)$  converges.
2. Let  $(a_n)$  be a sequence of real numbers. We say that  $(a_n)$  is *slack-convergent* if there is an  $a \in \mathbb{R}$  such that the following condition holds.  
For every  $\epsilon > 0$ , there is  $n_0 \in \mathbb{N}$  such that  $|a_n - a| \leq \epsilon$  for all  $n \geq n_0$ .  
Prove or disprove that a sequence is convergent (in the normal sense)  $\iff$  it is slack-convergent.  
  
(Additional) What happens if we change  $n \geq n_0$  to  $n > n_0$ ?
3. Let  $(a_n)$  be a sequence of real numbers. We say that  $(a_n)$  is *reciprocal-convergent* if there is an  $a \in \mathbb{R}$  such that the following condition holds.  
For every  $\epsilon > 0$ , there is  $n_0 \in \mathbb{N}$  such that  $|a_n - a| < 1/\epsilon$  for all  $n \geq n_0$ .  
Prove or disprove that a sequence is convergent (in the normal sense)  $\iff$  it is reciprocal-convergent.
4. Let  $(a_n)$  be a sequence of real numbers. We say that  $(a_n)$  is *natural-convergent* if the following condition holds.  
For every  $k \in \mathbb{N}$ ,  $\lim_{n \rightarrow \infty} |a_{n+k} - a_n| = 0$ .  
Prove or disprove that a sequence is convergent (in the normal sense)  $\iff$  it is natural-convergent.
5. Let  $(a_n)$  be a sequence of real numbers. We say that  $(a_n)$  is *weirdly-convergent* if there is an  $a \in \mathbb{R}$  such that the following condition holds.  
For every  $\epsilon > 0$ , there is  $n_0 \in \mathbb{N}$  such that  $|a_n - a| < \epsilon$  for infinitely many  $n \geq n_0$ .  
Prove or disprove that a sequence is convergent (in the normal sense)  $\iff$  it is weirdly-convergent.
6. Let  $(a_n)$  be a sequence of real numbers. We say that  $(a_n)$  is *reverse-convergent* if there is an  $a \in \mathbb{R}$  such that the following condition holds.  
For every  $n_0 \in \mathbb{N}$ , there is  $\epsilon > 0$  such that  $|a_n - a| < \epsilon$  for all  $n \geq n_0$ .  
Prove or disprove that a sequence is convergent (in the normal sense)  $\iff$  it is reverse-convergent.

For the question(s) in which the implication does not hold in both directions, does it hold in any? If yes, which?