

Calculus I Recap

Aryaman Maithani

<https://aryamanmaithani.github.io/tuts/ma-109>

IIT Bombay

Autumn Semester 2020-21

Start recording!

Definition 1 (Sequences)

Definition 2 (Convergence)

Definition 1 (Sequences)

A sequence in X

Definition 2 (Convergence)

Definition 1 (Sequences)

A sequence in X is a function $a : \mathbb{N} \rightarrow X$.

Definition 2 (Convergence)

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space.

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** .

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$.

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$,

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

for every $n > N$.

Definition 1 (Sequences)

A sequence **in** X is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in** X . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

for every $n > N$. L is said to be the *limit* of the sequence.

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

for every $n > N$. L is said to be the *limit* of the sequence.

In this case, we say that (a_n) converges

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

for every $n > N$. L is said to be the *limit* of the sequence.

In this case, we say that (a_n) converges **in X** .

Note the highlights.

Note the highlights. They are important.

Note the highlights. They are important. Consider $X = \mathbb{R}$

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$.

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier.

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$ and define $a_n = \frac{\lfloor 10^n \pi \rfloor}{10^n}$.

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$ and define $a_n = \frac{\lfloor 10^n \pi \rfloor}{10^n}$.

3.1, 3.14, 3.141, ...

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$ and define $a_n = \frac{\lfloor 10^n \pi \rfloor}{10^n}$.

$$3.1, 3.14, 3.141, \dots$$

The above is a sequence in \mathbb{Q} .

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$ and define $a_n = \frac{\lfloor 10^n \pi \rfloor}{10^n}$.

$$3.1, 3.14, 3.141, \dots$$

The above is a sequence in \mathbb{Q} . However, it does not converge in \mathbb{Q} .

Definition 3 (Cauchy Sequences)

Definition 3 (Cauchy Sequences)

Let X be a space.

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X .

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy*

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$,

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Proposition 1 (Convergence \implies Cauchy)

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Proposition 1 (Convergence \implies Cauchy)

If (a_n) is a convergent sequence

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Proposition 1 (Convergence \implies Cauchy)

If (a_n) is a convergent sequence in any space X ,

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Proposition 1 (Convergence \implies Cauchy)

If (a_n) is a convergent sequence in any space X , then (a_n) is Cauchy.

Definition 4 (Completeness)

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete*

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in** X .

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in** X .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in** X .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in** X .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples:

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in** X .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples: We saw some examples earlier.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in** X .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples: We saw some examples earlier. Go back and see that \mathbb{Q} and $(0, 1]$ are **not** complete.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in** X .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples: We saw some examples earlier. Go back and see that \mathbb{Q} and $(0, 1]$ are **not** complete.

Exercise: Show that \mathbb{N}, \mathbb{Z} are complete.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in** X .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples: We saw some examples earlier. Go back and see that \mathbb{Q} and $(0, 1]$ are **not** complete.

Exercise: Show that \mathbb{N}, \mathbb{Z} are complete. (What property do you really need? Can you generalise this?)

Now, we digress a bit to see what \mathbb{R} and completeness really means.

It is okay if you don't understand every single thing. It is more or less for you to know “okay, whatever we say works” even if you don't know the exact details why.

What is \mathbb{R} ?

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive,

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric,

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive,

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared*.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared*.

\mathbb{R} is an ordered field.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared*.

\mathbb{R} is an ordered field. All this means is that there is an order which is actually compatible with $+$ and \cdot .

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared*.

\mathbb{R} is an ordered field. All this means is that there is an order which is actually compatible with $+$ and \cdot . What does this mean?

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared*.

\mathbb{R} is an ordered field. All this means is that there is an order which is actually compatible with $+$ and \cdot . What does this mean?

$$x < y \implies x + z < y + z \text{ for all } x, y, z \in \mathbb{R},$$

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared*.

\mathbb{R} is an ordered field. All this means is that there is an order which is actually compatible with $+$ and \cdot . What does this mean?

$$x < y \implies x + z < y + z \text{ for all } x, y, z \in \mathbb{R},$$

$$x < y \implies x \cdot z < y \cdot z \text{ for all } x, y \in \mathbb{R} \text{ and } z \in \mathbb{R}_{>0}.$$

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual.

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R}

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R} which is bounded above

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R} which is bounded above has a least upper bound.

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R} which is bounded above has a least upper bound.

The least upper bound is called *supremum*.

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R} which is bounded above has a least upper bound.

The least upper bound is called *supremum*.

Note that **neither** of the above grey boxes is true if we replace \mathbb{R} by \mathbb{Q} .

What one must really ask at this point is:

What one must really ask at this point is: how do we know that \mathbb{R} exists?

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R}

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot and binary relation $<$

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot and binary relation $<$ which satisfies all the listed properties?

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot and binary relation $<$ which satisfies all the listed properties?

That is what I refer to as a non-trivial part.

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot and binary relation $<$ which satisfies all the listed properties?

That is what I refer to as a non-trivial part. It can be done but is not useful to us at the moment.

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing*

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing* if

$$a_{n+1} \geq a_n$$

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing* if

$$a_{n+1} \geq a_n$$

for all $n \in \mathbb{N}$.

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing* if

$$a_{n+1} \geq a_n$$

for all $n \in \mathbb{N}$.

Similarly, one defines a monotonically decreasing sequence.

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing* if

$$a_{n+1} \geq a_n$$

for all $n \in \mathbb{N}$.

Similarly, one defines a monotonically decreasing sequence. A sequence is said to be monotonic if it is either monotonically increasing or monotonically decreasing.

Definition 6 (Eventually monotonically increasing sequences)

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing*

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Theorem 2

An eventually monotonic sequence **in** \mathbb{R} which is bounded converges **in** \mathbb{R} .

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Theorem 2

An eventually monotonic sequence in \mathbb{R} which is bounded converges in \mathbb{R} .

Again, the above is not true if we take \mathbb{Q} instead of \mathbb{R} .

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Theorem 2

An eventually monotonic sequence in \mathbb{R} which is bounded converges in \mathbb{R} .

Again, the above is not true if we take \mathbb{Q} instead of \mathbb{R} . The π sequence shows this.

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Theorem 2

An eventually monotonic sequence in \mathbb{R} which is bounded converges in \mathbb{R} .

Again, the above is not true if we take \mathbb{Q} instead of \mathbb{R} . The π sequence shows this. In fact, the above is really a consequence of completeness.

We also saw series in the lectures. There's nothing much to be said about it.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.)

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205. Those taking a minor in Mathematics will also come across it in MA 403.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205. Those taking a minor in Mathematics will also come across it in MA 403. Of course, the ones in the Mathematics department will also see it in various courses.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205. Those taking a minor in Mathematics will also come across it in MA 403. Of course, the ones in the Mathematics department will also see it in various courses.

For us, all we need to know is that convergence of a series is just the convergence of the sequence of its *partial sums*.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205. Those taking a minor in Mathematics will also come across it in MA 403. Of course, the ones in the Mathematics department will also see it in various courses.

For us, all we need to know is that convergence of a series is just the convergence of the sequence of its *partial sums*. Thus, we are back in the case where we study sequences!

We then moved on to the definition of limits of functions defined on intervals.

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$.

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function.

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $l \in \mathbb{R}$.

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $l \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = l$$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $l \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = l$$

if for every $\epsilon > 0$,

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $l \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = l$$

if for every $\epsilon > 0$, there exists $\delta > 0$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $l \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = l$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $l \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = l$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - l| < \epsilon$$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $l \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = l$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - l| < \epsilon$$

for all $x \in (a, b)$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $l \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = l$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - l| < \epsilon$$

for all $x \in (a, b)$ such that $0 < |x - x_0| < \delta$.

Note in the above that we can still talk about limits at points at which the function is *not* defined.

Note in the above that we can still talk about limits at points at which the function is *not* defined.

If the thing in the previous slide does happen,

Note in the above that we can still talk about limits at points at which the function is *not* defined.

If the thing in the previous slide does happen, then we say that $f(x)$ tends to l as x tends to x_0 .

Note in the above that we can still talk about limits at points at which the function is *not* defined.

If the thing in the previous slide does happen, then we say that $f(x)$ tends to l as x tends to x_0 . Or that f has a limit l at x_0 .

Note in the above that we can still talk about limits at points at which the function is *not* defined.

If the thing in the previous slide does happen, then we say that $f(x)$ tends to l as x tends to x_0 . Or that f has a limit l at x_0 .

If no such l exists, then we say that f does not have any limit at x_0 .

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above.

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $l \in \mathbb{R}$.

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $l \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = l$$

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $l \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = l$$

if for every $\epsilon > 0$,

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $l \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = l$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $l \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = l$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

$$|f(x) - l| < \epsilon$$

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $l \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = l$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

$$|f(x) - l| < \epsilon$$

for all $x \in A$

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $l \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = l$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

$$|f(x) - l| < \epsilon$$

for all $x \in A$ such that $x > X$.

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $l \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = l$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

$$|f(x) - l| < \epsilon$$

for all $x \in A$ such that $x > X$.

Similarly, we have the limit at $-\infty$.

Stop recording. Start a new one.
Take doubts.

Start recording!

Last week, we had ended on limits. Today, we

Last week, we had ended on limits. Today, we *continue*

Last week, we had ended on limits. Today, we *continue* with

Last week, we had ended on limits. Today, we *continue* with *continuity*.

Last week, we had ended on limits. Today, we *continue* with *continuity*. Ba-dum-tss.

Last week, we had ended on limits. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Last week, we had ended on limits. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function

Last week, we had ended on limits. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$,

Last week, we had ended on limits. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be

Last week, we had ended on limits. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be *continuous at the point c*

Last week, we had ended on limits. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be *continuous at the point c* if (and only if)

$$\lim_{x \rightarrow c} f(x) = f(c).$$

Last week, we had ended on limits. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be *continuous at the point c* if (and only if)

$$\lim_{x \rightarrow c} f(x) = f(c).$$

We simply say “ f is continuous” if it is continuous at every point in the domain.

Last week, we had ended on limits. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be *continuous at the point c* if (and only if)

$$\lim_{x \rightarrow c} f(x) = f(c).$$

We simply say “ f is continuous” if it is continuous at every point in the domain. If f is not continuous at a point c *in the domain*, then we say that f is discontinuous at c .

We have the usual rules which tell us that
sum/product/composition of continuous functions is continuous.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c .

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. Let $u \in \mathbb{R}$ be between $f(a)$ and $f(b)$.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. Let $u \in \mathbb{R}$ be between $f(a)$ and $f(b)$. Then, there exists $c \in [a, b]$

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. Let $u \in \mathbb{R}$ be between $f(a)$ and $f(b)$. Then, there exists $c \in [a, b]$ such that $f(c) = u$.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. Let $u \in \mathbb{R}$ be between $f(a)$ and $f(b)$. Then, there exists $c \in [a, b]$ such that $f(c) = u$.

Note carefully that the domain is an interval.

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous.

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

for all $x \in [a, b]$.

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

for all $x \in [a, b]$.

Note very carefully that the above not only shows that the image of f is bounded

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

for all $x \in [a, b]$.

Note very carefully that the above not only shows that the image of f is bounded but also that the bounds are attained!

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

for all $x \in [a, b]$.

Note very carefully that the above not only shows that the image of f is bounded but also that the bounds are attained! Note that the domain was a closed and bounded interval.

Recall that a (non-empty) set which is bounded above can have many upper bounds.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$,

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above,

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above, then the supremum of the image is called the supremum of f on X .

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above, then the supremum of the image is called the supremum of f on X .

Analogous comments hold for infimum.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above, then the supremum of the image is called the supremum of f on X .

Analogous comments hold for infimum.

Thus, what the previous theorem told us was that not only is the image bounded but the supremum and infimum are actually attained.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above, then the supremum of the image is called the supremum of f on X .

Analogous comments hold for infimum.

Thus, what the previous theorem told us was that not only is the image bounded but the supremum and infimum are actually attained. (If the function is continuous and defined on a close and bounded interval, that is.)

Non-examples of the previous theorem:

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Consider $f : (0, 1) \rightarrow \mathbb{R}$

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = \frac{1}{x}.$$

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = \frac{1}{x}.$$

The image is not bounded above.

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = \frac{1}{x}.$$

The image is not bounded above. It is bounded below but the infimum is not attained.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$,

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$, we have $f(a_n) \rightarrow f(a)$.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$, we have $f(a_n) \rightarrow f(a)$.

This makes life simpler because it is sometimes easier to deal with sequences.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$, we have $f(a_n) \rightarrow f(a)$.

This makes life simpler because it is sometimes easier to deal with sequences. We had seen an example of this when we proved that a certain oscillatory function does not have a limit.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$, we have $f(a_n) \rightarrow f(a)$.

This makes life simpler because it is sometimes easier to deal with sequences. We had seen an example of this when we proved that a certain oscillatory function does not have a limit. Do you remember which?

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$.

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c*

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c* if the following limit exists:

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c* if the following limit exists:

$$\lim_{h \rightarrow 0} \frac{f(c + h) - f(c)}{h}.$$

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c* if the following limit exists:

$$\lim_{h \rightarrow 0} \frac{f(c + h) - f(c)}{h}.$$

In such a case, we call the value of the above limit the derivative of f at c

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c* if the following limit exists:

$$\lim_{h \rightarrow 0} \frac{f(c+h) - f(c)}{h}.$$

In such a case, we call the value of the above limit the derivative of f at c and denote it by $f'(c)$.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather

$$(fg)'(c) = f'(c)g(c) + f(c)g'(c).$$

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 .

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$,

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$, then we say that f has a *local maximum* at x_0 .

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$, then we say that f has a *local maximum* at x_0 .

Of course, we have an analogous definition for minimum.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$, then we say that f has a *local maximum* at x_0 .

Of course, we have an analogous definition for minimum. Note that here, we have that x_0 is an “interior point.”

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$, then we say that f has a *local maximum* at x_0 .

Of course, we have an analogous definition for minimum. Note that here, we have that x_0 is an “interior point.” That is, there is an interval *around* x_0 contained within the domain.

Theorem 5 (Fermat's Theorem)

Theorem 5 (Fermat's Theorem)

If $f : X \rightarrow \mathbb{R}$ is differentiable

Theorem 5 (Fermat's Theorem)

If $f : X \rightarrow \mathbb{R}$ is differentiable and has a local minimum or maximum at a point $x_0 \in X$,

Theorem 5 (Fermat's Theorem)

If $f : X \rightarrow \mathbb{R}$ is differentiable and has a local minimum or maximum at a point $x_0 \in X$, then $f'(x_0) = 0$.

Theorem 5 (Fermat's Theorem)

If $f : X \rightarrow \mathbb{R}$ is differentiable and has a local minimum or maximum at a point $x_0 \in X$, then $f'(x_0) = 0$.

Once again, note that this only talks about “interior points.”

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function.

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) .

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$,

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

Using the above, we have a more general result.

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

Using the above, we have a more general result.

Theorem 7 (Mean Value Theorem)

Let f be continuous and differentiable as above.

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

Using the above, we have a more general result.

Theorem 7 (Mean Value Theorem)

Let f be continuous and differentiable as above. There exists $c \in (a, b)$

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

Using the above, we have a more general result.

Theorem 7 (Mean Value Theorem)

Let f be continuous and differentiable as above. There exists $c \in (a, b)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function.

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) .

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$.

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

$$f'(x_0) = u.$$

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

$$f'(x_0) = u.$$

Note that the derivative of a (differentiable) function need not be continuous.

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

$$f'(x_0) = u.$$

Note that the derivative of a (differentiable) function need not be continuous. We shall see an example in the tutorial today, in fact.

We then saw a theorem which said “derivatives have IVP.” TO be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

$$f'(x_0) = u.$$

Note that the derivative of a (differentiable) function need not be continuous. We shall see an example in the tutorial today, in fact. However, the above theorem tells us how the derivative can't have “jump” discontinuity.

Stop recording. Start a new one.
Take doubts.