

Calculus I Recap

Aryaman Maithani

<https://aryamanmaithani.github.io/tuts/ma-109>

IIT Bombay

Autumn Semester **2020-21**

Week 1

Start recording!

Definition 1 (Sequences)

Definition 2 (Convergence)

Definition 1 (Sequences)

A sequence in X

Definition 2 (Convergence)

Definition 1 (Sequences)

A sequence in X is a function $a : \mathbb{N} \rightarrow X$.

Definition 2 (Convergence)

Definition 1 (Sequences)

A sequence in X is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Definition 1 (Sequences)

A sequence in X is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space.

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** .

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$.

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

Definition 1 (Sequences)

A sequence in X is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence in X . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$,

Definition 1 (Sequences)

A sequence in X is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence in X . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

for every $n > N$.

Definition 1 (Sequences)

A sequence in X is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence in X . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

for every $n > N$. L is said to be the *limit* of the sequence.

Definition 1 (Sequences)

A sequence in X is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence in X . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

for every $n > N$. L is said to be the *limit* of the sequence.

In this case, we say that (a_n) converges

Definition 1 (Sequences)

A sequence **in X** is a function $a : \mathbb{N} \rightarrow X$. We usually write a_n instead of $a(n)$.

Definition 2 (Convergence)

Let X be a space. Let (a_n) be a sequence **in X** . Let $L \in X$. We write

$$\lim_{n \rightarrow \infty} a_n = L$$

if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - L| < \epsilon$$

for every $n > N$. L is said to be the *limit* of the sequence.

In this case, we say that (a_n) converges **in X** .

Week 1

Note the highlights.

Week 1

Note the highlights. They are important.

Week 1

Note the highlights. They are important. Consider $X = \mathbb{R}$

Week 1

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

Week 1

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.
As we saw in class, (a_n) converges to $0 \in \mathbb{R}$.

Week 1

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier.

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$ and define $a_n = \frac{\lfloor 10^n \pi \rfloor}{10^n}$.

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$ and define $a_n = \frac{\lfloor 10^n \pi \rfloor}{10^n}$.

$$3.1, 3.14, 3.141, \dots$$

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$ and define $a_n = \frac{\lfloor 10^n \pi \rfloor}{10^n}$.

$$3.1, 3.14, 3.141, \dots$$

The above is a sequence in \mathbb{Q} .

Note the highlights. They are important. Consider $X = \mathbb{R}$ and the sequence $a_n := 1/n$.

As we saw in class, (a_n) converges to $0 \in \mathbb{R}$. Thus, (a_n) converges in \mathbb{R} .

However, consider $X = (0, 1]$ and (a_n) be as earlier. This sequence does not converge (in X) anymore.

Similarly, consider $X = \mathbb{Q}$ and define $a_n = \frac{\lfloor 10^n \pi \rfloor}{10^n}$.

$$3.1, 3.14, 3.141, \dots$$

The above is a sequence in \mathbb{Q} . However, it does not converge in \mathbb{Q} .

Definition 3 (Cauchy Sequences)

Definition 3 (Cauchy Sequences)

Let X be a space.

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X .

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy*

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$,

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be **Cauchy** if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Proposition 1 (Convergence \implies Cauchy)

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Proposition 1 (Convergence \implies Cauchy)

If (a_n) is a convergent sequence

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be *Cauchy* if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Proposition 1 (Convergence \implies Cauchy)

If (a_n) is a convergent sequence in any space X ,

Definition 3 (Cauchy Sequences)

Let X be a space. Let (a_n) be a sequence in X . (a_n) is said to be **Cauchy** if for every $\epsilon > 0$, there exists $N \in \mathbb{N}$ such that

$$|a_n - a_m| < \epsilon$$

for all $n, m > N$.

Proposition 1 (Convergence \implies Cauchy)

If (a_n) is a convergent sequence in any space X , then (a_n) is Cauchy.

Definition 4 (Completeness)

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete*

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples:

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples: We saw some examples earlier.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples: We saw some examples earlier. Go back and see that \mathbb{Q} and $(0, 1]$ are **not** complete.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples: We saw some examples earlier. Go back and see that \mathbb{Q} and $(0, 1]$ are **not** complete.

Exercise: Show that \mathbb{N}, \mathbb{Z} are complete.

Definition 4 (Completeness)

A space X is said to be *complete* if every Cauchy sequence in X converges **in X** .

Theorem 1 (\mathbb{R} is complete)

\mathbb{R} is complete.

This theorem is trivial and not trivial at the same time. You don't know what \mathbb{R} *truly* is. So you can't really prove this.

Non-examples: We saw some examples earlier. Go back and see that \mathbb{Q} and $(0, 1]$ are **not** complete.

Exercise: Show that \mathbb{N}, \mathbb{Z} are complete. (What property do you really need? Can you generalise this?)

Now, we digress a bit to see what \mathbb{R} and completeness really means.

It is okay if you don't understand every single thing. It is more or less for you to know "okay, whatever we say works" even if you don't know the exact details why.

What is \mathbb{R} ?

Week 1

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive,

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric,

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive,

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared.*

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared.*

\mathbb{R} is an ordered field.

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared.*

\mathbb{R} is an ordered field. All this means is that there is an order which is actually compatible with $+$ and \cdot .

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared.*

\mathbb{R} is an ordered field. All this means is that there is an order which is actually compatible with $+$ and \cdot . What does this mean?

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared.*

\mathbb{R} is an ordered field. All this means is that there is an order which is actually compatible with $+$ and \cdot . What does this mean?

$$x < y \implies x + z < y + z \text{ for all } x, y, z \in \mathbb{R},$$

What is \mathbb{R} ? Well, all one really needs is to know the following two slides about \mathbb{R} .

\mathbb{R} is a field. This means that the familiar properties of addition/multiplication are true. (Commutativity, associativity, existence of identity, inverses, and distributivity.)

\mathbb{R} is ordered. There is a binary operation \leq on \mathbb{R} which is reflexive, anti-symmetric, transitive, *and any two elements can be compared.*

\mathbb{R} is an ordered field. All this means is that there is an order which is actually compatible with $+$ and \cdot . What does this mean?

$$x < y \implies x + z < y + z \text{ for all } x, y, z \in \mathbb{R},$$

$$x < y \implies x \cdot z < y \cdot z \text{ for all } x, y \in \mathbb{R} \text{ and } z \in \mathbb{R}_{>0}.$$

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual.

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R}

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R} which is bounded above

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R} which is bounded above has a least upper bound.

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R} which is bounded above has a least upper bound.

The least upper bound is called *supremum*.

Note that all the properties earlier are also satisfied by \mathbb{Q} . Here's what sets \mathbb{R} apart:

\mathbb{R} is complete.

There's another way of defining completeness of \mathbb{R} , which coincides with the usual. It is the following:

Every non-empty subset of \mathbb{R} which is bounded above has a least upper bound.

The least upper bound is called *supremum*.

Note that **neither** of the above grey boxes is true if we replace \mathbb{R} by \mathbb{Q} .

What one must really ask at this point is:

What one must really ask at this point is: how do we know that \mathbb{R} exists?

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R}

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot .

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot and binary relation $<$

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot and binary relation $<$ which satisfies all the listed properties?

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot and binary relation $<$ which satisfies all the listed properties?

That is what I refer to as a non-trivial part.

What one must really ask at this point is: how do we know that \mathbb{R} exists?

That is, how do we know that there is some set \mathbb{R} with some operations $+$, \cdot and binary relation $<$ which satisfies all the listed properties?

That is what I refer to as a non-trivial part. It can be done but is not useful to us at the moment.

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing*

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing* if

$$a_{n+1} \geq a_n$$

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing* if

$$a_{n+1} \geq a_n$$

for all $n \in \mathbb{N}$.

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing* if

$$a_{n+1} \geq a_n$$

for all $n \in \mathbb{N}$.

Similarly, one defines a monotonically decreasing sequence.

Back to sequences now.

Definition 5 (Monotonically increasing sequences)

A sequence (a_n) is said to be *monotonically increasing* if

$$a_{n+1} \geq a_n$$

for all $n \in \mathbb{N}$.

Similarly, one defines a monotonically decreasing sequence. A sequence is said to be monotonic if it is either monotonically increasing or monotonically decreasing.

Definition 6 (Eventually monotonically increasing sequences)

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing*

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Theorem 2

An eventually monotonic sequence in \mathbb{R} which is bounded converges in \mathbb{R} .

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Theorem 2

An eventually monotonic sequence in \mathbb{R} which is bounded converges in \mathbb{R} .

Again, the above is not true if we take \mathbb{Q} instead of \mathbb{R} .

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Theorem 2

An eventually monotonic sequence in \mathbb{R} which is bounded converges in \mathbb{R} .

Again, the above is not true if we take \mathbb{Q} instead of \mathbb{R} . The π sequence shows this.

Definition 6 (Eventually monotonically increasing sequences)

A sequence (a_n) is said to be *eventually monotonically increasing* if there exists $N \in \mathbb{N}$ such that

$$a_{n+1} \geq a_n$$

for all $n \geq N$.

As earlier, we can define eventually monotonically decreasing sequences and simply, eventually monotonic sequences.

Theorem 2

An eventually monotonic sequence in \mathbb{R} which is bounded converges in \mathbb{R} .

Again, the above is not true if we take \mathbb{Q} instead of \mathbb{R} . The π sequence shows this. In fact, the above is really a consequence of completeness.

Week 1

We also saw series in the lectures. There's nothing much to be said about it.

Week 1

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.)

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205. Those taking a minor in Mathematics will also come across it in MA 403.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205. Those taking a minor in Mathematics will also come across it in MA 403. Of course, the ones in the Mathematics department will also see it in various courses.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205. Those taking a minor in Mathematics will also come across it in MA 403. Of course, the ones in the Mathematics department will also see it in various courses.

For us, all we need to know is that convergence of a series is just the convergence of the sequence of its *partial sums*.

We also saw series in the lectures. There's nothing much to be said about it. (As far as this course is concerned.) In reality, there is a lot more to be said about series and various tests for seeing if a series converges. Some of you will see this in future courses like MA 205. Those taking a minor in Mathematics will also come across it in MA 403. Of course, the ones in the Mathematics department will also see it in various courses.

For us, all we need to know is that convergence of a series is just the convergence of the sequence of its *partial sums*. Thus, we are back in the case where we study sequences!

Week 1

We then moved on to the definition of limits of functions defined on intervals.

Week 1

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$.

Week 1

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function.

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $L \in \mathbb{R}$.

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $L \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = L$$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $L \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = L$$

if for every $\epsilon > 0$,

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $L \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $L \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $L \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - L| < \epsilon$$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $L \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - L| < \epsilon$$

for all $x \in (a, b)$

We then moved on to the definition of limits of functions defined on intervals.

For the remainder, we fix $a, b \in \mathbb{R}$ such that $a < b$. (Just to recall, ∞ is not an element of \mathbb{R} .)

Definition 7 (Limit)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function. Let $x_0 \in [a, b]$ and $L \in \mathbb{R}$. Then, we write

$$\lim_{x \rightarrow x_0} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|f(x) - L| < \epsilon$$

for all $x \in (a, b)$ such that $0 < |x - x_0| < \delta$.

Note in the above that we can still talk about limits at points at which the function is *not* defined.

Note in the above that we can still talk about limits at points at which the function is *not* defined.

If the thing in the previous slide does happen,

Note in the above that we can still talk about limits at points at which the function is *not* defined.

If the thing in the previous slide does happen, then we say that $f(x)$ tends to l as x tends to x_0 .

Note in the above that we can still talk about limits at points at which the function is *not* defined.

If the thing in the previous slide does happen, then we say that $f(x)$ tends to l as x tends to x_0 . Or that f has a limit l at x_0 .

Note in the above that we can still talk about limits at points at which the function is *not* defined.

If the thing in the previous slide does happen, then we say that $f(x)$ tends to l as x tends to x_0 . Or that f has a limit l at x_0 .

If no such l exists, then we say that f does not have any limit at x_0 .

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above.

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$.

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = L$$

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = L$$

if for every $\epsilon > 0$,

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = L$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = L$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

$$|f(x) - L| < \epsilon$$

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = L$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

$$|f(x) - L| < \epsilon$$

for all $x \in A$

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = L$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

$$|f(x) - L| < \epsilon$$

for all $x \in A$ such that $x > X$.

We then also defined limit at $\pm\infty$.

Definition 8 (Limit at ∞)

Let $A \subset \mathbb{R}$ be a set which is not bounded above. Let $f : A \rightarrow \mathbb{R}$ be a function and let $L \in \mathbb{R}$. We say

$$\lim_{x \rightarrow \infty} f(x) = L$$

if for every $\epsilon > 0$, there exists $X \in \mathbb{R}$ such that

$$|f(x) - L| < \epsilon$$

for all $x \in A$ such that $x > X$.

Similarly, we have the limit at $-\infty$.

Stop recording. Start a new one.
Take doubts.

Start recording!

Week 2

Last week, we had *limited* ourselves to *limits*. Today, we

Week 2

Last week, we had *limited* ourselves to *limits*. Today, we *continue*

Week 2

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with

Week 2

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

Week 2

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*. Ba-dum-tss.

Week 2

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$,

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be *continuous at the point c*

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be *continuous at the point c* if (and only if)

$$\lim_{x \rightarrow c} f(x) = f(c).$$

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be *continuous at the point c* if (and only if)

$$\lim_{x \rightarrow c} f(x) = f(c).$$

We simply say “ f is continuous” if it is continuous at every point in the domain.

Last week, we had *limited* ourselves to *limits*. Today, we *continue* with *continuity*.

This is quite simple, using whatever we've already seen.

Definition 9 (Continuity)

If $f : [a, b] \rightarrow \mathbb{R}$ is a function and $c \in [a, b]$, then f is said to be *continuous at the point c* if (and only if)

$$\lim_{x \rightarrow c} f(x) = f(c).$$

We simply say “ f is continuous” if it is continuous at every point in the domain. If f is not continuous at a point c *in the domain*, then we say that f is discontinuous at c .

We have the usual rules which tell us that
sum/product/composition of continuous functions is continuous.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c .

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. Let $u \in \mathbb{R}$ be between $f(a)$ and $f(b)$.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. Let $u \in \mathbb{R}$ be between $f(a)$ and $f(b)$. Then, there exists $c \in [a, b]$

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. Let $u \in \mathbb{R}$ be between $f(a)$ and $f(b)$. Then, there exists $c \in [a, b]$ such that $f(c) = u$.

We have the usual rules which tell us that sum/product/composition of continuous functions is continuous. If f is continuous at c and $f(c) \neq 0$, then $1/f$ is continuous at c . We had also seen that the square root function is continuous. We now state an important property of continuous functions.

Definition 10 (Intermediate Value Property)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a continuous function. Let $u \in \mathbb{R}$ be between $f(a)$ and $f(b)$. Then, there exists $c \in [a, b]$ such that $f(c) = u$.

Note carefully that the domain is an interval.

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous.

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

for all $x \in [a, b]$.

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

for all $x \in [a, b]$.

Note very carefully that the above not only shows that the image of f is bounded

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

for all $x \in [a, b]$.

Note very carefully that the above not only shows that the image of f is bounded but also that the bounds are attained!

Now, we state another property, called the extreme value theorem.

Theorem 3 (Extreme value theorem)

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, there exist $x_1, x_2 \in [a, b]$ such that

$$f(x_1) \leq f(x) \leq f(x_2)$$

for all $x \in [a, b]$.

Note very carefully that the above not only shows that the image of f is bounded but also that the bounds are attained! Note that the domain was a closed and bounded interval.

Recall that a (non-empty) set which is bounded above can have many upper bounds.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*. Similarly, we had defined *infimum*.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$,

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above,

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above, then the supremum of the image is called the supremum of f on X .

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above, then the supremum of the image is called the supremum of f on X .

Analogous comments hold for infimum.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above, then the supremum of the image is called the supremum of f on X .

Analogous comments hold for infimum.

Thus, what the previous theorem told us was that not only is the image bounded but the supremum and infimum are actually attained.

Recall that a (non-empty) set which is bounded above can have many upper bounds. However, completeness of \mathbb{R} tells us that there is a *least* upper bound. We had called this the *supremum*.

Similarly, we had defined *infimum*.

By abuse of notation, given a function $f : X \rightarrow \mathbb{R}$, if the image $f(X) \subset \mathbb{R}$ is bounded above, then the supremum of the image is called the supremum of f on X .

Analogous comments hold for infimum.

Thus, what the previous theorem told us was that not only is the image bounded but the supremum and infimum are actually attained. (If the function is continuous and defined on a closed and bounded interval, that is.)

Non-examples of the previous theorem:

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Consider $f : (0, 1) \rightarrow \mathbb{R}$

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = \frac{1}{x}.$$

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = \frac{1}{x}.$$

The image is not bounded above.

Non-examples of the previous theorem:

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = x.$$

The image is bounded but the infimum/supremum are not attained.

Consider $f : (0, 1) \rightarrow \mathbb{R}$ defined by

$$f(x) = \frac{1}{x}.$$

The image is not bounded above. It is bounded below but the infimum is not attained.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$,

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$, we have $f(a_n) \rightarrow f(a)$.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$, we have $f(a_n) \rightarrow f(a)$.

This makes life simpler because it is sometimes easier to deal with sequences.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$, we have $f(a_n) \rightarrow f(a)$.

This makes life simpler because it is sometimes easier to deal with sequences. We had seen an example of this when we proved that a certain oscillatory function does not have a limit.

We saw one interesting result that helps simplify our life in some scenarios.

Theorem 4 (Sequential criterion)

Let $f : A \rightarrow \mathbb{R}$ be a function and let $a \in A$. Then, f is continuous at a iff given any sequence (a_n) in A such that $a_n \rightarrow a$, we have $f(a_n) \rightarrow f(a)$.

This makes life simpler because it is sometimes easier to deal with sequences. We had seen an example of this when we proved that a certain oscillatory function does not have a limit. Do you remember which?

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$.

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c*

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c* if the following limit exists:

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c* if the following limit exists:

$$\lim_{h \rightarrow 0} \frac{f(c + h) - f(c)}{h}.$$

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c* if the following limit exists:

$$\lim_{h \rightarrow 0} \frac{f(c + h) - f(c)}{h}.$$

In such a case, we call the value of the above limit the derivative of f at c

Next, we defined derivative. This was also not difficult.

Definition 11 (Derivative)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a function and let $c \in (a, b)$. f is said to be *differentiable at the point c* if the following limit exists:

$$\lim_{h \rightarrow 0} \frac{f(c + h) - f(c)}{h}.$$

In such a case, we call the value of the above limit the derivative of f at c and denote it by $f'(c)$.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather

$$(fg)'(c) = f'(c)g(c) + f(c)g'(c).$$

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 .

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$,

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$, then we say that f has a *local maximum* at x_0 .

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$, then we say that f has a *local maximum* at x_0 .

Of course, we have an analogous definition for minimum.

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$, then we say that f has a *local maximum* at x_0 .

Of course, we have an analogous definition for minimum. Note that here, we have that x_0 is an “interior point.”

We then have the usual rules about product/sum/composition of differentiable functions again being differentiable. Of course, we **don't** have the naïve product rule but rather $(fg)'(c) = f'(c)g(c) + f(c)g'(c)$. We then looked at minima/maxima.

Definition 12 (Local maximum)

Let $f : X \rightarrow \mathbb{R}$ be a function and let $x_0 \in X$. Suppose that there is an interval $(c, d) \subset X$ containing x_0 . If we have $f(x_0) \geq f(x)$ for all $x \in (c, d)$, then we say that f has a *local maximum* at x_0 .

Of course, we have an analogous definition for minimum. Note that here, we have that x_0 is an “interior point.” That is, there is an interval *around* x_0 contained within the domain.

Theorem 5 (Fermat's Theorem)

Theorem 5 (Fermat's Theorem)

If $f : X \rightarrow \mathbb{R}$ is differentiable

Theorem 5 (Fermat's Theorem)

If $f : X \rightarrow \mathbb{R}$ is differentiable and has a local minimum or maximum at a point $x_0 \in X$,

Theorem 5 (Fermat's Theorem)

If $f : X \rightarrow \mathbb{R}$ is differentiable and has a local minimum or maximum at a point $x_0 \in X$, then $f'(x_0) = 0$.

Theorem 5 (Fermat's Theorem)

If $f : X \rightarrow \mathbb{R}$ is differentiable and has a local minimum or maximum at a point $x_0 \in X$, then $f'(x_0) = 0$.

Once again, note that this only talks about “interior points.”

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function.

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) .

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$,

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

Using the above, we have a more general result.

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

Using the above, we have a more general result.

Theorem 7 (Mean Value Theorem)

Let f be continuous and differentiable as above.

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

Using the above, we have a more general result.

Theorem 7 (Mean Value Theorem)

Let f be continuous and differentiable as above. There exists $c \in (a, b)$

We then saw Rolle's Theorem. Note the hypothesis carefully.

Theorem 6 (Rolle's Theorem)

Suppose $f : [a, b] \rightarrow \mathbb{R}$ is a *continuous* function. Further, assume that it is differentiable on (a, b) . In this case, if $f(a) = f(b)$, then $f'(c) = 0$ for some $c \in (a, b)$.

Using the above, we have a more general result.

Theorem 7 (Mean Value Theorem)

Let f be continuous and differentiable as above. There exists $c \in (a, b)$ such that

$$f'(c) = \frac{f(b) - f(a)}{b - a}.$$

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function.

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) .

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$.

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

$$f'(x_0) = u.$$

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

$$f'(x_0) = u.$$

Note that the derivative of a (differentiable) function need not be continuous.

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

$$f'(x_0) = u.$$

Note that the derivative of a (differentiable) function need not be continuous. We shall see an example in the tutorial today, in fact.

We then saw a theorem which said “derivatives have IVP.” To be more precise:

Theorem 8 (Darboux's Theorem)

Let $f : (a, b) \rightarrow \mathbb{R}$ be a differentiable function. Let $c < d$ be points in (a, b) . Let u be between $f'(c)$ and $f'(d)$. Then, there exists $x_0 \in (c, d)$ such that

$$f'(x_0) = u.$$

Note that the derivative of a (differentiable) function need not be continuous. We shall see an example in the tutorial today, in fact. However, the above theorem tells us how the derivative can't have “jump” discontinuity.

Stop recording. Start a new one.
Take doubts.

Start recording!

Week 3

What did we see last week?

What did we see last week? Continuity,

Week 3

What did we see last week? Continuity, IVP,

Week 3

What did we see last week? Continuity, IVP, EVT,

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion,

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative,

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum,

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem,

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem, Rolle's and Mean Value Theorems,

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem, Rolle's and Mean Value Theorems, Darboux's Theorem.

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem, Rolle's and Mean Value Theorems, Darboux's Theorem. (Whew!)

We also saw an example of a function with non-continuous derivative.

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem, Rolle's and Mean Value Theorems, Darboux's Theorem. (Whew!)

We also saw an example of a function with non-continuous derivative. What was it?

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem, Rolle's and Mean Value Theorems, Darboux's Theorem. (Whew!)

We also saw an example of a function with non-continuous derivative. What was it?

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem, Rolle's and Mean Value Theorems, Darboux's Theorem. (Whew!)

We also saw an example of a function with non-continuous derivative. What was it?

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$f(x) := \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem, Rolle's and Mean Value Theorems, Darboux's Theorem. (Whew!)

We also saw an example of a function with non-continuous derivative. What was it?

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$f(x) := \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Keep this in mind.

Week 3

What did we see last week? Continuity, IVP, EVT, sequential criterion, derivative, (local) maximum and minimum, Fermat's not-Last Theorem, Rolle's and Mean Value Theorems, Darboux's Theorem. (Whew!)

We also saw an example of a function with non-continuous derivative. What was it?

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$f(x) := \begin{cases} x^2 \sin\left(\frac{1}{x}\right) & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Keep this in mind. please.

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

Theorem 9 (Second derivative test)

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

Theorem 9 (Second derivative test)

Assume that $f : [a, b] \rightarrow \mathbb{R}$ is continuous.

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

Theorem 9 (Second derivative test)

Assume that $f : [a, b] \rightarrow \mathbb{R}$ is continuous.

Suppose that $x_0 \in (a, b)$

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

Theorem 9 (Second derivative test)

Assume that $f : [a, b] \rightarrow \mathbb{R}$ is continuous.

Suppose that $x_0 \in (a, b)$ is such that $f'(x_0) = 0$ and $f''(x_0)$ exists.

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

Theorem 9 (Second derivative test)

Assume that $f : [a, b] \rightarrow \mathbb{R}$ is continuous.

Suppose that $x_0 \in (a, b)$ is such that $f'(x_0) = 0$ and $f''(x_0)$ exists.
Then,

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

Theorem 9 (Second derivative test)

Assume that $f : [a, b] \rightarrow \mathbb{R}$ is continuous.

Suppose that $x_0 \in (a, b)$ is such that $f'(x_0) = 0$ and $f''(x_0)$ exists.

Then,

- ① $f''(x_0) > 0 \implies f$ has a local minimum at x_0 ,

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

Theorem 9 (Second derivative test)

Assume that $f : [a, b] \rightarrow \mathbb{R}$ is continuous.

Suppose that $x_0 \in (a, b)$ is such that $f'(x_0) = 0$ and $f''(x_0)$ exists.

Then,

- ① $f''(x_0) > 0 \implies f$ has a local minimum at x_0 ,
- ② $f''(x_0) < 0 \implies f$ has a local maximum at x_0 .

We turn back to maximum and minimum and recall a theorem you must have seen in your previous life.

Theorem 9 (Second derivative test)

Assume that $f : [a, b] \rightarrow \mathbb{R}$ is continuous.

Suppose that $x_0 \in (a, b)$ is such that $f'(x_0) = 0$ and $f''(x_0)$ exists.

Then,

- ① $f''(x_0) > 0 \implies f$ has a local minimum at x_0 ,
- ② $f''(x_0) < 0 \implies f$ has a local maximum at x_0 .

If $f''(x_0) = 0$, then nothing can be concluded.

We now look at concavity and convexity.

Definition 13 (Convex)

We now look at concavity and convexity. For what follows, I will always denote an interval.

Definition 13 (Convex)

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

A function $f : I \rightarrow \mathbb{R}$ is said to be *convex* if

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

A function $f : I \rightarrow \mathbb{R}$ is said to be *convex* if for every $x_1, x_2 \in I$

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

A function $f : I \rightarrow \mathbb{R}$ is said to be *convex* if for every $x_1, x_2 \in I$ and every $t \in [0, 1]$,

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

A function $f : I \rightarrow \mathbb{R}$ is said to be *convex* if for every $x_1, x_2 \in I$ and every $t \in [0, 1]$, we have

$$f(tx_1 + (1 - t)x_2) \leq tf(x_1) + (1 - t)f(x_2).$$

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

A function $f : I \rightarrow \mathbb{R}$ is said to be *convex* if for every $x_1, x_2 \in I$ and every $t \in [0, 1]$, we have

$$f(tx_1 + (1 - t)x_2) \leq tf(x_1) + (1 - t)f(x_2).$$

More graphically, given any two points on the graph of the function,

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

A function $f : I \rightarrow \mathbb{R}$ is said to be *convex* if for every $x_1, x_2 \in I$ and every $t \in [0, 1]$, we have

$$f(tx_1 + (1 - t)x_2) \leq tf(x_1) + (1 - t)f(x_2).$$

More graphically, given any two points on the graph of the function, the line segment joining the two points

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

A function $f : I \rightarrow \mathbb{R}$ is said to be *convex* if for every $x_1, x_2 \in I$ and every $t \in [0, 1]$, we have

$$f(tx_1 + (1 - t)x_2) \leq tf(x_1) + (1 - t)f(x_2).$$

More graphically, given any two points on the graph of the function, the line segment joining the two points lies **above** the graph.

We now look at concavity and convexity. For what follows, I will always denote an interval. (It could be open/close/neither/unbounded.)

Definition 13 (Convex)

A function $f : I \rightarrow \mathbb{R}$ is said to be *convex* if for every $x_1, x_2 \in I$ and every $t \in [0, 1]$, we have

$$f(tx_1 + (1 - t)x_2) \leq tf(x_1) + (1 - t)f(x_2).$$

More graphically, given any two points on the graph of the function, the line segment joining the two points lies **above** the graph.

The definition of a *concave* function is obtained by replacing \leq with \geq and “**above**” with “**below**.”

Note that the definition does not even assume continuity.

Note that the definition does not even assume continuity. In particular, the function need not be differentiable,

Note that the definition does not even assume continuity. In particular, the function need not be differentiable, much less twice differentiable.

Note that the definition does not even assume continuity. In particular, the function need not be differentiable, much less twice differentiable.

However, if we do assume that it's differentiable, then we can say some things.

Note that the definition does not even assume continuity. In particular, the function need not be differentiable, much less twice differentiable.

However, if we do assume that it's differentiable, then we can say some things. If we assume twice differentiability, we can say some more things.

Note that the definition does not even assume continuity. In particular, the function need not be differentiable, much less twice differentiable.

However, if we do assume that it's differentiable, then we can say some things. If we assume twice differentiability, we can say some more things. I have put a summary of these on the next slide.

Note that the definition does not even assume continuity. In particular, the function need not be differentiable, much less twice differentiable.

However, if we do assume that it's differentiable, then we can say some things. If we assume twice differentiability, we can say some more things. I have put a summary of these on the next slide.

Read it some day.

Proposition 2

Suppose $f : I \rightarrow \mathbb{R}$ is differentiable. Then

- ① f' is increasing on $I \iff f$ is convex on I .
- ② f' is decreasing on $I \iff f$ is concave on I .
- ③ f' is strictly increasing on $I \iff f$ is strictly convex on I .
- ④ f' is strictly decreasing on $I \iff f$ is strictly concave on I .

Corollary 1

Suppose $f : I \rightarrow \mathbb{R}$ is **twice** differentiable. Then

- ① $f'' \geq 0$ on $I \iff f$ is convex on I .
- ② $f'' \leq 0$ on $I \iff f$ is concave on I .
- ③ $f'' > 0$ on $I \implies f$ is strictly convex on I .
- ④ $f'' < 0$ on $I \implies f$ is strictly concave on I .

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I .

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that **either**

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that **either**

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, **or**

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that **either**

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, **or**
- ② f is concave on $(x_0 - \delta, x_0)$ and convex on $(x_0, x_0 + \delta)$.

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that **either**

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, **or**
- ② f is concave on $(x_0 - \delta, x_0)$ and convex on $(x_0, x_0 + \delta)$.

As a crazy example, note that 0 is an inflection point of:

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that **either**

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, **or**
- ② f is concave on $(x_0 - \delta, x_0)$ and convex on $(x_0, x_0 + \delta)$.

As a crazy example, note that 0 is an inflection point of:

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that **either**

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, **or**
- ② f is concave on $(x_0 - \delta, x_0)$ and convex on $(x_0, x_0 + \delta)$.

As a crazy example, note that 0 is an inflection point of:

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$f(x) := \begin{cases} \frac{1}{x} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that either

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, or
- ② f is concave on $(x_0 - \delta, x_0)$ and convex on $(x_0, x_0 + \delta)$.

As a crazy example, note that 0 is an inflection point of:

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$f(x) := \begin{cases} \frac{1}{x} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Note that f is not even continuous at 0.

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that either

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, or
- ② f is concave on $(x_0 - \delta, x_0)$ and convex on $(x_0, x_0 + \delta)$.

As a crazy example, note that 0 is an inflection point of:

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$f(x) := \begin{cases} \frac{1}{x} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Note that f is not even continuous at 0. Let alone twice differentiable.

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that either

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, or
- ② f is concave on $(x_0 - \delta, x_0)$ and convex on $(x_0, x_0 + \delta)$.

As a crazy example, note that 0 is an inflection point of:

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$f(x) := \begin{cases} \frac{1}{x} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Note that f is not even continuous at 0. Let alone twice differentiable. Also note that every point is a point of inflection for an affine function $x \mapsto ax + b$.

Let's now talk about inflection points.

Definition 14 (Inflection point)

Let x_0 be an *interior point* of I . Then, x_0 is called an inflection for f if there exists $\delta > 0$ such that either

- ① f is convex on $(x_0 - \delta, x_0)$ and concave on $(x_0, x_0 + \delta)$, or
- ② f is concave on $(x_0 - \delta, x_0)$ and convex on $(x_0, x_0 + \delta)$.

As a crazy example, note that 0 is an inflection point of:

$f : \mathbb{R} \rightarrow \mathbb{R}$ defined as

$$f(x) := \begin{cases} \frac{1}{x} & x \neq 0, \\ 0 & x = 0. \end{cases}$$

Note that f is not even continuous at 0. Let alone twice differentiable. Also note that every point is a point of inflection for an affine function $x \mapsto ax + b$. (Even if $a = 0$.)

Week 3

Here's some more information being thrown at you.

Week 3

Here's some more information being thrown at you. Look at it some day.

Week 3

Here's some more information being thrown at you. Look at it some day. Let $x_0 \in I$ be an *interior point*, and $f : I \rightarrow \mathbb{R}$.

Here's some more information being thrown at you. Look at it some day. Let $x_0 \in I$ be an *interior point*, and $f : I \rightarrow \mathbb{R}$.

Theorem 10 (Derivative tests)

- ① **(First derivative test)** Suppose f is differentiable on $(x_0 - r, x_0) \cup (x_0, x_0 + r)$ for some $r > 0$. Then, x_0 is a point of inflection \iff there is $\delta > 0$ with $\delta < r$ such that f' is increasing on $(x_0 - \delta, x_0)$ and f' is decreasing on $(x_0, x_0 + \delta)$, or vice-versa.
- ② **(Second derivative test)** Suppose f is twice differentiable on $(x_0 - r, x_0) \cup (x_0, x_0 + r)$ for some $r > 0$. Then, x_0 is a point of inflection \iff there is $\delta > 0$ with $\delta < r$ such that $f'' \geq 0$ on $(x_0 - \delta, x_0)$ and $f'' \leq 0$ on $(x_0, x_0 + \delta)$, or vice-versa.

Thus, if f is twice differentiable, then x_0 is inflection point iff f'' changes sign. (Note that $f''(x_0)$ is not required to exist. Recall the crazy example.)

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 .

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f ,

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f , then $f''(x_0) = 0$.

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f , then $f''(x_0) = 0$.

In the above, we are not assuming the existence of f'' at other points.

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f , then $f''(x_0) = 0$.

In the above, we are not assuming the existence of f'' at other points. The following is now a **sufficient** condition.

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f , then $f''(x_0) = 0$.

In the above, we are not assuming the existence of f'' at other points. The following is now a **sufficient** condition.

Theorem 12 (A **third** derivative test)

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f , then $f''(x_0) = 0$.

In the above, we are not assuming the existence of f'' at other points. The following is now a **sufficient** condition.

Theorem 12 (A **third** derivative test)

Suppose f is **thrice** differentiable at x_0

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f , then $f''(x_0) = 0$.

In the above, we are not assuming the existence of f'' at other points. The following is now a **sufficient** condition.

Theorem 12 (A **third** derivative test)

Suppose f is **thrice** differentiable at x_0 such that $f''(x_0) = 0$

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f , then $f''(x_0) = 0$.

In the above, we are not assuming the existence of f'' at other points. The following is now a **sufficient** condition.

Theorem 12 (A **third** derivative test)

Suppose f is **thrice** differentiable at x_0 such that $f''(x_0) = 0$ and $f'''(x_0) \neq 0$.

The previous slide gives us a **necessary** condition for inflection point. We have the same notation as earlier.

Theorem 11 (Another second derivative test)

Suppose f is twice differentiable at x_0 . If x_0 is a point of inflection for f , then $f''(x_0) = 0$.

In the above, we are not assuming the existence of f'' at other points. The following is now a **sufficient** condition.

Theorem 12 (A **third** derivative test)

Suppose f is **thrice** differentiable at x_0 such that $f''(x_0) = 0$ and $f'''(x_0) \neq 0$. Then, x_0 is an inflection point for f .

Okay, that's enough about convex/concave/inflection points.

Okay, that's enough about convex/concave/inflection points.
Hopefully, any possible doubt about these is covered in the previous slides.

Okay, that's enough about convex/concave/inflection points. Hopefully, any possible doubt about these is covered in the previous slides. Read them some day and ask doubts, if any.

Okay, that's enough about convex/concave/inflection points. Hopefully, any possible doubt about these is covered in the previous slides. Read them some day and ask doubts, if any.

Once again, keep in mind the crazy example.

Okay, that's enough about convex/concave/inflection points. Hopefully, any possible doubt about these is covered in the previous slides. Read them some day and ask doubts, if any.

Once again, keep in mind the crazy example. The definitions of these concepts do not require any continuity or anything at the point.

Okay, that's enough about convex/concave/inflection points. Hopefully, any possible doubt about these is covered in the previous slides. Read them some day and ask doubts, if any.

Once again, keep in mind the crazy example. The definitions of these concepts do not require any continuity or anything at the point. However, we do have the theorem that a convex function on an open interval is continuous.

Okay, that's enough about convex/concave/inflection points. Hopefully, any possible doubt about these is covered in the previous slides. Read them some day and ask doubts, if any.

Once again, keep in mind the crazy example. The definitions of these concepts do not require any continuity or anything at the point. However, we do have the theorem that a convex function on an open interval is continuous. Proof:

[https://unapologetic.wordpress.com/2008/04/15/
convex-functions-are-continuous/](https://unapologetic.wordpress.com/2008/04/15/convex-functions-are-continuous/)

Okay, that's enough about convex/concave/inflection points. Hopefully, any possible doubt about these is covered in the previous slides. Read them some day and ask doubts, if any.

Once again, keep in mind the crazy example. The definitions of these concepts do not require any continuity or anything at the point. However, we do have the theorem that a convex function on an open interval is continuous. Proof:

[https://unapologetic.wordpress.com/2008/04/15/
convex-functions-are-continuous/](https://unapologetic.wordpress.com/2008/04/15/convex-functions-are-continuous/)

We also have the theorem that a convex function is differentiable at all but at most countably many points.

Okay, that's enough about convex/concave/inflection points. Hopefully, any possible doubt about these is covered in the previous slides. Read them some day and ask doubts, if any.

Once again, keep in mind the crazy example. The definitions of these concepts do not require any continuity or anything at the point. However, we do have the theorem that a convex function on an open interval is continuous. Proof:

<https://unapologetic.wordpress.com/2008/04/15/convex-functions-are-continuous/>

We also have the theorem that a convex function is differentiable at all but at most countably many points. Proof:

<https://math.stackexchange.com/questions/946311>

Let's now look at Taylor polynomials.

Definition 15 (Taylor polynomials)



Let's now look at Taylor polynomials. From now, I will be an **open** interval, a an interior point of I , and $f : I \rightarrow \mathbb{R}$ a function.

Definition 15 (Taylor polynomials)



Let's now look at Taylor polynomials. From now, I will be an **open** interval, a an interior point of I , and $f : I \rightarrow \mathbb{R}$ a function.

Definition 15 (Taylor polynomials)

Let f be n times differentiable



Let's now look at Taylor polynomials. From now, I will be an **open** interval, a an interior point of I , and $f : I \rightarrow \mathbb{R}$ a function.

Definition 15 (Taylor polynomials)

Let f be n times differentiable **at x_0** .



Let's now look at Taylor polynomials. From now, I will be an **open** interval, a an interior point of I , and $f : I \rightarrow \mathbb{R}$ a function.

Definition 15 (Taylor polynomials)

Let f be n times differentiable **at x_0** . We define the $n + 1$ Taylor polynomials as



Let's now look at Taylor polynomials. From now, I will be an **open** interval, a an interior point of I , and $f : I \rightarrow \mathbb{R}$ a function.

Definition 15 (Taylor polynomials)

Let f be n times differentiable **at x_0** . We define the $n + 1$ Taylor polynomials as

$$P_0(x) = f(x_0)$$



Let's now look at Taylor polynomials. From now, I will be an **open** interval, a an interior point of I , and $f : I \rightarrow \mathbb{R}$ a function.

Definition 15 (Taylor polynomials)

Let f be n times differentiable **at x_0** . We define the $n + 1$ Taylor polynomials as

$$P_0(x) = f(x_0)$$

$$P_1(x) = f(x_0) + \frac{f^{(1)}(x_0)}{1!}(x - x_0)$$



Let's now look at Taylor polynomials. From now, I will be an **open** interval, a an interior point of I , and $f : I \rightarrow \mathbb{R}$ a function.

Definition 15 (Taylor polynomials)

Let f be n times differentiable **at x_0** . We define the $n + 1$ Taylor polynomials as

$$P_0(x) = f(x_0)$$

$$P_1(x) = f(x_0) + \frac{f^{(1)}(x_0)}{1!}(x - x_0)$$

$$P_2(x) = f(x_0) + \frac{f^{(1)}(x_0)}{1!}(x - x_0) + \frac{f^{(2)}(x_0)}{2!}(x - x_0)^2$$



Let's now look at Taylor polynomials. From now, I will be an **open** interval, a an interior point of I , and $f : I \rightarrow \mathbb{R}$ a function.

Definition 15 (Taylor polynomials)

Let f be n times differentiable **at x_0** . We define the $n + 1$ Taylor polynomials as

$$P_0(x) = f(x_0)$$

$$P_1(x) = f(x_0) + \frac{f^{(1)}(x_0)}{1!}(x - x_0)$$

$$P_2(x) = f(x_0) + \frac{f^{(1)}(x_0)}{1!}(x - x_0) + \frac{f^{(2)}(x_0)}{2!}(x - x_0)^2$$

⋮

$$P_n(x) = f(x_0) + \frac{f^{(1)}(x_0)}{1!}(x - x_0) + \cdots + \frac{f^{(n)}(x_0)}{n!}(x - x_0)^n$$



Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have.

Theorem 13 (Taylor's theorem)

Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have. Also note that so far, we have just defined some polynomials.

Theorem 13 (Taylor's theorem)

Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have. Also note that so far, we have just defined some polynomials. We are yet to see how it actually connects with the function itself.

Theorem 13 (Taylor's theorem)

Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have. Also note that so far, we have just defined some polynomials. We are yet to see how it actually connects with the function itself. This is given by the following theorem.

Theorem 13 (Taylor's theorem)

Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have. Also note that so far, we have just defined some polynomials. We are yet to see how it actually connects with the function itself. This is given by the following theorem.

Theorem 13 (Taylor's theorem)

Suppose that f is $n+1$ times differentiable

Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have. Also note that so far, we have just defined some polynomials. We are yet to see how it actually connects with the function itself. This is given by the following theorem.

Theorem 13 (Taylor's theorem)

Suppose that f is $n+1$ times differentiable **on I** .

Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have. Also note that so far, we have just defined some polynomials. We are yet to see how it actually connects with the function itself. This is given by the following theorem.

Theorem 13 (Taylor's theorem)

Suppose that f is $n+1$ times differentiable **on I** . Suppose that $b \in I$.

Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have. Also note that so far, we have just defined some polynomials. We are yet to see how it actually connects with the function itself. This is given by the following theorem.

Theorem 13 (Taylor's theorem)

Suppose that f is $n+1$ times differentiable on I . Suppose that $b \in I$. Then, there exists $c \in (a, b) \cup (b, a)$ such that

Note that all the Taylor **polynomials** have only **finitely** many terms, as a polynomial should have. Also note that so far, we have just defined some polynomials. We are yet to see how it actually connects with the function itself. This is given by the following theorem.

Theorem 13 (Taylor's theorem)

Suppose that f is $n+1$ times differentiable on I . Suppose that $b \in I$. Then, there exists $c \in (a, b) \cup (b, a)$ such that

$$f(b) = P_n(b) + \frac{f^{(n+1)}(c)}{(n+1)!}(b-a)^{n+1},$$

where P_n is as in the previous slide.

Given a function f (which is $n + 1$ times differentiable) and a Taylor polynomial P_n ,

Given a function f (which is $n + 1$ times differentiable) and a Taylor polynomial P_n , we can define the n^{th} remainder as

Given a function f (which is $n + 1$ times differentiable) and a Taylor polynomial P_n , we can define the n^{th} remainder as

$$R_n(x) := f(x) - P_n(x), \quad \text{for } x \in I.$$

Given a function f (which is $n + 1$ times differentiable) and a Taylor polynomial P_n , we can define the n^{th} remainder as

$$R_n(x) := f(x) - P_n(x), \quad \text{for } x \in I.$$

By the previous theorem, we know that

$$R_n(x) = \frac{f^{(n+1)}(c_x)}{(n+1)!}(x-a)^{n+1},$$

Given a function f (which is $n + 1$ times differentiable) and a Taylor polynomial P_n , we can define the n^{th} remainder as

$$R_n(x) := f(x) - P_n(x), \quad \text{for } x \in I.$$

By the previous theorem, we know that

$$R_n(x) = \frac{f^{(n+1)}(c_x)}{(n+1)!}(x-a)^{n+1},$$

for some c_x between x and a .

Given a function f (which is $n + 1$ times differentiable) and a Taylor polynomial P_n , we can define the n^{th} remainder as

$$R_n(x) := f(x) - P_n(x), \quad \text{for } x \in I.$$

By the previous theorem, we know that

$$R_n(x) = \frac{f^{(n+1)}(c_x)}{(n+1)!}(x-a)^{n+1},$$

for some c_x between x and a .

Sometimes, assuming $f \in \mathcal{C}^\infty(I)$, we can bound $f^{(n+1)}(c_x)$ in a nice enough way to get that

Given a function f (which is $n + 1$ times differentiable) and a Taylor polynomial P_n , we can define the n^{th} remainder as

$$R_n(x) := f(x) - P_n(x), \quad \text{for } x \in I.$$

By the previous theorem, we know that

$$R_n(x) = \frac{f^{(n+1)}(c_x)}{(n+1)!}(x-a)^{n+1},$$

for some c_x between x and a .

Sometimes, assuming $f \in \mathcal{C}^\infty(I)$, we can bound $f^{(n+1)}(c_x)$ in a nice enough way to get that

$$R_n(x) \rightarrow 0$$

Given a function f (which is $n + 1$ times differentiable) and a Taylor polynomial P_n , we can define the n^{th} remainder as

$$R_n(x) := f(x) - P_n(x), \quad \text{for } x \in I.$$

By the previous theorem, we know that

$$R_n(x) = \frac{f^{(n+1)}(c_x)}{(n+1)!}(x-a)^{n+1},$$

for some c_x between x and a .

Sometimes, assuming $f \in \mathcal{C}^\infty(I)$, we can bound $f^{(n+1)}(c_x)$ in a nice enough way to get that

$$R_n(x) \rightarrow 0$$

for some $x \in I$.

If the previous thing happens,

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x .

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x . For some nice functions, we get an $R > 0$

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x . For some nice functions, we get an $R > 0$ such that the above happens for all $x \in (a - R, a + R)$.

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x . For some nice functions, we get an $R > 0$ such that the above happens for all $x \in (a - R, a + R)$. If such an R exists for all $a \in I$,

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x . For some nice functions, we get an $R > 0$ such that the above happens for all $x \in (a - R, a + R)$. If such an R exists for all $a \in I$, then f is said to be *analytic*.

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x . For some nice functions, we get an $R > 0$ such that the above happens for all $x \in (a - R, a + R)$. If such an R exists for all $a \in I$, then f is said to be *analytic*. (The R may depend on a .)

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x . For some nice functions, we get an $R > 0$ such that the above happens for all $x \in (a - R, a + R)$. If such an R exists for all $a \in I$, then f is said to be *analytic*. (The R may depend on a .)

Note that the Taylor **series**

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x . For some nice functions, we get an $R > 0$ such that the above happens for all $x \in (a - R, a + R)$. If such an R exists for all $a \in I$, then f is said to be *analytic*. (The R may depend on a .)

Note that the Taylor **series** about some point a may still converge but *not* to f .

If the previous thing happens, then we get that

$$f(x) = \lim_{n \rightarrow \infty} P_n(x)$$

for all such x .

Thus, we get

$$f(x) = \sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!} (x - a)^n$$

for all those x . For some nice functions, we get an $R > 0$ such that the above happens for all $x \in (a - R, a + R)$. If such an R exists for all $a \in I$, then f is said to be *analytic*. (The R may depend on a .)

Note that the Taylor **series** about some point a may still converge but *not* to f . Such a function is not called analytic.

Some final remarks:

Some final remarks:

The last thing written $\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$ is not a Taylor **polynomial**.

Some final remarks:

The last thing written $\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$ is not a Taylor **polynomial**.
It is the Taylor **series**.

Some final remarks:

The last thing written $\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$ is not a Taylor **polynomial**.

It is the Taylor **series**.

It may happen to not converge for *any* $x \neq a$.

Some final remarks:

The last thing written $\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$ is not a Taylor **polynomial**.

It is the Taylor **series**.

It may happen to not converge for *any* $x \neq a$. It may also happen to converge for all $x \in \mathbb{R}$.

Some final remarks:

The last thing written $\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$ is not a Taylor **polynomial**.

It is the Taylor **series**.

It may happen to not converge for *any* $x \neq a$. It may also happen to converge for all $x \in \mathbb{R}$.

Suppose that the series converges on some interval J such that $a \in J \subset I$.

Some final remarks:

The last thing written $\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$ is not a Taylor **polynomial**.

It is the Taylor **series**.

It may happen to not converge for *any* $x \neq a$. It may also happen to converge for all $x \in \mathbb{R}$.

Suppose that the series converges on some interval J such that $a \in J \subset I$. It is not necessary that the Taylor series converges to f on J .

Some final remarks:

The last thing written $\sum_{n=0}^{\infty} \frac{f^{(n)}(a)}{n!}(x - a)^n$ is not a Taylor **polynomial**.

It is the Taylor **series**.

It may happen to not converge for *any* $x \neq a$. It may also happen to converge for all $x \in \mathbb{R}$.

Suppose that the series converges on some interval J such that $a \in J \subset I$. It is not necessary that the Taylor series converges to f on J .

What was the example seen in class that illustrated this?

Stop recording. Start a new one.
Take doubts.

In the following, it will be tacitly assumed that $a, b \in \mathbb{R}$ with $a < b$.

Definition 16 (Partitions)

In the following, it will be tacitly assumed that $a, b \in \mathbb{R}$ with $a < b$.

Definition 16 (Partitions)

Given a closed interval $[a, b]$,

In the following, it will be tacitly assumed that $a, b \in \mathbb{R}$ with $a < b$.

Definition 16 (Partitions)

Given a closed interval $[a, b]$, a *partition* P of $[a, b]$

In the following, it will be tacitly assumed that $a, b \in \mathbb{R}$ with $a < b$.

Definition 16 (Partitions)

Given a closed interval $[a, b]$, a *partition* P of $[a, b]$ is a finite collection of points

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}.$$

In the following, it will be tacitly assumed that $a, b \in \mathbb{R}$ with $a < b$.

Definition 16 (Partitions)

Given a closed interval $[a, b]$, a *partition* P of $[a, b]$ is a finite collection of points

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}.$$

Note that a partition P is really just a subset of $[a, b]$

In the following, it will be tacitly assumed that $a, b \in \mathbb{R}$ with $a < b$.

Definition 16 (Partitions)

Given a closed interval $[a, b]$, a *partition* P of $[a, b]$ is a finite collection of points

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}.$$

Note that a partition P is really just a subset of $[a, b]$ with the requirement that it must be finite and contain a and b .

In the following, it will be tacitly assumed that $a, b \in \mathbb{R}$ with $a < b$.

Definition 16 (Partitions)

Given a closed interval $[a, b]$, a *partition* P of $[a, b]$ is a finite collection of points

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}.$$

Note that a partition P is really just a subset of $[a, b]$ with the requirement that it must be finite and contain a and b . It is customary to then list it in increasing order.

Definition 17 (Refinements)

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$,

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P if $P \subset P'$.

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P if $P \subset P'$.

The “ \subset ” makes sense because of our earlier remark about partitions just being subsets of $[a, b]$.

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P if $P \subset P'$.

The “ \subset ” makes sense because of our earlier remark about partitions just being subsets of $[a, b]$. In other words, it means that every point of P is also a point in P' .

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P if $P \subset P'$.

The “ \subset ” makes sense because of our earlier remark about partitions just being subsets of $[a, b]$. In other words, it means that every point of P is also a point in P' . Thus, we have “refined” the partition by further “chopping” it up.

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P if $P \subset P'$.

The “ \subset ” makes sense because of our earlier remark about partitions just being subsets of $[a, b]$. In other words, it means that every point of P is also a point in P' . Thus, we have “refined” the partition by further “chopping” it up.

Given two partitions P_1 and P_2 of $[a, b]$,

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P if $P \subset P'$.

The “ \subset ” makes sense because of our earlier remark about partitions just being subsets of $[a, b]$. In other words, it means that every point of P is also a point in P' . Thus, we have “refined” the partition by further “chopping” it up.

Given two partitions P_1 and P_2 of $[a, b]$, we see that $P = P_1 \cup P_2$ is also a partition of $[a, b]$.

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P if $P \subset P'$.

The “ \subset ” makes sense because of our earlier remark about partitions just being subsets of $[a, b]$. In other words, it means that every point of P is also a point in P' . Thus, we have “refined” the partition by further “chopping” it up.

Given two partitions P_1 and P_2 of $[a, b]$, we see that $P = P_1 \cup P_2$ is also a partition of $[a, b]$. Moreover, P is a refinement of both P_1 and P_2 .

Definition 17 (Refinements)

Given two partitions P and P' of $[a, b]$, we say that P' is a *refinement* of P if $P \subset P'$.

The “ \subset ” makes sense because of our earlier remark about partitions just being subsets of $[a, b]$. In other words, it means that every point of P is also a point in P' . Thus, we have “refined” the partition by further “chopping” it up.

Given two partitions P_1 and P_2 of $[a, b]$, we see that $P = P_1 \cup P_2$ is also a partition of $[a, b]$. Moreover, P is a refinement of both P_1 and P_2 . In other words, any two partitions have a common refinement.

Definition 18

Definition 18

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **bounded** function

Definition 18

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **bounded** function and

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}$$

a partition of $[a, b]$.

Definition 18

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **bounded** function and

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}$$

a partition of $[a, b]$.

We define the following quantities:

Definition 18

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **bounded** function and

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}$$

a partition of $[a, b]$.

We define the following quantities:

$$M_i := \sup_{x \in [x_{i-1}, x_i]} f(x) \quad \text{and} \quad m_i := \inf_{x \in [x_{i-1}, x_i]} f(x),$$

for $i = 1, \dots, n$.

Definition 18

Let $f : [a, b] \rightarrow \mathbb{R}$ be a **bounded** function and

$$P = \{a = x_0 < x_1 < \cdots < x_n = b\}$$

a partition of $[a, b]$.

We define the following quantities:

$$M_i := \sup_{x \in [x_{i-1}, x_i]} f(x) \quad \text{and} \quad m_i := \inf_{x \in [x_{i-1}, x_i]} f(x),$$

for $i = 1, \dots, n$.

Thus, m_i and M_i denote the infimum and supremum of f over the i -th interval, respectively.

Given everything as in the previous slide, we define lower/upper sums as following.

Definition 19 (Lower/Upper sum)

Given everything as in the previous slide, we define lower/upper sums as following.

Definition 19 (Lower/Upper sum)

The *lower sum* of f with respect to the partition P is defined as

Given everything as in the previous slide, we define lower/upper sums as following.

Definition 19 (Lower/Upper sum)

The *lower sum* of f with respect to the partition P is defined as

$$L(f, P) := \sum_{i=1}^n m_i(x_i - x_{i-1}).$$

Given everything as in the previous slide, we define lower/upper sums as following.

Definition 19 (Lower/Upper sum)

The *lower sum* of f with respect to the partition P is defined as

$$L(f, P) := \sum_{i=1}^n m_i(x_i - x_{i-1}).$$

The *upper sum* of f with respect to the partition P is defined as

Given everything as in the previous slide, we define lower/upper sums as following.

Definition 19 (Lower/Upper sum)

The *lower sum* of f with respect to the partition P is defined as

$$L(f, P) := \sum_{i=1}^n m_i(x_i - x_{i-1}).$$

The *upper sum* of f with respect to the partition P is defined as

$$U(f, P) := \sum_{i=1}^n M_i(x_i - x_{i-1}).$$

Given everything as in the previous slide, we define lower/upper sums as following.

Definition 19 (Lower/Upper sum)

The *lower sum* of f with respect to the partition P is defined as

$$L(f, P) := \sum_{i=1}^n m_i(x_i - x_{i-1}).$$

The *upper sum* of f with respect to the partition P is defined as

$$U(f, P) := \sum_{i=1}^n M_i(x_i - x_{i-1}).$$

In the above, note that we have both f and P in the notation.

Given everything as in the previous slide, we define lower/upper sums as following.

Definition 19 (Lower/Upper sum)

The *lower sum* of f with respect to the partition P is defined as

$$L(f, P) := \sum_{i=1}^n m_i(x_i - x_{i-1}).$$

The *upper sum* of f with respect to the partition P is defined as

$$U(f, P) := \sum_{i=1}^n M_i(x_i - x_{i-1}).$$

In the above, note that we have both f and P in the notation. This is crucial because the sums depend on the partition.

Using the earlier sums, we now define the upper and lower Darboux *integrals*. The notations are continuing from earlier.

Definition 20 (Lower/Upper Darboux integrals)

Using the earlier sums, we now define the upper and lower Darboux *integrals*. The notations are continuing from earlier.

Definition 20 (Lower/Upper Darboux integrals)

The *lower Darboux integral of f*

Using the earlier sums, we now define the upper and lower Darboux *integrals*. The notations are continuing from earlier.

Definition 20 (Lower/Upper Darboux integrals)

The *lower Darboux integral of f* is defined as

$$L(f) := \sup\{L(f, P) \mid P \text{ is a partition of } [a, b]\},$$

Using the earlier sums, we now define the upper and lower Darboux *integrals*. The notations are continuing from earlier.

Definition 20 (Lower/Upper Darboux integrals)

The *lower Darboux integral of f* is defined as

$$L(f) := \sup\{L(f, P) \mid P \text{ is a partition of } [a, b]\},$$

and the *upper Darboux integral of f*

Using the earlier sums, we now define the upper and lower Darboux *integrals*. The notations are continuing from earlier.

Definition 20 (Lower/Upper Darboux integrals)

The *lower Darboux integral of f* is defined as

$$L(f) := \sup\{L(f, P) \mid P \text{ is a partition of } [a, b]\},$$

and the *upper Darboux integral of f* is defined as

$$U(f) := \inf\{U(f, P) \mid P \text{ is a partition of } [a, b]\}.$$

Using the earlier sums, we now define the upper and lower Darboux *integrals*. The notations are continuing from earlier.

Definition 20 (Lower/Upper Darboux integrals)

The *lower Darboux integral of f* is defined as

$$L(f) := \sup\{L(f, P) \mid P \text{ is a partition of } [a, b]\},$$

and the *upper Darboux integral of f* is defined as

$$U(f) := \inf\{U(f, P) \mid P \text{ is a partition of } [a, b]\}.$$

Note that the sup / inf is over **all** the partitions P of $[a, b]$.

Using the earlier sums, we now define the upper and lower Darboux *integrals*. The notations are continuing from earlier.

Definition 20 (Lower/Upper Darboux integrals)

The *lower Darboux integral of f* is defined as

$$L(f) := \sup\{L(f, P) \mid P \text{ is a partition of } [a, b]\},$$

and the *upper Darboux integral of f* is defined as

$$U(f) := \inf\{U(f, P) \mid P \text{ is a partition of } [a, b]\}.$$

Note that the sup / inf is over **all** the partitions P of $[a, b]$.

Note that the notation now does not have any P .

Using the earlier sums, we now define the upper and lower Darboux *integrals*. The notations are continuing from earlier.

Definition 20 (Lower/Upper Darboux integrals)

The *lower Darboux integral of f* is defined as

$$L(f) := \sup\{L(f, P) \mid P \text{ is a partition of } [a, b]\},$$

and the *upper Darboux integral of f* is defined as

$$U(f) := \inf\{U(f, P) \mid P \text{ is a partition of } [a, b]\}.$$

Note that the sup / inf is over **all** the partitions P of $[a, b]$.

Note that the notation now does not have any P . This is because $L(f)$ and $U(f)$ don't depend on any specific partition.

Definition 21 (Darboux integrable)

Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$

Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable*

Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

In this case, we define

$$\int_a^b f(t)dt := U(f) = L(f).$$

Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

In this case, we define

$$\int_a^b f(t)dt := U(f) = L(f).$$

This value is called the Darboux integral.

Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

In this case, we define

$$\int_a^b f(t)dt := U(f) = L(f).$$

This value is called the Darboux integral.

Theorem 14 (Criteria for Darboux integrable)

Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

In this case, we define

$$\int_a^b f(t)dt := U(f) = L(f).$$

This value is called the Darboux integral.

Theorem 14 (Criteria for Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$



Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

In this case, we define

$$\int_a^b f(t)dt := U(f) = L(f).$$

This value is called the Darboux integral.

Theorem 14 (Criteria for Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is Darboux integrable if and only if



Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

In this case, we define

$$\int_a^b f(t)dt := U(f) = L(f).$$

This value is called the Darboux integral.

Theorem 14 (Criteria for Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is Darboux integrable if and only if for every $\epsilon > 0$,



Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

In this case, we define

$$\int_a^b f(t)dt := U(f) = L(f).$$

This value is called the Darboux integral.

Theorem 14 (Criteria for Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is Darboux integrable if and only if for every $\epsilon > 0$, there exists a partition P of $[a, b]$



Definition 21 (Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Darboux integrable* if $L(f) = U(f)$.

In this case, we define

$$\int_a^b f(t)dt := U(f) = L(f).$$

This value is called the Darboux integral.

Theorem 14 (Criteria for Darboux integrable)

A bounded function $f : [a, b] \rightarrow \mathbb{R}$ is Darboux integrable if and only if for every $\epsilon > 0$, there exists a partition P of $[a, b]$ such that

$$U(f, P) - L(f, P) < \epsilon.$$



A corollary of the previous is the following.

Corollary 2

A corollary of the previous is the following.

Corollary 2

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function.

A corollary of the previous is the following.

Corollary 2

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Suppose that (P_n) is a sequence of partitions of $[a, b]$

A corollary of the previous is the following.

Corollary 2

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Suppose that (P_n) is a sequence of partitions of $[a, b]$ such that

$$\lim_{n \rightarrow \infty} [U(f, P_n) - L(f, P_n)] = 0.$$

A corollary of the previous is the following.

Corollary 2

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Suppose that (P_n) is a sequence of partitions of $[a, b]$ such that

$$\lim_{n \rightarrow \infty} [U(f, P_n) - L(f, P_n)] = 0.$$

Then, f is Darboux integrable.

A corollary of the previous is the following.

Corollary 2

Let $f : [a, b] \rightarrow \mathbb{R}$ be a bounded function. Suppose that (P_n) is a sequence of partitions of $[a, b]$ such that

$$\lim_{n \rightarrow \infty} [U(f, P_n) - L(f, P_n)] = 0.$$

Then, f is Darboux integrable.

We now turn to the definition of Riemann integrals.

Some jargon.

Definition 22 (Norm of a partition)

Definition 23 (Tagged partition)

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$.

Definition 23 (Tagged partition)

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$. The *norm* of P is defined to be

Definition 23 (Tagged partition)

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$. The *norm* of P is defined to be

$$\|P\| := \max_{1 \leq i \leq n} [x_i - x_{i-1}].$$

Definition 23 (Tagged partition)

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$. The *norm* of P is defined to be

$$\|P\| := \max_{1 \leq i \leq n} [x_i - x_{i-1}].$$

In other words, it is the length of the largest sub-interval.

Definition 23 (Tagged partition)

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$. The *norm* of P is defined to be

$$\|P\| := \max_{1 \leq i \leq n} [x_i - x_{i-1}].$$

In other words, it is the length of the largest sub-interval.

Definition 23 (Tagged partition)

Given a partition P of $[a, b]$ as before,

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$. The *norm* of P is defined to be

$$\|P\| := \max_{1 \leq i \leq n} [x_i - x_{i-1}].$$

In other words, it is the length of the largest sub-interval.

Definition 23 (Tagged partition)

Given a partition P of $[a, b]$ as before, we get the intervals

$$I_i = [x_{i-1}, x_i]$$

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$. The *norm* of P is defined to be

$$\|P\| := \max_{1 \leq i \leq n} [x_i - x_{i-1}].$$

In other words, it is the length of the largest sub-interval.

Definition 23 (Tagged partition)

Given a partition P of $[a, b]$ as before, we get the intervals $I_i = [x_{i-1}, x_i]$ for $i = 1, \dots, n$. For each i , we pick a point $t_i \in I_i$.

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$. The *norm* of P is defined to be

$$\|P\| := \max_{1 \leq i \leq n} [x_i - x_{i-1}].$$

In other words, it is the length of the largest sub-interval.

Definition 23 (Tagged partition)

Given a partition P of $[a, b]$ as before, we get the intervals $I_i = [x_{i-1}, x_i]$ for $i = 1, \dots, n$. For each i , we pick a point $t_i \in I_i$. This collection of points together is denoted by t .

Some jargon.

Definition 22 (Norm of a partition)

Let $P = \{a = x_0 < \dots < x_n = b\}$ be a partition of $[a, b]$. The *norm* of P is defined to be

$$\|P\| := \max_{1 \leq i \leq n} [x_i - x_{i-1}].$$

In other words, it is the length of the largest sub-interval.

Definition 23 (Tagged partition)

Given a partition P of $[a, b]$ as before, we get the intervals $I_i = [x_{i-1}, x_i]$ for $i = 1, \dots, n$. For each i , we pick a point $t_i \in I_i$. This collection of points together is denoted by t . The pair (P, t) is called a *tagged partition* of $[a, b]$.

Definition 24 (Riemann sum)

Definition 24 (Riemann sum)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function.

Definition 24 (Riemann sum)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function. Let (P, t) be a tagged partition of $[a, b]$.

Definition 24 (Riemann sum)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function. Let (P, t) be a tagged partition of $[a, b]$. We define the *Riemann sum* associated to f and (P, t) by

Definition 24 (Riemann sum)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function. Let (P, t) be a tagged partition of $[a, b]$. We define the *Riemann sum* associated to f and (P, t) by

$$R(f, P, t) := \sum_{i=1}^n f(t_i)(x_i - x_{i-1}).$$

Definition 24 (Riemann sum)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function. Let (P, t) be a tagged partition of $[a, b]$. We define the *Riemann sum* associated to f and (P, t) by

$$R(f, P, t) := \sum_{i=1}^n f(t_i)(x_i - x_{i-1}).$$

Note that the notation here includes f , P , and t .

Definition 24 (Riemann sum)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function. Let (P, t) be a tagged partition of $[a, b]$. We define the *Riemann sum* associated to f and (P, t) by

$$R(f, P, t) := \sum_{i=1}^n f(t_i)(x_i - x_{i-1}).$$

Note that the notation here includes f , P , and t . Also note that here we didn't demand f be bounded.

On the next slide, we state two equivalent definitions of Riemann integrability.

Definition 25 (Riemann 1)

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable*

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$,

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for **all** tagged partitions (P, t)

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for all tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for all tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

A function $f : [a, b] \rightarrow \mathbb{R}$

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for all tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable*

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for all tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for all tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$,

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for **all** tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ and a partition P such that

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for all tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ and a partition P such that

$$|R(f, P', t') - R| < \epsilon$$

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for all tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ and a partition P such that

$$|R(f, P', t') - R| < \epsilon$$

for all tagged refinements (P', t') of P

Definition 25 (Riemann 1)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$|R(f, P, t) - R| < \epsilon$$

for all tagged partitions (P, t) such that $\|P\| < \delta$.

Definition 26 (Riemann 2)

A function $f : [a, b] \rightarrow \mathbb{R}$ is said to be *Riemann integrable* if for there exists $R \in \mathbb{R}$ such that for every $\epsilon > 0$, there exists $\delta > 0$ and a partition P such that

$$|R(f, P', t') - R| < \epsilon$$

for all tagged refinements (P', t') of P with $\|P'\| < \delta$.

Definition 27

Theorem 15 (Darboux and Riemann are friends)

Definition 27

In both the definitions on the earlier slide, the R is unique and it is called the *Riemann integral* of f over $[a, b]$.

Theorem 15 (Darboux and Riemann are friends)

Definition 27

In both the definitions on the earlier slide, the R is unique and it is called the *Riemann integral* of f over $[a, b]$.

Theorem 15 (Darboux and Riemann are friends)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function.

Definition 27

In both the definitions on the earlier slide, the R is unique and it is called the *Riemann integral* of f over $[a, b]$.

Theorem 15 (Darboux and Riemann are friends)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function.

If f is bounded and Darboux integrable, then f is also Riemann integrable.

Definition 27

In both the definitions on the earlier slide, the R is unique and it is called the *Riemann integral* of f over $[a, b]$.

Theorem 15 (Darboux and Riemann are friends)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function.

If f is bounded and Darboux integrable, then f is also Riemann integrable.

If f is Riemann integrable, then f is bounded and also Darboux integrable.

Definition 27

In both the definitions on the earlier slide, the R is unique and it is called the *Riemann integral* of f over $[a, b]$.

Theorem 15 (Darboux and Riemann are friends)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a function.

If f is bounded and Darboux integrable, then f is also Riemann integrable.

If f is Riemann integrable, then f is bounded and also Darboux integrable.

In both the cases above, the Darboux and Riemann integrals are the same.

Theorem 16 (Riemann sums approximating the integral)

Theorem 16 (Riemann sums approximating the integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be **Riemann integrable**. Suppose that (P_n, t_n) is a sequence of tagged partitions of $[a, b]$ such that $\|P_n\| \rightarrow 0$.

Theorem 16 (Riemann sums approximating the integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be **Riemann integrable**. Suppose that (P_n, t_n) is a sequence of tagged partitions of $[a, b]$ such that $\|P_n\| \rightarrow 0$. Then,

Theorem 16 (Riemann sums approximating the integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be **Riemann integrable**. Suppose that (P_n, t_n) is a sequence of tagged partitions of $[a, b]$ such that $\|P_n\| \rightarrow 0$. Then,

$$\lim_{n \rightarrow \infty} R(f, P_n, t_n) = \int_a^b f(x)dx.$$

Theorem 16 (Riemann sums approximating the integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be **Riemann integrable**. Suppose that (P_n, t_n) is a sequence of tagged partitions of $[a, b]$ such that $\|P_n\| \rightarrow 0$. Then,

$$\lim_{n \rightarrow \infty} R(f, P_n, t_n) = \int_a^b f(x)dx.$$

Note that we assumed f to be Riemann integrable to begin with.

Theorem 16 (Riemann sums approximating the integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be **Riemann integrable**. Suppose that (P_n, t_n) is a sequence of tagged partitions of $[a, b]$ such that $\|P_n\| \rightarrow 0$. Then,

$$\lim_{n \rightarrow \infty} R(f, P_n, t_n) = \int_a^b f(x) dx.$$

Note that we assumed f to be Riemann integrable to begin with. Thus, we cannot use the above theorem if we don't already know that f is Riemann integrable.

Theorem 16 (Riemann sums approximating the integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be **Riemann integrable**. Suppose that (P_n, t_n) is a sequence of tagged partitions of $[a, b]$ such that $\|P_n\| \rightarrow 0$. Then,

$$\lim_{n \rightarrow \infty} R(f, P_n, t_n) = \int_a^b f(x)dx.$$

Note that we assumed f to be Riemann integrable to begin with. Thus, we cannot use the above theorem if we don't already know that f is Riemann integrable. The next theorem helps us in determining when that happens.

Theorem 16 (Riemann sums approximating the integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be Riemann integrable. Suppose that (P_n, t_n) is a sequence of tagged partitions of $[a, b]$ such that $\|P_n\| \rightarrow 0$. Then,

$$\lim_{n \rightarrow \infty} R(f, P_n, t_n) = \int_a^b f(x) dx.$$

Note that we assumed f to be Riemann integrable to begin with. Thus, we cannot use the above theorem if we don't already know that f is Riemann integrable. The next theorem helps us in determining when that happens.

Theorem 17

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous.

Theorem 16 (Riemann sums approximating the integral)

Let $f : [a, b] \rightarrow \mathbb{R}$ be Riemann integrable. Suppose that (P_n, t_n) is a sequence of tagged partitions of $[a, b]$ such that $\|P_n\| \rightarrow 0$. Then,

$$\lim_{n \rightarrow \infty} R(f, P_n, t_n) = \int_a^b f(x) dx.$$

Note that we assumed f to be Riemann integrable to begin with. Thus, we cannot use the above theorem if we don't already know that f is Riemann integrable. The next theorem helps us in determining when that happens.

Theorem 17

Let $f : [a, b] \rightarrow \mathbb{R}$ be continuous. Then, f is Riemann integrable.

The converse of the previous theorem is not true.

The converse of the previous theorem is not true. In fact, the theorem is true even if we assume something less.

The converse of the previous theorem is not true. In fact, the theorem is true even if we assume something less. Namely, if f is bounded and is discontinuous on a finite set, then it is Riemann integrable.

The converse of the previous theorem is not true. In fact, the theorem is true even if we assume something less. Namely, if f is bounded and is discontinuous on a finite set, then it is Riemann integrable. The “finite” can even be replaced with “at most countable,” if you know what that means.

The converse of the previous theorem is not true. In fact, the theorem is true even if we assume something less. Namely, if f is bounded and is discontinuous on a finite set, then it is Riemann integrable. The “finite” can even be replaced with “at most countable,” if you know what that means.

The “at most countable” can actually be replaced with “measure zero.”

The converse of the previous theorem is not true. In fact, the theorem is true even if we assume something less. Namely, if f is bounded and is discontinuous on a finite set, then it is Riemann integrable. The “finite” can even be replaced with “at most countable,” if you know what that means.

The “at most countable” can actually be replaced with “measure zero.” At this point, the converse also becomes true!

The converse of the previous theorem is not true. In fact, the theorem is true even if we assume something less. Namely, if f is bounded and is discontinuous on a finite set, then it is Riemann integrable. The “finite” can even be replaced with “at most countable,” if you know what that means.

The “at most countable” can actually be replaced with “measure zero.” At this point, the converse also becomes true!

Now, we see how derivatives and integrals relate. These are the two parts of the Fundamental Theorem of Calculus.

Theorem 18 (FTC Part I)

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function,

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

for $x \in [a, b]$.

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

for $x \in [a, b]$.

Then, F is continuous.

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

for $x \in [a, b]$.

Then, F is continuous. Moreover, if f is continuous at some $c \in (a, b)$,

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

for $x \in [a, b]$.

Then, F is continuous. Moreover, if f is continuous at some $c \in (a, b)$, then F is differentiable at c and

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

for $x \in [a, b]$.

Then, F is continuous. Moreover, if f is continuous at some $c \in (a, b)$, then F is differentiable at c and

$$F'(c) = f(c).$$

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

for $x \in [a, b]$.

Then, F is continuous. Moreover, if f is continuous at some $c \in (a, b)$, then F is differentiable at c and

$$F'(c) = f(c).$$

In particular, if f is continuous,

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

for $x \in [a, b]$.

Then, F is continuous. Moreover, if f is continuous at some $c \in (a, b)$, then F is differentiable at c and

$$F'(c) = f(c).$$

In particular, if f is continuous, then Riemann integrability of f is guaranteed

Theorem 18 (FTC Part I)

Let $f : [a, b] \rightarrow \mathbb{R}$ be a Riemann integrable function, and let

$$F(x) := \int_a^x f(t)dt$$

for $x \in [a, b]$.

Then, F is continuous. Moreover, if f is continuous at some $c \in (a, b)$, then F is differentiable at c and

$$F'(c) = f(c).$$

In particular, if f is continuous, then Riemann integrability of f is guaranteed and the above equation is true for *all* $c \in (a, b)$.

Theorem 19 (FTC Part II)

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$ which is differentiable on (a, b)

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$ which is differentiable on (a, b) and satisfies $F' = f$ on (a, b) .

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$ which is differentiable on (a, b) and satisfies $F' = f$ on (a, b) . If f is Riemann integrable on $[a, b]$,

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$ which is differentiable on (a, b) and satisfies $F' = f$ on (a, b) . If f is Riemann integrable on $[a, b]$, then

$$\int_a^b f(t)dt = F(b) - F(a).$$

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$ which is differentiable on (a, b) and satisfies $F' = f$ on (a, b) . If f is Riemann integrable on $[a, b]$, then

$$\int_a^b f(t)dt = F(b) - F(a).$$

Note that the if is crucial.

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$ which is differentiable on (a, b) and satisfies $F' = f$ on (a, b) . If f is Riemann integrable on $[a, b]$, then

$$\int_a^b f(t)dt = F(b) - F(a).$$

Note that the if is crucial. It isn't necessary that the derivative of a function is Riemann integrable.

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$ which is differentiable on (a, b) and satisfies $F' = f$ on (a, b) . If f is Riemann integrable on $[a, b]$, then

$$\int_a^b f(t)dt = F(b) - F(a).$$

Note that the if is crucial. It isn't necessary that the derivative of a function is Riemann integrable. It needn't even be bounded.

Theorem 19 (FTC Part II)

Let $f : [a, b] \rightarrow \mathbb{R}$ be given and suppose there exists a continuous function $F : [a, b] \rightarrow \mathbb{R}$ which is differentiable on (a, b) and satisfies $F' = f$ on (a, b) . If f is Riemann integrable on $[a, b]$, then

$$\int_a^b f(t)dt = F(b) - F(a).$$

Note that the if is crucial. It isn't necessary that the derivative of a function is Riemann integrable. It needn't even be bounded. (But even if it is bounded, it needn't be Riemann integrable. Although an example of this is harder.)

Some pathological remarks:

- ① If a function is Riemann integrable, it doesn't mean that it is the derivative of a function. (That is, it needn't have an anti-derivative.)
- ② If a function has an anti-derivative, it doesn't mean that it is Riemann integrable. (That is, derivatives needn't be Riemann integrable.)

For the first, take $f : [0, 2] \rightarrow \mathbb{R}$ defined by $f(x) = \lfloor x \rfloor$. It cannot be the derivative of any function because it doesn't have IVP.
(Recall Theorem 8.)

For the second, consider the derivative of $F : [-1, 1] \rightarrow \mathbb{R}$ defined by $F(x) = x^2 \sin(1/x^2)$ for $x \neq 0$ and $F(0) = 0$. F' here isn't bounded.

Start recording!

Definition 28 (Limits)

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$.

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$. We write

$$\lim_{x \rightarrow c} f(x) = L$$

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$. We write

$$\lim_{x \rightarrow c} f(x) = L$$

if for every $\epsilon > 0$,

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$. We write

$$\lim_{x \rightarrow c} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$. We write

$$\lim_{x \rightarrow c} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$

$$\|f(x) - L\| < \epsilon$$

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$. We write

$$\lim_{x \rightarrow c} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$

$$\|f(x) - L\| < \epsilon$$

for all $x \in U$

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$. We write

$$\lim_{x \rightarrow c} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$\|f(x) - L\| < \epsilon$$

for all $x \in U$ such that $0 < \|x - c\| < \delta$.

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$. We write

$$\lim_{x \rightarrow c} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$\|f(x) - L\| < \epsilon$$

for all $x \in U$ such that $0 < \|x - c\| < \delta$.

Note that if $m = 1$, then $\|f(x) - L\|$ is just $|f(x) - L|$.

In whatever follows, $n, m \geq 1$ and U will be a subset of \mathbb{R}^n .

Definition 28 (Limits)

Let $f : U \rightarrow \mathbb{R}^m$ be a function and $c \in \mathbb{R}^n$ be a limit point of U .

Let $L \in \mathbb{R}^m$. We write

$$\lim_{x \rightarrow c} f(x) = L$$

if for every $\epsilon > 0$, there exists $\delta > 0$ such that

$$\|f(x) - L\| < \epsilon$$

for all $x \in U$ such that $0 < \|x - c\| < \delta$.

Note that if $m = 1$, then $\|f(x) - L\|$ is just $|f(x) - L|$. In fact, for $n = m = 1$, the definition above coincides with the earlier one.
(Definition 7.)

In the previous slide, we used the phrase “limit point.”

In the previous slide, we used the phrase “limit point.” The following is the definition.

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$.

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$. c is said to be a *limit point* of U

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$. c is said to be a *limit point* of U if for every $\delta > 0$,

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$. c is said to be a *limit point* of U if for every $\delta > 0$, there exists $x \in U$ such that

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$. c is said to be a *limit point* of U if for every $\delta > 0$, there exists $x \in U$ such that $0 < \|x - c\| < \delta$.

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$. c is said to be a *limit point* of U if for every $\delta > 0$, there exists $x \in U$ such that $0 < \|x - c\| < \delta$.

Note that a limit point of U can lie outside U .

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$. c is said to be a *limit point* of U if for every $\delta > 0$, there exists $x \in U$ such that $0 < \|x - c\| < \delta$.

Note that a limit point of U can lie outside U . Conversely, a point in U could still fail to be a limit point of U .

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$. c is said to be a *limit point* of U if for every $\delta > 0$, there exists $x \in U$ such that $0 < \|x - c\| < \delta$.

Note that a limit point of U can lie outside U . Conversely, a point in U could still fail to be a limit point of U .

To see this, consider $n = 1$ and $U = [0, 1) \cup \{2\}$.

In the previous slide, we used the phrase “limit point.” The following is the definition.

Definition 29 (Limit point)

Let $U \subset \mathbb{R}^n$ and $c \in \mathbb{R}^n$. c is said to be a *limit point* of U if for every $\delta > 0$, there exists $x \in U$ such that $0 < \|x - c\| < \delta$.

Note that a limit point of U can lie outside U . Conversely, a point in U could still fail to be a limit point of U .

To see this, consider $n = 1$ and $U = [0, 1) \cup \{2\}$. Then, 1 is a limit point of U while 2 is not.

As before, we can now define continuity easily.

Definition 30 (Continuity)

As before, we can now define continuity easily.

Definition 30 (Continuity)

If $f : U \rightarrow \mathbb{R}^m$ is a function

As before, we can now define continuity easily.

Definition 30 (Continuity)

If $f : U \rightarrow \mathbb{R}^m$ is a function and $c \in U$,

As before, we can now define continuity easily.

Definition 30 (Continuity)

If $f : U \rightarrow \mathbb{R}^m$ is a function and $c \in U$, then f is said to be

As before, we can now define continuity easily.

Definition 30 (Continuity)

If $f : U \rightarrow \mathbb{R}^m$ is a function and $c \in U$, then f is said to be *continuous at the point c*

As before, we can now define continuity easily.

Definition 30 (Continuity)

If $f : U \rightarrow \mathbb{R}^m$ is a function and $c \in U$, then f is said to be *continuous at the point c* if (and only if)

$$\lim_{x \rightarrow c} f(x) = f(c).$$

As before, we can now define continuity easily.

Definition 30 (Continuity)

If $f : U \rightarrow \mathbb{R}^m$ is a function and $c \in U$, then f is said to be *continuous at the point c* if (and only if)

$$\lim_{x \rightarrow c} f(x) = f(c).$$

As before, the case $n = m = 1$ recovers the original one.

Now, let us assume $n = 2$ and $m = 1$.

Definition 31 (Partial derivative)

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Let $f : U \rightarrow \mathbb{R}$ be a function

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Let $f : U \rightarrow \mathbb{R}$ be a function and $(a, b) \in U$ an interior point of U .

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Let $f : U \rightarrow \mathbb{R}$ be a function and $(a, b) \in U$ an interior point of U . The *partial derivative of f at (a, b) with respect to x_1* is defined by

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Let $f : U \rightarrow \mathbb{R}$ be a function and $(a, b) \in U$ an interior point of U . The *partial derivative of f at (a, b) with respect to x_1* is defined by

$$\frac{\partial f}{\partial x_1}(a, b) := \lim_{x_1 \rightarrow a} \frac{f(x_1, b) - f(a, b)}{x_1 - a},$$

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Let $f : U \rightarrow \mathbb{R}$ be a function and $(a, b) \in U$ an interior point of U . The *partial derivative of f at (a, b) with respect to x_1* is defined by

$$\frac{\partial f}{\partial x_1}(a, b) := \lim_{x_1 \rightarrow a} \frac{f(x_1, b) - f(a, b)}{x_1 - a},$$

provided that the limit exists.

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Let $f : U \rightarrow \mathbb{R}$ be a function and $(a, b) \in U$ an interior point of U . The *partial derivative of f at (a, b) with respect to x_1* is defined by

$$\frac{\partial f}{\partial x_1}(a, b) := \lim_{x_1 \rightarrow a} \frac{f(x_1, b) - f(a, b)}{x_1 - a},$$

provided that the limit exists.

The partial derivative with respect to x_2 is defined similarly.

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Let $f : U \rightarrow \mathbb{R}$ be a function and $(a, b) \in U$ an interior point of U . The *partial derivative of f at (a, b) with respect to x_1* is defined by

$$\frac{\partial f}{\partial x_1}(a, b) := \lim_{x_1 \rightarrow a} \frac{f(x_1, b) - f(a, b)}{x_1 - a},$$

provided that the limit exists.

The partial derivative with respect to x_2 is defined similarly.

Note that the limit above is an ordinary one-variable limit of a real function, as we had seen earlier.

Now, let us assume $n = 2$ and $m = 1$. That is, $U \subset \mathbb{R}^2$ and we look at functions of the form $f : U \rightarrow \mathbb{R}$.

Definition 31 (Partial derivative)

Let $f : U \rightarrow \mathbb{R}$ be a function and $(a, b) \in U$ an interior point of U . The *partial derivative of f at (a, b) with respect to x_1* is defined by

$$\frac{\partial f}{\partial x_1}(a, b) := \lim_{x_1 \rightarrow a} \frac{f(x_1, b) - f(a, b)}{x_1 - a},$$

provided that the limit exists.

The partial derivative with respect to x_2 is defined similarly.

Note that the limit above is an ordinary one-variable limit of a real function, as we had seen earlier. Also note that b is fixed in the numerator.

In the previous slide, we used the phrase “interior point.”

In the previous slide, we used the phrase “interior point.” The following is the definition.

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$ and $c \in U$.

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$ and $c \in U$. c is said to be an *interior point* of U

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$ and $c \in U$. c is said to be an *interior point* of U if there exists $\delta > 0$

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$ and $c \in U$. c is said to be an *interior point* of U if there exists $\delta > 0$ such that for every $x \in \mathbb{R}^n$ with $\|x - c\| < \delta$,

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$ and $c \in U$. c is said to be an *interior point* of U if there exists $\delta > 0$ such that for every $x \in \mathbb{R}^n$ with $\|x - c\| < \delta$, we have $x \in U$.

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$ and $c \in U$. c is said to be an *interior point* of U if there exists $\delta > 0$ such that for every $x \in \mathbb{R}^n$ with $\|x - c\| < \delta$, we have $x \in U$.

The above says that not only is $c \in U$ but also that there is a “ball” around c contained in U .

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$ and $c \in U$. c is said to be an *interior point* of U if there exists $\delta > 0$ such that for every $x \in \mathbb{R}^n$ with $\|x - c\| < \delta$, we have $x \in U$.

The above says that not only is $c \in U$ but also that there is a “ball” around c contained in U .

Note that the above says “there exists” and not “for every.”

In the previous slide, we used the phrase “interior point.” The following is the definition.

Definition 32 (Interior point)

Let $U \subset \mathbb{R}^n$ and $c \in U$. c is said to be an *interior point* of U if there exists $\delta > 0$ such that for every $x \in \mathbb{R}^n$ with $\|x - c\| < \delta$, we have $x \in U$.

The above says that not only is $c \in U$ but also that there is a “ball” around c contained in U .

Note that the above says “there exists” and not “for every.” Compare this with the definition of “limit point.”

Definition 33 (Directional derivatives)

Definition 33 (Directional derivatives)

Let $f : U \rightarrow \mathbb{R}$ be a function

Definition 33 (Directional derivatives)

Let $f : U \rightarrow \mathbb{R}$ be a function and $v = (v_1, v_2) \in \mathbb{R}^2$ a **unit vector**.

Definition 33 (Directional derivatives)

Let $f : U \rightarrow \mathbb{R}$ be a function and $v = (v_1, v_2) \in \mathbb{R}^2$ a **unit vector**.
Let $x = (x_1, x_2) \in U$ be an interior point of U .

Definition 33 (Directional derivatives)

Let $f : U \rightarrow \mathbb{R}$ be a function and $v = (v_1, v_2) \in \mathbb{R}^2$ a **unit vector**.

Let $x = (x_1, x_2) \in U$ be an interior point of U . The *directional derivative of f in the direction v at x* is defined as

Definition 33 (Directional derivatives)

Let $f : U \rightarrow \mathbb{R}$ be a function and $v = (v_1, v_2) \in \mathbb{R}^2$ a **unit vector**.

Let $x = (x_1, x_2) \in U$ be an interior point of U . The *directional derivative of f in the direction v at x* is defined as

$$\nabla_v f(x) = \lim_{t \rightarrow 0} \frac{f(x_1 + tv_1, x_2 + tv_2) - f(x_1, x_2)}{t},$$

provided it exists.

Definition 33 (Directional derivatives)

Let $f : U \rightarrow \mathbb{R}$ be a function and $v = (v_1, v_2) \in \mathbb{R}^2$ a **unit vector**. Let $x = (x_1, x_2) \in U$ be an interior point of U . The *directional derivative of f in the direction v at x* is defined as

$$\nabla_v f(x) = \lim_{t \rightarrow 0} \frac{f(x_1 + tv_1, x_2 + tv_2) - f(x_1, x_2)}{t},$$

provided it exists.

As before, this is an ordinary limit.

Definition 33 (Directional derivatives)

Let $f : U \rightarrow \mathbb{R}$ be a function and $v = (v_1, v_2) \in \mathbb{R}^2$ a **unit vector**.

Let $x = (x_1, x_2) \in U$ be an interior point of U . The *directional derivative of f in the direction v at x* is defined as

$$\nabla_v f(x) = \lim_{t \rightarrow 0} \frac{f(x_1 + tv_1, x_2 + tv_2) - f(x_1, x_2)}{t},$$

provided it exists.

As before, this is an ordinary limit. Taking $v = (1, 0)$ and $(0, 1)$ recovers the usual the partial derivatives with respect to x_1 and x_2 , respectively.

Definition 34 (Differentiability)

Definition 34 (Differentiability)

Let $f : U \rightarrow \mathbb{R}$ be a function and (x_0, y_0) be an interior point of U .

Definition 34 (Differentiability)

Let $f : U \rightarrow \mathbb{R}$ be a function and (x_0, y_0) be an interior point of U .
 f is said to be *differentiable at (x_0, y_0)*

Definition 34 (Differentiability)

Let $f : U \rightarrow \mathbb{R}$ be a function and (x_0, y_0) be an interior point of U .
 f is said to be *differentiable at (x_0, y_0)* if there exists a 1×2 matrix A

Definition 34 (Differentiability)

Let $f : U \rightarrow \mathbb{R}$ be a function and (x_0, y_0) be an interior point of U . f is said to be *differentiable at (x_0, y_0)* if there exists a 1×2 matrix A such that

$$\lim_{(h,k) \rightarrow (0,0)} \frac{\left| f(x_0 + h, y_0 + k) - f(x_0, y_0) - A \begin{bmatrix} h \\ k \end{bmatrix} \right|}{\|(h, k)\|} = 0.$$

Definition 34 (Differentiability)

Let $f : U \rightarrow \mathbb{R}$ be a function and (x_0, y_0) be an interior point of U . f is said to be *differentiable at (x_0, y_0)* if there exists a 1×2 matrix A such that

$$\lim_{(h,k) \rightarrow (0,0)} \frac{\left| f(x_0 + h, y_0 + k) - f(x_0, y_0) - A \begin{bmatrix} h \\ k \end{bmatrix} \right|}{\|(h, k)\|} = 0.$$

In this case, we write $Df(x_0, y_0) = A$

Definition 34 (Differentiability)

Let $f : U \rightarrow \mathbb{R}$ be a function and (x_0, y_0) be an interior point of U . f is said to be *differentiable at (x_0, y_0)* if there exists a 1×2 matrix A such that

$$\lim_{(h,k) \rightarrow (0,0)} \frac{\left| f(x_0 + h, y_0 + k) - f(x_0, y_0) - A \begin{bmatrix} h \\ k \end{bmatrix} \right|}{\|(h, k)\|} = 0.$$

In this case, we write $Df(x_0, y_0) = A$ and call A the *total derivative* of f at (x_0, y_0) .

In your slides, we had seen originally seen a different definition.

Theorem 20

In your slides, we had seen originally seen a different definition.
That happens to be equivalent to the above.

Theorem 20

In your slides, we had seen originally seen a different definition.
That happens to be equivalent to the above. We have the
following theorem.

Theorem 20

In your slides, we had seen originally seen a different definition. That happens to be equivalent to the above. We have the following theorem. We keep the same notations as earlier.

Theorem 20

In your slides, we had seen originally seen a different definition. That happens to be equivalent to the above. We have the following theorem. We keep the same notations as earlier.

Theorem 20

Suppose that f is differentiable at (x_0, y_0) .

In your slides, we had seen originally seen a different definition. That happens to be equivalent to the above. We have the following theorem. We keep the same notations as earlier.

Theorem 20

Suppose that f is differentiable at (x_0, y_0) . Then, both the partial derivatives of f at (x_0, y_0) exist

In your slides, we had seen originally seen a different definition. That happens to be equivalent to the above. We have the following theorem. We keep the same notations as earlier.

Theorem 20

Suppose that f is differentiable at (x_0, y_0) . Then, both the partial derivatives of f at (x_0, y_0) exist and

$$Df(x_0, y_0) = \begin{bmatrix} \frac{\partial f}{\partial x_1}(x_0, y_0) & \frac{\partial f}{\partial x_2}(x_0, y_0) \end{bmatrix}.$$

In your slides, we had seen originally seen a different definition. That happens to be equivalent to the above. We have the following theorem. We keep the same notations as earlier.

Theorem 20

Suppose that f is differentiable at (x_0, y_0) . Then, both the partial derivatives of f at (x_0, y_0) exist and

$$Df(x_0, y_0) = \begin{bmatrix} \frac{\partial f}{\partial x_1}(x_0, y_0) & \frac{\partial f}{\partial x_2}(x_0, y_0) \end{bmatrix}.$$

The above matrix is also called the *gradient* and denoted by $\nabla f(x_0, y_0)$.

Stop recording. Start a new one.
Take doubts.