

# Complex Analysis TSC - 1

Aryaman Maithani

<https://aryamanmaithani.github.io/tuts/ma-205>

IIT Bombay

18th September, 2020

Hi,

Hi, welcome to this

Hi, welcome to this

*complex*

Hi, welcome to this

*complex* discussion.

Hi, welcome to this

*complex* discussion.

Here are some “guidelines” for this TSC -

Hi, welcome to this

*complex* discussion.

Here are some “guidelines” for this TSC -

- 1 Unmute your mic at any time and ask your doubt.

Hi, welcome to this

*complex* discussion.

Here are some “guidelines” for this TSC -

- 1 Unmute your mic at any time and ask your doubt.
- 2 I will not be checking chat often (or maybe at all), so posting it there might not be helpful.



Hi, welcome to this

*complex* discussion.

Here are some “guidelines” for this TSC -

- 1 Unmute your mic at any time and ask your doubt.
- 2 I will not be checking chat often (or maybe at all), so posting it there might not be helpful.

You can find a link to this document on [bit.ly/ma-205](https://bit.ly/ma-205). Both with and without pauses. You may keep it open alongside for quick reference.

# Warning

This is primarily going to be a quick recap of the facts important.

# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

In particular, this will *not* be a substitute for all the lectures done so far.

# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

In particular, this will *not* be a substitute for all the lectures done so far.

I will also not be going through the proofs.

# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

In particular, this will *not* be a substitute for all the lectures done so far.

I will also not be going through the proofs. We can discuss these finer things at the end, if time permits.

# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

In particular, this will *not* be a substitute for all the lectures done so far.

I will also not be going through the proofs. We can discuss these finer things at the end, if time permits.

Though I'm not a fan of this - this session is pretty much going to cover things important from the point of view of an exam.

# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

In particular, this will *not* be a substitute for all the lectures done so far.

I will also not be going through the proofs. We can discuss these finer things at the end, if time permits.

Though I'm not a fan of this - this session is pretty much going to cover things important from the point of view of an exam. I may also skip things from the lectures if I think that they are not important.



# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

In particular, this will *not* be a substitute for all the lectures done so far.

I will also not be going through the proofs. We can discuss these finer things at the end, if time permits.

Though I'm not a fan of this - this session is pretty much going to cover things important from the point of view of an exam. I may also skip things from the lectures if I think that they are not important. They *might* turn out to be important, though.

# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

In particular, this will *not* be a substitute for all the lectures done so far.

I will also not be going through the proofs. We can discuss these finer things at the end, if time permits.

Though I'm not a fan of this - this session is pretty much going to cover things important from the point of view of an exam. I may also skip things from the lectures if I think that they are not important. They *might* turn out to be important, though.

Of course, I will not  
mathematically incorrect.

say anything which is

# Warning

This is primarily going to be a quick recap of the facts important. It is, of course, not possible to go through everything in just 2 hours.

In particular, this will *not* be a substitute for all the lectures done so far.

I will also not be going through the proofs. We can discuss these finer things at the end, if time permits.

Though I'm not a fan of this - this session is pretty much going to cover things important from the point of view of an exam. I may also skip things from the lectures if I think that they are not important. They *might* turn out to be important, though.

Of course, I will not (intentionally) say anything which is mathematically incorrect.

# Lecture 1

## Definition 1 (Some notation)

Given  $z_0 \in \mathbb{C}$  and  $\delta > 0$ , the  $\delta$ -neighbourhood of  $z_0$ , denoted by  $B_\delta(z_0)$  is the set

$$B_\delta(z_0) := \{z \in \mathbb{C} : |z - z_0| < \delta\}.$$

# Lecture 1

## Definition 1 (Some notation)

Given  $z_0 \in \mathbb{C}$  and  $\delta > 0$ , the  $\delta$ -neighbourhood of  $z_0$ , denoted by  $B_\delta(z_0)$  is the set

$$B_\delta(z_0) := \{z \in \mathbb{C} : |z - z_0| < \delta\}.$$

## Definition 2 (Open sets)

A set  $U \subset \mathbb{C}$  is said to be open if:  
for every  $z_0 \in \mathbb{C}$ , there exists *some*  $\delta > 0$  such that

$$B_\delta(z_0) \subset U.$$

# Lecture 1

## Definition 1 (Some notation)

Given  $z_0 \in \mathbb{C}$  and  $\delta > 0$ , the  $\delta$ -neighbourhood of  $z_0$ , denoted by  $B_\delta(z_0)$  is the set

$$B_\delta(z_0) := \{z \in \mathbb{C} : |z - z_0| < \delta\}.$$

## Definition 2 (Open sets)

A set  $U \subset \mathbb{C}$  is said to be open if:  
for every  $z_0 \in \mathbb{C}$ , there exists *some*  $\delta > 0$  such that

$$B_\delta(z_0) \subset U.$$

## Definition 3 (Path-connected sets)

A set  $P \subset \mathbb{C}$  is said to be path-connected if any two points in  $P$  can be joined by a path in  $P$ . (A continuous function from  $[0, 1]$  to  $P$ .)

## Definition 4 (Differentiable)

Let  $\Omega \subset \mathbb{C}$  be open. Let

$$f : \Omega \rightarrow \mathbb{C}$$

be a function.

## Definition 4 (Differentiable)

Let  $\Omega \subset \mathbb{C}$  be **open**. Let

$$f : \Omega \rightarrow \mathbb{C}$$

be a function.



## Definition 4 (Differentiable)

Let  $\Omega \subset \mathbb{C}$  be **open**. Let

$$f : \Omega \rightarrow \mathbb{C}$$

be a function. Let  $z_0 \in \Omega$ .

## Definition 4 (Differentiable)

Let  $\Omega \subset \mathbb{C}$  be **open**. Let

$$f : \Omega \rightarrow \mathbb{C}$$

be a function. Let  $z_0 \in \Omega$ .  $f$  is said to be *differentiable* at  $z_0$  if

## Definition 4 (Differentiable)

Let  $\Omega \subset \mathbb{C}$  be **open**. Let

$$f : \Omega \rightarrow \mathbb{C}$$

be a function. Let  $z_0 \in \Omega$ .  $f$  is said to be *differentiable* at  $z_0$  if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

## Definition 4 (Differentiable)

Let  $\Omega \subset \mathbb{C}$  be **open**. Let

$$f : \Omega \rightarrow \mathbb{C}$$

be a function. Let  $z_0 \in \Omega$ .  $f$  is said to be *differentiable* at  $z_0$  if

$$\lim_{z \rightarrow z_0} \frac{f(z) - f(z_0)}{z - z_0}$$

exists. In this case, it is denoted by  $f'(z_0)$ .

## Definition 5 (Holomorphic)

A function  $f$  is said to be holomorphic on an open set  $\Omega$  if it is differentiable at every  $z_0 \in \Omega$ .

A function  $f$  is said to be holomorphic at  $z_0$  if it is holomorphic on some neighbourhood of  $z_0$ .

## Definition 5 (Holomorphic)

A function  $f$  is said to be holomorphic on an open set  $\Omega$  if it is differentiable at every  $z_0 \in \Omega$ .

A function  $f$  is said to be holomorphic at  $z_0$  if it is holomorphic on some neighbourhood of  $z_0$ .

## Remark 1

A function may be differentiable at  $z_0$  but not holomorphic at  $z_0$ .

## Definition 5 (Holomorphic)

A function  $f$  is said to be holomorphic on an open set  $\Omega$  if it is differentiable at every  $z_0 \in \Omega$ .

A function  $f$  is said to be holomorphic at  $z_0$  if it is holomorphic on some neighbourhood of  $z_0$ .

## Remark 1

A function may be differentiable at  $z_0$  but not holomorphic at  $z_0$ . For example,  $f(z) = |z|^2$  is differentiable only at 0.

## Definition 5 (Holomorphic)

A function  $f$  is said to be holomorphic on an open set  $\Omega$  if it is differentiable at every  $z_0 \in \Omega$ .

A function  $f$  is said to be holomorphic at  $z_0$  if it is holomorphic on some neighbourhood of  $z_0$ .

## Remark 1

A function may be differentiable at  $z_0$  but not holomorphic at  $z_0$ . For example,  $f(z) = |z|^2$  is differentiable only at 0. Thus, it is differentiable at 0 but holomorphic nowhere.



## Definition 5 (Holomorphic)

A function  $f$  is said to be holomorphic on an open set  $\Omega$  if it is differentiable at every  $z_0 \in \Omega$ .

A function  $f$  is said to be holomorphic at  $z_0$  if it is holomorphic on some neighbourhood of  $z_0$ .

## Remark 1

A function may be differentiable at  $z_0$  but not holomorphic at  $z_0$ . For example,  $f(z) = |z|^2$  is differentiable only at 0. Thus, it is differentiable at 0 but holomorphic nowhere.

For sets, however, there is no difference.

# End of Lecture 1

Any questions?

From this point on,  $\Omega$  will always denote an open subset of  $\mathbb{C}$ .  
Whenever I write some complex number  $z$  as  $z = x + iy$ , it will be assumed that  $x, y \in \mathbb{R}$ .  
Similarly for  $f(z) = u(z) + iv(z)$ .

## Lecture 2: CR Equations

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. We can decompose  $f$  as

$$f(z) = u(z) + \iota v(z),$$

where  $u, v : \Omega \rightarrow \mathbb{R}$  are real valued functions.

## Lecture 2: CR Equations

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. We can decompose  $f$  as

$$f(z) = u(z) + \iota v(z),$$

where  $u, v : \Omega \rightarrow \mathbb{R}$  are real valued functions.

The idea now is to consider  $u$  and  $v$  as functions of two variables. We can do so by simply considering  $u(x, y) = u(x + \iota y)$  and similarly for  $v$ .

# Lecture 2: CR Equations

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. We can decompose  $f$  as

$$f(z) = u(z) + \iota v(z),$$

where  $u, v : \Omega \rightarrow \mathbb{R}$  are real valued functions.

The idea now is to consider  $u$  and  $v$  as functions of two variables. We can do so by simply considering  $u(x, y) = u(x + \iota y)$  and similarly for  $v$ . Now, if we know that  $f$  is holomorphic, then we have the following result.

# Lecture 2: CR Equations

## Theorem 1 (CR equations)

Let  $f : \Omega \rightarrow \mathbb{C}$  be differentiable at a point  $z_0 \in \Omega$ . Let  $z_0 = x_0 + iy_0$ .

# Lecture 2: CR Equations

## Theorem 1 (CR equations)

Let  $f : \Omega \rightarrow \mathbb{C}$  be differentiable at **a point**  $z_0 \in \Omega$ . Let

$z_0 = x_0 + iy_0$ .

Then, we have

$$u_x(x_0, y_0) = v_y(x_0, y_0) \quad \text{and} \quad u_y(x_0, y_0) = -v_x(x_0, y_0).$$



# Lecture 2: CR Equations

## Theorem 1 (CR equations)

Let  $f : \Omega \rightarrow \mathbb{C}$  be differentiable at a point  $z_0 \in \Omega$ . Let  $z_0 = x_0 + \iota y_0$ .

Then, we have

$$u_x(x_0, y_0) = v_y(x_0, y_0) \quad \text{and} \quad u_y(x_0, y_0) = -v_x(x_0, y_0).$$

Moreover, we have

$$f'(z_0) = u_x(x_0, y_0) + \iota v_x(x_0, y_0).$$

# Lecture 2: CR Equations

## Theorem 1 (CR equations)

Let  $f : \Omega \rightarrow \mathbb{C}$  be differentiable at a point  $z_0 \in \Omega$ . Let

$$z_0 = x_0 + \iota y_0.$$

Then, we have

$$u_x(x_0, y_0) = v_y(x_0, y_0) \quad \text{and} \quad u_y(x_0, y_0) = -v_x(x_0, y_0).$$

Moreover, we have

$$f'(z_0) = u_x(x_0, y_0) + \iota v_x(x_0, y_0).$$

Note the subscript is  $x$  for both in the above.

# Lecture 2: CR Equations

## Theorem 1 (CR equations)

Let  $f : \Omega \rightarrow \mathbb{C}$  be differentiable at a point  $z_0 \in \Omega$ . Let

$$z_0 = x_0 + \iota y_0.$$

Then, we have

$$u_x(x_0, y_0) = v_y(x_0, y_0) \quad \text{and} \quad u_y(x_0, y_0) = -v_x(x_0, y_0).$$

Moreover, we have

$$f'(z_0) = u_x(x_0, y_0) + \iota v_x(x_0, y_0).$$

Note the subscript is  $x$  for both in the above.

Also note that all the equalities are only at the point  $z_0$ .

# Lecture 2: CR Equations

## Theorem 1 (CR equations)

Let  $f : \Omega \rightarrow \mathbb{C}$  be differentiable at **a point**  $z_0 \in \Omega$ . Let

$$z_0 = x_0 + \iota y_0.$$

Then, we have

$$u_x(x_0, y_0) = v_y(x_0, y_0) \quad \text{and} \quad u_y(x_0, y_0) = -v_x(x_0, y_0).$$

Moreover, we have

$$f'(z_0) = u_x(x_0, y_0) + \iota v_x(x_0, y_0).$$

Note the subscript is  $x$  for both in the above.

Also note that all the equalities are only at **the point**  $z_0$ . In particular, we are only assuming differentiability at  $z_0$ .

# Lecture 2: CR Equations

Converse? What is the converse?

## Lecture 2: CR Equations

Converse? What is the converse? **No**. The converse is **not** true.

## Lecture 2: CR Equations

Converse? What is the converse? **No**. The converse is **not** true.  
An example for you to check is

$$f(z) := \begin{cases} \frac{\bar{z}^2}{z} & z \neq 0, \\ 0 & z = 0. \end{cases}$$

## Lecture 2: CR Equations

Converse? What is the converse? **No**. The converse is **not** true.  
An example for you to check is

$$f(z) := \begin{cases} \frac{\bar{z}^2}{z} & z \neq 0, \\ 0 & z = 0. \end{cases}$$

Check that  $u$  and  $v$  satisfy the CR equations at  $(0,0)$  but  $f$  is not differentiable at  $0 + 0i$ . (Page 23 of slides.)



## Lecture 2: CR Equations

We recall MA 105 now.

### Definition 6 (Total derivative)

If  $f : \Omega \rightarrow \mathbb{C}$  is a function, we may view it as a function

$$f : \Omega \rightarrow \mathbb{R}^2.$$

Recall that  $f$  is said to be real differentiable at  $(x_0, y_0) \in \Omega \subset \mathbb{R}^2$  if

## Lecture 2: CR Equations

We recall MA 105 now.

### Definition 6 (Total derivative)

If  $f : \Omega \rightarrow \mathbb{C}$  is a function, we may view it as a function

$$f : \Omega \rightarrow \mathbb{R}^2.$$

Recall that  $f$  is said to be real differentiable at  $(x_0, y_0) \in \Omega \subset \mathbb{R}^2$  if there exists a  $2 \times 2$  real matrix  $A$  such that

# Lecture 2: CR Equations

We recall MA 105 now.

## Definition 6 (Total derivative)

If  $f : \Omega \rightarrow \mathbb{C}$  is a function, we may view it as a function

$$f : \Omega \rightarrow \mathbb{R}^2.$$

Recall that  $f$  is said to be real differentiable at  $(x_0, y_0) \in \Omega \subset \mathbb{R}^2$  if there exists a  $2 \times 2$  real matrix  $A$  such that

$$\lim_{(h,k) \rightarrow (0,0)} \frac{\left\| f(x_0 + h, y_0 + k) - f(x_0, y_0) - A \begin{bmatrix} h \\ k \end{bmatrix} \right\|}{\|(h, k)\|} = 0.$$

## Lecture 2: CR Equations

We recall MA 105 now.

### Definition 6 (Total derivative)

If  $f : \Omega \rightarrow \mathbb{C}$  is a function, we may view it as a function

$$f : \Omega \rightarrow \mathbb{R}^2.$$

Recall that  $f$  is said to be real differentiable at  $(x_0, y_0) \in \Omega \subset \mathbb{R}^2$  if there exists a  $2 \times 2$  real matrix  $A$  such that

$$\lim_{(h,k) \rightarrow (0,0)} \frac{\left\| f(x_0 + h, y_0 + k) - f(x_0, y_0) - A \begin{bmatrix} h \\ k \end{bmatrix} \right\|}{\|(h, k)\|} = 0.$$

The matrix  $A$  was called the *total derivative of  $f$  at  $(x_0, y_0)$* .

# Lecture 2: CR Equations

## Theorem 2

If  $f$  is (complex) differentiable at a point  $z_0 = x_0 + iy_0$ , then  $f$  is real differentiable at  $(x_0, y_0)$ .

### Theorem 2

If  $f$  is (complex) differentiable at a point  $z_0 = x_0 + iy_0$ , then  $f$  is real differentiable at  $(x_0, y_0)$ .

Once again, this is only talking about differentiability at a point.

### Theorem 2

If  $f$  is (complex) differentiable at a point  $z_0 = x_0 + iy_0$ , then  $f$  is real differentiable at  $(x_0, y_0)$ .

Once again, this is only talking about differentiability at a point. The converse is again **not** true.

# Lecture 2: CR Equations

## Theorem 2

If  $f$  is (complex) differentiable at a point  $z_0 = x_0 + iy_0$ , then  $f$  is real differentiable at  $(x_0, y_0)$ .

Once again, this is only talking about differentiability at a point. The converse is again **not** true.

Thus, we have seen two sufficient conditions for complex differentiability so far. Neither is individually sufficient.



# Lecture 2: CR Equations

## Theorem 2

If  $f$  is (complex) differentiable at a point  $z_0 = x_0 + iy_0$ , then  $f$  is real differentiable at  $(x_0, y_0)$ .

Once again, this is only talking about differentiability at a point. The converse is again **not** true.

Thus, we have seen two sufficient conditions for complex differentiability so far. Neither is individually sufficient. However, together, they are.

## Theorem 3

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function and let  $z_0 \in \mathbb{C}$ . If

## Theorem 3

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function and let  $z_0 \in \mathbb{C}$ . If the CR equations hold **at the point**  $(x_0, y_0)$  and

## Theorem 3

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function and let  $z_0 \in \mathbb{C}$ . If the CR equations hold **at the point**  $(x_0, y_0)$  and if  $f$  is real differentiable **at the point**  $(x_0, y_0)$ , then

## Theorem 3

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function and let  $z_0 \in \mathbb{C}$ . If the CR equations hold **at the point**  $(x_0, y_0)$  and if  $f$  is real differentiable **at the point**  $(x_0, y_0)$ , then  $f$  is complex differentiable **at the point**  $z_0$ .

# Lecture 2: CR Equations

## Definition 7 (Harmonic functions)

Let  $u : \Omega \rightarrow \mathbb{R}$  be a twice continuously differentiable function.  $u$  is said to be *harmonic* if  $u_{xx} + u_{yy} = 0$ .

# Lecture 2: CR Equations

## Definition 7 (Harmonic functions)

Let  $u : \Omega \rightarrow \mathbb{R}$  be a twice continuously differentiable function.  $u$  is said to be *harmonic* if  $u_{xx} + u_{yy} = 0$ .

## Proposition 1

The real and imaginary parts of a holomorphic function are harmonic.

# Lecture 2: CR Equations

## Definition 7 (Harmonic functions)

Let  $u : \Omega \rightarrow \mathbb{R}$  be a twice continuously differentiable function.  $u$  is said to be *harmonic* if  $u_{xx} + u_{yy} = 0$ .

## Proposition 1

The real and imaginary parts of a holomorphic function are harmonic.

Suppose  $u$  and  $v$  are harmonic on  $\Omega$ .  $v$  is said to be a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic on  $\Omega$ .



# Lecture 2: CR Equations

## Definition 7 (Harmonic functions)

Let  $u : \Omega \rightarrow \mathbb{R}^2$  be a twice continuously differentiable function.  $u$  is said to be *harmonic* if  $u_{xx} + u_{yy} = 0$ .

## Proposition 1

The real and imaginary parts of a holomorphic function are harmonic.

Suppose  $u$  and  $v$  are harmonic on  $\Omega$ .  $v$  is said to be a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic on  $\Omega$ .

If  $v$  is a harmonic conjugate of  $u$ , then  $-u$  is a harmonic conjugate of  $v$ .

# Lecture 2: CR Equations

## Definition 7 (Harmonic functions)

Let  $u : \Omega \rightarrow \mathbb{R}$  be a twice continuously differentiable function.  $u$  is said to be *harmonic* if  $u_{xx} + u_{yy} = 0$ .

## Proposition 1

The real and imaginary parts of a holomorphic function are harmonic.

Suppose  $u$  and  $v$  are harmonic on  $\Omega$ .  $v$  is said to be a harmonic conjugate of  $u$  if  $f = u + iv$  is holomorphic on  $\Omega$ .

If  $v$  is a harmonic conjugate of  $u$ , then  $-u$  is a harmonic conjugate of  $v$ .

Check the second last slide of this lecture to find the algorithm for finding a harmonic conjugate.

# End of Lecture 2

Any questions?

# Lecture 3: Power Series

## Definition 8 (Convergence of series)

A series of the form

$$\sum_{n=0}^{\infty} a_n$$

of complex numbers is said to converge if the sequence of partial sums

$$s_n = \sum_{k=0}^n a_k$$

converges (to a finite complex number).

# Lecture 3: Power Series

## Definition 8 (Convergence of series)

A series of the form

$$\sum_{n=0}^{\infty} a_n$$

of complex numbers is said to converge if the sequence of partial sums

$$s_n = \sum_{k=0}^n a_k$$

converges (to a finite complex number).

The sequence of partial sums is just the following sequence:

$$a_0, a_0 + a_1, a_0 + a_1 + a_2, \dots$$

# Lecture 3: Power Series

## Definition 8 (Convergence of series)

A series of the form

$$\sum_{n=0}^{\infty} a_n$$

of complex numbers is said to converge if the sequence of partial sums

$$s_n = \sum_{k=0}^n a_k$$

converges (to a finite complex number).

The sequence of partial sums is just the following sequence:

$$a_0, a_0 + a_1, a_0 + a_1 + a_2, \dots$$

“Divergent” is simply used to mean “not convergent.”

# Lecture 3: Power Series

## Definition 8 (Convergence of series)

A series of the form

$$\sum_{n=0}^{\infty} a_n$$

of complex numbers is said to converge if the sequence of partial sums

$$s_n = \sum_{k=0}^n a_k$$

converges (to a finite complex number).

The sequence of partial sums is just the following sequence:

$$a_0, a_0 + a_1, a_0 + a_1 + a_2, \dots$$

“Divergent” is simply used to mean “not convergent.”

Check that  $\sum (-1)^n$  and  $\sum n$  both diverge.

# Lecture 3: Power Series

## Definition 9 (limsup)

Given a sequence  $(x_n)$  of **real numbers**, we may define a new sequence  $(y_n)$  as



## Definition 9 (limsup)

Given a sequence  $(x_n)$  of **real numbers**, we may define a new sequence  $(y_n)$  as

$$y_n = \sup\{x_m : m \geq n\}.$$

## Definition 9 (limsup)

Given a sequence  $(x_n)$  of **real numbers**, we may define a new sequence  $(y_n)$  as

$$y_n = \sup\{x_m : m \geq n\}.$$

The limit of this sequence always exists and we define

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n.$$

# Lecture 3: Power Series

## Definition 9 (limsup)

Given a sequence  $(x_n)$  of **real numbers**, we may define a new sequence  $(y_n)$  as

$$y_n = \sup\{x_m : m \geq n\}.$$

The limit of this sequence always exists and we define

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n.$$

## Remark 2

Each  $y_n$  might be  $\infty$ . That is allowed.

# Lecture 3: Power Series

## Definition 9 (limsup)

Given a sequence  $(x_n)$  of **real numbers**, we may define a new sequence  $(y_n)$  as

$$y_n = \sup\{x_m : m \geq n\}.$$

The limit of this sequence always exists and we define

$$\limsup_{n \rightarrow \infty} x_n = \lim_{n \rightarrow \infty} y_n.$$

## Remark 2

Each  $y_n$  might be  $\infty$ . That is allowed.

The limsup might be  $\infty$ . This is also allowed.

# Lecture 3: Power Series

We will be interested in discussing radius of convergence of *power series*. We all know what that is. It is a series of the form

$$\sum_{n=0}^{\infty} a_n (z - z_0)^n \quad (*)$$

where  $z_0 \in \mathbb{C}$  and each  $a_n \in \mathbb{C}$ .

# Lecture 3: Power Series

We will be interested in discussing radius of convergence of *power series*. We all know what that is. It is a series of the form

$$\sum_{n=0}^{\infty} a_n (z - z_0)^n \quad (*)$$

where  $z_0 \in \mathbb{C}$  and each  $a_n \in \mathbb{C}$ .

What is the radius of convergence, though?

# Lecture 3: Power Series

We will be interested in discussing radius of convergence of *power series*. We all know what that is. It is a series of the form

$$\sum_{n=0}^{\infty} a_n (z - z_0)^n \quad (*)$$

where  $z_0 \in \mathbb{C}$  and each  $a_n \in \mathbb{C}$ .

What is the radius of convergence, though? (The definition, that is.)

## Theorem 4 (Radius of convergence)

Given any power series as  $(*)$ , there exists  $R \in [0, \infty]$  such that

- 1  $(*)$  converges for any  $z$  with  $|z - z_0| < R$

This  $R$  is called the radius of convergence.

# Lecture 3: Power Series

We will be interested in discussing radius of convergence of *power series*. We all know what that is. It is a series of the form

$$\sum_{n=0}^{\infty} a_n(z - z_0)^n \quad (*)$$

where  $z_0 \in \mathbb{C}$  and each  $a_n \in \mathbb{C}$ .

What is the radius of convergence, though? (The definition, that is.)

## Theorem 4 (Radius of convergence)

Given any power series as  $(*)$ , there exists  $R \in [0, \infty]$  such that

- 1  $(*)$  converges for any  $z$  with  $|z - z_0| < R$ , and
- 2  $(*)$  diverges for any  $z$  with  $|z - z_0| > R$ .

This  $R$  is called the radius of convergence.



# Lecture 3: Power Series

We will be interested in discussing radius of convergence of *power series*. We all know what that is. It is a series of the form

$$\sum_{n=0}^{\infty} a_n(z - z_0)^n \quad (*)$$

where  $z_0 \in \mathbb{C}$  and each  $a_n \in \mathbb{C}$ .

What is the radius of convergence, though? (The definition, that is.)

## Theorem 4 (Radius of convergence)

Given any power series as  $(*)$ , there exists  $R \in [0, \infty]$  such that

- ①  $(*)$  converges for any  $z$  with  $|z - z_0| < R$ , and
- ②  $(*)$  diverges for any  $z$  with  $|z - z_0| > R$ .

This  $R$  is called the radius of convergence.

Note the **brackets**.

# Lecture 3: Power Series

We would now like to be able to calculate the radius of convergence.

## Theorem 5 (Root test)

# Lecture 3: Power Series

We would now like to be able to calculate the radius of convergence.

## Theorem 5 (Root test)

Let  $(*)$  be as earlier. Define

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n}.$$

# Lecture 3: Power Series

We would now like to be able to calculate the radius of convergence.

## Theorem 5 (Root test)

Let  $(*)$  be as earlier. Define

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n}.$$

Then,  $R = \alpha^{-1}$  is the radius of convergence.

# Lecture 3: Power Series

We would now like to be able to calculate the radius of convergence.

## Theorem 5 (Root test)

Let  $(*)$  be as earlier. Define

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n}.$$

Then,  $R = \alpha^{-1}$  is the radius of convergence.

This test *always works*. We had no assumptions of any kind on  $(*)$ .

# Lecture 3: Power Series

We would now like to be able to calculate the radius of convergence.

## Theorem 5 (Root test)

Let  $(*)$  be as earlier. Define

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n}.$$

Then,  $R = \alpha^{-1}$  is the radius of convergence.

This test *always works*. We had no assumptions of any kind on  $(*)$ . Note that  $^{-1}$ .

# Lecture 3: Power Series

We would now like to be able to calculate the radius of convergence.

## Theorem 5 (Root test)

Let  $(*)$  be as earlier. Define

$$\alpha = \limsup_{n \rightarrow \infty} \sqrt[n]{a_n}.$$

Then,  $R = \alpha^{-1}$  is the radius of convergence.

This test *always works*. We had no assumptions of any kind on  $(*)$ . Note that  $^{-1}$ .

If  $\alpha = 0$ , then  $R = \infty$  and vice-versa.

# Lecture 3: Power Series

We have another test. This is simpler (to calculate) but mightn't always work.

## Theorem 6 (Ratio test)

Let  $(*)$  be as earlier.



# Lecture 3: Power Series

We have another test. This is simpler (to calculate) but mightn't always work.

## Theorem 6 (Ratio test)

Let  $(*)$  be as earlier.

Assume that the limit

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$$

exists.

# Lecture 3: Power Series

We have another test. This is simpler (to calculate) but mightn't always work.

## Theorem 6 (Ratio test)

Let  $(*)$  be as earlier.

Assume that the limit

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$$

exists.

Then,  $R$  is the radius of convergence.

# Lecture 3: Power Series

We have another test. This is simpler (to calculate) but mightn't always work.

## Theorem 6 (Ratio test)

Let  $(*)$  be as earlier.

Assume that the limit

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$$

exists.

Then,  $R$  is the radius of convergence.

Note that here we assume that the limit does exist. This may not always be true.

# Lecture 3: Power Series

We have another test. This is simpler (to calculate) but mightn't always work.

## Theorem 6 (Ratio test)

Let  $(*)$  be as earlier.

Assume that the limit

$$R = \lim_{n \rightarrow \infty} \left| \frac{a_n}{a_{n+1}} \right|$$

exists.

Then,  $R$  is the radius of convergence.

Note that here we assume that the limit does exist. This may not always be true.

Note that I'm not taking any inverse here but also note the way the ratio is taken. We have  $a_n/a_{n+1}$ .

# Lecture 3: Power Series

Differentiability of power series is what one should expect.

## Theorem 7 (Differentiability)

Let  $\sum_{n=0}^{\infty} a_n z^n$  be a power series with radius of convergence  $R > 0$ . On the **open disc** of radius  $R$ , let  $f(z)$  denote this sum. Then, on this disc, we have

$$f'(z) =$$

# Lecture 3: Power Series

Differentiability of power series is what one should expect.

## Theorem 7 (Differentiability)

Let  $\sum_{n=0}^{\infty} a_n z^n$  be a power series with radius of convergence  $R > 0$ . On the **open disc** of radius  $R$ , let  $f(z)$  denote this sum. Then, on this disc, we have

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}.$$

# Lecture 3: Power Series

Differentiability of power series is what one should expect.

## Theorem 7 (Differentiability)

Let  $\sum_{n=0}^{\infty} a_n z^n$  be a power series with radius of convergence  $R > 0$ . On the **open disc** of radius  $R$ , let  $f(z)$  denote this sum. Then, on this disc, we have

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}.$$

Note that this is again a power series with the same radius of convergence. Thus, we may repeat the process indefinitely.

# Lecture 3: Power Series

Differentiability of power series is what one should expect.

## Theorem 7 (Differentiability)

Let  $\sum_{n=0}^{\infty} a_n z^n$  be a power series with radius of convergence  $R > 0$ . On the **open disc** of radius  $R$ , let  $f(z)$  denote this sum. Then, on this disc, we have

$$f'(z) = \sum_{n=1}^{\infty} n a_n z^{n-1}.$$

Note that this is again a power series with the same radius of convergence. Thus, we may repeat the process indefinitely. In other words, power series are infinite differentiable.



# End of Lecture 3

Any questions?

# Lecture 4: Exponential function

I shall just recall the facts from the lecture.

## Definition 10 (Exponential function)

The power series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

converges on all of  $\mathbb{C}$ . This sum is denoted by  $\exp(z)$ .

## Theorem 8 (Facts)

# Lecture 4: Exponential function

I shall just recall the facts from the lecture.

## Definition 10 (Exponential function)

The power series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

converges on all of  $\mathbb{C}$ . This sum is denoted by  $\exp(z)$ .

## Theorem 8 (Facts)

①  $\exp'(z) = \exp(z),$

# Lecture 4: Exponential function

I shall just recall the facts from the lecture.

## Definition 10 (Exponential function)

The power series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

converges on all of  $\mathbb{C}$ . This sum is denoted by  $\exp(z)$ .

## Theorem 8 (Facts)

- 1  $\exp'(z) = \exp(z)$ ,
- 2  $\exp(bz) = b \exp(bz)$ , for  $b \in \mathbb{C}$ ,

# Lecture 4: Exponential function

I shall just recall the facts from the lecture.

## Definition 10 (Exponential function)

The power series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

converges on all of  $\mathbb{C}$ . This sum is denoted by  $\exp(z)$ .

## Theorem 8 (Facts)

- 1  $\exp'(z) = \exp(z)$ ,
- 2  $\exp(bz) = b \exp(bz)$ , for  $b \in \mathbb{C}$ ,
- 3  $\exp(z) \cdot \exp(-z) = 1$  for all  $z \in \mathbb{C}$ ,

# Lecture 4: Exponential function

I shall just recall the facts from the lecture.

## Definition 10 (Exponential function)

The power series

$$\sum_{n=0}^{\infty} \frac{z^n}{n!}$$

converges on all of  $\mathbb{C}$ . This sum is denoted by  $\exp(z)$ .

## Theorem 8 (Facts)

- 1  $\exp'(z) = \exp(z)$ ,
- 2  $\exp(bz) = b \exp(bz)$ , for  $b \in \mathbb{C}$ ,
- 3  $\exp(z) \cdot \exp(-z) = 1$  for all  $z \in \mathbb{C}$ ,
- 4  $\exp(z)$  is always nonzero.

# Lecture 4: Exponential function

Now, we some “converse” facts.

## Theorem 9 (Characterisations)

# Lecture 4: Exponential function

Now, we some “converse” facts.

## Theorem 9 (Characterisations)

- 1 If  $f'(z) = bf(z)$ , then  $f(z) = a \exp(bz)$  for some  $a, b \in \mathbb{C}$ ,



# Lecture 4: Exponential function

Now, we some “converse” facts.

## Theorem 9 (Characterisations)

- 1 If  $f'(z) = bf(z)$ , then  $f(z) = a \exp(bz)$  for some  $a, b \in \mathbb{C}$ ,
- 2 If  $f' = f$  and  $f(0) = 1$ , then  $f(z) = \exp(z)$ .

# Lecture 4: Exponential function

Now, we some “converse” facts.

## Theorem 9 (Characterisations)

- 1 If  $f'(z) = bf(z)$ , then  $f(z) = a \exp(bz)$  for some  $a, b \in \mathbb{C}$ ,
- 2 If  $f' = f$  and  $f(0) = 1$ , then  $f(z) = \exp(z)$ .

## Theorem 10 (Final fact)

Let  $z, w \in \mathbb{C}$ , then

$$\exp(z + w) = \exp(z) \cdot \exp(w).$$

# Lecture 4: Exponential function

## Definition 11 (Domain)

A subset  $\Omega \subset \mathbb{C}$  is said to be a *domain* if it is open and path-connected.

# Lecture 4: Exponential function

## Definition 11 (Domain)

A subset  $\Omega \subset \mathbb{C}$  is said to be a *domain* if it is open and path-connected.

More discussion - informal-tut.

# Lecture 4: Exponential function

## Definition 11 (Domain)

A subset  $\Omega \subset \mathbb{C}$  is said to be a *domain* if it is open and path-connected.

More discussion - informal-tut.

We had one very nice result on the zeroes of a analytic functions.

# Lecture 4: Exponential function

## Definition 11 (Domain)

A subset  $\Omega \subset \mathbb{C}$  is said to be a *domain* if it is open and path-connected.

More discussion - informal-tut.

We had one very nice result on the zeroes of a analytic functions.

## Theorem 11 (Zeroes are isolated)

Let  $\Omega$  be a **domain** and  $f : \Omega \rightarrow \mathbb{C}$  be a **non-constant** holomorphic function.

# Lecture 4: Exponential function

## Definition 11 (Domain)

A subset  $\Omega \subset \mathbb{C}$  is said to be a *domain* if it is open and path-connected.

More discussion - informal-tut.

We had one very nice result on the zeroes of a analytic functions.

## Theorem 11 (Zeroes are isolated)

Let  $\Omega$  be a **domain** and  $f : \Omega \rightarrow \mathbb{C}$  be a **non-constant** holomorphic function. Let  $z_0 \in \Omega$  be such that  $f(z_0) = 0$ .

# Lecture 4: Exponential function

## Definition 11 (Domain)

A subset  $\Omega \subset \mathbb{C}$  is said to be a *domain* if it is open and path-connected.

More discussion - informal-tut.

We had one very nice result on the zeroes of a analytic functions.

## Theorem 11 (Zeroes are isolated)

Let  $\Omega$  be a **domain** and  $f : \Omega \rightarrow \mathbb{C}$  be a **non-constant** holomorphic function. Let  $z_0 \in \Omega$  be such that  $f(z_0) = 0$ . Then, there exists  $\delta > 0$  such that  $f$  has no other zero in  $B_\delta(z_0)$ .



# Lecture 4: Exponential function

## Definition 11 (Domain)

A subset  $\Omega \subset \mathbb{C}$  is said to be a *domain* if it is open and path-connected.

More discussion - informal-tut.

We had one very nice result on the zeroes of a analytic functions.

## Theorem 11 (Zeroes are isolated)

Let  $\Omega$  be a **domain** and  $f : \Omega \rightarrow \mathbb{C}$  be a **non-constant** holomorphic function. Let  $z_0 \in \Omega$  be such that  $f(z_0) = 0$ . Then, there exists  $\delta > 0$  such that  $f$  has no other zero in  $B_\delta(z_0)$ .

The above is saying that around every zero of  $f$ , we can draw a (sufficiently small) circle such that  $f$  has no other zero in that disc.

# Lecture 4: Exponential function

## Definition 11 (Domain)

A subset  $\Omega \subset \mathbb{C}$  is said to be a *domain* if it is open and path-connected.

More discussion - informal-tut.

We had one very nice result on the zeroes of a analytic functions.

## Theorem 11 (Zeroes are isolated)

Let  $\Omega$  be a **domain** and  $f : \Omega \rightarrow \mathbb{C}$  be a **non-constant** holomorphic function. Let  $z_0 \in \Omega$  be such that  $f(z_0) = 0$ . Then, there exists  $\delta > 0$  such that  $f$  has no other zero in  $B_\delta(z_0)$ .

The above is saying that around every zero of  $f$ , we can draw a (sufficiently small) circle such that  $f$  has no other zero in that disc. This is the same as saying that the set of zeroes is *discrete*.

# End of Lecture 4

Any questions?

# Lecture 5: Integration

## Definition 12

Let  $f : [a, b] \rightarrow \mathbb{C}$  be a piecewise continuous function. Writing  $f = u + \iota v$  as usual, we define

$$\int_a^b f(t)dt := \int_a^b u(t)dt + \iota \int_a^b v(t)dt.$$

# Lecture 5: Integration

## Definition 12

Let  $f : [a, b] \rightarrow \mathbb{C}$  be a piecewise continuous function. Writing  $f = u + \iota v$  as usual, we define

$$\int_a^b f(t)dt := \int_a^b u(t)dt + \iota \int_a^b v(t)dt.$$

This is naturally what one would have wanted to define.

# Lecture 5: Integration

## Definition 12

Let  $f : [a, b] \rightarrow \mathbb{C}$  be a piecewise continuous function. Writing  $f = u + \iota v$  as usual, we define

$$\int_a^b f(t)dt := \int_a^b u(t)dt + \iota \int_a^b v(t)dt.$$

This is naturally what one would have wanted to define. Now, we define integration over a *contour*.

# Lecture 5: Integration

## Definition 12

Let  $f : [a, b] \rightarrow \mathbb{C}$  be a piecewise continuous function. Writing  $f = u + \iota v$  as usual, we define

$$\int_a^b f(t) dt := \int_a^b u(t) dt + \iota \int_a^b v(t) dt.$$

This is naturally what one would have wanted to define. Now, we define integration over a *contour*. (What is a contour?)

# Lecture 5: Integration

## Definition 12

Let  $f : [a, b] \rightarrow \mathbb{C}$  be a piecewise continuous function. Writing  $f = u + \iota v$  as usual, we define

$$\int_a^b f(t) dt := \int_a^b u(t) dt + \iota \int_a^b v(t) dt.$$

This is naturally what one would have wanted to define. Now, we define integration over a *contour*. (What is a contour?)

## Definition 13

Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function. Let  $\gamma : [a, b] \rightarrow \Omega$  be a contour. We define

$$\int_{\gamma} f(z) dz := \int_a^b f(\gamma(t)) \gamma'(t) dt.$$



# Lecture 5: Integration

## Definition 12

Let  $f : [a, b] \rightarrow \mathbb{C}$  be a piecewise continuous function. Writing  $f = u + \iota v$  as usual, we define

$$\int_a^b f(t) dt := \int_a^b u(t) dt + \iota \int_a^b v(t) dt.$$

This is naturally what one would have wanted to define. Now, we define integration over a *contour*. (What is a contour?)

## Definition 13

Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function. Let  $\gamma : [a, b] \rightarrow \Omega$  be a contour. We define

$$\int_{\gamma} f(z) dz := \int_a^b f(\gamma(t)) \gamma'(t) dt.$$

# Lecture 5: Integration

We have a useful inequality called the **ML** inequality.

## Theorem 12 (**ML** Inequality)

# Lecture 5: Integration

We have a useful inequality called the **ML** inequality.

## Theorem 12 (**ML** Inequality)

Let  $\gamma$  be a contour of length  **$L$**

# Lecture 5: Integration

We have a useful inequality called the **ML** inequality.

## Theorem 12 (**ML** Inequality)

Let  $\gamma$  be a contour of length  $L$  and  $f$  be a continuous function defined on the image of  $\gamma$ .

# Lecture 5: Integration

We have a useful inequality called the **ML** inequality.

## Theorem 12 (**ML** Inequality)

Let  $\gamma$  be a contour of length **L** and  $f$  be a continuous function defined on the image of  $\gamma$ .

Suppose that

$$|f(\gamma(t))| \leq \mathbf{M}, \quad \text{for all } t \in [a, b].$$

# Lecture 5: Integration

We have a useful inequality called the  $ML$  inequality.

## Theorem 12 ( $ML$ Inequality)

Let  $\gamma$  be a contour of length  $L$  and  $f$  be a continuous function defined on the image of  $\gamma$ .

Suppose that

$$|f(\gamma(t))| \leq M, \quad \text{for all } t \in [a, b].$$

Then, we have

$$\left| \int_{\gamma} f(z) dz \right| \leq ML.$$

# Lecture 5: Integration

## Theorem 13 (Primitives and integrals)

Suppose  $f : \Omega \rightarrow \mathbb{C}$  has a *primitive* on  $\Omega$ .

# Lecture 5: Integration

## Theorem 13 (Primitives and integrals)

Suppose  $f : \Omega \rightarrow \mathbb{C}$  has a *primitive* on  $\Omega$ . That is, there exists a function  $F : \Omega \rightarrow \mathbb{C}$  such that  $F' = f$ . (The complex derivative.)



# Lecture 5: Integration

## Theorem 13 (Primitives and integrals)

Suppose  $f : \Omega \rightarrow \mathbb{C}$  has a *primitive* on  $\Omega$ . That is, there exists a function  $F : \Omega \rightarrow \mathbb{C}$  such that  $F' = f$ . (The complex derivative.) Then, we have

# Lecture 5: Integration

## Theorem 13 (Primitives and integrals)

Suppose  $f : \Omega \rightarrow \mathbb{C}$  has a *primitive* on  $\Omega$ . That is, there exists a function  $F : \Omega \rightarrow \mathbb{C}$  such that  $F' = f$ . (The complex derivative.) Then, we have

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

# Lecture 5: Integration

## Theorem 13 (Primitives and integrals)

Suppose  $f : \Omega \rightarrow \mathbb{C}$  has a *primitive* on  $\Omega$ . That is, there exists a function  $F : \Omega \rightarrow \mathbb{C}$  such that  $F' = f$ . (The complex derivative.) Then, we have

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

If  $\gamma$  is *closed*, that is, if  $\gamma(b) = \gamma(a)$ , then

# Lecture 5: Integration

## Theorem 13 (Primitives and integrals)

Suppose  $f : \Omega \rightarrow \mathbb{C}$  has a *primitive* on  $\Omega$ . That is, there exists a function  $F : \Omega \rightarrow \mathbb{C}$  such that  $F' = f$ . (The complex derivative.) Then, we have

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

If  $\gamma$  is *closed*, that is, if  $\gamma(b) = \gamma(a)$ , then

$$\int_{\gamma} f(z) dz = 0.$$

# Lecture 5: Integration

## Theorem 13 (Primitives and integrals)

Suppose  $f : \Omega \rightarrow \mathbb{C}$  has a *primitive* on  $\Omega$ . That is, there exists a function  $F : \Omega \rightarrow \mathbb{C}$  such that  $F' = f$ . (The complex derivative.) Then, we have

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

If  $\gamma$  is *closed*, that is, if  $\gamma(b) = \gamma(a)$ , then

$$\int_{\gamma} f(z) dz = 0.$$

Existence of a primitive is a strong condition, by the way.

# Lecture 5: Integration

## Theorem 13 (Primitives and integrals)

Suppose  $f : \Omega \rightarrow \mathbb{C}$  has a *primitive* on  $\Omega$ . That is, there exists a function  $F : \Omega \rightarrow \mathbb{C}$  such that  $F' = f$ . (The complex derivative.) Then, we have

$$\int_{\gamma} f(z) dz = F(\gamma(b)) - F(\gamma(a)).$$

If  $\gamma$  is *closed*, that is, if  $\gamma(b) = \gamma(a)$ , then

$$\int_{\gamma} f(z) dz = 0.$$

Existence of a primitive is a strong condition, by the way. A holomorphic function need not have a primitive on all of  $\Omega$ .

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple,



Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple, closed contour

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple, closed contour and let  $f$  be a holomorphic function defined

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple, closed contour and let  $f$  be a holomorphic function defined on an open set  $\Omega$  containing  $\gamma$

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple, closed contour and let  $f$  be a holomorphic function defined on an open set  $\Omega$  containing  $\gamma$  as well as its interior.

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple, closed contour and let  $f$  be a holomorphic function defined on an open set  $\Omega$  containing  $\gamma$  **as well as its interior**. Then,

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple, closed contour and let  $f$  be a holomorphic function defined on an open set  $\Omega$  containing  $\gamma$  **as well as its interior**. Then,

$$\int_{\gamma} f(z) dz = 0.$$

# Lecture 5: Integration

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple, closed contour and let  $f$  be a holomorphic function defined on an open set  $\Omega$  containing  $\gamma$  **as well as its interior**. Then,

$$\int_{\gamma} f(z) dz = 0.$$

If  $\Omega$  is simply-connected, then the interior condition is automatically met.

# Lecture 5: Integration

Now, we come to Cauchy's theorem.

## Theorem 14 (Cauchy's Theorem)

Let  $\gamma$  be a simple, closed contour and let  $f$  be a holomorphic function defined on an open set  $\Omega$  containing  $\gamma$  **as well as its interior**. Then,

$$\int_{\gamma} f(z) dz = 0.$$

If  $\Omega$  is simply-connected, then the interior condition is automatically met. This gives us the next result.



## Theorem 15 (“General” Cauchy Theorem)

Let  $\Omega$  be a simply-connected domain. Let  $\gamma : [a, b] \rightarrow \mathbb{C}$  be a simple, closed contour and  $f : \Omega \rightarrow \mathbb{C}$  holomorphic. Then,

$$\int_{\gamma} f(z) dz = 0.$$

# End of Lecture 5

Any questions?

## Theorem 16 (Cauchy Integral Formula)

Let  $f$  be holomorphic everywhere on an open set  $\Omega$ .

## Theorem 16 (Cauchy Integral Formula)

Let  $f$  be holomorphic everywhere on an open set  $\Omega$ . Let  $\gamma$  be a simple closed curve in  $\Omega$ , oriented positively.

## Theorem 16 (Cauchy Integral Formula)

Let  $f$  be holomorphic everywhere on an open set  $\Omega$ . Let  $\gamma$  be a simple closed curve in  $\Omega$ , oriented positively. If  $z_0$  is interior to  $\gamma$

## Theorem 16 (Cauchy Integral Formula)

Let  $f$  be holomorphic everywhere on an open set  $\Omega$ . Let  $\gamma$  be a simple closed curve in  $\Omega$ , oriented positively. If  $z_0$  is interior to  $\gamma$  and  $\Omega$  contains the interior of  $\gamma$ , then

## Theorem 16 (Cauchy Integral Formula)

Let  $f$  be holomorphic everywhere on an open set  $\Omega$ . Let  $\gamma$  be a simple closed curve in  $\Omega$ , oriented positively. If  $z_0$  is interior to  $\gamma$  and  $\Omega$  contains the interior of  $\gamma$ , then

$$f(z_0) = \frac{1}{2\pi i} \int_{\gamma} \frac{f(z)}{z - z_0} dz$$

## Lecture 6: CIF and Consequences

We then saw a consequence of CIF which I state as a theorem below.

### Theorem 17 (Holomorphic $\implies$ Analytic)

Let  $\Omega \subset \mathbb{C}$  be open and  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. Pick any  $z_0 \in \Omega$ .



# Lecture 6: CIF and Consequences

We then saw a consequence of CIF which I state as a theorem below.

## Theorem 17 (Holomorphic $\implies$ Analytic)

Let  $\Omega \subset \mathbb{C}$  be open and  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. Pick any  $z_0 \in \Omega$ . Let  $R > 0$  be the largest such that  $B_R(z_0) \subset \Omega$ .

# Lecture 6: CIF and Consequences

We then saw a consequence of CIF which I state as a theorem below.

## Theorem 17 (Holomorphic $\implies$ Analytic)

Let  $\Omega \subset \mathbb{C}$  be open and  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. Pick any  $z_0 \in \Omega$ . Let  $R > 0$  be the largest such that  $B_R(z_0) \subset \Omega$ . (The case  $R = \infty$  is allowed. That just means  $\Omega = \mathbb{C}$ .)

## Lecture 6: CIF and Consequences

We then saw a consequence of CIF which I state as a theorem below.

### Theorem 17 (Holomorphic $\implies$ Analytic)

Let  $\Omega \subset \mathbb{C}$  be open and  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. Pick any  $z_0 \in \Omega$ . Let  $R > 0$  be the largest such that  $B_R(z_0) \subset \Omega$ . (The case  $R = \infty$  is allowed. That just means  $\Omega = \mathbb{C}$ .) Then, on the disc  $B_R(z_0)$ , we may write  $f(z)$  as

# Lecture 6: CIF and Consequences

We then saw a consequence of CIF which I state as a theorem below.

## Theorem 17 (Holomorphic $\implies$ Analytic)

Let  $\Omega \subset \mathbb{C}$  be open and  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. Pick any  $z_0 \in \Omega$ . Let  $R > 0$  be the largest such that  $B_R(z_0) \subset \Omega$ . (The case  $R = \infty$  is allowed. That just means  $\Omega = \mathbb{C}$ .) Then, on the disc  $B_R(z_0)$ , we may write  $f(z)$  as

$$f(z) = \sum_{n=0}^{\infty} a_n z^n,$$

# Lecture 6: CIF and Consequences

We then saw a consequence of CIF which I state as a theorem below.

## Theorem 17 (Holomorphic $\implies$ Analytic)

Let  $\Omega \subset \mathbb{C}$  be open and  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. Pick any  $z_0 \in \Omega$ . Let  $R > 0$  be the largest such that  $B_R(z_0) \subset \Omega$ .

(The case  $R = \infty$  is allowed. That just means  $\Omega = \mathbb{C}$ .)

Then, on the disc  $B_R(z_0)$ , we may write  $f(z)$  as

$$f(z) = \sum_{n=0}^{\infty} a_n z^n,$$

where each  $a_n$  is given by

## Lecture 6: CIF and Consequences

We then saw a consequence of CIF which I state as a theorem below.

### Theorem 17 (Holomorphic $\implies$ Analytic)

Let  $\Omega \subset \mathbb{C}$  be open and  $f : \Omega \rightarrow \mathbb{C}$  be holomorphic. Pick any  $z_0 \in \Omega$ . Let  $R > 0$  be the largest such that  $B_R(z_0) \subset \Omega$ .

(The case  $R = \infty$  is allowed. That just means  $\Omega = \mathbb{C}$ .)

Then, on the disc  $B_R(z_0)$ , we may write  $f(z)$  as

$$f(z) = \sum_{n=0}^{\infty} a_n z^n,$$

where each  $a_n$  is given by

$$a_n = \frac{1}{2\pi i} \int_{|w-z_0|=r} \frac{f(w)}{(w-z_0)^{n+1}} dw.$$

## Lecture 6: CIF and Consequences

The above also gives us (what I call) the “generalised” Cauchy Integral Formula.

### Theorem 18 (“Generalised” CIF)

## Lecture 6: CIF and Consequences

The above also gives us (what I call) the “generalised” Cauchy Integral Formula.

### Theorem 18 (“Generalised” CIF)

$$\int_{|w-z_0|=r} \frac{f(w)}{(w-z_0)^{n+1}} dw = \frac{2\pi i}{n!} f^{(n)}(z_0),$$

where  $f$  is a function which is holomorphic on an open disc  $B_R(z_0)$  and  $r < R$ .



## Lecture 6: CIF and Consequences

The above also gives us (what I call) the “generalised” Cauchy Integral Formula.

### Theorem 18 (“Generalised” CIF)

$$\int_{|w-z_0|=r} \frac{f(w)}{(w-z_0)^{n+1}} dw = \frac{2\pi i}{n!} f^{(n)}(z_0),$$

where  $f$  is a function which is holomorphic on an open disc  $B_R(z_0)$  and  $r < R$ .

### Remark 3

Note that, as usual, we require  $f$  to be holomorphic within the circle as well.

# Lecture 7: CIF and Consequences

## Theorem 19 (Cauchy's estimate)

## Theorem 20 (Liouville's Theorem)

# Lecture 7: CIF and Consequences

## Theorem 19 (Cauchy's estimate)

Suppose that  $f$  is holomorphic on  $|z - z_0| < R$

## Theorem 20 (Liouville's Theorem)

# Lecture 7: CIF and Consequences

## Theorem 19 (Cauchy's estimate)

Suppose that  $f$  is holomorphic on  $|z - z_0| < R$  and bounded by  $M > 0$  on this disc. Then,

## Theorem 20 (Liouville's Theorem)

# Lecture 7: CIF and Consequences

## Theorem 19 (Cauchy's estimate)

Suppose that  $f$  is holomorphic on  $|z - z_0| < R$  and bounded by  $M > 0$  on this disc. Then,

$$\left| f^{(n)}(z_0) \right| \leq \frac{n!M}{R^n}.$$

## Theorem 20 (Liouville's Theorem)

# Lecture 7: CIF and Consequences

## Theorem 19 (Cauchy's estimate)

Suppose that  $f$  is holomorphic on  $|z - z_0| < R$  and bounded by  $M > 0$  on this disc. Then,

$$\left| f^{(n)}(z_0) \right| \leq \frac{n!M}{R^n}.$$

An easy application of this give us:

## Theorem 20 (Liouville's Theorem)

# Lecture 7: CIF and Consequences

## Theorem 19 (Cauchy's estimate)

Suppose that  $f$  is holomorphic on  $|z - z_0| < R$  and bounded by  $M > 0$  on this disc. Then,

$$\left| f^{(n)}(z_0) \right| \leq \frac{n!M}{R^n}.$$

An easy application of this give us:

## Theorem 20 (Liouville's Theorem)

Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be holomorphic.

# Lecture 7: CIF and Consequences

## Theorem 19 (Cauchy's estimate)

Suppose that  $f$  is holomorphic on  $|z - z_0| < R$  and bounded by  $M > 0$  on this disc. Then,

$$\left| f^{(n)}(z_0) \right| \leq \frac{n!M}{R^n}.$$

An easy application of this give us:

## Theorem 20 (Liouville's Theorem)

Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be holomorphic. If  $f$  is bounded, then



# Lecture 7: CIF and Consequences

## Theorem 19 (Cauchy's estimate)

Suppose that  $f$  is holomorphic on  $|z - z_0| < R$  and bounded by  $M > 0$  on this disc. Then,

$$\left| f^{(n)}(z_0) \right| \leq \frac{n!M}{R^n}.$$

An easy application of this give us:

## Theorem 20 (Liouville's Theorem)

Let  $f : \mathbb{C} \rightarrow \mathbb{C}$  be holomorphic. If  $f$  is bounded, then  $f$  is constant!

# End of Lectures 6 and 7

Any questions?

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

# Logarithm

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

Let  $\Omega \subset \mathbb{C}$  be a **domain**.

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

Let  $\Omega \subset \mathbb{C}$  be a **domain**. Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function such that

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

Let  $\Omega \subset \mathbb{C}$  be a **domain**. Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function such that

$$\exp(f(z)) = z, \quad \text{for all } z \in \Omega.$$

# Logarithm

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

Let  $\Omega \subset \mathbb{C}$  be a **domain**. Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function such that

$$\exp(f(z)) = z, \quad \text{for all } z \in \Omega.$$

Then,  $f$  is called a *branch of the logarithm*.

# Logarithm

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

Let  $\Omega \subset \mathbb{C}$  be a **domain**. Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function such that

$$\exp(f(z)) = z, \quad \text{for all } z \in \Omega.$$

Then,  $f$  is called a *branch of the logarithm*.

## Theorem 21 (Uniqueness of branches)

Assume that  $f, g : \Omega \rightarrow \mathbb{C}$  are two branches of the logarithm.



# Logarithm

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

Let  $\Omega \subset \mathbb{C}$  be a **domain**. Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function such that

$$\exp(f(z)) = z, \quad \text{for all } z \in \Omega.$$

Then,  $f$  is called a *branch of the logarithm*.

## Theorem 21 (Uniqueness of branches)

Assume that  $f, g : \Omega \rightarrow \mathbb{C}$  are two branches of the logarithm. Then,  $f - g$  is a constant function.

# Logarithm

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

Let  $\Omega \subset \mathbb{C}$  be a **domain**. Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function such that

$$\exp(f(z)) = z, \quad \text{for all } z \in \Omega.$$

Then,  $f$  is called a *branch of the logarithm*.

## Theorem 21 (Uniqueness of branches)

Assume that  $f, g : \Omega \rightarrow \mathbb{C}$  are two branches of the logarithm. Then,  $f - g$  is a constant function. Moreover, this constant is an integer multiple of  $2\pi i$ .

# Logarithm

We discuss logarithm a bit.

## Definition 14 (Branch of the logarithm)

Let  $\Omega \subset \mathbb{C}$  be a **domain**. Let  $f : \Omega \rightarrow \mathbb{C}$  be a continuous function such that

$$\exp(f(z)) = z, \quad \text{for all } z \in \Omega.$$

Then,  $f$  is called a *branch of the logarithm*.

## Theorem 21 (Uniqueness of branches)

Assume that  $f, g : \Omega \rightarrow \mathbb{C}$  are two branches of the logarithm. Then,  $f - g$  is a constant function. Moreover, this constant is an integer multiple of  $2\pi i$ .

The last theorem also assumed that  $\Omega$  is a **domain**.

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ .

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$  and  $0 \notin \Omega$ .

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$  and  $0 \notin \Omega$ .

There exists a unique function  $F : \Omega \rightarrow \mathbb{C}$  such that



The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$  and  $0 \notin \Omega$ .

There exists a unique function  $F : \Omega \rightarrow \mathbb{C}$  such that

①  $F(1) = 0,$

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$  and  $0 \notin \Omega$ .

There exists a unique function  $F : \Omega \rightarrow \mathbb{C}$  such that

- ①  $F(1) = 0$ ,
- ②  $F'(z) = 1/z$ ,

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$  and  $0 \notin \Omega$ .

There exists a unique function  $F : \Omega \rightarrow \mathbb{C}$  such that

- ①  $F(1) = 0$ ,
- ②  $F'(z) = 1/z$ ,
- ③  $\exp(F(z)) = z$  for all  $z \in \Omega$ ,

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$  and  $0 \notin \Omega$ .

There exists a unique function  $F : \Omega \rightarrow \mathbb{C}$  such that

- ①  $F(1) = 0$ ,
- ②  $F'(z) = 1/z$ ,
- ③  $\exp(F(z)) = z$  for all  $z \in \Omega$ ,
- ④  $F(r) = \log(r)$  for all  $r \in \Omega \cap \mathbb{R}$ .

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$  and  $0 \notin \Omega$ .

There exists a unique function  $F : \Omega \rightarrow \mathbb{C}$  such that

- ①  $F(1) = 0$ ,
- ②  $F'(z) = 1/z$ ,
- ③  $\exp(F(z)) = z$  for all  $z \in \Omega$ ,
- ④  $F(r) = \log(r)$  for all  $r \in \Omega \cap \mathbb{R}$ .

The log in the last point is the usual log for real numbers as seen in 105.

The previous theorem talked about uniqueness of branches (up to a constant) assuming the existence of such a branch. Now, we see when a branch is actually possible.

## Theorem 22 (Existence of a branch)

Let  $\Omega$  be a **simply-connected** domain in  $\mathbb{C}$ . Assume that  $1 \in \Omega$  and  $0 \notin \Omega$ .

There exists a unique function  $F : \Omega \rightarrow \mathbb{C}$  such that

- ①  $F(1) = 0$ ,
- ②  $F'(z) = 1/z$ ,
- ③  $\exp(F(z)) = z$  for all  $z \in \Omega$ ,
- ④  $F(r) = \log(r)$  for all  $r \in \Omega \cap \mathbb{R}$ .

The log in the last point is the usual log for real numbers as seen in 105. The above  $F$  is then denoted by  $\log$ .

# Lecture 8: Singularities

## Definition 15 (Singularities)

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. A point  $z_0 \in \mathbb{C}$  is said to be a singularity of  $f$  if

## Definition 16 (Isolated singularity)

# Lecture 8: Singularities

## Definition 15 (Singularities)

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. A point  $z_0 \in \mathbb{C}$  is said to be a singularity of  $f$  if

- 1  $z_0 \notin \Omega$ , i.e.,  $f$  is not defined at  $z_0$ , or

## Definition 16 (Isolated singularity)



# Lecture 8: Singularities

## Definition 15 (Singularities)

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. A point  $z_0 \in \mathbb{C}$  is said to be a singularity of  $f$  if

- 1  $z_0 \notin \Omega$ , i.e.,  $f$  is not defined at  $z_0$ , or
- 2  $z_0 \in \Omega$  and  $f$  is not holomorphic at  $z_0$ .

## Definition 16 (Isolated singularity)

# Lecture 8: Singularities

## Definition 15 (Singularities)

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. A point  $z_0 \in \mathbb{C}$  is said to be a singularity of  $f$  if

- ①  $z_0 \notin \Omega$ , i.e.,  $f$  is not defined at  $z_0$ , or
- ②  $z_0 \in \Omega$  and  $f$  is not holomorphic at  $z_0$ .

## Definition 16 (Isolated singularity)

A singularity  $z_0 \in \mathbb{C}$  is said to be *isolated* if

# Lecture 8: Singularities

## Definition 15 (Singularities)

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. A point  $z_0 \in \mathbb{C}$  is said to be a singularity of  $f$  if

- 1  $z_0 \notin \Omega$ , i.e.,  $f$  is not defined at  $z_0$ , or
- 2  $z_0 \in \Omega$  and  $f$  is not holomorphic at  $z_0$ .

## Definition 16 (Isolated singularity)

A singularity  $z_0 \in \mathbb{C}$  is said to be *isolated* if there exists *some*  $\delta > 0$  such that

# Lecture 8: Singularities

## Definition 15 (Singularities)

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. A point  $z_0 \in \mathbb{C}$  is said to be a singularity of  $f$  if

- 1  $z_0 \notin \Omega$ , i.e.,  $f$  is not defined at  $z_0$ , or
- 2  $z_0 \in \Omega$  and  $f$  is not holomorphic at  $z_0$ .

## Definition 16 (Isolated singularity)

A singularity  $z_0 \in \mathbb{C}$  is said to be *isolated* if there exists *some*  $\delta > 0$  such that  $f$  is holomorphic on  $B_\delta(z_0) \setminus \{z_0\}$ .

# Lecture 8: Singularities

## Definition 15 (Singularities)

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. A point  $z_0 \in \mathbb{C}$  is said to be a singularity of  $f$  if

- 1  $z_0 \notin \Omega$ , i.e.,  $f$  is not defined at  $z_0$ , or
- 2  $z_0 \in \Omega$  and  $f$  is not holomorphic at  $z_0$ .

## Definition 16 (Isolated singularity)

A singularity  $z_0 \in \mathbb{C}$  is said to be *isolated* if there exists *some*  $\delta > 0$  such that  $f$  is holomorphic on  $B_\delta(z_0) \setminus \{z_0\}$ .

The above is saying that “ $f$  is holomorphic on some *punctured disc* around  $z_0$ .”

# Lecture 8: Singularities

## Definition 15 (Singularities)

Let  $f : \Omega \rightarrow \mathbb{C}$  be a function. A point  $z_0 \in \mathbb{C}$  is said to be a singularity of  $f$  if

- 1  $z_0 \notin \Omega$ , i.e.,  $f$  is not defined at  $z_0$ , or
- 2  $z_0 \in \Omega$  and  $f$  is not holomorphic at  $z_0$ .

## Definition 16 (Isolated singularity)

A singularity  $z_0 \in \mathbb{C}$  is said to be *isolated* if there exists *some*  $\delta > 0$  such that  $f$  is holomorphic on  $B_\delta(z_0) \setminus \{z_0\}$ .

The above is saying that “ $f$  is holomorphic on some *punctured disc* around  $z_0$ .”

Compare this “isolation” with what we saw earlier when we said that “zeroes are isolated.”

# Lecture 8: Singularities

## Definition 17 (Non-isolated singularity)

A singularity which is not an isolated singularity is called a non-isolated singularity.

# Lecture 8: Singularities

## Definition 17 (Non-isolated singularity)

A singularity which is not an isolated singularity is called a non-isolated singularity.

*The floor is made of floor.*



# Lecture 8: Singularities

## Definition 17 (Non-isolated singularity)

A singularity which is not an isolated singularity is called a non-isolated singularity.

*The floor is made of floor.*

Note that if  $f$  has only finitely many singularities, then all the singularities are isolated.

# Lecture 8: Singularities

## Definition 17 (Non-isolated singularity)

A singularity which is not an isolated singularity is called a non-isolated singularity.

*The floor is made of floor.*

Note that if  $f$  has only finitely many singularities, then all the singularities are isolated.

We classify **isolated** singularities into three types:

# Lecture 8: Singularities

## Definition 17 (Non-isolated singularity)

A singularity which is not an isolated singularity is called a non-isolated singularity.

*The floor is made of floor.*

Note that if  $f$  has only finitely many singularities, then all the singularities are isolated.

We classify **isolated** singularities into three types:

- 1 Removable singularities,

# Lecture 8: Singularities

## Definition 17 (Non-isolated singularity)

A singularity which is not an isolated singularity is called a non-isolated singularity.

*The floor is made of floor.*

Note that if  $f$  has only finitely many singularities, then all the singularities are isolated.

We classify **isolated** singularities into three types:

- 1 Removable singularities,
- 2 Poles,

# Lecture 8: Singularities

## Definition 17 (Non-isolated singularity)

A singularity which is not an isolated singularity is called a non-isolated singularity.

*The floor is made of floor.*

Note that if  $f$  has only finitely many singularities, then all the singularities are isolated.

We classify **isolated** singularities into three types:

- 1 Removable singularities,
- 2 Poles,
- 3 Essential singularities.

# Lecture 8: Singularities

## Definition 17 (Non-isolated singularity)

A singularity which is not an isolated singularity is called a non-isolated singularity.

*The floor is made of floor.*

Note that if  $f$  has only finitely many singularities, then all the singularities are isolated.

We classify **isolated** singularities into three types:

- 1 Removable singularities,
- 2 Poles,
- 3 Essential singularities.

## Remark 4

The above classification is only for **isolated** singularities.

# Lecture 8: Singularities

## Definition 18 (Removable singularity)

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable.

## Theorem 23 (Riemann's Removable Singularity Theorem)

# Lecture 8: Singularities

## Definition 18 (Removable singularity)

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable.

These are characterised by the following theorem.

## Theorem 23 (Riemann's Removable Singularity Theorem)



# Lecture 8: Singularities

## Definition 18 (Removable singularity)

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable.

These are characterised by the following theorem.

## Theorem 23 (Riemann's Removable Singularity Theorem)

$z_0$  is a removable singularity of  $f$  iff

# Lecture 8: Singularities

## Definition 18 (Removable singularity)

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable.

These are characterised by the following theorem.

## Theorem 23 (Riemann's Removable Singularity Theorem)

$z_0$  is a removable singularity of  $f$  iff  $\lim_{z \rightarrow z_0} f(z)$  exists.

# Lecture 8: Singularities

## Definition 18 (Removable singularity)

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable.

These are characterised by the following theorem.

## Theorem 23 (Riemann's Removable Singularity Theorem)

$z_0$  is a removable singularity of  $f$  iff  $\lim_{z \rightarrow z_0} f(z)$  exists.

In the above, we mean that it exists as a (finite) complex number.

# Lecture 8: Singularities

## Definition 18 (Removable singularity)

If an isolated singularity can be removed by defining the function by assigning a certain value at that point, we say that the singularity is removable.

These are characterised by the following theorem.

## Theorem 23 (Riemann's Removable Singularity Theorem)

$z_0$  is a removable singularity of  $f$  iff  $\lim_{z \rightarrow z_0} f(z)$  exists.

In the above, we mean that it exists as a (finite) complex number.

$$f(z) = \frac{\sin z}{z}$$

defined on  $\mathbb{C} \setminus \{0\}$  has 0 as a removable singularity.

# Lecture 8: Singularities

## Definition 19 (Pole)

An isolated singularity  $z_0$  is said to be a pole if

# Lecture 8: Singularities

## Definition 19 (Pole)

An isolated singularity  $z_0$  is said to be a pole if  $|f(z)| \rightarrow \infty$  as

# Lecture 8: Singularities

## Definition 19 (Pole)

An isolated singularity  $z_0$  is said to be a pole if  $|f(z)| \rightarrow \infty$  as  $z \rightarrow z_0$ .

# Lecture 8: Singularities

## Definition 19 (Pole)

An isolated singularity  $z_0$  is said to be a pole if  $|f(z)| \rightarrow \infty$  as  $z \rightarrow z_0$ .

## Theorem 24

An isolated singularity  $z_0$  is a pole of  $f$  iff

$$\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0.$$



# Lecture 8: Singularities

## Definition 19 (Pole)

An isolated singularity  $z_0$  is said to be a pole if  $|f(z)| \rightarrow \infty$  as  $z \rightarrow z_0$ .

## Theorem 24

An isolated singularity  $z_0$  is a pole of  $f$  iff

$$\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0.$$

## Theorem 25 (Order of a pole)

If  $z_0$  is a pole of  $f$ , then there exists an integer  $m > 0$  such that

$$f(z) = (z - z_0)^{-m} f_1(z)$$

# Lecture 8: Singularities

## Definition 19 (Pole)

An isolated singularity  $z_0$  is said to be a pole if  $|f(z)| \rightarrow \infty$  as  $z \rightarrow z_0$ .

## Theorem 24

An isolated singularity  $z_0$  is a pole of  $f$  iff

$$\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0.$$

## Theorem 25 (Order of a pole)

If  $z_0$  is a pole of  $f$ , then there exists an integer  $m > 0$  such that

$$f(z) = (z - z_0)^{-m} f_1(z)$$

on a punctured neighbourhood of  $z_0$ , for some function  $f_1$  which is holomorphic on the complete neighbourhood.

# Lecture 8: Singularities

## Definition 19 (Pole)

An isolated singularity  $z_0$  is said to be a pole if  $|f(z)| \rightarrow \infty$  as  $z \rightarrow z_0$ .

## Theorem 24

An isolated singularity  $z_0$  is a pole of  $f$  iff

$$\lim_{z \rightarrow z_0} \frac{1}{f(z)} = 0.$$

## Theorem 25 (Order of a pole)

If  $z_0$  is a pole of  $f$ , then there exists an integer  $m > 0$  such that

$$f(z) = (z - z_0)^{-m} f_1(z)$$

on a punctured neighbourhood of  $z_0$ , for some function  $f_1$  which is holomorphic on the complete neighbourhood.

# Lecture 8: Singularities

## Definition 20 (Essential singularity)

An isolated singularity is called an essential singularity if it is neither a removable singularity nor a pole.

## Theorem 26 (Casorati-Weierstrass Theorem)

# Lecture 8: Singularities

## Definition 20 (Essential singularity)

An isolated singularity is called an essential singularity if it is neither a removable singularity nor a pole.

## Theorem 26 (Casorati-Weierstrass Theorem)

If  $z_0$  is an isolated singularity, then it is essential iff

# Lecture 8: Singularities

## Definition 20 (Essential singularity)

An isolated singularity is called an essential singularity if it is neither a removable singularity nor a pole.

## Theorem 26 (Casorati-Weierstrass Theorem)

If  $z_0$  is an isolated singularity, then it is essential iff the values of  $f$  come arbitrarily close to every complex number in a neighborhood of  $z_0$ .

# End of Lecture 8

Any questions?

## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ .



## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ . Consider an annulus of the form

$$A = \{z : r < |z - z_0| < R\},$$

where  $0 < r < R \leq \infty$ .

## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ . Consider an annulus of the form

$$A = \{z : r < |z - z_0| < R\},$$

where  $0 < r < R \leq \infty$ . Assume that  $f$  is holomorphic on this open annulus  $A$ .

# Lecture 9: Laurent Series

## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ . Consider an annulus of the form

$$A = \{z : r < |z - z_0| < R\},$$

where  $0 < r < R \leq \infty$ . Assume that  $f$  is holomorphic on this open annulus  $A$ . Then, CIF takes the form

# Lecture 9: Laurent Series

## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ . Consider an annulus of the form

$$A = \{z : r < |z - z_0| < R\},$$

where  $0 < r < R \leq \infty$ . Assume that  $f$  is holomorphic on this open annulus  $A$ . Then, CIF takes the form

$$f(z) = \sum_{n=0}^{\infty} a_n (z - z_0)^{-n} + \sum_{n=1}^{\infty} b_n (z - z_0)^n,$$

## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ . Consider an annulus of the form

$$A = \{z : r < |z - z_0| < R\},$$

where  $0 < r < R \leq \infty$ . Assume that  $f$  is holomorphic on this open annulus  $A$ . Then, CIF takes the form

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=R'} \frac{f(w)}{z-w} dw, \quad ,$$

## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ . Consider an annulus of the form

$$A = \{z : r < |z - z_0| < R\},$$

where  $0 < r < R \leq \infty$ . Assume that  $f$  is holomorphic on this open annulus  $A$ . Then, CIF takes the form

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=R'} \frac{f(w)}{z-w} dw - \frac{1}{2\pi i} \int_{|w-z_0|=r'} \frac{f(w)}{z-w} dw,$$

# Lecture 9: Laurent Series

## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ . Consider an annulus of the form

$$A = \{z : r < |z - z_0| < R\},$$

where  $0 < r < R \leq \infty$ . Assume that  $f$  is holomorphic on this open annulus  $A$ . Then, CIF takes the form

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=R'} \frac{f(w)}{z-w} dw - \frac{1}{2\pi i} \int_{|w-z_0|=r'} \frac{f(w)}{z-w} dw,$$

where  $r < r' < R' < R$ .

# Lecture 9: Laurent Series

## Theorem 27 (Modified CIF)

Suppose that  $z_0$  is an isolated singularity of  $f$ . Consider an annulus of the form

$$A = \{z : r < |z - z_0| < R\},$$

where  $0 < r < R \leq \infty$ . Assume that  $f$  is holomorphic on this open annulus  $A$ . Then, CIF takes the form

$$f(z) = \frac{1}{2\pi i} \int_{|w-z_0|=R'} \frac{f(w)}{z-w} dw - \frac{1}{2\pi i} \int_{|w-z_0|=r'} \frac{f(w)}{z-w} dw,$$

where  $r < r' < R' < R$ .

Just like how the usual CIF gave us the power series, this CIF gives us the Laurent series.



# Lecture 9: Laurent Series

## Theorem 28 (Laurent Series)

With the same setup as earlier, for  $z \in A$ , we can write  $f(z)$  as

# Lecture 9: Laurent Series

## Theorem 28 (Laurent Series)

With the same setup as earlier, for  $z \in A$ , we can write  $f(z)$  as

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n,$$

# Lecture 9: Laurent Series

## Theorem 28 (Laurent Series)

With the same setup as earlier, for  $z \in A$ , we can write  $f(z)$  as

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n,$$

where each  $a_n$  is given, as before, by

# Lecture 9: Laurent Series

## Theorem 28 (Laurent Series)

With the same setup as earlier, for  $z \in A$ , we can write  $f(z)$  as

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n,$$

where each  $a_n$  is given, as before, by

$$a_n = \frac{1}{2\pi i} \int_{|z-w|=r_0} \frac{f(w)}{(z-w)^{n+1}} dw,$$

## Theorem 28 (Laurent Series)

With the same setup as earlier, for  $z \in A$ , we can write  $f(z)$  as

$$f(z) = \sum_{n=-\infty}^{\infty} a_n (z - z_0)^n,$$

where each  $a_n$  is given, as before, by

$$a_n = \frac{1}{2\pi i} \int_{|z-w|=r_0} \frac{f(w)}{(z-w)^{n+1}} dw,$$

where  $r < r_0 < R$ .

# Lecture 9: Laurent Series

## Theorem 28 (Laurent Series)

With the same setup as earlier, for  $z \in A$ , we can write  $f(z)$  as

$$f(z) = \sum_{n=-\infty}^{\infty} a_n(z - z_0)^n,$$

where each  $a_n$  is given, as before, by

$$a_n = \frac{1}{2\pi i} \int_{|z-w|=r_0} \frac{f(w)}{(z-w)^{n+1}} dw,$$

where  $r < r_0 < R$ .

Note that the above is valid for  $n < 0$  as well.

# Lecture 9: Laurent Series

The most interesting coefficient is the  $-1^{\text{st}}$  one.

## Lecture 9: Laurent Series

The most interesting coefficient is the  $-1^{\text{st}}$  one. When we integrate a Laurent series along a circle centered at  $z_0$ , only  $a_{-1}$  remains (with a factor of  $2\pi i$ ).



## Lecture 9: Laurent Series

The most interesting coefficient is the  $-1^{\text{st}}$  one. When we integrate a Laurent series along a circle centered at  $z_0$ , only  $a_{-1}$  remains (with a factor of  $2\pi i$ ). This is given by

$$a_{-1} = \frac{1}{2\pi i} \int_{|z-z_0|=r_0} f(w)dw.$$

## Lecture 9: Laurent Series

The most interesting coefficient is the  $-1^{\text{st}}$  one. When we integrate a Laurent series along a circle centered at  $z_0$ , only  $a_{-1}$  remains (with a factor of  $2\pi i$ ). This is given by

$$a_{-1} = \frac{1}{2\pi i} \int_{|z-z_0|=r_0} f(w)dw.$$

This is what is usually called the *residue* and written as

## Lecture 9: Laurent Series

The most interesting coefficient is the  $-1^{\text{st}}$  one. When we integrate a Laurent series along a circle centered at  $z_0$ , only  $a_{-1}$  remains (with a factor of  $2\pi i$ ). This is given by

$$a_{-1} = \frac{1}{2\pi i} \int_{|z-z_0|=r_0} f(w)dw.$$

This is what is usually called the *residue* and written as

$$a_{-1} = \text{Res}(f; z_0).$$

With residues, calculation of integrals becomes easier.

## Theorem 29 (Cauchy's Residue Theorem)

Suppose  $f$  is given and has finitely many singularities  $z_1, \dots, z_n$  within a **simple** closed contour  $\gamma$ .

# Lecture 9: Laurent Series

With residues, calculation of integrals becomes easier.

## Theorem 29 (Cauchy's Residue Theorem)

Suppose  $f$  is given and has finitely many singularities  $z_1, \dots, z_n$  within a **simple** closed contour  $\gamma$ . Then, we have

$$\int_{\gamma} f(z) dz = 2\pi i \sum_{i=1}^n \text{Res}(f; z_i).$$

# Lecture 9: Laurent Series

With residues, calculation of integrals becomes easier.

## Theorem 29 (Cauchy's Residue Theorem)

Suppose  $f$  is given and has finitely many singularities  $z_1, \dots, z_n$  within a **simple** closed contour  $\gamma$ . Then, we have

$$\int_{\gamma} f(z) dz = 2\pi i \sum_{i=1}^n \text{Res}(f; z_i).$$

Note that the above is implicitly implying that  $f$  is holomorphic at all other points within  $\gamma$ .

# Lecture 9: Laurent Series

## Definition 21 (Principal part of a Laurent Series)

Given a Laurent series as earlier, its principal part is

## Theorem 30 (Isolated singularities and their principal parts)

# Lecture 9: Laurent Series

## Definition 21 (Principal part of a Laurent Series)

Given a Laurent series as earlier, its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

## Theorem 30 (Isolated singularities and their principal parts)



# Lecture 9: Laurent Series

## Definition 21 (Principal part of a Laurent Series)

Given a Laurent series as earlier, its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

## Theorem 30 (Isolated singularities and their principal parts)

The isolated singularity  $z_0$  is

# Lecture 9: Laurent Series

## Definition 21 (Principal part of a Laurent Series)

Given a Laurent series as earlier, its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

## Theorem 30 (Isolated singularities and their principal parts)

The isolated singularity  $z_0$  is

- ① removable iff the principal part has no terms,

# Lecture 9: Laurent Series

## Definition 21 (Principal part of a Laurent Series)

Given a Laurent series as earlier, its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

## Theorem 30 (Isolated singularities and their principal parts)

The isolated singularity  $z_0$  is

- ① removable iff the principal part has no terms,
- ② a pole iff the principal part has finitely many (and at least one) terms, and

# Lecture 9: Laurent Series

## Definition 21 (Principal part of a Laurent Series)

Given a Laurent series as earlier, its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

## Theorem 30 (Isolated singularities and their principal parts)

The isolated singularity  $z_0$  is

- ① removable iff the principal part has no terms,
- ② a pole iff the principal part has finitely many (and at least one) terms, and
- ③ essential iff the principal part has infinitely many terms.

# Lecture 9: Laurent Series

## Definition 21 (Principal part of a Laurent Series)

Given a Laurent series as earlier, its principal part is

$$\sum_{n=-\infty}^{-1} a_n(z - z_0)^n.$$

## Theorem 30 (Isolated singularities and their principal parts)

The isolated singularity  $z_0$  is

- ① removable iff the principal part has no terms,
- ② a pole iff the principal part has finitely many (and at least one) terms, and
- ③ essential iff the principal part has infinitely many terms.

In particular, the residue at a removable singularity is 0.

# Lecture 9: Laurent Series

Now, we see how one can calculate residue at a pole.

# Lecture 9: Laurent Series

Now, we see how one can calculate residue at a pole.

By the previous theorem, we know that  $f$  can be written as

# Lecture 9: Laurent Series

Now, we see how one can calculate residue at a pole.

By the previous theorem, we know that  $f$  can be written as

$$f(z) = \frac{a_{-m}}{(z - z_0)^m} + \cdots + \frac{a_{-1}}{z - z_0} + a_0 + a_1(z - z_0) + \cdots ,$$



# Lecture 9: Laurent Series

Now, we see how one can calculate residue at a pole.

By the previous theorem, we know that  $f$  can be written as

$$f(z) = \frac{a_{-m}}{(z - z_0)^m} + \cdots + \frac{a_{-1}}{z - z_0} + a_0 + a_1(z - z_0) + \cdots ,$$

for some integer  $m > 0$ .

# Lecture 9: Laurent Series

Now, we see how one can calculate residue at a pole.

By the previous theorem, we know that  $f$  can be written as

$$f(z) = \frac{a_{-m}}{(z - z_0)^m} + \cdots + \frac{a_{-1}}{z - z_0} + a_0 + a_1(z - z_0) + \cdots,$$

for some integer  $m > 0$ .

Thus,

$$g(z) = (z - z_0)^m f(z)$$

is holomorphic at  $z_0$  and

## Lecture 9: Laurent Series

Now, we see how one can calculate residue at a pole.

By the previous theorem, we know that  $f$  can be written as

$$f(z) = \frac{a_{-m}}{(z - z_0)^m} + \cdots + \frac{a_{-1}}{z - z_0} + a_0 + a_1(z - z_0) + \cdots,$$

for some integer  $m > 0$ .

Thus,

$$g(z) = (z - z_0)^m f(z)$$

is holomorphic at  $z_0$  and

$$a_{-1} = \frac{1}{(m-1)!} g^{(m-1)}(z_0).$$

# The End

Doubts?