

Extending Riemann maps to the boundary

Aryaman Maithani

IIT Bombay

16th March 2022

Notations and Conventions

- 1 \mathbb{D} will denote the open unit disc. $S^1 := \partial\mathbb{D}$ is the unit circle.

Notations and Conventions

- 1 \mathbb{D} will denote the open unit disc. $S^1 := \partial\mathbb{D}$ is the unit circle.
- 2 \mathcal{O}^∞ denotes the set of bounded holomorphic functions on \mathbb{D} .

Notations and Conventions

- ① \mathbb{D} will denote the open unit disc. $S^1 := \partial\mathbb{D}$ is the unit circle.
- ② \mathcal{O}^∞ denotes the set of bounded holomorphic functions on \mathbb{D} .
- ③ Ω will always denote a nonempty, open, bounded, and simply-connected subset of \mathbb{C} .

Notations and Conventions

- 1 \mathbb{D} will denote the open unit disc. $S^1 := \partial\mathbb{D}$ is the unit circle.
- 2 \mathcal{O}^∞ denotes the set of bounded holomorphic functions on \mathbb{D} .
- 3 Ω will always denote a nonempty, open, bounded, and simply-connected subset of \mathbb{C} .
- 4 Recall that a Riemann mapping of Ω onto \mathbb{D} is simply a biholomorphism $\Omega \rightarrow \mathbb{D}$.

Notations and Conventions

- ① \mathbb{D} will denote the open unit disc. $S^1 := \partial\mathbb{D}$ is the unit circle.
- ② \mathcal{O}^∞ denotes the set of bounded holomorphic functions on \mathbb{D} .
- ③ Ω will always denote a nonempty, open, bounded, and simply-connected subset of \mathbb{C} .
- ④ Recall that a Riemann mapping of Ω onto \mathbb{D} is simply a biholomorphism $\Omega \rightarrow \mathbb{D}$.
- ⑤ A curve shall mean a continuous function with domain $[0, 1]$. Typically, γ will be a curve such that $\gamma([0, 1)) \subseteq \Omega$ and $\gamma(1) \in \partial\Omega$. Similarly, Γ will be a curve such that $\Gamma([0, 1)) \subseteq \mathbb{D}$ and $\Gamma(1) \in \partial\mathbb{D}$.

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} .

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$.

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

The obvious way to extend f is

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

The obvious way to extend f is via sequences.

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

The obvious way to extend f is via sequences. In fact, if an extension exists, this *is* how it must be obtained.

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

The obvious way to extend f is via sequences. In fact, if an extension exists, this *is* how it must be obtained. In particular, this extension is unique

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

The obvious way to extend f is via sequences. In fact, if an extension exists, this *is* how it must be obtained. In particular, this extension is unique and we must have $f(\overline{\Omega}) \subseteq \overline{\mathbb{D}}$.

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

The obvious way to extend f is via sequences. In fact, if an extension exists, this *is* how it must be obtained. In particular, this extension is unique and we must have $f(\overline{\Omega}) \subseteq \overline{\mathbb{D}}$. This must force $f(\overline{\Omega}) = \overline{\mathbb{D}}$.

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

The obvious way to extend f is via sequences. In fact, if an extension exists, this *is* how it must be obtained. In particular, this extension is unique and we must have $f(\overline{\Omega}) \subseteq \overline{\mathbb{D}}$. This must force $f(\overline{\Omega}) = \overline{\mathbb{D}}$. (Why?)

Introduction

Let Ω be a bounded simply-connected domain in \mathbb{C} . By the Riemann Mapping Theorem, we know that there exists a biholomorphism $f : \Omega \rightarrow \mathbb{D}$. The following is a natural question.

Question

Can f be continuously extended up to $\overline{\Omega}$?

The obvious way to extend f is via sequences. In fact, if an extension exists, this *is* how it must be obtained. In particular, this extension is unique and we must have $f(\overline{\Omega}) \subseteq \overline{\mathbb{D}}$. This must force $f(\overline{\Omega}) = \overline{\mathbb{D}}$. (Why?)

Thus, it also makes sense to ask whether *the* extension is a homeomorphism onto $\overline{\mathbb{D}}$.

Spoilers

What we shall see is the following:

Spoilers

What we shall see is the following: By imposing a *simple* topological restriction on Ω ,

Spoilers

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$.

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$. Moreover, this will be a *homeomorphism*.

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$. Moreover, this will be a *homeomorphism*.

Remark 1

The last line is not difficult to see.

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$. Moreover, this will be a *homeomorphism*.

Remark 1

The last line is not difficult to see. Indeed, once we have continuously extended f to $\tilde{f} : \overline{\Omega} \rightarrow \overline{\mathbb{D}}$,

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$. Moreover, this will be a *homeomorphism*.

Remark 1

The last line is not difficult to see. Indeed, once we have continuously extended f to $\tilde{f} : \overline{\Omega} \rightarrow \overline{\mathbb{D}}$, we have

$$\mathbb{D} \subseteq \tilde{f}(\overline{\Omega}) \subseteq \overline{\mathbb{D}}.$$

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$. Moreover, this will be a *homeomorphism*.

Remark 1

The last line is not difficult to see. Indeed, once we have continuously extended f to $\tilde{f} : \overline{\Omega} \rightarrow \overline{\mathbb{D}}$, we have

$$\mathbb{D} \subseteq \tilde{f}(\overline{\Omega}) \subseteq \overline{\mathbb{D}}.$$

As $\tilde{f}(\overline{\Omega})$ is compact,

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$. Moreover, this will be a *homeomorphism*.

Remark 1

The last line is not difficult to see. Indeed, once we have continuously extended f to $\tilde{f} : \overline{\Omega} \rightarrow \overline{\mathbb{D}}$, we have

$$\mathbb{D} \subseteq \tilde{f}(\overline{\Omega}) \subseteq \overline{\mathbb{D}}.$$

As $\tilde{f}(\overline{\Omega})$ is compact, we have $\tilde{f}(\overline{\Omega}) = \overline{\mathbb{D}}$.

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$. Moreover, this will be a *homeomorphism*.

Remark 1

The last line is not difficult to see. Indeed, once we have continuously extended f to $\tilde{f} : \overline{\Omega} \rightarrow \overline{\mathbb{D}}$, we have

$$\mathbb{D} \subseteq \tilde{f}(\overline{\Omega}) \subseteq \overline{\mathbb{D}}.$$

As $\tilde{f}(\overline{\Omega})$ is compact, we have $\tilde{f}(\overline{\Omega}) = \overline{\mathbb{D}}$.

Furthermore, if \tilde{f} is an injection,

What we shall see is the following: By imposing a *simple* topological restriction on Ω , one gets that *any* biholomorphism $\Omega \rightarrow \mathbb{D}$ can be extended to a continuous *injective* map $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$. Moreover, this will be a *homeomorphism*.

Remark 1

The last line is not difficult to see. Indeed, once we have continuously extended f to $\tilde{f} : \overline{\Omega} \rightarrow \overline{\mathbb{D}}$, we have

$$\mathbb{D} \subseteq \tilde{f}(\overline{\Omega}) \subseteq \overline{\mathbb{D}}.$$

As $\tilde{f}(\overline{\Omega})$ is compact, we have $\tilde{f}(\overline{\Omega}) = \overline{\mathbb{D}}$.

Furthermore, if \tilde{f} is an injection, then compactness again tells us that \tilde{f} is a homeomorphism (as \tilde{f} is a bijection).

Definition 2

Simple Boundary Points

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

Simple Boundary Points

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω

Simple Boundary Points

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω such that $\alpha_n \rightarrow \beta$ as $n \rightarrow \infty$,

Simple Boundary Points

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω such that $\alpha_n \rightarrow \beta$ as $n \rightarrow \infty$, there exists a curve γ

Simple Boundary Points

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω such that $\alpha_n \rightarrow \beta$ as $n \rightarrow \infty$, there exists a curve γ and a strictly increasing sequence $(t_n)_n$ in $(0, 1)$

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω such that $\alpha_n \rightarrow \beta$ as $n \rightarrow \infty$, there exists a curve γ and a strictly increasing sequence $(t_n)_n$ in $(0, 1)$ such that

$$t_n \rightarrow 1,$$

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω such that $\alpha_n \rightarrow \beta$ as $n \rightarrow \infty$, there exists a curve γ and a strictly increasing sequence $(t_n)_n$ in $(0, 1)$ such that

$$t_n \rightarrow 1, \gamma(t_n) = \alpha_n \ (n = 1, 2, \dots),$$

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω such that $\alpha_n \rightarrow \beta$ as $n \rightarrow \infty$, there exists a curve γ and a strictly increasing sequence $(t_n)_n$ in $(0, 1)$ such that

$$t_n \rightarrow 1, \gamma(t_n) = \alpha_n \ (n = 1, 2, \dots), \gamma([0, 1)) \in \Omega.$$

Simple Boundary Points

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω such that $\alpha_n \rightarrow \beta$ as $n \rightarrow \infty$, there exists a curve γ and a strictly increasing sequence $(t_n)_n$ in $(0, 1)$ such that

$$t_n \rightarrow 1, \gamma(t_n) = \alpha_n \ (n = 1, 2, \dots), \gamma([0, 1)) \in \Omega.$$

$\gamma(1) = \beta$ follows by continuity.

Simple Boundary Points

Definition 2

A boundary point β of Ω is called a **simple boundary point** if β has the following property:

For every sequence $(\alpha_n)_n$ in Ω such that $\alpha_n \rightarrow \beta$ as $n \rightarrow \infty$, there exists a curve γ and a strictly increasing sequence $(t_n)_n$ in $(0, 1)$ such that

$$t_n \rightarrow 1, \gamma(t_n) = \alpha_n \ (n = 1, 2, \dots), \gamma([0, 1)) \in \Omega.$$

$\gamma(1) = \beta$ follows by continuity.

In words: there is a curve in Ω which passes through α_n and ends at β .


(Non-)Examples

(Non-)Examples

- 1 Every boundary point of \mathbb{D} is a simple boundary point.


(Non-)Examples

- ① Every boundary point of \mathbb{D} is a simple boundary point.
- ② Let $\Omega := \mathbb{D} \setminus [0, 1)$.



(Non-)Examples


- 1 Every boundary point of \mathbb{D} is a simple boundary point.
- 2 Let $\Omega := \mathbb{D} \setminus [0, 1)$.



The boundary points of Ω lying on the real axis are not simple.

(Non-)Examples


- 1 Every boundary point of \mathbb{D} is a simple boundary point.
- 2 Let $\Omega := \mathbb{D} \setminus [0, 1)$.



The boundary points of Ω lying on the real axis are not simple. Note that Ω is indeed bounded and simply-connected and thus, biholomorphic to \mathbb{D} .

(Non-)Examples


- 1 Every boundary point of \mathbb{D} is a simple boundary point.
- 2 Let $\Omega := \mathbb{D} \setminus [0, 1)$.



The boundary points of Ω lying on the real axis are not simple. Note that Ω is indeed bounded and simply-connected and thus, biholomorphic to \mathbb{D} . However, $\partial\Omega$ is clearly not homeomorphic to $\partial\mathbb{D}$ and

(Non-)Examples

- 1 Every boundary point of \mathbb{D} is a simple boundary point.
- 2 Let $\Omega := \mathbb{D} \setminus [0, 1)$.



The boundary points of Ω lying on the real axis are not simple. Note that Ω is indeed bounded and simply-connected and thus, biholomorphic to \mathbb{D} . However, $\partial\Omega$ is clearly not homeomorphic to $\partial\mathbb{D}$ and thus, no biholomorphism can be extended to a homeomorphism $\overline{\Omega} \rightarrow \overline{\mathbb{D}}$.

Theorem 3 (Helper Theorem)

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- 1 If β is a simple boundary point of Ω ,

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- 1 If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$.

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- 1 If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- 1 If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.
- 2 If β_1 and β_2 are distinct simple boundary points of Ω

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- ① If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.
- ② If β_1 and β_2 are distinct simple boundary points of Ω and if f is continuously extended to $\Omega \cup \{\beta_1, \beta_2\}$,

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- ① If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.
- ② If β_1 and β_2 are distinct simple boundary points of Ω and if f is continuously extended to $\Omega \cup \{\beta_1, \beta_2\}$, then $f(\beta_1) \neq f(\beta_2)$.

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- ① If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.
- ② If β_1 and β_2 are distinct simple boundary points of Ω and if f is continuously extended to $\Omega \cup \{\beta_1, \beta_2\}$, then $f(\beta_1) \neq f(\beta_2)$.

We give the proof after proving the main theorem assuming the above.

As remarked, the extension in ① is unique and would have to be attained as follows:

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- ① If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.
- ② If β_1 and β_2 are distinct simple boundary points of Ω and if f is continuously extended to $\Omega \cup \{\beta_1, \beta_2\}$, then $f(\beta_1) \neq f(\beta_2)$.

We give the proof after proving the main theorem assuming the above.

As remarked, the extension in ① is unique and would have to be attained as follows: given a sequence $(\alpha_n)_n$ in Ω converging to β ,

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- ① If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.
- ② If β_1 and β_2 are distinct simple boundary points of Ω and if f is continuously extended to $\Omega \cup \{\beta_1, \beta_2\}$, then $f(\beta_1) \neq f(\beta_2)$.

We give the proof after proving the main theorem assuming the above.

As remarked, the extension in ① is unique and would have to be attained as follows: given a sequence $(\alpha_n)_n$ in Ω converging to β , we have $f(\beta) := \lim f(\alpha_n)$.

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- ① If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.
- ② If β_1 and β_2 are distinct simple boundary points of Ω and if f is continuously extended to $\Omega \cup \{\beta_1, \beta_2\}$, then $f(\beta_1) \neq f(\beta_2)$.

We give the proof after proving the main theorem assuming the above.

As remarked, the extension in ① is unique and would have to be attained as follows: given a sequence $(\alpha_n)_n$ in Ω converging to β , we have $f(\beta) := \lim f(\alpha_n)$. Once we show that this limit (exists and) is independent of the sequence (α_n) ,

Theorem 3 (Helper Theorem)

Let Ω be a bounded simply-connected domain, and let f be a Riemann mapping of Ω onto \mathbb{D} .

- ① If β is a simple boundary point of Ω , then f has a continuous extension to $\Omega \cup \{\beta\}$. If f is so extended, then $|f(\beta)| = 1$.
- ② If β_1 and β_2 are distinct simple boundary points of Ω and if f is continuously extended to $\Omega \cup \{\beta_1, \beta_2\}$, then $f(\beta_1) \neq f(\beta_2)$.

We give the proof after proving the main theorem assuming the above.

As remarked, the extension in ① is unique and would have to be attained as follows: given a sequence $(\alpha_n)_n$ in Ω converging to β , we have $f(\beta) := \lim f(\alpha_n)$. Once we show that this limit (exists and) is independent of the sequence (α_n) , we would have shown continuity.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple,

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D}

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it,

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②,

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$. As remarked earlier, this would finish the proof.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto \overline{D} .

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$. As remarked earlier, this would finish the proof.

To this end, let $(z_n)_n$ be an arbitrary sequence in $\overline{\Omega}$ that converges to z .

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto $\overline{\mathbb{D}}$.

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$. As remarked earlier, this would finish the proof.

To this end, let $(z_n)_n$ be an arbitrary sequence in $\overline{\Omega}$ that converges to z . We can pick a corresponding sequence $(\alpha_n)_n$ in Ω such that

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto $\overline{\mathbb{D}}$.

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$. As remarked earlier, this would finish the proof.

To this end, let $(z_n)_n$ be an arbitrary sequence in $\overline{\Omega}$ that converges to z . We can pick a corresponding sequence $(\alpha_n)_n$ in Ω such that

$$|\alpha_n - z_n| < 1/n$$

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto $\overline{\mathbb{D}}$.

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$. As remarked earlier, this would finish the proof.

To this end, let $(z_n)_n$ be an arbitrary sequence in $\overline{\Omega}$ that converges to z . We can pick a corresponding sequence $(\alpha_n)_n$ in Ω such that $|\alpha_n - z_n| < 1/n$ and $|f(\alpha_n) - f(z_n)| < 1/n$.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto $\overline{\mathbb{D}}$.

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By 2, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$. As remarked earlier, this would finish the proof.

To this end, let $(z_n)_n$ be an arbitrary sequence in $\overline{\Omega}$ that converges to z . We can pick a corresponding sequence $(\alpha_n)_n$ in Ω such that $|\alpha_n - z_n| < 1/n$ and $|f(\alpha_n) - f(z_n)| < 1/n$. Thus, $\alpha_n \rightarrow z$ and hence, $f(\alpha_n) \rightarrow f(z)$.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto $\overline{\mathbb{D}}$.

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$. As remarked earlier, this would finish the proof.

To this end, let $(z_n)_n$ be an arbitrary sequence in $\overline{\Omega}$ that converges to z . We can pick a corresponding sequence $(\alpha_n)_n$ in Ω such that $|\alpha_n - z_n| < 1/n$ and $|f(\alpha_n) - f(z_n)| < 1/n$. Thus, $\alpha_n \rightarrow z$ and hence, $f(\alpha_n) \rightarrow f(z)$. In turn, $f(z_n) \rightarrow f(z)$, as desired.

The Main Extension Theorem

Theorem 4

If Ω is a bounded simply-connected domain and if every boundary point of Ω is simple, then every Riemann mapping of Ω onto \mathbb{D} extends to a homeomorphism of $\overline{\Omega}$ onto $\overline{\mathbb{D}}$.

Proof.

Let $f : \Omega \rightarrow \mathbb{D}$ be a biholomorphism. By the Helper Theorem and the remark following it, we see that we may extend f to $\overline{\Omega}$ using sequences. By ②, it follows that f so extended is one-one. We now check that it is continuous on $\overline{\Omega}$. As remarked earlier, this would finish the proof.

To this end, let $(z_n)_n$ be an arbitrary sequence in $\overline{\Omega}$ that converges to z . We can pick a corresponding sequence $(\alpha_n)_n$ in Ω such that $|\alpha_n - z_n| < 1/n$ and $|f(\alpha_n) - f(z_n)| < 1/n$. Thus, $\alpha_n \rightarrow z$ and hence, $f(\alpha_n) \rightarrow f(z)$. In turn, $f(z_n) \rightarrow f(z)$, as desired. □

A Purely Topological Corollary

Recall that a Jordan curve is the image of an injective map $S^1 \rightarrow \mathbb{C}$.

A Purely Topological Corollary

Recall that a Jordan curve is the image of an injective map $S^1 \rightarrow \mathbb{C}$.

Corollary 5

A Purely Topological Corollary

Recall that a Jordan curve is the image of an injective map $S^1 \rightarrow \mathbb{C}$.

Corollary 5

If every boundary point of a bounded simply-connected region Ω is simple,

A Purely Topological Corollary

Recall that a Jordan curve is the image of an injective map $S^1 \rightarrow \mathbb{C}$.

Corollary 5

If every boundary point of a bounded simply-connected region Ω is simple, then the boundary of Ω is a Jordan curve,

A Purely Topological Corollary

Recall that a Jordan curve is the image of an injective map $S^1 \rightarrow \mathbb{C}$.

Corollary 5

If every boundary point of a bounded simply-connected region Ω is simple, then the boundary of Ω is a Jordan curve, and $\overline{\Omega}$ is homeomorphic to $\overline{\mathbb{D}}$.

A Purely Topological Corollary

Recall that a Jordan curve is the image of an injective map $S^1 \rightarrow \mathbb{C}$.

Corollary 5

If every boundary point of a bounded simply-connected region Ω is simple, then the boundary of Ω is a Jordan curve, and $\overline{\Omega}$ is homeomorphic to $\overline{\mathbb{D}}$.

In fact, the converse is true too:

A Purely Topological Corollary

Recall that a Jordan curve is the image of an injective map $S^1 \rightarrow \mathbb{C}$.

Corollary 5

If every boundary point of a bounded simply-connected region Ω is simple, then the boundary of Ω is a Jordan curve, and $\overline{\Omega}$ is homeomorphic to $\overline{\mathbb{D}}$.

In fact, the converse is true too: If the boundary of Ω is a Jordan curve, then every boundary point of Ω is simple.

Theorem 6 (Radial Limit Theorem)

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$,

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If $g^*(e^{i\theta}) = 0$

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If $g^*(e^{i\theta}) = 0$ for almost all $e^{i\theta}$ on some arc $J \subseteq S^1$,

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If $g^*(e^{i\theta}) = 0$ for almost all $e^{i\theta}$ on some arc $J \subseteq S^1$, then $g(z) = 0$ for every $z \in \mathbb{D}$.

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If $g^*(e^{i\theta}) = 0$ for almost all $e^{i\theta}$ on some arc $J \subseteq S^1$, then $g(z) = 0$ for every $z \in \mathbb{D}$.

Theorem 7 (Lindelöf's Theorem)

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If $g^*(e^{i\theta}) = 0$ for almost all $e^{i\theta}$ on some arc $J \subseteq S^1$, then $g(z) = 0$ for every $z \in \mathbb{D}$.

Theorem 7 (Lindelöf's Theorem)

Suppose Γ is a curve such that $\Gamma([0, 1)) \subseteq \mathbb{D}$ and $\Gamma(1) = 1$.

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If $g^*(e^{i\theta}) = 0$ for almost all $e^{i\theta}$ on some arc $J \subseteq S^1$, then $g(z) = 0$ for every $z \in \mathbb{D}$.

Theorem 7 (Lindelöf's Theorem)

Suppose Γ is a curve such that $\Gamma([0, 1)) \subseteq \mathbb{D}$ and $\Gamma(1) = 1$.

If $g \in \mathcal{O}^\infty$ and

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If $g^*(e^{i\theta}) = 0$ for almost all $e^{i\theta}$ on some arc $J \subseteq S^1$, then $g(z) = 0$ for every $z \in \mathbb{D}$.

Theorem 7 (Lindelöf's Theorem)

Suppose Γ is a curve such that $\Gamma([0, 1)) \subseteq \mathbb{D}$ and $\Gamma(1) = 1$.

If $g \in \mathcal{O}^\infty$ and

$$\lim_{t \rightarrow 1^-} g(\Gamma(t)) = L,$$

Theorem 6 (Radial Limit Theorem)

To every $g \in \mathcal{O}^\infty$ corresponds a function $g^* \in L^\infty(S^1)$, defined almost everywhere by

$$g^*(e^{i\theta}) = \lim_{r \rightarrow 1} g(re^{i\theta}).$$

If $g^*(e^{i\theta}) = 0$ for almost all $e^{i\theta}$ on some arc $J \subseteq S^1$, then $g(z) = 0$ for every $z \in \mathbb{D}$.

Theorem 7 (Lindelöf's Theorem)

Suppose Γ is a curve such that $\Gamma([0, 1)) \subseteq \mathbb{D}$ and $\Gamma(1) = 1$.

If $g \in \mathcal{O}^\infty$ and

$$\lim_{t \rightarrow 1^-} g(\Gamma(t)) = L,$$

then g has radial limit L at 1.

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

- 1 Suppose that f cannot be extended to β .

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta,$$

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_n) \rightarrow w_1,$$

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2,$$

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point,

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$.

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$.
Let $g = f^{-1}$

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$.

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω .

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$,

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r)

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$.

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$. Thus, $|\Gamma(t)| \rightarrow 1$ as $t \rightarrow 1$.

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$. Thus, $|\Gamma(t)| \rightarrow 1$ as $t \rightarrow 1$. In particular, $|w_1| = |w_2| = 1$.

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$. Thus, $|\Gamma(t)| \rightarrow 1$ as $t \rightarrow 1$. In particular, $|w_1| = |w_2| = 1$.

Let J be one of the open arcs of $S^1 \setminus \{w_1, w_2\}$

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$. Thus, $|\Gamma(t)| \rightarrow 1$ as $t \rightarrow 1$. In particular, $|w_1| = |w_2| = 1$.

Let J be one of the open arcs of $S^1 \setminus \{w_1, w_2\}$ such that every radius of \mathbb{D} which ends at a point of J

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$. Thus, $|\Gamma(t)| \rightarrow 1$ as $t \rightarrow 1$. In particular, $|w_1| = |w_2| = 1$.

Let J be one of the open arcs of $S^1 \setminus \{w_1, w_2\}$ such that every radius of \mathbb{D} which ends at a point of J intersects the range of Γ in a set which has a limit point on S^1 .

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$. Thus, $|\Gamma(t)| \rightarrow 1$ as $t \rightarrow 1$. In particular, $|w_1| = |w_2| = 1$.

Let J be one of the open arcs of $S^1 \setminus \{w_1, w_2\}$ such that every radius of \mathbb{D} which ends at a point of J intersects the range of Γ in a set which has a limit point on S^1 . By the Radial Limit Theorem, g has radial limits a.e. on S^1 .

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$. Thus, $|\Gamma(t)| \rightarrow 1$ as $t \rightarrow 1$. In particular, $|w_1| = |w_2| = 1$.

Let J be one of the open arcs of $S^1 \setminus \{w_1, w_2\}$ such that every radius of \mathbb{D} which ends at a point of J intersects the range of Γ in a set which has a limit point on S^1 . By the Radial Limit Theorem, g has radial limits a.e. on S^1 since $g \in \mathcal{O}^\infty$

Proof of Helper Theorem

Now, we prove the earlier Helper Theorem assuming the earlier two results.

① Suppose that f cannot be extended to β . Then, there exists a sequence $(\alpha_n)_n$ in Ω and points $w_1, w_2 \in \overline{\mathbb{D}}$ such that

$$\alpha_n \rightarrow \beta, f(\alpha_{2n}) \rightarrow w_1, f(\alpha_{2n+1}) \rightarrow w_2, w_1 \neq w_2.$$

Choose γ as given by β being a simple boundary point, and put $\Gamma := f \circ \gamma$. Let $g = f^{-1}$ and put $K_r := g(\overline{D}(0; r))$ for $0 < r < 1$. Then K_r is a compact subset of Ω . Since $\gamma(t) \rightarrow \beta$ as $t \rightarrow 1$, there exists $t^* < 1$ (depending on r) such that $\gamma(t) \notin K_r$ if $t^* < t < 1$. Thus, $|\Gamma(t)| \rightarrow 1$ as $t \rightarrow 1$. In particular, $|w_1| = |w_2| = 1$.

Let J be one of the open arcs of $S^1 \setminus \{w_1, w_2\}$ such that every radius of \mathbb{D} which ends at a point of J intersects the range of Γ in a set which has a limit point on S^1 . By the Radial Limit Theorem, g has radial limits a.e. on S^1 since $g \in \mathcal{O}^\infty$ (as Ω is bounded).

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 .

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist,

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) =$$

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

since $g(\Gamma(t)) = \gamma(t) \rightarrow \beta$ as $t \rightarrow 1$.

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

since $g(\Gamma(t)) = \gamma(t) \rightarrow \beta$ as $t \rightarrow 1$. Thus, by the Radial Limit Theorem again,

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

since $g(\Gamma(t)) = \gamma(t) \rightarrow \beta$ as $t \rightarrow 1$. Thus, by the Radial Limit Theorem again, applied to $g - \beta$,

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

since $g(\Gamma(t)) = \gamma(t) \rightarrow \beta$ as $t \rightarrow 1$. Thus, by the Radial Limit Theorem again, applied to $g - \beta$, we see that $g \equiv \beta$ on \mathbb{D} , contradicting that

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

since $g(\Gamma(t)) = \gamma(t) \rightarrow \beta$ as $t \rightarrow 1$. Thus, by the Radial Limit Theorem again, applied to $g - \beta$, we see that $g \equiv \beta$ on \mathbb{D} , contradicting that g is an injection.

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

since $g(\Gamma(t)) = \gamma(t) \rightarrow \beta$ as $t \rightarrow 1$. Thus, by the Radial Limit Theorem again, applied to $g - \beta$, we see that $g \equiv \beta$ on \mathbb{D} , contradicting that g is an injection.

Thus, we have shown that $w_1 = w_2$

Continuing the Proof

We have that $g \circ \Gamma = \gamma$ and that g has radial limits a.e. on S^1 . Now, for whichever $e^{it} \in J$ the radial limit does exist, we must have

$$\lim_{r \rightarrow 1} g(re^{it}) = \beta,$$

since $g(\Gamma(t)) = \gamma(t) \rightarrow \beta$ as $t \rightarrow 1$. Thus, by the Radial Limit Theorem again, applied to $g - \beta$, we see that $g \equiv \beta$ on \mathbb{D} , contradicting that g is an injection.

Thus, we have shown that $w_1 = w_2$ and $|w_1| = 1$.

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points.

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) = f(\beta_2)$.

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) = f(\beta_2)$. We may assume $f(\beta_i) = 1$.

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) = f(\beta_2)$. We may assume $f(\beta_i) = 1$. Let γ_i be curves with $\gamma_i([0, 1)) \subseteq \Omega$,

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) = f(\beta_2)$. We may assume $f(\beta_i) = 1$. Let γ_i be curves with $\gamma_i([0, 1)) \subseteq \Omega$, and $\gamma_i(1) = \beta_i$,

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) = f(\beta_2)$. We may assume $f(\beta_i) = 1$. Let γ_i be curves with $\gamma_i([0, 1)) \subseteq \Omega$, and $\gamma_i(1) = \beta_i$, and let $\Gamma_i := f \circ \gamma_i$.

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) = f(\beta_2)$. We may assume $f(\beta_i) = 1$. Let γ_i be curves with $\gamma_i([0, 1)) \subseteq \Omega$, and $\gamma_i(1) = \beta_i$, and let $\Gamma_i := f \circ \gamma_i$. Then, each Γ_i satisfies the condition of Lindelöf's Theorem with

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) \neq f(\beta_2)$. We may assume $f(\beta_1) = 1$. Let γ_i be curves with $\gamma_i([0, 1)) \subseteq \Omega$, and $\gamma_i(1) = \beta_i$, and let $\Gamma_i := f \circ \gamma_i$. Then, each Γ_i satisfies the condition of Lindelöf's Theorem with

$$\lim_{t \rightarrow 1} g(\Gamma_i(t)) = \lim_{t \rightarrow 1} \gamma_i(t) = \beta_i.$$

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) \neq f(\beta_2)$. We may assume $f(\beta_1) = 1$. Let γ_i be curves with $\gamma_i([0, 1)) \subseteq \Omega$, and $\gamma_i(1) = \beta_i$, and let $\Gamma_i := f \circ \gamma_i$. Then, each Γ_i satisfies the condition of Lindelöf's Theorem with

$$\lim_{t \rightarrow 1} g(\Gamma_i(t)) = \lim_{t \rightarrow 1} \gamma_i(t) = \beta_i.$$

Thus, the radial limit of g at 1 is

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) \neq f(\beta_2)$. We may assume $f(\beta_1) = 1$. Let γ_i be curves with $\gamma_i([0, 1)) \subseteq \Omega$, and $\gamma_i(1) = \beta_i$, and let $\Gamma_i := f \circ \gamma_i$. Then, each Γ_i satisfies the condition of Lindelöf's Theorem with

$$\lim_{t \rightarrow 1} g(\Gamma_i(t)) = \lim_{t \rightarrow 1} \gamma_i(t) = \beta_i.$$

Thus, the radial limit of g at 1 is both β_1 and β_2

Continuing the Proof

② Now, we need to prove that an extension takes different values at different boundary points. Let β_1, β_2 be simple boundary points with $f(\beta_1) \neq f(\beta_2)$. We may assume $f(\beta_1) = 1$. Let γ_i be curves with $\gamma_i([0, 1)) \subseteq \Omega$, and $\gamma_i(1) = \beta_i$, and let $\Gamma_i := f \circ \gamma_i$. Then, each Γ_i satisfies the condition of Lindelöf's Theorem with

$$\lim_{t \rightarrow 1} g(\Gamma_i(t)) = \lim_{t \rightarrow 1} \gamma_i(t) = \beta_i.$$

Thus, the radial limit of g at 1 is both β_1 and β_2 and hence, $\beta_1 = \beta_2$. \square