

## Problem Set 5 - pre-REU 2025

### Problem set on Invariant Theory

Problems marked with <sup>L</sup> are problems on linear algebra.

1. Show that the following sets of matrices form a group under matrix multiplication. In each case, justify why the matrices are indeed invertible.
  - (a) The set of  $n \times n$  matrices with determinant one.
  - (b) The set of  $n \times n$  matrices  $M$  satisfying  $MM^T = I_n$ .
  - (c) The set of  $2n \times 2n$  matrices  $M$  satisfying  $M\Omega M^T = \Omega$ , where  $\Omega$  is the  $2n \times 2n$  block matrix given as  $\Omega = \begin{pmatrix} 0 & I_n \\ -I_n & 0 \end{pmatrix}$ .
2. Let  $R$  be a ring (such as the ring of polynomials). Let  $f: R \rightarrow R$  be a function satisfying
  - $f(1) = 1$ ,
  - $f(x+y) = f(x) + f(y)$  for all  $x, y \in R$ ,
  - $f(xy) = f(x)f(y)$  for all  $x, y \in R$ .

Let  $S$  be the set of fixed points of  $R$ , i.e.,  $S := \{r \in R : f(r) = r\}$ . Show that  $S$  is closed under addition, multiplication, and contains 1.

3. From class, we know that any element of the orthogonal group looks like

$$M = \begin{bmatrix} \cos(\theta) & -\varepsilon \sin(\theta) \\ \sin(\theta) & \varepsilon \cos(\theta) \end{bmatrix}$$

for some  $\theta \in \mathbb{R}$  and some  $\varepsilon \in \{1, -1\}$ . Using this description, check that for the action of  $G = O_2(\mathbb{R})$  on  $R = \mathbb{R}[x, y]$ , we have  $x^2 + y^2 \in R^G$ . (As before, the action is via  $\begin{bmatrix} x \\ y \end{bmatrix} \mapsto M \begin{bmatrix} x \\ y \end{bmatrix}$ .)

4. Let  $G = GL_2(\mathbb{R})$  act on  $R = \mathbb{R}[X_{2 \times m}]$  in the usual way. Show that  $R^G = \mathbb{R}$ , i.e., the only invariant polynomials are the constants.  
*Hint:* Think about what the matrix  $\begin{pmatrix} 2 & 0 \\ 0 & 2 \end{pmatrix}$  does.  
 Start out with  $m = 1$  or  $2$  to get an idea.

- <sup>L</sup>5. Let  $A, B$  be  $n \times n$  matrices such that  $\exists P \in GL_n(\mathbb{R})$  with  $PAP^{-1} = B$ . Show that  $A$  and  $B$  have the same characteristic polynomial, i.e., show that  $\det(A - \lambda I) = \det(B - \lambda I)$ .
- 5.5. Recall that for the action of  $G = GL_n(\mathbb{R})$  on  $R = \mathbb{R}[X_{n \times n}]$  given by conjugation,  $R^G$  is generated by the coefficients of the characteristic polynomial. Interpret the previous problem in this context.
- <sup>L</sup>6. Let  $A$  be an  $n \times m$  matrix with columns  $\mathbf{a}_1, \dots, \mathbf{a}_m \in \mathbb{R}^n$ . Show that the  $(i, j)$ th entry of  $A^T A$  is  $\mathbf{a}_i \cdot \mathbf{a}_j$ .
- <sup>L</sup>6.5. Recall that for the action of  $G = O_n(\mathbb{R})$  on  $R = \mathbb{R}[X_{n \times m}]$  given by left multiplication,  $R^G$  is generated by the entries of  $X^T X$ . Using the previous problem, how does this description of  $R^G$  fit in with our geometrical definition of  $O_n(\mathbb{R})$ ?

7. Consider the action of  $G = \mathrm{SL}_2(\mathbb{R})$  on  $R = \mathbb{R}[X_{2 \times 2}]$  by left multiplication. For ease of notation, assume that the variables are denoted and arranged as

$$\begin{bmatrix} x_1 & x_2 \\ y_1 & y_2 \end{bmatrix}.$$

Show that  $x_1y_2 - x_2y_1 \in R^G$ . More generally, show that if  $\mathrm{SL}_2$  is acting on the polynomial with variables

$$\begin{bmatrix} x_1 & x_2 & \cdots & x_m \\ y_1 & y_2 & \cdots & y_m \end{bmatrix},$$

then  $x_iy_j - x_jy_i \in R^G$  for all  $1 \leq i < j \leq m$ .

*Hint for the last part:* If  $A$  and  $B$  are matrices of compatible sizes, think about how the columns of  $AB$  look. In particular, the  $i$ -th column of  $AB$  is the product of  $A$  and the  $i$ -th column of  $B$ .

- 7.5. Generalise the previous to  $\mathrm{SL}_n(\mathbb{R})$  acting on  $\mathbb{R}[X_{n \times m}]$ .